


**Universal form of thermodynamic uncertainty relation for Langevin dynamics**Jae Sung Lee ,\* Jong-Min Park , and Hyunggyu Park <sup>†</sup>*School of Physics and Quantum Universe Center, Korea Institute for Advanced Study, Seoul 02455, Korea* (Received 16 June 2021; accepted 19 October 2021; published 11 November 2021)

The thermodynamic uncertainty relation (TUR) provides a stricter bound for entropy production (EP) than that of the thermodynamic second law. This stricter bound can be utilized to infer the EP and derive other tradeoff relations. Though the validity of the TUR has been verified in various stochastic systems, its application to general Langevin dynamics has not been successfully unified, especially for underdamped Langevin dynamics, where odd parity variables in time-reversal operation such as velocity get involved. Previous TURs for underdamped Langevin dynamics are neither experimentally accessible nor reduced to the original form of the overdamped Langevin dynamics in the zero-mass limit. Here, we find a TUR for underdamped Langevin dynamics with an arbitrary time-dependent protocol, which is operationally accessible when all mechanical forces are controllable. We show that the original TUR is a consequence of our underdamped TUR in the zero-mass limit. This indicates that the TUR formulation presented here can be regarded as the universal form of the TUR for general Langevin dynamics.

DOI: [10.1103/PhysRevE.104.L052102](https://doi.org/10.1103/PhysRevE.104.L052102)

*Introduction.* Thermodynamic processes and accompanying entropy production (EP) are constrained by the thermodynamic second law, stating that the EP is always nonnegative. Beyond the second law, a new bound was discovered in 2015 [1], called the thermodynamic uncertainty relation (TUR) expressed in terms of the TUR factor  $\mathcal{Q}$  as

$$\mathcal{Q} \equiv \frac{\text{Var}[\Theta]}{\langle \Theta \rangle^2} \Delta S^{\text{tot}} \geq 2k_B, \quad (1)$$

with a time-accumulated current  $\Theta$ , its steady-state average  $\langle \Theta \rangle$  and variance  $\text{Var}[\Theta]$ , the Boltzmann constant  $k_B$ , and the average total EP  $\Delta S^{\text{tot}}$ . This is basically a tradeoff relation between the fluctuation magnitude and the thermodynamic cost of a stochastic system given as an inequality with the universal lower bound. As the variance is always positive, the TUR sets a positive lower bound of the EP, thus provides a tighter bound than the second law. This bound can be utilized for inferring the EP by measuring a certain current statistics in a nonequilibrium process [2–4]. Moreover, a recent debate on thermodynamic tradeoff relations among the efficiency, power, and reversibility of a heat engine [5–12] has also been investigated based on the TUR bound [13].

After the first discovery in 2015 [1], the validity of the TUR has been rigorously proved for a variety of stochastic systems [14–26]. First, it was shown that the TUR in the original form, Eq. (1), holds for a continuous-time Markov jump process [14,15] and the overdamped Langevin dynamics in the steady state [16]. Later, TURs for these two stochastic systems with an arbitrary initial state [17,18] and an arbitrary time-dependent driving [19] have been found. The TUR for a discrete-time Markov process was first discovered only in

an exponential form [20], but later the linearized version was also found [18]. We note that the TUR for general stochastic systems was found in an exponential form recently [21,22]. However, the exponential form is not practically useful in a sense that the physical meaning of the cost function is hard to interpret and its bound can be quite loose.

Compared to other stochastic systems, studies on the TUR for underdamped Langevin systems have made little progress. In contrast to the overdamped Langevin systems, the odd-parity variables like velocity come into play in the underdamped dynamics, and the probability current is divided into two parts: the reversible and the irreversible currents. As only the latter contributes to the EP [27,28], the thermodynamic cost function could not be simply written in terms of the EP only, but also includes some kinetic quantities such as dynamical activity, which are not easily accessible in experiments [29,30]. This significantly degrades the applicability of the TUR for inferring the EP in the underdamped Langevin dynamics. In addition, the link between the TURs for the overdamped and underdamped Langevin dynamics has been missing. Mathematically, the overdamped dynamics is usually attained in the zero-mass limit of the underdamped dynamics. However, the zero-mass limit of the previous TURs for the underdamped dynamics becomes meaningless as the dynamic activity (thus, the cost function) diverges [30]. This clearly reveals the lack of systematic understanding of the thermodynamic tradeoff relation at a more fundamental level of description. Moreover, due to this difficulty, the TUR for the underdamped Langevin dynamics with an arbitrary time-dependent driving force in a linearized form has not been found.

In this study, we derive a TUR for general underdamped Langevin systems with an arbitrary time-dependent protocol, including velocity-dependent forces like a magnetic Lorentz force breaking the time reversal symmetry. The cost function

\*jslee@kias.re.kr

<sup>†</sup>hgpark@kias.re.kr

of this TUR is expressed in terms of the EP without any kinetic quantity and an initial-state-dependent term which is negligible for the long observation-time limit. Furthermore, this TUR returns back to the original TUR of the overdamped dynamics [Eq. (1)] in the zero-mass limit when the driving forces and the current weight function do not include odd variables. Thus, our TUR can be regarded as the universal form of the TUR for general Langevin dynamics.

*Model and main results.* We consider a  $N$ -dimensional underdamped Langevin system driven by a force  $\mathbf{F}(\mathbf{x}, \mathbf{v}, t) = (F_1, \dots, F_N)$ , where  $\mathbf{x} = (x_1, \dots, x_N)$  and  $\mathbf{v} = (v_1, \dots, v_N)$  are the position and velocity vectors of the system, respectively. Dynamics of the  $i$ th component of the system ( $x_i, v_i$ ) is in contact with a thermal reservoir with temperature  $T_i$ . Then, the dynamics can be described by the following equation:

$$\dot{x}_i = v_i, \quad m_i \dot{v}_i = F_i(\mathbf{x}, \mathbf{v}, t) - \gamma_i v_i + \xi_i, \quad (2)$$

where  $m_i$ ,  $\gamma_i$ , and  $\xi_i$  are the  $i$ th mass, dissipation coefficient, and Gaussian white noise satisfying  $\langle \xi_i(t) \xi_j(t') \rangle = 2k_B \gamma_i T_i \delta_{ij} \delta(t - t')$  with zero mean, respectively. For convenience, we set the Boltzmann constant  $k_B = 1$  in the following discussion. A general time-dependent force  $F_i$  consists of two parts: reversible  $F_i^{\text{rev}}$  and irreversible  $F_i^{\text{ir}}$  forces, that is,  $F_i(\mathbf{x}, \mathbf{v}, t) = F_i^{\text{rev}}(\mathbf{x}, \mathbf{v}, t) + F_i^{\text{ir}}(\mathbf{x}, \mathbf{v}, t)$  with  $F_i^{\text{rev}}(\mathbf{x}, \mathbf{v}, t) = F_i^{\text{rev}\dagger}(\mathbf{x}, -\mathbf{v}, t)$  and  $F_i^{\text{ir}}(\mathbf{x}, \mathbf{v}, t) = -F_i^{\text{ir}\dagger}(\mathbf{x}, -\mathbf{v}, t)$ , where the “ $\dagger$ ” operation reverses signs of all odd parameters in the time reversal [28,30]. Without loss of generality, we can set

$$F_i^{\text{rev}}(\mathbf{x}, \mathbf{v}, t) = s f_i^{\text{rev}}(r\mathbf{x}, \mathbf{v}, \omega t), \\ F_i^{\text{ir}}(\mathbf{x}, \mathbf{v}, t) = f_i^{\text{ir}}(r\mathbf{x}, \mathbf{v}, \omega t), \quad (3)$$

where  $s, r, \omega$  are the scaling parameters for force, position, and time, respectively. Note that  $s$  is multiplied to the reversible force only, which is one of the key manipulations for deriving the TUR. We consider  $\Gamma = [\mathbf{x}_t, \mathbf{v}_t]_{t=0}^{t=\tau}$ , which denotes a trajectory of the system from  $t = 0$  to  $t = \tau$ , and a  $\Gamma$ -dependent current  $\Theta$  which has the following form:

$$\Theta_\tau(\Lambda) \equiv \int_0^\tau dt \Lambda(\mathbf{x}_t, \mathbf{v}_t, t; s, r, \omega) \cdot \mathbf{v}_t, \quad (4)$$

with the weight function vector

$$\Lambda = s \chi(r\mathbf{x}_t, \mathbf{v}_t, \omega t). \quad (5)$$

Note that the same scale parameter  $s$  is used for the weight function and the reversible force for later convenience.

Then, our first main result is the following *underdamped* TUR in terms of the underdamped TUR factor  $\mathcal{Q}^u$  as

$$\mathcal{Q}^u \equiv \frac{\text{Var}[\Theta_\tau]}{\Omega_\tau^2} (\Delta S_\tau^{\text{tot}} + \mathcal{I}) \geq 2, \quad (6)$$

where  $\Omega_\tau$  is defined as

$$\Omega_\tau \equiv \hat{h}_\tau(\Theta_\tau), \quad \text{where } \hat{h}_\tau \equiv \tau \partial_\tau - s \partial_s - r \partial_r - \omega \partial_\omega, \quad (7)$$

and  $\mathcal{I}$  is an initial-state-dependent term defined in Eq. (26) which depends on the dynamic details but becomes negligible in the large- $\tau$  limit. Equation (6) holds for processes with arbitrary time-dependent driving from an arbitrary initial state. This underdamped TUR resembles the overdamped TUR recently found in [19] with additional parameters  $s$  and  $r$ .

In principle,  $\Omega_\tau$  is operationally accessible by measuring the response of  $\langle \Theta_\tau \rangle$  with respect to a slight change of the observation time  $\tau$ , the reversible force magnitude  $s$ , the system scale  $r$ , and the driving speed  $\omega$ . However, this requires a full control of all mechanical forces, which is usually not available in experiments for complex systems with uncontrollable interaction forces, while it is still possible in numerical simulations. Thus, our TUR is experimentally accessible only for simple systems where all forces can be controlled. Note that the EP can be readily inferred from numerical simulations (or real experiments for simple systems) by measuring a proper current or a set of currents [31]. We emphasize that our underdamped TUR does not contain any kinetic term like dynamical activity. Furthermore, this TUR provides a much tighter bound, compared to the previous underdamped TURs [21,22,29,30], which will be explicitly shown in the examples below.

Another fascinating part of our underdamped TUR is that the overdamped TUR, Eq. (1), arises naturally by taking the zero-mass limit, in case of no velocity-dependent force. For simplicity, we consider a steady-state TUR without any time-dependent protocol and no time dependence in the weight function  $\Lambda$  of a current of interest ( $\omega = 0$ ), that is,

$$F_i = s f_i^{\text{rev}}(r\mathbf{x}) \quad \text{and} \quad \Lambda = s \chi(r\mathbf{x}_t). \quad (8)$$

Then, in the zero mass limit,  $\Omega_\tau$  and  $\mathcal{I}$  in Eq. (6) becomes

$$\Omega_\tau = -\langle \Theta_\tau \rangle \quad \text{and} \quad \mathcal{I} = 0 \quad (9)$$

in the steady state, which leads to the original TUR [Eq. (1)]. This is our second main result. The proofs of Eqs. (6) and (9) will be sketched later, and full details are given in Supplemental Material (SM) [32].

*Examples.* To illustrate the usefulness and validity of our main results, we concentrate on steady-state processes where  $\mathbf{F}$  and  $\Lambda$  have no explicit time dependence in the following examples. With these conditions, the underdamped TUR is simplified with

$$\Omega_\tau = \Omega_\tau^{\text{ss}} \equiv (1 - s \partial_s - r \partial_r) \langle \Theta_\tau \rangle, \quad (10)$$

in the steady state. We demonstrate this result in two examples below and an additional one on a molecular refrigerator in SM [32].

*Example 1: Free diffusion with drift.* Consider a displacement current in the free diffusion process of a Brownian particle with mass  $m$ , driven by a constant external force  $\mathbf{F}$ . We set  $s f \mathbf{e}_1 = \mathbf{F}^{\text{rev}}$ , where  $f$  is a constant and  $\mathbf{e}_1$  is the unit vector along the  $x_1$  axis. We choose the weight function  $\Lambda = s \mathbf{e}_1$ , yielding  $\Theta_\tau(\Lambda)$  as displacement at  $t = \tau$  from the initial position at  $t = 0$  along the  $x_1$  axis. Note that  $s$  is a scale parameter, which will be set to be unity after the whole calculation. This model was studied recently as a paradigmatic example for a conjecture of the underdamped TUR in one dimension [33].

As the steady-state velocity is  $\langle v_1 \rangle^{\text{ss}} = s f / \gamma_1$ , we get  $\langle \Theta_\tau \rangle = \tau s \langle v_1 \rangle^{\text{ss}} = \tau s^2 f / \gamma_1$ . Consequently, we obtain

$$\Omega_\tau^{\text{ss}} = (1 - s \partial_s) \langle \Theta_\tau \rangle = -\langle \Theta_\tau \rangle. \quad (11)$$




FIG. 1.  $Q^u$  (blue filled symbol) and  $Q$  (red open symbol) as a function of the observation time  $\tau$  for free diffusion with drift. Solid, dashed, and dash-dotted curves represent analytic results for  $m = 1$ , 0.1, and 0.01, respectively. The black dotted line indicates the lower bound, i.e., 2. Other system parameters are set to be unity,  $f = \gamma_1 = T_1 = 1$ .

Using Eqs. (6) and (11), the underdamped TUR for the free diffusion process with drift at  $s = 1$  becomes

$$Q^u = \frac{\text{Var}[\Theta_\tau]}{\langle \Theta_\tau \rangle^2} (\Delta S_\tau^{\text{tot}} + \mathcal{I}^{\text{fr}}) \geq 2, \quad (12)$$

where  $\Delta S_\tau^{\text{tot}} = \tau f^2 / (T_1 \gamma_1)$  and  $\mathcal{I}^{\text{fr}} = 2mf^2 / (T_1 \gamma_1^2)$ . Calculations of  $\Delta S_\tau^{\text{tot}}$ ,  $\mathcal{I}^{\text{fr}}$ , and  $\text{Var}[\Theta_\tau]$  are presented in SM [32]. Note that  $\mathcal{I}^{\text{fr}}$  vanishes in the zero-mass limit, confirming that our underdamped TUR in Eq. (12) returns back to the original TUR form in the overdamped limit. For a finite mass, the original TUR is recovered only when  $\mathcal{I}^{\text{fr}}$  is negligible in the large- $\tau$  limit.

Figure 1 shows analytic (curves) and numerical (dots) plots of  $Q^u$  and  $Q$  for various values of  $m$  as a function of  $\tau$ . The analytic expressions are presented in SM [32] and the numerical data are obtained by averaging over  $10^7$  trajectories from the Langevin equation. As expected from our underdamped TUR,  $Q^u$  is always above the lower bound of 2 for any observation time period  $\tau$ , and approaches the bound either in the zero-mass limit or in the large- $\tau$  limit. The conventional TUR factor  $Q$  approaches the bound from below (violations of the original TUR) in these limits. This example clearly demonstrates the importance of the initial-state dependent term  $\mathcal{I}$  in the underdamped dynamics for a finite  $\tau$ , which usually vanishes in the overdamped limit. The free-diffusion bound conjecture [33] also involves  $\tau$  in the lower bound, though it differs from our rigorous bound (see SM [32] for discussions).

*Example 2: Charged particle in a magnetic field.* The next example is the motion of a charged Brownian particle under a magnetic field  $B$  in a two-dimensional space [30,34]. The particle is trapped in a harmonic potential with stiffness  $k$  and driven by a nonconservative rotational force. Then, the total force is given by  $\mathbf{F} = \mathbf{F}^{\text{nc}} + \mathbf{F}^{\text{mag}} + \mathbf{F}^{\text{har}}$  with the nonconservative rotational force  $\mathbf{F}^{\text{nc}} = s\kappa(rx_2, -rx_1)$ , the Lorentz force induced by the magnetic field  $\mathbf{F}^{\text{mag}} = sB(v_2, -v_1)$ , and

the harmonic force  $\mathbf{F}^{\text{har}} = -s\kappa(rx_1, rx_2)$ . By regarding the magnetic field  $B$  as an odd-parity parameter, we treat the whole force  $\mathbf{F}$  as a reversible one. The opposite choice is also possible [30,35,36]. Here, we consider the case  $\gamma_1 = \gamma_2 \equiv \gamma$ ,  $m_1 = m_2 \equiv m$ , and  $T_1 = T_2 \equiv T$ . We are interested in the work current done by the nonconservative force, thus  $\mathbf{\Lambda} = \mathbf{F}^{\text{nc}}$ . By replacing the parameters as  $\kappa \rightarrow sr\kappa$ ,  $B \rightarrow sB$ , and  $k \rightarrow srk$  from the result of Ref. [34], the steady-state work current can be written as

$$\langle \Theta_\tau \rangle = \frac{2\tau r\kappa^2 T}{\gamma k/s + \kappa B - r\kappa^2 m/\gamma}, \quad (13)$$

with the stability condition  $\gamma k/s + \kappa B - r\kappa^2 m/\gamma > 0$ . Then, we obtain from Eq. (10)

$$\Omega_\tau^{\text{ss}} = -\frac{\gamma k/s + r\kappa^2 m/\gamma}{\gamma k/s + \kappa B - r\kappa^2 m/\gamma} \langle \Theta_\tau \rangle. \quad (14)$$

With dimensionless parameters  $B_0 = B/\gamma$ ,  $\kappa_0 = \kappa/k$ , and  $m_0 = mk/\gamma^2$ , the underdamped TUR at  $s = r = 1$  can be written as

$$Q^u = \frac{\text{Var}[\Theta_\tau]}{g^{\text{mag}} \langle \Theta_\tau \rangle^2} (\Delta S_\tau^{\text{tot}} + \mathcal{I}^{\text{mag}}) \geq 2, \quad (15)$$

with

$$g^{\text{mag}} = \left( \frac{1 + \kappa_0^2 m_0}{1 + \kappa_0 B_0 - \kappa_0^2 m_0} \right)^2, \quad (16)$$

$$\mathcal{I}^{\text{mag}} = \frac{2\kappa_0^2 [B_0^2 - 2\kappa_0 m_0 B_0 + 2m_0(1 + \kappa_0^2 m_0)]}{(1 + \kappa_0 B_0 - \kappa_0^2 m_0)^2}. \quad (17)$$

The derivation of  $\mathcal{I}^{\text{mag}}$  is shown in SM [32], and  $\text{Var}[\Theta_\tau]$  can be also calculated for any finite  $\tau$  by solving rather complex matrix differential equations numerically (not shown here, but see Ref. [37] for a sketch of derivations). The EP is given by the Clausius EP with the odd-parity choice of  $B$  [30,36], thus we obtain  $\Delta S_\tau^{\text{tot}} = \langle \Theta_\tau \rangle / T$  as the average heat current is equal to the average work current in the steady state.

In Fig. 2, we plot  $Q^u$  evaluated at various values of parameters against  $B_0$ . The parameter values of  $m_0$ ,  $\kappa_0$ , and  $\tau$  are randomly selected from the uniform distribution with ranges of  $[0, 1]$ ,  $[0, 10]$ , and  $[0, 10]$ , respectively, with fixed  $\gamma = k = T = 1$ . All points stay above the lower bound of 2, which turns out to be a very tight one for any value of  $B_0$ . In the large- $\tau$  limit,  $\mathcal{I}^{\text{mag}}$  is negligible and  $\text{Var}[\Theta_\tau]$  takes a simple form [34]. Then, the conventional TUR factor becomes

$$Q = \frac{\text{Var}[\Theta_\tau]}{\langle \Theta_\tau \rangle^2} \Delta S_\tau^{\text{tot}} = 2 \frac{1 + \kappa_0^2(1 + 3m_0) + \kappa_0^3 m_0 B_0}{(1 + \kappa_0 B_0 - \kappa_0^2 m_0)^2}, \quad (18)$$

which is larger than  $2g^{\text{mag}}$  under the stability condition  $1 + \kappa_0 B_0 - \kappa_0^2 m_0 > 0$ , which confirms our underdamped TUR, but can be smaller than the conventional bound of 2 for  $\kappa_0 B_0 > 0$ . The previous bound including dynamical activity [29,30] is very loose compared to our bound here (see Fig. S1 in Ref. [32]). It is interesting to note that, in the equilibrium limit ( $\kappa \rightarrow 0$ ),  $g^{\text{mag}} \approx 1$  and  $Q^u \simeq Q$  approaches 2 for large  $\tau$ .

In the zero-mass limit ( $m_0 = 0$ ), we get  $g^{\text{mag}} = 1/(1 + \kappa_0 B_0)^2$  and  $\mathcal{I}^{\text{mag}} = 2\kappa_0^2 B_0^2 / (1 + \kappa_0 B_0)^2$ . Thus, the original TUR is restored when  $B_0 = 0$  (no velocity-dependent force).




FIG. 2. Plot for  $Q^u$  of the charged particle in a magnetic field evaluated at various values of the system parameters and the observations time against  $B_0$ . The dashed line indicates the lower bound of 2.

With nonzero  $B_0$ , the broken time-reversal symmetry due to the Lorentz force is known to lower the TUR bound even in the overdamped limit [34,37]. Very recently, its lower bound for the conventional TUR factor  $\mathcal{Q}$  is rigorously obtained as  $2/(1+B_0^2)$  for general nonlinear forces with a finite  $\tau$  [37]. Our underdamped TUR also gives a lower bound for  $\mathcal{Q}$  from Eq. (15), which may be tighter than the above rigorous bound for the overdamped limit, depending on the parameter values.

*Sketch of TUR derivation.* Here we provide a brief sketch and essence of the TUR derivation. Full details are given in SM [32]. The Fokker-Planck (FP) equation of the probability distribution function  $P_t = P(\mathbf{x}, \mathbf{v}, t; s, r, \omega)$  for the Langevin equation, Eq. (2), can be written as

$$\partial_t P_t = \mathcal{L}P_t = \sum_i (\mathcal{L}_i^{\text{rev}} + \mathcal{L}_i^{\text{ir}})P_t, \quad (19)$$

where the FP operator  $\mathcal{L}$  is split into the reversible and irreversible parts as

$$\mathcal{L}_i^{\text{rev}} = -\left[ \partial_{x_i} v_i + \left( \frac{s}{m_i} \right) \partial_{v_i} f_i^{\text{rev}}(\mathbf{r}\mathbf{x}, \mathbf{v}, \omega t) \right], \quad (20)$$

$$\mathcal{L}_i^{\text{ir}} = -\frac{1}{m_i} \partial_{v_i} \left[ f_i^{\text{ir}}(\mathbf{r}\mathbf{x}, \mathbf{v}, \omega t) - \gamma_i v_i - \frac{\gamma_i T_i}{m_i} \partial_{v_i} \right]. \quad (21)$$

Now, we consider a *modified* dynamics, satisfying the following FP equation parametrized by  $\theta$ :

$$\partial_t P_{t,\theta} = \sum_i [\mathcal{L}_i^{\text{rev}} + (1+\theta)\mathcal{L}_i^{\text{ir}}]P_{t,\theta}, \quad (22)$$

which is called the  $\theta$  process. Then, it is straightforward to show that its solution is given by

$$P_{t,\theta} = P_\theta(\mathbf{x}, \mathbf{v}, t; s, r, \omega) = (1+\theta)^N P(\mathbf{x}_\theta, \mathbf{v}, t_\theta; s_\theta, r_\theta, \omega_\theta) \quad (23)$$

with the scaled variables and parameters as

$$\begin{aligned} \mathbf{x}_\theta &= (1+\theta)\mathbf{x}, & t_\theta &= (1+\theta)t, \\ s_\theta &= \frac{s}{1+\theta}, & r_\theta &= \frac{r}{1+\theta}, & \omega_\theta &= \frac{\omega}{1+\theta}, \end{aligned} \quad (24)$$

and the normalization factor  $(1+\theta)^N$ . Note from Eq. (23) that the initial distribution  $P_{0,\theta}$  at  $t=0$  is  $\theta$  dependent. This modification in the FP equation is equivalent to adding an extra force  $\theta\mathcal{Y}_i$  to the original process as

$$F_{i,\theta}(\mathbf{x}, \mathbf{v}, t) = F_i(\mathbf{x}, \mathbf{v}, t) + \theta\mathcal{Y}_i(\mathbf{x}_\theta, \mathbf{v}, t_\theta; s_\theta, r_\theta, \omega_\theta), \quad (25)$$

where  $\mathcal{Y}_i = J_i^{\text{ir}}/P_t$  with the irreversible current  $J_i^{\text{ir}}$  of the original process given by  $J_i^{\text{ir}} = \frac{1}{m_i} [f_i^{\text{ir}} - \gamma_i v_i - (\gamma_i T_i/m_i) \partial_{v_i}] P_t$ .

With this setup, we take a similar derivation procedure in Refs. [16,19], using the Cramér-Rao inequality [38,39], leading to the main result of Eq. (6) with

$$\begin{aligned} \Delta S_\tau^{\text{tot}} &= \sum_{i=1}^N \int_0^\tau dt \left\langle \frac{(m_i J_i^{\text{ir}})^2}{\gamma_i T_i P_t^2} \right\rangle, \\ \mathcal{I} &= 2 \int d\mathbf{x}_0 d\mathbf{v}_0 \frac{(\partial_\theta P_{0,\theta})^2|_{\theta=0}}{P_0} = 2\langle (N + \hat{h}' \ln P_0)^2 \rangle, \end{aligned} \quad (26)$$

where  $\hat{h}' \equiv \mathbf{x} \cdot \nabla - s\partial_s - r\partial_r - \omega\partial_\omega$ . Note that  $\mathcal{I}$  is determined by the derivatives of the initial Shannon entropy.

In order to find the TUR in the overdamped limit, we consider the case where the force and the current weight function are velocity-independent (thus,  $f_i^{\text{ir}} = 0$ ) as

$$F_i = s f_i^{\text{rev}}(\mathbf{r}\mathbf{x}, \omega t), \quad \Lambda_i = s \chi_i(\mathbf{r}\mathbf{x}, \omega t). \quad (27)$$

The corresponding overdamped FP equation of the probability distribution function  $\rho_t = \rho(\mathbf{x}, t; s, r, \omega)$  in the zero-mass limit is given as

$$\partial_t \rho_t = \sum_i \mathcal{L}_i^o \rho_t, \quad (28)$$

where the overdamped FP operator  $\mathcal{L}_i^o$  is given as

$$\mathcal{L}_i^o = -\frac{1}{\gamma_i} \partial_{x_i} [s f_i^{\text{rev}}(\mathbf{r}\mathbf{x}, \omega t) - T_i \partial_{x_i}]. \quad (29)$$

The overdamped limit of the underdamped  $\theta$  process can be obtained formally by the standard small-mass expansion method using the Brinkman's hierarchy [27]. In the presence of a velocity-dependent force such as a magnetic Lorentz force, the overdamped limit could become quite subtle [37], which is not considered here. However, with no irreversible force ( $f_i^{\text{ir}} = 0$ ), it can be easily seen from Eqs. (22) and (21) that the  $\theta$  process is simply given by the original process with the replacement of  $\gamma_i$  by  $(1+\theta)\gamma_i$ . Thus, we can immediately write down the FP equation for the  $\theta$  process in the overdamped limit as

$$\partial_t \rho_{t,\theta} = \left( \frac{1}{1+\theta} \right) \mathcal{L}_i^o \rho_{t,\theta}. \quad (30)$$

This is exactly the same as the *virtual-perturbation* FP equation in Refs. [16,19,31,40] with the relation of  $1+\epsilon = 1/(1+\theta)$  (the perturbation parameter  $\epsilon$ ). This clearly shows that the  $\theta$  process of Eq. (22) in the underdamped dynamics is a natural extension of that in the overdamped dynamics. The original TUR with Eq. (9) and its extension to the case with a time-dependent protocol [19] can be easily obtained (see details in SM [32]).

*Conclusion.* We derived the TUR for general underdamped Langevin systems with an arbitrary time-dependent driving



from an arbitrary initial state, including velocity-dependent forces. In contrast to the previously reported one, our result is experimentally accessible when all forces are controllable and its lower bound is much tighter. Therefore, this bound can be utilized to facilitate inferring the EP by measuring a current statistics and its response to a slight change of various system parameters. Furthermore, the original TUR for the overdamped Langevin dynamics can be understood as its zero-mass limit. This implies that our underdamped TUR provides a universal form of the tradeoff relation for general Langevin systems. It would be interesting to extend our result to systems with non-Markovian environmental noises such as

active-matter systems, which are known to be described by effective underdamped Langevin dynamics [41,42].

### ACKNOWLEDGMENTS

The authors acknowledge the Korea Institute for Advanced Study for providing computing resources (KIAS Center for Advanced Computation Linux Cluster System). This research was supported by the NRF Grant No. 2017R1D1A1B06035497 (H.P.) and the KIAS individual Grants No. PG013604 (H.P.), No. PG074002 (J.M.P.), and No. PG064901 (J.S.L.) at the Korea Institute for Advanced Study.

- 
- [1] A. C. Barato and U. Seifert, *Phys. Rev. Lett.* **114**, 158101 (2015).  
 [2] J. Li, J. M. Horowitz, T. R. Gingrich, and N. Fakhri, *Nat. Commun.* **10**, 1666 (2019).  
 [3] S. K. Manikandan, D. Gupta, and S. Krishnamurthy, *Phys. Rev. Lett.* **124**, 120603 (2020).  
 [4] T. R. Gingrich, G. M. Rotskoff, and J. M. Horowitz, *J. Phys. A: Math. Theor.* **50**, 184004 (2017).  
 [5] G. Benenti, K. Saito, and G. Casati, *Phys. Rev. Lett.* **106**, 230602 (2011).  
 [6] K. Brandner, K. Saito, and U. Seifert, *Phys. Rev. Lett.* **110**, 070603 (2013).  
 [7] K. Proesmans and C. Van den Broeck, *Phys. Rev. Lett.* **115**, 090601 (2015).  
 [8] M. Campisi and R. Fazio, *Nat. Commun.* **7**, 11895 (2016).  
 [9] N. Shiraishi, K. Saito, and H. Tasaki, *Phys. Rev. Lett.* **117**, 190601 (2016).  
 [10] V. Holubec and A. Ryabov, *Phys. Rev. Lett.* **121**, 120601 (2018).  
 [11] J. S. Lee and H. Park, *Sci. Rep.* **7**, 10725 (2017).  
 [12] J. S. Lee, S. H. Lee, J. Um, and H. Park, *J. Korean Phys. Soc.* **75**, 948 (2019).  
 [13] P. Pietzonka and U. Seifert, *Phys. Rev. Lett.* **120**, 190602 (2018).  
 [14] T. R. Gingrich, J. M. Horowitz, N. Perunov, and J. L. England, *Phys. Rev. Lett.* **116**, 120601 (2016).  
 [15] J. M. Horowitz and T. R. Gingrich, *Phys. Rev. E* **96**, 020103(R) (2017).  
 [16] Y. Hasegawa and T. Van Vu, *Phys. Rev. E* **99**, 062126 (2019).  
 [17] A. Dechant and S.-I. Sasa, *J. Stat. Mech: Theory Exp.* (2018) 063209.  
 [18] K. Liu, Z. Gong, and M. Ueda, *Phys. Rev. Lett.* **125**, 140602 (2020).  
 [19] T. Koyuk and U. Seifert, *Phys. Rev. Lett.* **125**, 260604 (2020).  
 [20] K. Proesmans and C. Van den Broeck, *Europhys. Lett.* **119**, 20001 (2017).  
 [21] P. P. Potts and P. Samuelsson, *Phys. Rev. E* **100**, 052137 (2019).  
 [22] K. Proesmans and J. M. Horowitz, *J. Stat. Mech.: Theory Exp.* (2019) 054005.  
 [23] A. C. Barato, R. Chetrite, A. Faggionato, and D. Gabrielli, *New J. Phys.* **20**, 103023 (2018).  
 [24] L. P. Fischer, P. Pietzonka, and U. Seifert, *Phys. Rev. E* **97**, 022143 (2018).  
 [25] Y. Hasegawa and T. Van Vu, *Phys. Rev. Lett.* **123**, 110602 (2019).  
 [26] K. Macieszczak, K. Brandner, and J. P. Garrahan, *Phys. Rev. Lett.* **121**, 130601 (2018).  
 [27] H. Risken and H. Haken, *The Fokker-Planck Equation: Methods of Solution and Applications* 2nd ed. (Springer, Berlin, 1989).  
 [28] A. Dechant and S.-I. Sasa, *Phys. Rev. E* **97**, 062101 (2018).  
 [29] T. Van Vu and Y. Hasegawa, *Phys. Rev. E* **100**, 032130 (2019).  
 [30] J. S. Lee, J.-M. Park, and H. Park, *Phys. Rev. E* **100**, 062132 (2019).  
 [31] A. Dechant, *J. Phys. A: Math. Theor.* **52**, 035001 (2019).  
 [32] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevE.104.L052102> for details, which includes Refs. [43–49].  
 [33] L. P. Fischer, H.-M. Chun, and U. Seifert, *Phys. Rev. E* **102**, 012120 (2020).  
 [34] H.-M. Chun, L. P. Fischer, and U. Seifert, *Phys. Rev. E* **99**, 042128 (2019).  
 [35] C. Kwon, J. Yeo, H. K. Lee, and H. Park, *J. Korean Phys. Soc.* **68**, 633 (2016).  
 [36] H.-M. Chun and J. D. Noh, *J. Stat. Mech: Theory Exp.* (2018) 023208.  
 [37] J.-M. Park and H. Park, *Phys. Rev. Research* **3**, 043005 (2021).  
 [38] H. Cramér, *Mathematical Methods of Statistics* (Princeton University Press, Princeton, NJ, 1999), Vol. 9.  
 [39] C. R. Rao, *Bull. Calcutta Math. Soc.* **37**, 81 (1945).  
 [40] A. Dechant and S.-I. Sasa, *Phys. Rev. Research* **3**, L042012 (2021).  
 [41] D. Mandal, K. Klymko, and M. R. DeWeese, *Phys. Rev. Lett.* **119**, 258001 (2017).  
 [42] E. Fodor, C. Nardini, M. E. Cates, J. Tailleur, P. Visco, and F. van Wijland, *Phys. Rev. Lett.* **117**, 038103 (2016).  
 [43] P. F. Cohadon, A. Heidmann, and M. Pinard, *Phys. Rev. Lett.* **83**, 3174 (1999).  
 [44] M. Pinard, P. F. Cohadon, T. Briant, and A. Heidmann, *Phys. Rev. A* **63**, 013808 (2000).  
 [45] J. Mertz, O. Marti, and J. Mlynek, *Appl. Phys. Lett.* **62**, 2344 (1993).  
 [46] G. Jourdan, G. Torricelli, J. Chevrier, and F. Comin, *Nanotechnology* **18**, 475502 (2007).  
 [47] S. Liang, D. Medich, D. M. Czajkowsky, S. Sheng, J.-Y. Yuan, and Z. Shao, *Ultramicroscopy* **84**, 119 (2000).  
 [48] K. H. Kim and H. Qian, *Phys. Rev. Lett.* **93**, 120602 (2004).  
 [49] L. Onsager and S. Machlup, *Phys. Rev.* **91**, 1505 (1953).