

Symmetries of many-body systems imply distance-dependent potentials

Jonathan Utterson ^{*} and Radek Erban [†]*Mathematical Institute, University of Oxford, Radcliffe Observatory Quarter, Woodstock Road, Oxford OX2 6GG, United Kingdom*

(Received 31 March 2023; accepted 21 June 2023; published 21 July 2023)

Considering an interatomic potential $U(\mathbf{q})$, where $\mathbf{q} = [\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N] \in \mathbb{R}^{3N}$ is a vector describing positions $\mathbf{q}_i \in \mathbb{R}^3$, it is shown that U can be defined as a function of the interatomic distance variables $r_{ij} = |\mathbf{q}_i - \mathbf{q}_j|$ provided the potential U satisfies some symmetry assumptions. Moreover, the potential U can be defined as a function of a proper subset of the distance variables r_{ij} , provided $N > 5$, with the number of distance variables used scaling linearly with the number of atoms N .

DOI: [10.1103/PhysRevE.108.014122](https://doi.org/10.1103/PhysRevE.108.014122)

I. INTRODUCTION

The theory of classical interatomic potentials has been developed for decades; a review of this research area is provided by Murrell *et al.* [1] and more recently by Ackland [2]. The basis of molecular modeling is dependent on creating a suitable potential energy function that defines the free-energy surface and dynamics of the system accurately while also balancing computational feasibility. One must compromise by reducing the degrees of freedom with some method of coarse graining [3]. A key way to do this is by explicitly constructing a potential energy function that reduces the complexity of the system. Many such function choices can naturally arise for a given system [4]. Pair potentials are commonly used to approximate potential energy contributions, though caution must be taken to use these appropriately [5]. Despite this, effective pair potentials in many classical circumstances have had fair degrees of success for decades in simulations of liquids [6–11].

To obtain more accurate results from thermodynamic calculations, many-body contributions are considered in the potential energy function [12,13]. An example potential incorporating two-body and three-body terms is the Stillinger-Weber potential [14], which accurately incorporates the geometry of silicon, meaning that, not only do the pairwise bonds between the silicon atoms matter, but so do the triangular substructures connecting neighboring atoms [15]. The embedded-atom-method potentials [16] incorporate an effective pairwise potential and a density-dependent contribution without explicitly using the geometric features. Progressing from pair potentials to those incorporating three-body terms and four-body terms, the most general interatomic potential

considered is a sum of all of these contributions, which can also include the single-body terms that arise when an external field is present. The n -body terms are explicitly evaluated given the coordinates of the N atoms, which can be thought of as vertices of a polygon (if coplanar) or a polyhedron. These n -body terms in the potential are then thought of as contributions arising from the n -gon substructures of the shape formed by the vertices. This forms the basis of fragmentation methods used in *ab initio* quantum chemistry; a summary and a closed-form expression for energy were presented by Richard *et al.* [17]. Tandem to this, cluster descriptions of many-body configurations [18] can also be used in conjunction with n -body expansions of the many-body potential [19], which differs from the previous method as this relies on the ordering of vertices as opposed to their position.

Nonreciprocal interactions, where pairwise forces do not obey Newton's third law [20], are applicable to colloidal physics [21], plasma physics [22,23], and active transport [24]. The statistical mechanics framework used to analyze such a system that exhibits nonreciprocity relies on defining an interatomic potential; Ivlev *et al.* [25] provided some pioneering analysis in this area. A question that has not been answered is characterizing when a general interatomic potential displays nonreciprocal interactions. For example, if the potential depends purely on pairwise distances, then reciprocity is a consequence; so one possible way to approach this problem is to study under what symmetries can we conclude that a general position-dependent potential function can be written as a function that depends purely on distances. Separately, this is a fundamental question that underpins classical potential theory and it is addressed in this paper in Theorems 1 and 2. We should note that thought into symmetries of a potential was undertaken by Kinghorn and Adamowicz [26]. It was used to analyze a specific functional form of potential developed, while in Sec. II we consider a general potential with the goal of understanding when distances are appropriate variables used to describe the potential function. The potential function U considered in this paper has translational, rotational, and reflectional symmetries as formulated in Definition 1, where we study systems of identical particles (atoms or, more generally, coarse-grained particles).

^{*}utterson@maths.ox.ac.uk[†]erban@maths.ox.ac.uk

We present proofs of Theorems 1 and 2 in Sec. III, where we also show that we require only a relatively small subset of distances to uniquely determine the potential as stated in Theorem 2. Limitations of this description are discussed in Sec. IV, where we also present some generalizations of Theorems 1 and 2 to mixtures of atoms of different types.

II. RESULTS

The configuration of a system of N atoms at positions \mathbf{q}_i , $i = 1, 2, \dots, N$, is defined as a $3N$ -dimensional vector $\mathbf{q} = (\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N) \in \mathbb{R}^{3N}$. We note that these can provisionally be thought of as vertices of an N -gon, or an N -polyhedron, assuming that $\mathbf{q}_i \neq \mathbf{q}_j$ for $i \neq j$. The lengths of edges are distances between atoms, which we define as

$$r_{ij} = |\mathbf{q}_j - \mathbf{q}_i| \quad \text{for } i, j = 1, 2, \dots, N. \quad (1)$$

In this paper we study potential functions $U : \mathbb{R}^{3N} \rightarrow \mathbb{R}$ called central potential functions, which satisfy certain symmetries as specified in Definition 1. These symmetries are (i) translational invariance, (ii) rotational invariance, (iii) reflectional invariance, and (iv) parity for i and j identical atoms. An example of potential satisfying the assumptions in Definition 1 is

$$U(\mathbf{q}) = \sum_{i < j}^N \Psi_2(r_{ij}) + \sum_{i < j < k}^N \Psi_3(r_{ij}, r_{ik}, r_{jk}), \quad (2)$$

where $\Psi_2 : [0, \infty) \rightarrow \mathbb{R}$ and $\Psi_3 : [0, \infty)^3 \rightarrow \mathbb{R}$ are two-body and three-body potentials which depend on distances between atoms.

Definition 1. A function $U : \mathbb{R}^{3N} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ is called a central potential function provided it takes finite values on the subset

$$\Omega = \{\mathbf{q} \in \mathbb{R}^{3N} \mid \mathbf{q} = \{\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N\} \text{ with } \mathbf{q}_i \neq \mathbf{q}_j \text{ for } i \neq j\}$$

and for any $\mathbf{q} \in \Omega$ it satisfies

- (i) $U(\mathbf{q}_1 + \mathbf{c}, \mathbf{q}_2 + \mathbf{c}, \dots, \mathbf{q}_N + \mathbf{c}) = U(\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N)$ for all translations $\mathbf{c} \in \mathbb{R}^3$,
- (ii) $U(R\mathbf{q}_1, R\mathbf{q}_2, \dots, R\mathbf{q}_N) = U(\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N)$ for all rotations $R \in \text{SO}(3)$,
- (iii) $U(Q\mathbf{q}_1, Q\mathbf{q}_2, \dots, Q\mathbf{q}_N) = U(\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N)$ for all reflections Q satisfying that all points \mathbf{q}_i , $i = 1, 2, \dots, N$, lie on one side of the plane of reflection,
- (iv) $U(\mathbf{q}_1, \dots, \mathbf{q}_i, \dots, \mathbf{q}_j, \dots, \mathbf{q}_N) = U(\mathbf{q}_1, \dots, \mathbf{q}_j, \dots, \mathbf{q}_i, \dots, \mathbf{q}_N)$ for any $i, j = 1, 2, \dots, N$.

The symmetries considered in Definition 1 are satisfied by other generalizations of the example potential (2), which include n -body terms depending only on the distances (1) between atoms. In fact, the symmetries (i)–(iv) imply that the potential $U : \mathbb{R}^{3N} \rightarrow \mathbb{R}$ can be written as a function of distances. We have the following theorem, which we prove in Sec. III.

Theorem 1. A central potential function $U : \mathbb{R}^{3N} \rightarrow \mathbb{R}$ can be written as

$$\phi : [0, \infty)^{N(N-1)/2} \rightarrow \mathbb{R},$$

where the $N(N-1)/2$ inputs are interpreted as the set of all pairwise distances (1) between atoms.

Considering $N = 2$, Theorem 1 states that a central potential function U of six variables can be written as a function ϕ of one variable r_{12} . Consequently, Theorem 1 reduces the dimensionality of the potential U for any $N < 7$. If $N = 7$ then we have $3N = N(N-1)/2 = 21$ and the 21-dimensional state space \mathbb{R}^{3N} corresponds to the 21 distance variables (1). Since the dimension of the state space scales as $O(N)$ and the number of distances scales as $O(N^2)$, Theorem 1 can be further improved by considering only a subset of the distance variables (1). In Sec. III we also prove the following result.

Theorem 2. Let $N \geq 4$. Then a central potential function $U : \mathbb{R}^{3N} \rightarrow \mathbb{R}$ can be written as

$$\phi : [0, \infty)^{4N-10} \rightarrow \mathbb{R},$$

where the $4N - 10$ inputs are a subset of the set of all pairwise distances (1).

Considering $N = 4$ and $N = 5$, we have $4N - 10 = 6$ and $4N - 10 = 10$, respectively. In particular, Theorems 1 and 2 state the same conclusion for $N = 4$ and $N = 5$. Theorem 2 improves the result of Theorem 1 for $N > 5$. We will prove Theorems 1 and 2 together in Sec. III by considering the cases $N = 2, N = 3, N = 4, N = 5$, and $N > 5$.

Applying Theorem 2 to our example potential (2), we observe that it reduces the number of independent variables for $N > 5$. In particular, while the function ϕ constructed in the proof of Theorem 2 depends only on distances (1), it is not given in the form (2).

In addition to central potential functions satisfying conditions in Definition 1, there are potentials to which Theorems 1 and 2 are not applicable. For example, if the potential U corresponds to an external nonuniform field Ψ_1 , then we have

$$U(\mathbf{q}) = \sum_{i=1}^N \Psi_1(\mathbf{q}_i)$$

and U will not satisfy the conditions in Definition 1 nor will it be possible to write as a function of pairwise distances (1). Assuming that there is no external field present and that we have a system of N identical atoms interacting [i.e., U satisfies condition (iv) in Definition 1], then we can formally write it as a sum of the n -body interactions for $2 \leq n \leq N$ in the form

$$U(\mathbf{q}) = \sum_{i < j}^N U_2(\mathbf{q}_i, \mathbf{q}_j) + \sum_{i < j < k}^N U_3(\mathbf{q}_i, \mathbf{q}_j, \mathbf{q}_k) + \dots + U_N(\mathbf{q}_1, \dots, \mathbf{q}_N), \quad (3)$$

where we can naturally think about n polyhedrons of atoms as the input to the potential function, but these are fixed in space and a natural assumption is that given this input, it should not matter where we fix this polyhedron [leading to translational invariance (i)] or how we orient this polyhedron [rotational invariance (ii)]. One slightly more subtle assumption is that we should be allowed to reflect our polyhedron in any plane that keeps the polyhedron on one side [reflectional symmetry (iii)].

One difference between symmetries (i) and (ii) and symmetries (iii) and (iv) is that the former are continuous symmetries whereas the reflection invariance (iii) and parity (iv) are not. Noether's theorem [27] states that each continuous symmetry gives rise to a corresponding conserved

quantity (in a closed system). In particular, translational invariance (i) leads to conserved linear momentum (which is a consequence of reciprocity of forces) and rotational invariance (ii) gives rise to conserved angular momentum. In the next section we provide a proof that functions obeying symmetries (i)–(iv) should only rely on distances and we also show that a proper subset of pairwise distances for $N > 5$ can be used to describe the potential function U .

III. PROOFS OF THEOREMS 1 AND 2

We prove Theorems 1 and 2 together by considering the cases $N = 2$, $N = 3$, $N = 4$, and $N = 5$, followed by an inductive argument for $N > 5$. We define displacement vectors by

$$\mathbf{\Delta}_{ij} = \mathbf{q}_j - \mathbf{q}_i \quad \text{for } i, j = 1, 2, \dots, N, \quad (4)$$

i.e., we have $r_{ij} = |\mathbf{\Delta}_{ij}|$.

A. Case $N = 2$

We define the function $\phi : [0, \infty) \rightarrow \mathbb{R}$ by

$$\phi(s) = U(\mathbf{0}, s\hat{\mathbf{k}}) = U(0, 0, 0, 0, 0, s), \quad (5)$$

where $\hat{\mathbf{k}}$ is a unit vector in the direction of the positive z axis and $\mathbf{0} = [0, 0, 0]$. Given atom positions $\mathbf{q}_1, \mathbf{q}_2 \in \mathbb{R}^3$, we translate the configuration to position atom 1 at the origin. Using symmetry (i) in Definition 1, we have $U(\mathbf{q}_1, \mathbf{q}_2) = U(\mathbf{0}, \mathbf{\Delta}_{12})$. We then rotate the axes using rotation $R_1 \in \text{SO}(3)$ such that the displacement vector connecting the two atoms is aligned with the positive z axis, giving $R_1\mathbf{\Delta}_{12} = r_{12}\hat{\mathbf{k}}$, while maintaining $R_1\mathbf{0} = \mathbf{0}$. Using symmetry (ii) in Definition 1, we have

$$U(\mathbf{q}_1, \mathbf{q}_2) = U(\mathbf{0}, \mathbf{\Delta}_{12}) = U(\mathbf{0}, r_{12}\hat{\mathbf{k}}) = \phi(r_{12}),$$

where the last equality follows from our definition (5). This concludes the proof of Theorem 1 for $N = 2$.

B. Case $N = 3$

Given atom positions $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3 \in \mathbb{R}^3$, we consider the function $U(\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3)$. Using symmetry (i) in Definition 1, we translate the configuration to position atom 1 at the origin and consequently we have

$$U(\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3) = U(\mathbf{0}, \mathbf{\Delta}_{12}, \mathbf{\Delta}_{13}).$$

Given that we have three axes to rotate around, we can always find a rotation R_1 such that $R_1\mathbf{\Delta}_{12} = r_{12}\hat{\mathbf{k}}$, as we did in the $N = 2$ case. Using symmetry (ii), we have

$$U(\mathbf{0}, \mathbf{\Delta}_{12}, \mathbf{\Delta}_{13}) = U(\mathbf{0}, r_{12}\hat{\mathbf{k}}, R_1\mathbf{\Delta}_{13}).$$

We note that we can find a rotation R_2 about the z axis which rotates the triangle defined by the transformed atom positions $\mathbf{0}$, $r_{12}\hat{\mathbf{k}}$, and $R_1\mathbf{\Delta}_{13}$ to a planar triangle in the $x-z$ plane, as demonstrated in Fig. 1. Using symmetry (ii) again, we have

$$U(\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3) = U(\mathbf{0}, r_{12}\hat{\mathbf{k}}, R_2R_1\mathbf{\Delta}_{13}).$$

However, the key point is that $R_2R_1\mathbf{\Delta}_{13}$ is uniquely defined by the triangle with lengths r_{12} , r_{13} , and r_{23} , the angles of which




FIG. 1. Schematic of the constructive method in aid of the proof for the case $N = 3$.

can be calculated using the cosine rule, i.e., $R_2R_1\mathbf{\Delta}_{13}$ can be expressed as

$$\left[\sqrt{r_{13}^2 - \left(\frac{r_{13}^2 + r_{12}^2 - r_{23}^2}{2r_{12}} \right)^2}, 0, \frac{r_{13}^2 + r_{12}^2 - r_{23}^2}{2r_{12}} \right]. \quad (6)$$

Therefore, there exists a function $\phi : [0, \infty)^3 \rightarrow \mathbb{R}$ such that $U(\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3) = \phi(r_{12}, r_{13}, r_{23})$ for any \mathbf{q}_1 , \mathbf{q}_2 , and \mathbf{q}_3 , confirming Theorem 1 for $N = 3$.

C. Case $N = 4$

Given atom positions $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3, \mathbf{q}_4 \in \mathbb{R}^3$, these can be thought of as defining the vertices of a tetrahedron (or if coplanar a quadrilateral). Following similar steps as in the case $N = 3$ in Sec. III B, we translate atom 1 to the origin, apply rotation R_1 to orient the displacement vector $\mathbf{\Delta}_{12}$ with the positive z axis, and then do a second rotation R_2 that fixes the triangle formed by the vertices of atoms 1, 2, and 3 in the $x-z$ plane. As in Sec. III B, we have

$$U(\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3, \mathbf{q}_4) = U(\mathbf{0}, r_{12}\hat{\mathbf{k}}, R_2R_1\mathbf{\Delta}_{13}, R_2R_1\mathbf{\Delta}_{14}).$$

Using Eq. (6), we know that $R_2R_1\mathbf{\Delta}_{13}$ is determined entirely by distances r_{12} , r_{13} , and r_{23} . All that remains to be shown is that $R_2R_1\mathbf{\Delta}_{14}$ is determined by pairwise distances. We note that the triangle formed by atoms 1, 2, and 3 (denoted by ABC in the lower part of our illustration of the proof in Fig. 2) is uniquely determined (after orienting one side with the positive z axis). Consequently, this fixes the side BC . On the other hand, the triangle BCD is uniquely determined (as one side BC is fixed) by distances r_{23} , r_{24} , and r_{34} . These can be thought of as two triangles which can rotate around a hinge BC , so to determine the vector $R_2R_1\mathbf{\Delta}_{14}$ we necessarily need the final distance r_{14} that gives the angle between the planes containing triangles ABC and BCD (two configurations are illustrated in Fig. 2). If triangles ABC and BCD are coplanar, the set of all pairwise distances, with this orientation, will give a unique




FIG. 2. Schematic of the constructive method in aid of the proof for $N = 4$. For clarity we have highlighted only the additional three displacement vectors, though the triangle formed by vertices $\{1, 2, 3\}$ lying in the x - z plane is the same as in Fig. 1.

description of $R_2R_1\Delta_{14}$. If these triangles are not coplanar, this final distance gives two possible vectors for $R_2R_1\Delta_{14}$. These correspond to a unique $R_2R_1\Delta_{14}$ and the copy obtained by reflection in the plane containing triangle ABC . However, by property (iii) we know that if we reflect in the plane containing ABC with a matrix denoted by Q , then

$$U(\mathbf{0}, r_{12}\hat{\mathbf{k}}, R_2R_1\Delta_{13}, R_2R_1\Delta_{14}) = U(\mathbf{0}, r_{12}\hat{\mathbf{k}}, R_2R_1\Delta_{13}, QR_2R_1\Delta_{14}).$$

Therefore, there exists a function $\phi : [0, \infty)^6 \rightarrow \mathbb{R}$ such that $U(\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3, \mathbf{q}_4) = \phi(r_{12}, r_{13}, r_{14}, r_{23}, r_{24}, r_{34})$ for any $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3$, and \mathbf{q}_4 , confirming Theorem 1 for $N = 4$.

D. Case $N = 5$

To proceed in this case, we note that any N -vertex polyhedron can be made by adding a single vertex to an $N - 1$ polyhedron or polygon (in the case where all other points

are coplanar). The task at hand, as in the case $N = 4$ in Sec. III C, is being able to determine the displacement vectors once we have translated and rotated the configuration such that $R_2R_1\Delta_{12} = r_{12}\hat{\mathbf{k}}$ is aligned with the positive z axis.

An $N = 5$ polyhedron can be constructed from adding a vertex onto either a preexisting $N = 4$ polyhedron (at most three points are coplanar) or an $N = 4$ polygon (where all points are coplanar). In the first case, we may take any three vertices on the preexisting polyhedron; we call these vertices the transformed positions of atoms 2, 3, and 4 [by property (iv) in Definition 1]. If we know r_{25} , then this fifth vertex must lie on a sphere of radius r_{25} , with the transformed position of atom 2 as the center; we denote this by S_2 . Similarly, we construct S_3 and S_4 as spheres of radii r_{35} and r_{45} , respectively. This is illustrated in Fig. 3(a). The fifth vertex lies at the intersection of three spheres S_2, S_3 , and S_4 , which contains at most two points. If it contains exactly two points, then we need another distance r_{15} (which is the distance from the vertex in the preexisting polyhedron that was not used as a center of spheres S_2, S_3 , or S_4) to determine which of those two positions is correct [see Fig. 3(b)]. In this way, four more distances are used to specify all of the vertices of the $N = 5$ polyhedron. Therefore, there exists a function $\phi : [0, \infty)^{10} \rightarrow \mathbb{R}$ such that $U(\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3, \mathbf{q}_4, \mathbf{q}_5) = \phi(r_{12}, r_{13}, r_{14}, r_{15}, r_{23}, r_{24}, r_{25}, r_{34}, r_{35}, r_{45})$ for any $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3, \mathbf{q}_4$, and \mathbf{q}_5 .

To arrive at this conclusion, we used an assumption that four points are not coplanar. If this is not the case, then we need fewer distances for the specific configuration. For example, if the four preexisting vertices are coplanar, utilizing the sphere approach for any three of those vertices will result again in two possible positions for vertex 5; however, using the pairwise distance between this vertex and the new vertex gives no information, as the fourth point lies on the plane of symmetry formed by the spheres. In this case we use property (iii), considering the reflective symmetry about this plane to argue that we have determined all displacement vectors with this orientation uniquely up to a reflection in the plane containing vertices 1, 2, 3, and 4. In this case, we do




FIG. 3. (a) The intersection of three spheres, based on three known centers and radii, is used to position an additional vertex. (b) A fourth vertex, chosen noncoplanar to the three vertices used to construct the spheres previously, is used to uniquely determine the fourth vertex position.

not need the fourth distance mentioned above, and evaluating $U(\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3, \mathbf{q}_4, \mathbf{q}_5)$ is possible with the nine pairwise distances. The tenth distance is also not needed if the intersection of spheres S_2 , S_3 , and S_4 is exactly equal to one point (when vertices 2, 3, and 4 are collinear). Thus we have proven Theorems 1 and 2 in the case $N = 5$.

E. Case $N > 5$

We inductively prove that a setup similar to that in the $N = 5$ case in Sec. III D works by constructing polyhedra of higher order by the addition of a new vertex, that is, the $N - 1$ case requires the set of distances D_{N-1} to evaluate $U(\mathbf{q}_1, \dots, \mathbf{q}_{N-1})$, where $|D_{N-1}| = 4(N - 1) - 10$, which is true for the base case of $N = 5$. The most general case to consider is when we have an $N - 1$ polyhedron before we introduce the new vertex. In this case, any three of the $N - 1$ vertices can be chosen, say, i , j , and k . The three distances r_{iN} , r_{jN} , and r_{kN} are used to create three intersecting spheres and two potential positions for vertex N . We use vertex l , which is not coplanar to i , j , and k , and distance r_{lN} determines this position uniquely. Therefore, the required set of distances to evaluate $U(\mathbf{q}_1, \dots, \mathbf{q}_N)$ is $D_N = D_{N-1} \cup \{r_{iN}, r_{jN}, r_{kN}, r_{lN}\}$, i.e., we need $4N - 10$ pairwise distances. If there are at least four coplanar points, then we only need three additional pairwise distances (so we would only need $4N - 11$ pairwise distances). Since the inductive step holds for all N and it works for the base case of $N = 5$, this concludes our proofs of Theorems 1 and 2 for all N .

IV. DISCUSSION AND CONCLUSIONS

Theorems 1 and 2 show that symmetries of the many-body system imply that the potential U can be written in a form which only depends on pairwise distances between atoms. Since U is a function of the $3N$ -dimensional state space, Theorem 1 provides a nonlinear transformation of U from a function of $3N$ variables \mathbf{q} into a function of $N(N - 1)/2$ distance variables, which is not optimal, as it is shown in Theorem 2 where the number of distance variables scales linearly with N .

Considering the example potential (2), which depends on all $N(N - 1)/2$ distance variables, Theorem 2 provides a reduction of the number of distance variables to $O(N)$. However, if we use the resulting form of the potential ϕ this does not directly translate to $O(N)$ complexity of evaluating ϕ . To illustrate this, let us consider our example potential (2) with $\Psi_2(r_{ij}) = 1/r_{ij}$ and $\Psi_3 \equiv 0$, giving

$$U(\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N) = \sum_{\{i,j\} \in \mathcal{S}} \frac{1}{r_{ij}} + \sum_{\{i,j\} \notin \mathcal{S}} \frac{1}{r_{ij}}, \quad (7)$$

where r_{ij} is defined by (1) and \mathcal{S} is the set of pairs of indices $\{i, j\}$ corresponding to the subset of distances which is used to define ϕ in Theorem 2.

Theorem 2 shows that the number of elements in the set \mathcal{S} scales as $O(N)$, i.e., the number of terms in the first sum on the right-hand side of (7) is $O(N)$, while the number of terms in the second sum on the right-hand side of (7) scales as

$O(N^2)$. Considering $\{i, j\} \notin \mathcal{S}$, we can find $k \in \{1, 2, \dots, N\}$ such that $\{i, k\} \in \mathcal{S}$ and $\{j, k\} \in \mathcal{S}$. In particular, the distance r_{ij} for $\{i, j\} \notin \mathcal{S}$ can be expressed in terms of distances r_{ik} and r_{jk} using the cosine rule. Therefore, we can find an explicit form of the potential U as a function of $O(N)$ distances corresponding to the indices in the set \mathcal{S} . However, the second term in the form (7) will contain summations over $O(N^2)$ terms, that is, Theorem 2 does not reduce the $O(N^2)$ complexity of calculations of ϕ . It has been included to illustrate that the number of distance variables needed scales linearly with N in the same way the dimension of the phase space scales linearly with N . Theorem 2 provides a nonlinear transformation of U from a function of $3N$ variables \mathbf{q} into a function ϕ of $4N - 10$ distance variables for $N \geq 4$. While it improves the $O(N^2)$ scaling of Theorem 1, this result does not reduce the number of independent variables for $N \geq 11$. If $N = 10$, then $3N = 4N - 10$ and the number of distance variables used to define ϕ in Theorem 2 is the same as the number of independent variables defining U , while Theorem 2 provides a reduction in dimensionality for $N < 10$.

Theorem 1 has been formulated as an implication, stating that symmetries (i)–(iv) of a central potential function in Definition 1 imply that the potential can be written as a function of pairwise distances (1). However, translations, rotations, and reflections are Euclidean isometries, preserving pairwise distances between atoms, so a partial inverse of Theorem 1 also holds, i.e., any potential given as a function of pairwise distances satisfies symmetry assumptions (i)–(iii). Property (iv) states that we consider systems of identical particles in this paper. In particular, symmetries (i)–(iv) are both necessary and sufficient conditions for a potential to be expressed as a function of pairwise distances for systems of identical atoms.

We can generalize Theorems 1 and 2 to mixtures of particles, i.e., for systems when symmetry (iv) in Definition 1 does not hold. Then properties (i)–(iii) of potential U imply that it can be expressed as a function of pairwise distances. If we further reduce the number of symmetries the potential U has, then we can find potential functions which cannot be expressed as a function of pairwise distances. For example,

$$U(\mathbf{q}) = U(\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_N) = \mathbf{d} \cdot (\mathbf{q}_2 - \mathbf{q}_1)$$

for any nonzero constant vector \mathbf{d} satisfies the translational symmetry (i), but not the rotational symmetry (ii). An example of a potential function satisfying the rotational symmetry (ii) but not the translational symmetry (i) is $U(\mathbf{q}) = |\mathbf{q}_1|$. In fact, symmetries (i)–(iii) are both necessary and sufficient conditions for a potential to be expressed as a function of pairwise distances (for systems of nonidentical particles).

ACKNOWLEDGMENT

This work was supported by the Engineering and Physical Sciences Research Council (Grant No. EP/V047469/1) and by the Royal Society (Grant No. RGF\EA\180058).

- [1] J. N. Murrell, S. Carter, S. C. Farantos, P. Huxley, and A. J. C. Varandas, *Molecular Potential Energy Functions* (Wiley, New York, 1984).
- [2] G. J. Ackland, in *Comprehensive Nuclear Materials*, edited by R. J. M. Konings (Elsevier, Amsterdam, 2012), pp. 267–291.
- [3] W. G. Noid, Perspective: Coarse-grained models for biomolecular systems, *J. Chem. Phys.* **139**, 090901 (2013).
- [4] M. A. van der Hoef and P. Madden, Three-body dispersion contributions to the thermodynamic properties and effective pair interactions in liquid argon, *J. Chem. Phys.* **111**, 1520 (1999).
- [5] A. A. Louis, Beware of density dependent pair potentials, *J. Phys.: Condens. Matter* **14**, 9187 (2002).
- [6] M. Sprik, in *Computer Simulation in Chemical Physics*, edited by M. P. Allen and D. J. Tildesley, NATO Advanced Studies Institute, Series C: Mathematical and Physical Sciences (Springer, Dordrecht, 1993), Vol. 397.
- [7] C.-L. Zhao, D.-X. Zhao, C.-C. Bei, X.-N. Meng, S. Li, and Z.-Z. Yang, Seven-site effective pair potential for simulating liquid water, *J. Phys. Chem. B* **123**, 4594 (2019).
- [8] J.-M. Bomont and J.-L. Bretonnet, An effective pair potential for thermodynamics and structural properties of liquid mercury, *J. Chem. Phys.* **124**, 054504 (2006).
- [9] J. Utterson and R. Erban, On standardised moments of force distribution in simple liquids, *Phys. Chem. Chem. Phys.* **24**, 5646 (2022).
- [10] K. O. Waseda, Structure and effective pair potential of liquid silicon, *Jpn. J. Appl. Phys.* **35**, 151 (1996).
- [11] R. Erban and S. J. Chapman, *Stochastic Modelling of Reaction-Diffusion Processes* (Cambridge University Press, Cambridge, 2020).
- [12] S. J. Plimpton and A. P. Thompson, Computational aspects of many-body potentials, *MRS Bull.* **37**, 513 (2012).
- [13] G. A. Cisneros, K. T. Wikfeldt, L. Ojamäe, J. Lu, Y. Xu, H. Torabifard, A. P. Bartók, G. Csányi, V. Molinero, and F. Paesani, Modeling molecular interactions in water: From pairwise to many-body potential energy functions, *Chem. Rev.* **116**, 7501 (2016).
- [14] F. H. Stillinger and T. A. Weber, Computer simulation of local order in condensed phases of silicon, *Phys. Rev. B* **31**, 5262 (1985).
- [15] R. Biswas and D. R. Hamann, New classical models for silicon structural energies, *Phys. Rev. B* **36**, 6434 (1987).
- [16] M. S. Daw and M. I. Baskes, Embedded-atom method: Derivation and application to impurities, surfaces, and other defects in metals, *Phys. Rev. B* **29**, 6443 (1984).
- [17] R. M. Richard, K. U. Lao, and J. M. Herbert, Understanding the many-body expansion for large systems. I. Precision considerations, *J. Chem. Phys.* **141**, 014108 (2014).
- [18] J. M. Sanchez, F. Ducastelle, and D. Gratias, Generalized cluster description of multicomponent systems, *Physica A* **128**, 334 (1984).
- [19] R. Drautz, M. Fähnle, and J. M. Sanchez, General relations between many-body potentials and cluster expansions in multicomponent systems, *J. Phys.: Condens. Matter* **16**, 3843 (2004).
- [20] S. A. M. Loos, S. Arabha, A. Rajabpour, A. Hassanali, and É. Roldán, Nonreciprocal forces enable cold-to-hot heat transfer between nanoparticles, *Sci. Rep.* **13**, 4517 (2023).
- [21] V. N. Tsytovich, Dust plasma crystals, drops, and clouds, *Phys. Usp.* **40**, 53 (1997).
- [22] E. A. Lisin, O. F. Petrov, E. A. Sametov, O. S. Vaulina, K. B. Stratsenko, M. M. Vasiliev, J. Carmona-Reyes, and T. W. Hyde, Experimental study of the nonreciprocal effective interactions between microparticles in an anisotropic plasma, *Sci. Rep.* **10**, 13653 (2020).
- [23] S. A. Khrapak, A. V. Ivlev, and G. Morfill, Interaction potential of microparticles in a plasma: Role of collisions with plasma particles, *Phys. Rev. E* **64**, 046403 (2001).
- [24] R. Soto and R. Golestanian, Self-Assembly of Catalytically Active Colloidal Molecules: Tailoring Activity Through Surface Chemistry, *Phys. Rev. Lett.* **112**, 068301 (2014).
- [25] A. V. Ivlev, J. Bartnick, M. Heinen, C. R. Du, V. Nosenko, and H. Lowen, Statistical Mechanics where Newton’s Third Law is Broken, *Phys. Rev. X* **5**, 011035 (2015).
- [26] D. B. Kinghorn and L. Adamowicz, A new N -body potential and basis set for adiabatic and non-adiabatic variational energy calculations, *J. Chem. Phys.* **106**, 8760 (1997).
- [27] V. I. Arnold, *Mathematical Methods of Classical Mechanics*, Graduate Texts in Mathematics Vol. 60 (Springer, Berlin, 1989).