

Space-time evolution of a beam-plasma instability in strongly correlated plasmas

Z. C. Tao

*Department of Physics, Boston College, Chestnut Hill, Massachusetts 02167
and Department of Physics, University of Western Ontario, London, Ontario, Canada N6A 3K7**

A. K. Ram and A. Bers

Plasma Fusion Center, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

G. Kalman

Department of Physics, Boston College, Chestnut Hill, Massachusetts 02167

(Received 4 May 1993)

We examine the interaction of an electron beam penetrating a strongly correlated plasma. Conditions are established for this unstable interaction to lead to an absolute instability. We propose the absolute instability as a means for probing the strongly correlated plasmas.

PACS number(s): 52.35.Qz, 52.40.Mj, 52.25.Mq

The collective-mode structure of a strongly correlated (strongly coupled) plasma has characteristic features [1], distinguishing it from the mode structure of a weakly correlated system, which is well described by the Vlasov or random-phase approximation. Most importantly, the dispersion ($\partial\omega/\partial k$) of the plasmon mode is diminished and becomes *negative* at a critical value of the coupling parameter. This latter represents the ratio of the potential to kinetic energy and is conveniently given either as $\Gamma = e^2/k_B Ta$ (a being the Wigner-Seitz radius) or as the plasma parameter $\gamma = 1/4\pi N_D$ defined through the number of electrons in the Debye cube $N_D = n/\kappa_D^3$, where $\kappa_D = \sqrt{4\pi\beta ne^2}$ is the Debye wave number. Strongly coupled plasmas occur under various physical circumstances. Here we consider strongly coupled electron plasmas and study the high-frequency dynamics of a classical plasma. Our description of the physical scenario and our conclusions remain valid both for strongly coupled ionic plasmas and for strongly coupled degenerate electron plasmas as well. (A conversion of the coupling parameters pertaining to degenerate and classical plasmas, respectively, can be easily effected [2].)

The critical value of the plasma parameter γ has been determined by computer simulations [3], by theoretical calculations [4,5], and by experiments on alkali metals [6], converging to the value $\gamma = 30$ –50. When an electron beam penetrates a plasma, a beam-plasma instability develops. In this Rapid Communication we point out that, due to the change of the dispersion, the space-time evolution of the beam-plasma instability changes character as the correlations become sufficiently strong.

The physical reason for the negative dispersion in the strongly correlated system is that the correlations induce a quasilocalization of the particles in a pattern that exhibits a short-range order. It is the interaction of the beam particle with the ordered structure that is ultimately responsible for the change of the character of the instability.

The space-time evolution of the instability in an ob-

server frame moving with velocity V , in the nonrelativistic limit, is described [7] by the Green function

$$G(x', t) = \int_L \frac{d\omega'}{2\pi} \int_F \frac{dk}{2\pi} e^{i(kx' - \omega't)} \frac{1}{\epsilon_v(k, \omega')} \\ = \int_L \frac{d\omega}{2\pi} \int_F \frac{dk}{2\pi} e^{i(kx - \omega t)} \frac{1}{\epsilon(k, \omega + kV)}. \quad (1)$$

Here $x' = x + Vt$; $\epsilon(k, \omega)$ is the dielectric function for the system under consideration. L and F are the Laplace and Fourier contours in the ω and k planes, respectively. The behavior of the Green function $G(x', t)$ is totally determined by the analytic nature of the dielectric function $\epsilon(k, \omega)$. The time asymptotic form of the Green function gives the pulse shape in the laboratory frame. It is known [7] that the pulse shape in the laboratory frame can be deduced from the analysis of the pinch-point (k_0, ω_0) condition:

$$\epsilon(k_0, \omega_0 + k_0 V) = 0, \quad \frac{\partial}{\partial k} \epsilon(k_0, \omega_0 + k_0 V) = 0. \quad (2)$$

We consider the interaction of an electron beam penetrating a strongly correlated plasma. We model the present beam-plasma system by two plasmas in relative motion v_0 with different temperatures $1/\beta_b$ and $1/\beta$, and different densities n_b and n . Hence, these two plasmas have two different plasma parameters γ_b and γ . The dielectric function for the present beam-plasma system is given by

$$\epsilon(k, \omega) = 1 - \frac{\varphi(k)\chi_0(k, \omega)}{1 + \mathcal{G}(k, \omega)\varphi(k)\chi_0(k, \omega)} \\ - \frac{\varphi(k)\chi_{0b}(k, \omega - kv_0)}{1 + \mathcal{G}_b(k, \omega - kv_0)\varphi(k)\chi_{0b}(k, \omega - kv_0)}, \quad (3)$$

with $\chi_0(k, \omega)$ and $\mathcal{G}(k, \omega)$ denoting the noninteracting Vlasov density response function and the dynamical local field, respectively; $\varphi(k) = 4\pi e^2/k^2$ is the Coulomb potential. The dynamical local field describes the correlation

contribution to the dielectric function, as formulated in Refs. [1,2,5,8]. In the present paper, we will only consider a special case where the electron beam is weakly correlated $\gamma_b \ll 1$ and its temperature is much less than that of the strongly correlated plasma $\beta/\beta_b \ll 1$. Then the dielectric function reduces to

$$\epsilon(k, \omega) = 1 - \frac{b}{(\omega - kv_0)^2} - \frac{\varphi(k)\chi_0(k, \omega)}{1 + \mathcal{G}(k, \omega)\varphi(k)\chi_0(k, \omega)}. \quad (4)$$

Here $b = n_b/n$, and k , v , and ω are normalized to κ_D , the electron thermal velocity $v_T = \sqrt{1/m\beta}$, and the plasma frequency $\omega_p = \sqrt{4\pi ne^2/m}$, respectively. The zeros of this dielectric function are given by

$$\begin{aligned} (\omega - kv_0)^2 [k^2 - Z(\omega/k) + \mathcal{G}(k, \omega)Z(\omega/k)] \\ = b [k^2 + \mathcal{G}(k, \omega)Z(\omega/k)], \end{aligned} \quad (5)$$

with Z denoting the plasma dispersion function [9]. In the long-wavelength limit, i.e., $k/\kappa_D \ll 1$, the above equation becomes

$$\begin{aligned} (\omega - kv_0)^2 [\omega^2 - 1 - 2\alpha k^2 + 2i\eta\omega k^2] \\ = b [\omega^2 - (2\alpha - 3)k^2 + 2i\eta\omega k^2], \end{aligned} \quad (6)$$

where α and η essentially determine the plasmon group velocity and damping. α and η as a function of the plasma parameter γ have been calculated [4,5]; the results presented here (Fig. 1) are based on the calculation of Ref. [5]. The warm Vlasov plasma is obtained in the limit $\gamma \rightarrow 0$ where $\alpha = 1.5$ and $\eta = 0$. As γ increases, the plasmon group velocity decreases. Above the critical value $\gamma \gtrsim 50$, the plasmon group velocity becomes negative. The plasmon damping increases as γ increases.

The interaction between the beam waves and the plasma waves in the strongly coupled plasma, as shown in Fig. 2, is described by Eq. (6). As can be seen, in the strongly correlated plasma where $\gamma > \gamma_{\text{crit}}$, the plasmon group velocity can become opposite of that of the electron beam waves, thus giving rise to the possibility of an absolute instability. The absolute instability results from the phase resonant interaction between the forward beam waves and backward plasma waves. We now establish the conditions for the beam-plasma instability to become absolute.




FIG. 1. The plasmon group velocity α and damping η as a function of the plasma parameter γ .




FIG. 2. $\text{Re}\omega$ vs $\text{Re}k$, as given by Eq. (6). A and B denote $\text{Re}\omega$ for the weakly ($\gamma=0$) and strongly ($\gamma=87$) correlated plasmas; C gives the beam wave.

We begin with an analysis of the solutions to Eq. (6) for the real k . The results are presented in the complex ω plane in Fig. 3. The branch with $\omega_i > 0$ indicates that the beam-plasma system exhibits an instability. The maximum growth of this instability occurs when condition $|kv_0| \simeq 1$ is met. Note that Eq. (6) is an adequate representation to Eq. (5) only in the long-wavelength limit. Hence, in order for our above analysis and the following to be valid, the velocity of the electron beam should be much greater than the thermal velocity of the strongly correlated plasma. The electron number, on the other hand, should be much smaller than that of the strongly correlated plasma so as not to disturb the strongly correlated plasma too much. The beam parameters $b = 0.001$ and $v_0 = 10$ are chosen to meet above conditions.

We employ the pinch-point analysis [7] to determine the nature of the space-time evolution for the present beam-plasma system. The pinch-point analysis consists of mapping the complex ω plane into the complex k plane. Through this analysis, the solutions to Eq. (2) can be easily determined. It is obvious from Eqs. (2), (5), and (6) that the asymptotic form of the Green function is influenced by the plasmon group velocity α and damping η . The damping in the strongly correlated situation is almost exclusively collisional, moderated by the increased tendency of the particles to localize and to avoid their neighbors; Landau damping is negligible. We have performed the pinch-point analysis for the present beam-




FIG. 3. Numerical solutions to Eq. (6) in the ω plane for $0.0 < k < 0.2$ for $\gamma=87$.




FIG. 4. Convective-absolute boundary in the α - η plane, together with the results for α and η as a function of γ (dashed curve).

plasma system: it is found that lowering α moves the trailing edge of the pulse shape to the negative direction, while increasing η moves the trailing edge toward the opposite direction. Hence, under favorable conditions, the trailing edge of the pulse shape lies in the negative side of the origin, and the beam-plasma instability becomes absolute [7]. In Fig. 4, we present the condition for $v_0 = 10$ and $b = 0.001$. In the α - η plane, these conditions form a boundary line separating convective instability from absolute instability for the present beam-plasma system.

In the strongly correlated plasma, α and η are determined by the plasma parameter γ . Plotting α and η as a function of γ yields another curve in Fig. 4. We find that the present curve intersects with the boundary curve which separates convective from absolute instabilities at the point corresponding to $\gamma \sim 60$. Calculations based on other analyses of the plasmon dispersion [1,4] would give somewhat lower, but not substantially different results. Therefore, for $\gamma > 60$, the beam-plasma instability in strongly correlated plasmas is absolute for the present choice of the parameters.

As mentioned earlier, the pinch-point analysis not only distinguishes convective from absolute space-time evolution, but also yields the asymptotic form of the Green function $G(x', t)$ or the pulse shape. We present the pulse shape for the beam-plasma system in the weakly and strongly correlated plasmas in Fig. 5. As can be seen, the trailing edges of the pulse shape in the weakly correlated plasma such as $\gamma = 0$ and in the strongly correlated plasma such as $\gamma = 87$ lie on the opposite sides of the origin, revealing the nature of the space-time evolution for the corresponding beam-plasma system. The leading edges, however, are in the same place, despite the differences between these two different plasmas. As demonstrated in a previous study [11], the leading edges always reside in the same place as long as the beam plasma is cold. It has been shown that the thermal motion in the electron beam would slow down the leading edge. Hence, this would not interfere with the nature of the




FIG. 5. The pulse shape for the beam-plasma system. Solid and dashed curves give the pulse shape in the strongly ($\gamma = 87$) and weakly ($\gamma = 0$) correlated plasmas.

space-time evolution of the present beam-plasma system.

The distinguishing feature between an absolute and a convective instability is the bandwidth of the observed frequency spectrum. For an absolute instability it has been shown [10] that an absolute instability corresponds to a narrow-band emission centered around the pinch-point frequency in the laboratory frame. Meanwhile, a convective instability corresponds to a broadband emission. We propose the difference in the emission spectrum to be used as a means to probe and identify strongly correlated plasmas. Depending on the actual nature of the plasma to be experimentally studied, different ways of producing, injecting, and designing the electron beam can be envisioned. In particular, a degenerate electron gas in a solid would serve as an ideal target [6]: electron beams could be injected into such a plasma by strong electric fields. As emphasized in our analysis for the probing purposes, the electron number density in the electron beam should be much smaller than the density of the main strongly correlated plasma, so that the latter would not be disturbed by the former too much. We expect that the present paper will be a useful guide to the planning and interpretation of such probing experiments.

To summarize, we find that particle correlations profoundly affect the manner the beam-plasma system evolves with time. As the plasma parameter increases, the nature of the beam-plasma instability is changed from convective to absolute, at a critical value of the plasma parameter. This phenomenon may provide a useful experimental approach to probe and identify strongly correlated plasmas.

This work was partially supported by the National Science Foundation, Grants No. ECS-87-13337, No. PHY-91-15714, and No. ECS-88-22475, and the Department of Energy, Grant No. DE-FG02-91ER-54109. Z.C.T. acknowledges the support of his colleagues at UWO during the final stage of the work.

*Present address. Electronic address: tao@uwovax.uwo.ca

- [1] G. Kalman, in *Condensed Matter Theories 7*, edited by A. N. Proto and J. L. Aliaga (Plenum, New York, 1991).
- [2] Z. C. Tao and G. Kalman, *Phys. Rev. A* **43**, 973 (1990).

- [3] J. P. Hansen, E. L. Pollock, and I. R. McDonald, *Phys. Rev.* **11A**, 1025 (1975).

- [4] P. Carini, G. Kalman, and K. I. Golden, *Phys. Lett.* **78A**, 450 (1980); P. Carini and G. Kalman, *ibid.* **105A**, 229

- (1984); M. Minella, G. Kalman, K. I. Golden, and P. Carini (unpublished).
- [5] Z. C. Tao, Ph.D. thesis, Boston College, 1990 (unpublished).
- [6] A. VonFelde, J. Sprosser-Prou, and J. Fink, *Phys. Rev. B* **40**, 10 181 (1989).
- [7] A. Bers, in *Handbook of Plasma Physics*, edited by M. W. Rosenbluth and R. Z. Sagdeev, Basic Plasma Physics Vol. 1 (North-Holland, Amsterdam, 1983), Chap. 3.2
- [8] S. Ichimaru, *Rev. Mod. Phys.* **54**, 1017 (1982).
- [9] B. D. Fried and S. Conte, *The Plasma Dispersion Function* (Academic, New York, 1961).
- [10] A. K. Ram and A. Bers, in *Physics of Space Plasmas (1990)*, edited by T. Chang, G. B. Crew, and J. R. Jasperse, SPI Conference Proceedings and Reprint Series, No. 10 (Scientific, Cambridge, MA, 1991), pp. 351–365.
- [11] G. Francis, A. K. Ram, and A. Bers, *Phys. Fluids* **29**, 255 (1986).