

## Decoherence produces coherent states: An explicit proof for harmonic chains

Max Tegmark\*

*Max-Planck-Institut für Physik, Föhringer Ring 6, D-80805 München, Germany*

Harold S. Shapiro†

*Department of Mathematics, Royal Institute of Technology, S-11044 Stockholm, Sweden*

(Received 10 February 1994)

We study the behavior of infinite systems of coupled harmonic oscillators as the time  $t \rightarrow \infty$ , and generalize the central limit theorem (CLT) to show that their reduced Wigner distributions become Gaussian under quite general conditions. This shows that generalized coherent states tend to be produced naturally. A sufficient condition for this to happen is shown to be that the spectral function is analytic and nonlinear. For a chain of coupled oscillators, the nonlinearity requirement means that waves must be dispersive, so that localized wave packets become suppressed. Virtually all harmonic heat-bath models in the literature satisfy this constraint, and we have good reason to believe that coherent states and their generalizations are not merely a useful analytical tool, but that nature is indeed full of them. Standard proofs of the CLT rely heavily on the fact that probability densities are non-negative. Although the CLT is generally not applicable if the densities are allowed to take negative values, we show that a CLT does indeed hold for a special class of such functions. We find that, intriguingly, nature has arranged things so that all Wigner functions belong to this class.

PACS number(s): 05.30.Ch, 02.50.-r, 03.65.-w

### I. INTRODUCTION

The phenomenon of decoherence and the useful quantum states known as coherent states have been extensively studied quite separately, both being interesting in their own right, and the linguistic similarity of the names may be no more than a coincidence. Yet, it is becoming increasingly clear that the link between decoherence and coherent states is quite a close one—see [1] (Zurek, Habib, and Paz, 1993, hereafter ZHP) and references therein. ZHP give an excellent and up-to-date discussion of this link, and indicate that decoherence may indeed produce coherent states, since it is shown that the latter tend to be the most robust states when subjected to interactions with other systems. This link appears to have been first pointed out by Kübler and Zeh [2]. In this paper, we will in a sense complete this justification of the use of coherent states and their generalizations, by explicitly proving that they are created under quite generic circumstances.

#### A. Decoherence

Decoherence refers to some of the changes in a system that are due to its interaction with its environment. Such effects may include suppression of off-diagonal elements in the spatial density matrix (which makes the system appear more “classical”) and increase in entropy. Decoherence is now widely recognized as a key to the relationship

between the quantum and classical realms of physics (see Ref. [3] and references therein). Sources of decoherence discussed in the literature include scattering ([4–7] and others) and quantum gravity (for instance, Refs. [8,9]), but most of the literature has focused on systems with quadratic Hamiltonians, typically coupled harmonic oscillators in a chain or some other simple configuration. One reason for this is that systems with quadratic Hamiltonians are just about the only quantum systems whose time evolution can be found analytically. Hence they have provided useful and tractable models. This is why harmonic chains will be the model of choice in the present paper as well.

Before the interest in decoherence, the main motivation for studying harmonic chains was the pursuit of a dynamical basis for equilibrium statistical mechanics. An excellent summary of the early developments in this area is given in Ref. [10]. A recent summary of subsequent work is given in Ref. [11] (Tegmark and Yeh, 1994, hereafter TY), and Ref. [12] gives a more comprehensive review. In decoherence applications, the basic calculational procedure is identical to that in the statistical mechanics applications mentioned above: the idea is to study the time evolution of some small subset of the oscillators, called the *system*, by taking a partial trace over the rest of the oscillators, called the *heat bath* or the *environment*. In statistical mechanics applications, the goal is to investigate whether the system exhibits standard thermodynamic features such as Brownian motion and an approach to thermal equilibrium. In decoherence applications, the emphasis is on the behavior of the reduced density matrix of the system and on the extent to which certain quantum phase correlations are destroyed.

\*Electronic address: max@physics.berkeley.edu

†Electronic address: shapiro@math.kth.se

### B. Generalized coherent states

For historical reasons, states whose Wigner functions [13–15] are Gaussian have been given many different names. The single-oscillator ground state is a Gaussian centered on the origin. When translated in the  $q$  and  $p$  directions in phase space, it is usually called a coherent state. When rescaled so that it is shortened in the  $q$  direction and elongated in the  $p$  direction (or vice versa), it is known as a squeezed state. When subjected to the most general linear canonical transformation (translated, squeezed, and rotated), it is sometimes known as a tiltedly squeezed state. When expanded, it is called a thermal state, and is no longer pure. The translated ground state of a many-oscillator system is sometimes called a multimode coherent state, and so on. Thus the most general state with a Gaussian Wigner function might be termed a multimode tiltedly squeezed mixed state. We will simply refer to all these states as *generalized coherent states*, or *Gaussian states* for short.

As is indicated by the profusion of names for them, Gaussian states have been intensely studied in many areas of physics, from quantum optics to statistical mechanics. One reason for this is (just as with harmonic chains) analytic tractability: if a state is Gaussian at some given time, it will always remain Gaussian if the Hamiltonian of the system is quadratic, so it is sufficient to compute the time evolution of the mean and the covariance matrix, which specify the Gaussian uniquely. Another reason for their popularity is that coherent states, invented by Schrödinger in 1926 [16] and further developed by Glauber [17], have been seen as a clue to understanding the classical limit of quantum mechanics. This is because they, as opposed to, for instance, energy eigenstates, exhibit fairly “classical” behavior.

### C. The connection

Another Gaussian distribution, the Maxwell-Boltzmann velocity distribution, is well known to arise dynamically from the interactions of many independent particles, along the lines of the central limit theorem (CLT). Thus, in the spirit of ZHP, a natural question to ask is whether generalized coherent states also tend to be produced dynamically, from interactions within many-body systems. In this paper, we will address this question in a case where much of the necessary mathematical machinery is already in place: the case where the many-body system is a harmonic chain.

The paper is organized as follows: In Sec. II, we review some basic results about classical and quantum harmonic chains and establish some notation. In Sec. III, we prove the main result of the paper for the classical case. In Sec. IV, we show that the same result is true for the quantum-mechanical case as well. Finally, in Sec. V, we give a more heuristic and qualitative discussion of what happens for finite systems and for chains lacking translational invariance. Some necessary mathematical results are proven in the Appendices: In Appendix B we place a constraint on the dispersion relationship, and in Appendix C we prove a generalization of the central limit

theorem for the case where the “probability density” can take negative values.

## II. THE GENERAL HARMONIC CHAIN

In this section, we establish some notation and review some basic results about classical and quantum harmonic chains and cyclic matrices.

As our quantum system, let us take  $2N+1$  coupled harmonic oscillators of equal mass, labeled  $-N, \dots, -1, 0, 1, \dots, N$ . Denoting a point in the  $2(2N+1)$ -dimensional phase space by

$$\mathbf{z} = \begin{bmatrix} \mathbf{q} \\ \mathbf{p} \end{bmatrix} \quad (1)$$

and the corresponding operators by

$$\hat{\mathbf{z}} = \begin{bmatrix} \hat{\mathbf{q}} \\ \hat{\mathbf{p}} \end{bmatrix}, \quad (2)$$

we can write the Hamiltonian as

$$\hat{H} = \frac{1}{2m} \hat{\mathbf{p}}^T \hat{\mathbf{p}} + \frac{m\omega_0^2}{2} \hat{\mathbf{q}}^T A \hat{\mathbf{q}}, \quad (3)$$

where the time-independent matrix  $A$  is symmetric and positive definite. Throughout this paper, we will use units where  $m = \omega_0 = \hbar = 1$ . The number of oscillators can be either finite or infinite, but we will limit ourselves to the infinite case except in Sec. V.

At any given time, we will specify the (pure or mixed) state of the system by its Wigner function  $W(\mathbf{z})$ . It is well known that since the Hamiltonian is quadratic, the equation of motion for the Wigner function is identical to that of the Liouville function in classical statistical mechanics and has the solution

$$W_t(\mathbf{z}) = W_0([U(t)]^{-1}\mathbf{z}), \quad (4)$$

where the time-evolution matrix  $U$  is given by

$$U(t) = \begin{bmatrix} X & Y \\ Z & X \end{bmatrix} = \begin{bmatrix} \cos A^{1/2}t & A^{-1/2}\sin A^{1/2}t \\ -A^{1/2}\sin A^{1/2}t & \cos A^{1/2}t \end{bmatrix}. \quad (5)$$

Here and throughout this paper, the action of a function on a symmetric matrix is defined as the corresponding real-valued function acting on its eigenvalues: Since all symmetric matrices  $A$  can be diagonalized as

$$A = R \Lambda R^T,$$

where  $R$  is orthogonal and  $\Lambda = \text{diag}\{d_i\}$  is diagonal and real, we can extend any mapping  $f$  on the real line to symmetric matrices by defining

$$f(R \text{diag}\{d_i\} R^T) \equiv R \text{diag}\{f(d_i)\} R^T. \quad (6)$$

It is easy to see that this definition is consistent with power series expansions whenever the latter converge. For example,

$$\cos(A^{1/2}) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} A^n.$$

By a *Gaussian state* in  $n$  dimensions (we will often have  $n < 2N + 1$  further on, when dealing with reduced Wigner functions), we will mean a state whose Wigner function is Gaussian, i.e., is of the form

$$W(\mathbf{z}) = (2\pi)^{-n} (\det C)^{-1/2} \exp\left[-\frac{1}{2}(\mathbf{z} - \boldsymbol{\mu})^T C^{-1}(\mathbf{z} - \boldsymbol{\mu})\right]. \quad (7)$$

Here the mean vector  $\boldsymbol{\mu}$  and the covariance matrix  $C$  satisfy

$$\boldsymbol{\mu} = \langle \hat{\mathbf{z}} \rangle, \quad (8)$$

$$C = \begin{pmatrix} \langle \hat{\mathbf{q}} \hat{\mathbf{q}}^T \rangle & \frac{1}{2} \langle \hat{\mathbf{q}} \hat{\mathbf{p}}^T + \hat{\mathbf{p}} \hat{\mathbf{q}}^T \rangle \\ \frac{1}{2} \langle \hat{\mathbf{q}} \hat{\mathbf{p}}^T + \hat{\mathbf{p}} \hat{\mathbf{q}}^T \rangle & \langle \hat{\mathbf{p}} \hat{\mathbf{p}}^T \rangle \end{pmatrix} - \boldsymbol{\mu} \boldsymbol{\mu}^T.$$

(The symmetric ordering is necessary since  $\hat{\mathbf{q}}$  and  $\hat{\mathbf{p}}$  do not commute.) The Wigner function being Gaussian is equivalent to the density matrix being Gaussian in the position (or momentum) representation.

By a *time-independent state*, we will mean a state with a time-independent Wigner function (or, equivalently, with a time-independent density matrix). In TY it is shown that a necessary but not sufficient condition for a state to be time independent is that

$$\boldsymbol{\mu} = 0, \quad (9)$$

$$C = \begin{pmatrix} D & 0 \\ 0 & AD \end{pmatrix},$$

where  $D$  is some constant, symmetric, positive definite matrix that commutes with  $A$ . If the state is Gaussian, then this is evidently also a sufficient condition, since the Wigner function is completely specified by  $\boldsymbol{\mu}$  and  $C$ . We will assume that all states have  $\boldsymbol{\mu} = 0$ . This in no way reduces the generality of our treatment, as the time evolution of  $\boldsymbol{\mu}$  and the time evolution of the shape of the Wigner function (about its center  $\boldsymbol{\mu}$ ) are totally independent (see TY). Thus assuming  $\boldsymbol{\mu} = 0$  is much like assuming that the center of mass is at rest at the origin when studying the motion of a blob of jello in the absence of external forces.

As is conventional, we will assume that the harmonic chain is translationally invariant. This is equivalent to the potential matrix  $A$  being *cyclic* (such matrices are often called *circulant* in the mathematics literature [18]), i.e., that each row is a cyclic permutation of the row above it:  $A_{i+1, j+1} = A_{ij}$ , understood (mod  $n$ ) for an  $n \times n$  matrix. Since  $A$  is also symmetric, this means that we can write  $A_{ij} = a_{|i-j|}$  and interpret the system as a chain of harmonic oscillators where the coupling between any two oscillators depends only on the separation between them. (If  $N$  is finite, we can interpret the system as oscillators arranged in a ring rather than a line.) Using (6), we can write any function of a (cyclic or noncyclic) matrix  $A$  as

$$f(A)_{mn} = \sum_k R_{mk} R_{nk} f(d_k). \quad (10)$$

Cyclic matrices have the great advantage that they all

commute. This is because they can all be diagonalized by the same matrix  $R$ , the discrete Fourier matrix. Physically, this means that plane waves form a complete set of solutions. If  $A$  is symmetric, positive-definite, cyclic, and infinite-dimensional, then Eq. (10) reduces to [19]

$$f(A)_{mn} = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\theta f[\lambda^2(\theta)] \cos[(m-n)\theta], \quad (11)$$

where the spectral function  $\lambda^2(\theta)$  is the function whose Fourier coefficients are row zero of  $A$ . The spectral function can be interpreted as a dispersion relationship,  $\lambda$  being the frequency of a wave with wave number  $\theta$ . Note that  $f(A)$  is cyclic as well, i.e., its components depend only on the distance to the diagonal.

A cyclic potential frequently discussed in the literature is the *nearest neighbor potential*, the case where each mass is coupled only to a fixed spring and to its nearest neighbor:

$$\hat{H} = \sum_{k=-\infty}^{\infty} \left[ \frac{1}{2} \hat{p}_k^2 + \frac{1}{2} \hat{q}_k^2 + \frac{\gamma^2}{2} (\hat{q}_{i+1} - \hat{q}_i)^2 \right], \quad (12)$$

i.e.,  $A_{kk} = 1 + 2\gamma^2$ ,  $A_{k, k\pm 1} = -\gamma^2$  and all other elements of  $A$  vanish. For this special case, the spectral function is

$$\lambda^2(\theta) = 1 + 4\gamma^2 \sin^2 \frac{\theta}{2}. \quad (13)$$

### III. THE FINITE CLASSICAL CHAIN

In this section, we will investigate the circumstances under which states become Gaussian in classical statistical mechanics. Here the positions and momenta at time  $t$  are specified by  $\mathbf{z}(t)$ , which is a vector of random variables. These random variables are given by the initial random variables as

$$\mathbf{z}(t) = U(t)\mathbf{z}(0),$$

and we wish to study the circumstances under which the probability distribution of  $\mathbf{z}(t)$  becomes a multivariate Gaussian as  $t \rightarrow \infty$ .

According to Eq. (5), the position of oscillator  $m$  at time  $t$  is given by the initial data as

$$q_m(t) = \sum_{n=-\infty}^{\infty} \xi_{mn}(t), \quad (14)$$

where we have defined the random variables

$$\xi_{mn}(t) \equiv X_{mn} q_n(0) + Y_{mn} p_n(0). \quad (15)$$

(The above expression is to be understood without any summation.) Using the Liapunov version of the central limit theorem (e.g., Ref. [20]), we see that the distribution of  $q_m(t)$  becomes Gaussian as  $t \rightarrow \infty$  if the Liapunov condition

$$\frac{M^{(3)}}{M^{(2)}} \rightarrow 0 \quad \text{as } t \rightarrow \infty \quad (16)$$

is satisfied, where we have defined

$$M^{(k)}(t) = \left[ \sum_{n=-\infty}^{\infty} \langle |\xi_{mn}(t)|^k \rangle \right]^{1/k}. \quad (17)$$

If we make the physically reasonable assumption about the second and third moments of the initial data that  $\langle z_k(0)^2 \rangle^{1/2} > \sigma$  and  $\langle |z_k(0)|^3 \rangle^{1/3} < \kappa$  for some positive constants  $\sigma$  and  $\kappa$ , then the Liapunov condition reduces to the requirement that

$$\frac{\left[ \sum_{n=-\infty}^{\infty} |X_{mn}|^3 + |Y_{mn}|^3 \right]^{1/3}}{\left[ \sum_{n=-\infty}^{\infty} X_{mn}^2 + Y_{mn}^2 \right]^{1/2}} \rightarrow 0 \text{ as } t \rightarrow \infty. \quad (18)$$

(For the quantum case to be treated in the next section, the assumption of a minimum standard deviation follows directly from the Heisenberg uncertainty principle, if we simply assume that the standard deviations are bounded from above.)

If  $A$  is cyclic, then (5) and (11) yield

$$X_{mn} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \cos[\lambda(\theta)t] \cos[(m-n)\theta] d\theta, \quad (19)$$

$$Y_{mn} = \frac{1}{2\pi} \int_{-\pi}^{\pi} [\lambda(\theta)]^{-1} \sin[\lambda(\theta)t] \cos[(m-n)\theta] d\theta.$$

Now Parseval's theorem gives

$$\sum_{n=-\infty}^{\infty} X_{mn}^2 + Y_{mn}^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} (\cos^2[\lambda(\theta)t] + [\lambda(\theta)]^{-2} \sin^2[\lambda(\theta)t]) d\theta, \quad (20)$$

which approaches some positive constant  $c$  as  $t \rightarrow \infty$ . Thus, the Liapunov condition is satisfied if and only if the numerator of Eq. (18) approaches zero as  $t \rightarrow \infty$ . But

$$\sum_{n=-\infty}^{\infty} |X_{mn}|^3 + |Y_{mn}|^3 \leq \left[ \sup_n |X_{mn}| + |Y_{mn}| \right] \sum_{n=-\infty}^{\infty} X_{mn}^2 + Y_{mn}^2, \quad (21)$$

so it suffices to show that this supremum approaches zero. This is not merely a sufficient condition but also a necessary condition for  $q_m(t)$  to become Gaussian, since a nonzero supremum means that there is some  $\xi_{mn}$  that makes a finite contribution to the sum (14). This would imply that the distribution of  $q_m(t)$  depends on the details of the distribution of  $\xi_{mn}$  and thus cannot in general be Gaussian. In Appendix A, we show that this supremum does indeed approach zero under quite general conditions, namely, for any spectral function  $\lambda$  that is analytic on the entire interval  $[-\pi, \pi]$  and in addition is nonlinear.

In conclusion, we have shown that the probability distribution of  $q_m(t)$  becomes Gaussian as  $t \rightarrow \infty$  if the spectral function  $\lambda$  is nonlinear and analytic on  $[-\pi, \pi]$  and if the initial probability distributions of all positions and momenta are independent and have bounded second and third moments. The proof that  $p_m(t)$  becomes Gaussian is completely analogous.

The assumption that all random variables are independent can be relaxed to assuming that no dependence exists at oscillator separations larger than some fixed in-

teger  $M$ . More precisely [21,22], it is sufficient that there exists an  $M$  such that  $n-m > M$  implies that the two infinite sets

$$(\dots, q(0)_{m-1}, p(0)_{m-1}, q(0)_m, p(0)_m)$$

and

$$(q(0)_n, p(0)_n, q(0)_{n+1}, p(0)_{n+1}, \dots)$$

are independent. Finally, our proof can readily be generalized by using the multivariate CLT [23,24] to show that all finite multivariate distributions become Gaussian.

#### IV. THE INFINITE QUANTUM CHAIN

In this section we will see that all the results of the previous section can be generalized to the quantum-mechanical case. Much of the mathematics remains the same, but the interpretation changes. The big mathematical difference is that a Wigner function can take negative values, whereas a classical probability distribution cannot. A generalization of the central limit theorem for Wigner distributions is proved in Appendix C.

By analogy with reduced density matrices, all expectation values of the  $n$ th oscillator can be calculated from the  $n$ th single oscillator *reduced Wigner function* [14,15]

$$W^{(n)}(q_n, p_n) \equiv \int_{(n)} W(\mathbf{q}, \mathbf{p}), \quad (22)$$

where the integral is to be taken over all variables *except*  $x_n$  and  $p_n$ . This is analogous to the way the marginal probability distribution for  $(x_n, p_n)$  is calculated in classical statistical mechanics. The only difference is that the Wigner function can take negative values and cannot be interpreted as a probability distribution. In Sec. III, we gave necessary and sufficient conditions for when various marginal distributions become Gaussian as  $t \rightarrow \infty$ . Here we will pursue the quantum analog and give conditions for when various reduced Wigner functions become Gaussian.

Fourier transforming Eq. (22) with respect to all variables yields

$$\hat{W}^{(n)}(q_n, p_n) = \hat{W}(0, \dots, 0, q_n, 0, \dots, 0, p_n, 0, \dots, 0), \quad (23)$$

i.e., the Fourier transformed Wigner function (also known as the *characteristic function*) with all variables except  $q_n$  and  $p_n$  set equal to zero. This expression is often more useful than (22), as it contains no integrals. Fourier transforming Eq. (4) and using the fact that  $\det U = 1$  yields

$$\hat{W}_t(\mathbf{z}) = \hat{W}_0([U(t)]^T \mathbf{z}), \quad (24)$$

where  $U^T$  denotes the transpose of  $U$ .

Let us first assume that the oscillators are not entangled initially, so that the Wigner function for the initial state is completely separable, i.e., of the form

$$W_0(\mathbf{q}, \mathbf{p}) = \prod_n W_0^{(n)}(q_n, p_n) \quad (25)$$

for some set of reduced Wigner functions  $W_0^{(n)}$ . Substi-

tuting Eq. (25) into Eqs. (23) and (24) yields

$$\hat{W}_t^{(m)}(q_m, p_m) = \prod_n \hat{W}_0^{(n)}(X_{mn}q_m + Z_{mn}p_m, Y_{mn}p_m + X_{mn}q_m), \quad (26)$$

where the matrices  $X$ ,  $Y$ , and  $Z$  are those defined in Eq. (5), and no summation is implied. Thus the reduced Wigner function is obtained by Fourier transforming the initial reduced Wigner functions, multiplying them together, rescaling their arguments appropriately, and performing an inverse Fourier transform on the result. This is exactly how we would compute the probability density for a weighted sum of independent two-dimensional random variables, which is the classical case that we investigated in the previous section. The standard versions of the CLT all make heavy use of the assumption that probability densities are non-negative. Thus in order to show that the reduced Wigner function becomes Gaussian, we need a Liapunov type CLT for “random variables” whose “probability densities” are allowed negative values, a subject which to our knowledge has not been previously studied. We leave the full mathematical details of such a study for a future paper, but prove such a generalized CLT in Appendix C for the special case where all the “random variables” are identically distributed. It appears highly plausible that the standard Liapunov proof can be appropriately generalized employing similar techniques.

In conclusion, this would show that any one-particle reduced Wigner function  $\hat{W}_t^{(n)}(q_n, p_n)$  becomes Gaussian as  $t \rightarrow \infty$  if the spectral function  $\lambda$  is nonlinear and analytic on  $[-\pi, \pi]$  and if the initial states of all oscillators satisfy the condition that certain expectation values be bounded as described in Sec. III. Specifically, the expectation values of all linear, quadratic and cubic combinations of  $\hat{p}_n$  and  $\hat{q}_n$  should be bounded from above by some constant independent of  $n$ . Then as shown in theorem III in Appendix C, the other moment constraints will be automatically satisfied because of the Heisenberg uncertainty relationship.

Just as in the classical case, the assumption that no initial correlations exist between different oscillators can be relaxed to assuming that the joint Wigner functions are separable for oscillator separations larger than some fixed integer  $M$ . The generalization to the reduced Wigner function for more than one particle is also completely analogous.

Our result shows that virtually all harmonic chains treated in the literature will produce Gaussian states as  $t \rightarrow \infty$ , since they tend to have spectral functions that are both analytic and nonlinear. Some well-known examples of such harmonic chains are the above-mentioned nearest neighbor model [25,10] and the Ford, Kac, and Mazur (FKM) model [19]. Since the FKM model has been shown to be equivalent to the independent-oscillator heat bath model [12], the latter will also produce Gaussian states under quite general conditions.

An interesting mathematical problem is to generalize our results to arbitrary quadratic systems, by giving conditions for when they produce Gaussian states. It is our

belief that Gaussian states will be seen to be produced under quite generic circumstances, and thus are ubiquitous whenever there is interaction between a very large number of systems.

In TY, it is shown that if a harmonic chain starts out with an arbitrary cyclic covariance matrix

$$C = \begin{pmatrix} E & G \\ G & F \end{pmatrix},$$

then

$$C \rightarrow \begin{pmatrix} D & 0 \\ 0 & AD \end{pmatrix} \text{ as } t \rightarrow \infty, \quad (27)$$

where

$$D = \frac{1}{2}[E + A^{-1}F].$$

If the spectral function is nonlinear and analytic as discussed above, the convergence will not merely be pointwise as shown in the TY, but indeed uniform. Since a Gaussian is uniquely specified by its mean vector  $\mu$  and its covariance matrix  $C$ , we thus know not only that the harmonic chain approaches a Gaussian state, but also exactly which Gaussian state. As we would expect, the only information that is preserved about the initial data is the second moments, i.e., the covariance matrix, whereas all fine details of the Wigner function and all information about higher moments are lost. Note that the initial data enter only in the combination  $E + A^{-1}F$ , so all information about  $G$  (initial position-momentum correlations) is lost as well.

Without loss of generality, we assumed that the mean vector  $\mu = 0$  in the above treatment. The effect of relaxing this assumption is discussed in TY. It is seen that whereas the covariance matrix still converges to the value given above (and from what we have shown, all higher central moments converge to the values required by Gaussianity), the mean vector  $\mu$  does *not* converge towards a constant, but keeps oscillating forever.

## V. DISCUSSION

In this paper, we have shown that any part of a generic harmonic chain will evolve into a Gaussian state as  $t \rightarrow \infty$ . Given that the spectral function is mathematically well-behaved (analytic on the interval  $[-\pi, \pi]$ ), “generic” is to be interpreted as forbidding two special cases:

(1) The spectral function is linear.

(2) Fine-tuned long-range correlations exist in the initial data.

We will now attempt to give a more intuitive and physical interpretation of these two conditions (which apply for infinite chains), as well as qualitatively discuss what happens if  $N$  is large but finite.

The gist of the CLT as we have used it is that a weighted average of infinitely many independent random variables approaches Gaussianity as  $t \rightarrow \infty$  if all weights become infinitesimal. Very loosely speaking, a sum of infinitely many infinitesimally small independent random contributions is Gaussian. In terms of our harmonic

chains, information about the initial data must be mixed, and mixed so thoroughly that the state of any subsystem of the chain at  $t=0$  will have only an infinitesimal impact on the state of any subsystem of the chain as  $t \rightarrow \infty$ . Physically, what can go wrong? In the extreme case  $A \propto I$ , which corresponds to the oscillators being completely uncoupled, there is no mixing of information whatsoever and the CLT fails miserably. Now one might think that as long as an oscillator is coupled to at least one other oscillator (and thus indirectly to an infinite number of oscillators through it, by translational invariance), the CLT should always apply, and Gaussians should be obtained for any cyclic potential matrix *except*  $A \propto I$ . This is false. If the spectral function is linear (or, dropping the analyticity requirement, if it is linear on any finite interval), then a wave-packet composed only of wave numbers in this interval will simply travel down the chain without dispersing, retaining its initial shape forever. Thus the initial data at one point will have a non-infinitesimal impact on the state somewhere else, even at arbitrarily late times. This is reflected as  $U_{mn} \rightarrow 0$  as  $t \rightarrow \infty$  for any fixed  $m$  and  $n$  as shown in TY, while  $\sup_{m,n} |U_{mn}|$  remains bounded away from zero, as elements of order unity merely propagate further and further away from the diagonal, at a linear rate. In summary, the key is that the propagation of waves must be *dispersive*, i.e., the dispersion relationship must be non-linear. This will ensure that all localized wave packets gradually get destroyed.

The second constraint, that on the initial data, is closely related to the second law of thermodynamics: although for most initial data, the entropy of isolated gas in a container will not decrease, there is a small set of rather contrived initial data for which it will, and time will appear to run backwards for a while. The easiest way to obtain such initial data is to let a low-entropy state evolve into a high-entropy state and then reverse all velocities. The situation with our harmonic chains is completely analogous: If an uncorrelated state is allowed to evolve, the entropy of the subsystems will increase as each oscillator becomes increasingly correlated with ever more distance neighbors. If we now replace  $W(\mathbf{q}, \mathbf{p})$  by exactly  $W(\mathbf{q}, -\mathbf{p})$  (approximately will not suffice), the system will evolve back into the uncorrelated (and perhaps non-Gaussian) system we started with. Apparent time reversal is always caused by such long-range correlations, and since we used a version of the CLT that bans such correlations, such troubles are avoided altogether. Of course, after the uncorrelated initial state has been obtained, new correlations begin to arise again, and the subsystems eventually approach Gaussianity. An interesting problem is to investigate whether, in this vein, our result can be proven to hold for any cyclic initial conditions whatsoever.

The result that subsystems become Gaussian as  $t \rightarrow \infty$  holds strictly only for infinite chains. So what happens when  $N$  is finite but very large? If the waves are dispersive, then the discussion of finite  $N$  in TY can readily be extended to show that  $\max_{m,n} U_{mn}$  will evolve as follows when  $N$  is large.

(i) During an initial transition period whose duration is

of the order of the dynamical time scale  $\omega_0^{-1}$ , it decays from its initial value of order unity to a value of order  $N^{-1/2}$ .

(ii) After that, it oscillates around this value with an oscillation amplitude of the same order.

(iii) Since the time evolution of  $U_{mn}$  is almost periodic, some components must return to values of order unity an infinite number of times. This happens approximately once every Poincaré recurrence time. However, as shown by Ref. [26], the Poincaré time scale is generally enormous compared to the dynamical time scale, since it tends to grow exponentially with  $N$  for systems of this type.

In a discussion of density matrices [27], Feynman writes the following: "When we solve a quantum-mechanical problem, what we really do is divide the universe into two parts—the system in which we are interested and the rest of the universe. We then usually act as if the system in which we are interested comprised the entire universe." In this spirit we summarize our harmonic chain result: The effect of "the rest of the universe" is to make our subsystem approach a generalized coherent state. Since most systems in the real world are coupled to their environment, this gives us even more reason to believe that nature is indeed full of generalized coherent states.

#### ACKNOWLEDGMENTS

The authors would like to thank Emory Bunn, Leehwa Yeh, and Wojtek Zurek for useful comments on the manuscript.

#### APPENDIX A

In this appendix we give a condition for when  $\sup_{m,n} |U(t)_{mn}| \rightarrow 0$  as  $t \rightarrow \infty$ . This rests on theorem (I), which is proved in Appendix B. Since according to Eqs. (5) and (11),

$$\begin{aligned} X_{mn} &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \cos[\lambda(\theta)t] \cos[(m-n)\theta] d\theta, \\ Y_{mn} &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \sin[\lambda(\theta)t] \cos[(m-n)\theta] [\lambda(\theta)]^{-1} d\theta, \\ Z_{mn} &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \sin[\lambda(\theta)t] \cos[(m-n)\theta] \lambda(\theta) d\theta, \end{aligned}$$

we wish to show that

$$\sup_k \left| \int_{-\pi}^{\pi} e^{i\lambda(\theta)t} e^{-ik\theta} g(\theta) d\theta \right| \rightarrow 0 \text{ as } t \rightarrow \infty,$$

where  $k = \pm(m-n)$  is any integer,  $g(\theta) = [\lambda(\theta)]^\nu$ , and  $\nu = 0$ ,  $\nu = -1$ , and  $\nu = 1$ , respectively. Setting  $f(\theta) = \lambda(\theta)$ , theorem I shows that  $\sup_{m,n} |X(t)_{mn}| \rightarrow 0$  as  $t \rightarrow \infty$  if  $\lambda$  is a nonlinear analytic function on the entire interval  $[-\pi, \pi]$ . The same holds for  $Z(t)_{mn}$ , i.e., the  $\nu = 1$  case. Since  $A$  is positive definite,  $\lambda$  is bounded from below by some positive constant, so  $\lambda^{-1}$  is also analytic and  $\sup_{m,n} |Y(t)_{mn}| \rightarrow 0$  as  $t \rightarrow \infty$  follows under the same conditions. In summary,  $\sup_{m,n} |U(t)_{mn}| \rightarrow 0$  as  $t \rightarrow \infty$  for any bounded nonlinear analytic spectral function  $\lambda$ .

It is noteworthy that the nonlinearity requirement is

crucial to ensure that the convergence to zero is uniform, independent of  $m$  and  $n$ . By simply changing variables and using Riemann-Lebesgue's Lemma, it is readily seen that  $U(t)_{mn}$  will approach zero as  $t \rightarrow \infty$  for any fixed  $m$  and  $n$ , even if  $\lambda$  is linear. However, as was discussed in Sec. V, this alone is not sufficient for producing Gaussian states.

APPENDIX B

In this Appendix, which is purely mathematical, we prove the basic convergence theorem upon which the conclusion of the paper rests. The theorem is of course a version of "Van der Corput's Lemma" [28] in the theory of oscillatory integrals, but the uniformity with respect to  $k$  (which we believe is new) requires quite delicate handling.

*Theorem I:* Let  $f$  be a function analytic on a neighborhood of the closed bounded interval  $I$  of the real axis, and real valued on  $I$ . Assume  $f$  is not a polynomial of degree  $\leq 1$ . Then, for any  $g \in C^1(I)$ , we have

$$\sup_{k \in \mathbf{R}} \left| \int_I e^{if(x)t} e^{-ikx} g(x) dx \right| \rightarrow 0 \text{ as } t \rightarrow \pm \infty. \quad (B1)$$

Actually, as the proof will show, the left-hand side is  $O(|t|^{-\sigma})$  for some  $\sigma > 0$ .

In the proof, we may restrict attention to  $t > 0$ , as the other case then follows if we replace  $f$  by  $-f$ . By hypothesis, there is an open simply connected domain  $D$  containing  $I$  such that  $f$  is analytic on a neighborhood of the closure  $\bar{D}$  of  $D$ .

*Lemma (B1):* There is an integer  $l$  such that, for every  $w \in \mathbf{C}$ ,  $f'(z) - w = 0$  has at most  $l$  roots (counting multiplicities) in  $\bar{D}$ .

*Proof:* This is a simple exercise in complex analysis.

*Lemma (B2):* If  $F$  is of class  $C^2(J)$  and real valued for some closed bounded interval  $J \subset \mathbf{R}$ ,  $g \in C^1(J)$ , and  $F$  is strictly monotone on  $J$ , then for  $t > 0$  we have

$$\left| \int_J e^{iF(x)t} g(x) dx \right| \leq \frac{C_1}{\delta^2 t}, \quad (B2)$$

where  $\delta$  denotes the smaller of 1 and  $\min |F'(x)|$  for  $x \in J$ . Here  $C_1$  is a constant depending only on  $g$  and the number  $\max |F''(x)|$ ,  $x \in J$ .

*Proof:* This is a standard estimate of "Van der Corput type" (see, for instance, Ref. [28]). This is a rather primitive version, the proof being a straightforward variable change  $y = F(x)$  followed by partial integration. With stronger hypotheses one can get  $\delta$  rather than  $\delta^2$  in the denominator, but this is not required for our purposes.

In the following, a number of constants whose precise values are not essential will arise. Constants denoted  $C_1, C_2, \dots$ , will all be independent of  $k$  (later  $u$ ), depending only on the functions  $f$  and  $g$  and the geometric entities  $I, D$ .

*Proof of theorem:* We must estimate the integral

$$\int_I e^{if(x)t} e^{-ikx} g(x) dx = \int_I e^{iF(x)t} g(x) dx,$$

where  $F(x) = f(x) - ux$ , and  $u = k/t$  is a real parameter. By Lemma (B1), the number of complex zeroes  $z$  of  $f'(z) - u = 0$  in  $\bar{D}$  is bounded by an integer  $l$  independent of  $u$ . Denote the distinct zeroes by  $z_j = z_j(u)$ ;  $j = 1, \dots, s$ , with corresponding multiplicities  $m_1, \dots, m_s$  and  $\sum_{j=1}^s m_j \leq l$ . Now, fix  $\epsilon > 0$  and let  $\Delta_j$  denote an open disk of radius  $\epsilon$  centered at  $z_j$ . Then,  $I \setminus \cup_{j=1}^s \Delta_j$  consists of a union of  $r \leq l + 1$  pairwise disjoint closed intervals  $J_i$ , on each of which  $F$  is strictly monotone. Moreover we have the estimate

$$|F'(x)| \geq C_2 \epsilon^l \quad (B3)$$

for all  $x$  in these intervals. We will show this after completion of the argument. By (B2) we have for small  $\epsilon$

$$\left| \int_{J_i} e^{iF(x)t} g(x) dx \right| \leq \frac{C_3}{\epsilon^{2l} t}.$$

Summing over  $i$ , and noting that  $I \setminus \cup J_i$  has length  $\leq 2l\epsilon$ , we get

$$\left| \int_I e^{iF(x)t} g(x) dx \right| \leq C_4 [\epsilon + (\epsilon^{2l} t)^{-1}]. \quad (B4)$$

For fixed (large)  $t$ , choose here  $\epsilon = t^{-1/(2l+1)}$  and we see that the left-hand term in (B4) is bounded by  $C_5 t^{-1/(2l+1)}$ . This concludes the proof of the theorem.

We now supply the proof for the estimate (B3). Let us define the polynomial

$$P(z) = P(z, u) \equiv \prod_{j=1}^s [z - z_j(u)]^{m_j}.$$

It is clear that there is some constant  $C_6$  such that  $P(z, u) < C_6$  for all  $z \in \bar{D}$  and for all  $u$ . Now, consider the function  $P(z, u)/(f'(z) - u)$ . It is analytic in  $\bar{D}$ . Moreover, for some constant  $C_7$ ,

$$\max_{x \in I} \frac{|P(x, u)|}{|f'(x) - u|} \leq C_7. \quad (B5)$$

[We will return to the proof of (B5) shortly.] Thus, for  $x \in I$ ,

$$|f'(x) - u| \geq C_7^{-1} |P(x, u)| \geq C_2 \epsilon^l$$

for some constant  $C_2$  when  $x \in I \setminus (\cup \Delta_j)$ . Thus all that remains in order to prove (B3) is to show that (B5) holds. This can be done as follows. Let  $\Gamma_1, \Gamma_2, \dots, \Gamma_{l+1}$  be pairwise disjoint simple closed curves in  $D$ , each of which encloses  $I$ .

*Lemma (B3):* There is a positive constant  $C_8$  such that for any  $u$ ,

$$\min_{z \in \Gamma_j} |f'(z) - u| \geq C_8$$

holds for at least one value of  $j$ .

*Proof:* Let us define  $\varphi_j(u) \equiv \min |f'(z) - u|$ ,  $z \in \Gamma_j$ . It is

easy to see that  $\varphi_j$  is continuous. Hence, so is

$$\varphi(u) \equiv \max_{1 \leq j \leq l+1} \varphi_j(u).$$

Moreover,  $\varphi(u) > 0$ , because if  $\varphi(u) = 0$  for some  $u$ , then all  $\varphi_j(u)$  are zero, so  $f'(z) - u$  vanishes at least once on each  $\Gamma_j$  and thus has at least  $l+1$  zeroes, a contradiction. Since  $\varphi(u)$  is continuous, positive and obviously  $\rightarrow \infty$  as  $|u| \rightarrow \infty$ , it attains a positive minimum value  $C_8$ . We thus have that for every  $u$ , there is at least one  $j = j(u)$  such that  $\varphi_j(u) \leq C_8$ , which proves the Lemma.

By the maximum modulus theorem, for any  $u \in \mathbb{C}$ ,

$$\max_{z \in \Gamma} \frac{|P(z, u)|}{|f'(z) - u|} \leq \max_{z \in \Gamma_j} \frac{|P(z, u)|}{|f'(z) - u|},$$

where we choose  $j = j(u)$  as in Lemma (B3). Thus on the right-hand side, the numerator is bounded from above by  $C_6$  and the denominator is bounded from below by  $C_8$ , so the entire expression is  $\leq C_7 \equiv C_6/C_8$ . This completes the proof.

### APPENDIX C

In this Appendix, we prove a generalized version of the central limit theorem (CLT) that holds for Wigner distributions. Although the CLT *cannot* be generalized to arbitrary quasiprobability densities that are allowed to take negative values, we show that repeated convolutions do indeed lead to Gaussianity for a special class of such densities. We find that, for some reason, nature has arranged things so that all Wigner functions belong to this class.

Given a function  $f$  on  $\mathbb{R}^d$ , its zeroth, first, and second moments are defined as

$$M^{(0)} \equiv \int f(x) d^d x,$$

$$M_i^{(1)} \equiv \int f(x) x_i d^d x,$$

$$M_{ij}^{(2)} \equiv \int f(x) x_i x_j d^d x$$

if the moments exist, i.e., if these integrals are convergent in the Lebesgue sense. In probability theory, the second central moment matrix  $V_{ij} \equiv M_{ij}^{(2)} - M_i^{(1)} M_j^{(1)}$  is usually called the *covariance matrix*. Let us define a *quasiprobability density* on  $\mathbb{R}^d$  as a real-valued function  $f$  having the following properties:

- (i)  $M^{(0)} = 1$ .
- (ii) The first moments  $M_i^{(1)}$  exist.
- (iii) The second moments exist and the covariance matrix is strictly positive definite.
- (iv) For reasons that will become clear later, we will also make the technical assumption that  $f$  is an  $L^2$  function, i.e., square integrable.

If  $f$  has the additional property that it is non-negative, i.e., that it is a probability density, then the basic version of the CLT states that if we define  $f_n$  to be  $f$  convolved with itself  $n$  times and translated and rescaled so as to have the same first and second moments as  $f$ , then  $f_n$  approaches a Gaussian  $g$  as  $n \rightarrow \infty$ . The convergence is usually shown to be in the weak topology of measures, which in our context means that integral of  $f_n$  times any bounded test function tends to the corresponding integral

for  $g$ . We wish to investigate under which circumstances  $f_n$  approaches a Gaussian if we drop the assumption of non-negativity.

Without loss of generality, we may assume that  $M_i^{(1)} = 0$  and that  $V_{ij} = \delta_{ij}$ , the identity matrix, as the general case can be obtained from this by a simple change of variables. By Fourier transforming and using the convolution theorem, one then obtains the standard expression

$$\hat{f}_n(\mathbf{k}) = [\hat{f}(n^{-1/2} \mathbf{k})]^n. \quad (C1)$$

Our problem decomposes into two parts.

(A) To give conditions for when  $\hat{f}_n(\mathbf{k}) \rightarrow \hat{g}(\mathbf{k}) = e^{-k^2/2}$  as  $n \rightarrow \infty$ .

(B) To show that this convergence to Gaussianity on the Fourier side really implies that  $f_n \rightarrow g$  in some meaningful sense.

It is important to note that (B) is not merely an unphysical mathematical detail. This is illustrated by the following counterexample: Take  $d=1$  and chose  $\hat{f}(k)$  to be any smooth, symmetric  $L^2$  function such that  $\hat{f}(0) = 1$ ,  $\hat{f}'(0) = 0$ ,  $\hat{f}''(0) = -1$ , and  $\hat{f}(k_*) > 1$  for some constant  $k_* > 0$ . An example of such a function is  $\hat{f}(k) = (1+k^4)e^{-k^2/2}$ . It is easy to see that its inverse Fourier transform  $f$  will have all the properties of a quasiprobability density. It is also easy to show that  $f_n(k) \rightarrow g(k)$  as  $n \rightarrow \infty$  pointwise, for any fixed  $k$ , since  $\hat{f}(k) = 1 - k^2/2 + O(k^3)$  follows from our assumptions, and

$$\left[1 - \frac{k^2}{2n}\right]^n \rightarrow e^{-k^2/2} \quad \text{as } n \rightarrow \infty.$$

Yet Eq. (C1) clearly shows that the part of the curve that exceeds unity will grow ever larger as  $n$  increases. Pointwise convergence is obtained merely because the growing  $|\hat{f}_n| > 1$  hump keeps shifting out to higher and higher frequencies  $k$ . Thus as  $n$  grows large,  $\hat{f}_n$  may look quite Gaussian on the interval  $|k| \ll n^{1/2} k_*$ , but there will be exponentially growing bumps of height  $\hat{f}(k_*)^n$  at  $k = \pm n^{1/2} k_*$ . Inverse Fourier transforming, this means  $f_n$  will behave like a sum of a Gaussian and violent noise, whose frequency and amplitude increase without bounds as  $n \rightarrow \infty$ .

We will refer to a quasiprobability density as *proper* if the absolute value of its Fourier transform takes its maximum only at the origin. Thus  $f$  is proper if  $|\hat{f}(\mathbf{k})| \leq 1$ , with equality only for  $\mathbf{k} = 0$ . If a quasiprobability density never takes negative values (and hence is a probability density in the conventional sense), then it is easy to show that it will automatically be proper. The ‘‘ultraviolet catastrophe’’ described above shows that a necessary condition for a CLT to hold is that  $|\hat{f}|$  never exceeds unity. Thus being proper is a necessary condition, except perhaps for the borderline case where  $|\hat{f}(\mathbf{k})| \leq 1$  but actually equals unity for some  $\mathbf{k} \neq 0$ . In what follows, we will show that being proper is also a sufficient condition. We will also see that, interestingly, all Wigner quasiprobability densities are proper.

In what follows, the function  $g$  will always denote the  $d$ -dimensional Gaussian



$$g(\mathbf{x}) \equiv \frac{1}{(2\pi)^{d/2}} e^{-x^2/2}.$$

Unless otherwise indicated, all integrals below are to be taken over all space.  $\|\cdot\|_2$  will denote the  $L^2$  norm in  $\mathbb{R}^d$ , defined by

$$\|f\|_2 \equiv \left[ \int |f(\mathbf{x})|^2 d^d x \right]^{1/2}.$$

**Lemma (C1):** If  $f$  is a proper quasiprobability density, then for any  $\epsilon > 0$ , there exists a  $\delta > 0$  such that  $|\hat{f}(\mathbf{k})| \leq 1 - \delta$  for all  $|\mathbf{k}| > \epsilon$ .

*Proof:* If  $f$  is integrable, then  $\hat{f}(\mathbf{k}) \rightarrow 0$  as  $|\mathbf{k}| \rightarrow \infty$  by Riemann-Lebesgue's Lemma, so the continuous function  $|\hat{f}(\mathbf{k})|$  attains some maximum value  $M_\epsilon$  on the set  $\{k: |\mathbf{k}| \geq \epsilon\}$ .  $M_\epsilon < 1$  since  $f$  is proper, so we can choose  $\delta = 1 - M_\epsilon$ . Alternatively, if we do not wish to assume that  $f$  is integrable, it is straightforward to show that  $\hat{f}(\mathbf{k}) \rightarrow 0$  as  $|\mathbf{k}| \rightarrow \infty$  if  $f$  is any Wigner function.

**Lemma (C2):** The norms  $\|\hat{f}_n\|_2$  are bounded by a constant independent of  $n$ .

*Proof:* For small  $\mathbf{k}$ ,  $\hat{f}$  has the asymptotic behavior  $\hat{f}(\mathbf{k}) = 1 - k^2/2 + O(k^3)$ . Thus it is easy to see that given any constant  $p < 1$ , there exists an  $\epsilon_p$  such that

$$|\hat{f}(\mathbf{k})|^2 \leq e^{-pk^2}$$

for all  $|\mathbf{k}| \leq \epsilon_p$ . For all other  $\mathbf{k}$ , we have

$$|\hat{f}(\mathbf{k})|^2 \leq (1 - \delta_p)^2$$

for some  $\delta_p > 0$  by Lemma (C1). Combining these two bounds, we obtain

$$\begin{aligned} \|\hat{f}_n\|_2^2 &= \int |\hat{f}_n(\mathbf{k})|^2 d^d k \\ &\leq \int_{|\mathbf{k}| \leq \sqrt{n}\epsilon_p} e^{-pk^2} d^d k \\ &\quad + (1 - \delta_p)^{2(n-1)} \int_{|\mathbf{k}| > \sqrt{n}\epsilon_p} |\hat{f}(n^{-1/2}\mathbf{k})|^2 d^d k. \end{aligned}$$

Extending both integrals to all of space and changing variables in the second one, we get

$$\|\hat{f}_n\|_2^2 \leq \|e^{-pk^2/2}\|_2^2 + n^{d/2} (1 - \delta_p)^{2(n-1)} \|\hat{f}\|_2^2. \tag{C2}$$

Since the last term  $\rightarrow 0$  as  $n \rightarrow \infty$ , the left hand side is bounded by a constant independent of  $n$ .

**Lemma (C3):**  $\hat{f}_n(\mathbf{k}) \rightarrow \hat{g}(\mathbf{k})$  pointwise as  $n \rightarrow \infty$ .

*Proof:* This step is identical to that in proofs of the classical CLT (see, for instance, Ref. [20]), so we omit it.

**Lemma (C4):**  $\hat{f}_n \rightarrow \hat{g}$  in weak  $L^2$  topology as  $n \rightarrow \infty$ .

*Proof:* By a standard result in functional analysis [29], weak  $L^2$ -convergence (that  $\hat{f}_n - \hat{g}$  integrated against any  $L^2$  test function approaches zero) follows from the pointwise convergence [Lemma (C3)] and bounded norms [Lemma (C2)].

**Lemma (C5):**  $\|\hat{f}_n\|_2 \rightarrow \|\hat{g}\|_2$  as  $n \rightarrow \infty$ .

*Proof:* Letting  $p \uparrow 1$  in Eq. (C2) and invoking Fatou's Lemma,

$$\limsup_{n \rightarrow \infty} \|\hat{f}_n\|_2 \leq \|\hat{g}\|_2.$$

But since  $\hat{f}_n \rightarrow \hat{g}$  in weak  $L^2$  topology, we have

$$\|\hat{g}\|_2 \leq \liminf_{n \rightarrow \infty} \|\hat{f}_n\|_2.$$

The two preceding inequalities imply  $\limsup \|\hat{f}_n\|_2 \leq \liminf \|\hat{f}_n\|_2$ . Since the reverse is true always,  $\liminf = \limsup$ , which implies that  $\lim \|\hat{f}_n\|_2$  exists and equals  $\|\hat{g}\|_2$ .

**Theorem II:** If  $f$  is a proper quasiprobability density, then

$$\int |f_n(\mathbf{x}) - g(\mathbf{x})|^2 d^d x \rightarrow 0 \text{ as } n \rightarrow \infty,$$

i.e.,  $f_n$  approaches a Gaussian in  $L^2$  norm.

*Proof:* Because of the Plancherel Theorem ( $L^2$  unitarity of the Fourier transform), this is equivalent to

$$\|\hat{f}_n - \hat{g}\|_2 \rightarrow 0 \text{ as } n \rightarrow \infty,$$

i.e., that  $\hat{f}_n \rightarrow \hat{g}$  in strong  $L^2$  topology. But by a standard functional analysis result [29], this follows from weak  $L^2$  convergence [Lemma (C4)] combined with convergence of the norm [Lemma (C5)], so the proof is complete.

Thus we have shown that  $f_n$  approaches Gaussianity in the strong  $L^2$  sense. Note that in Lemma 1, we used the technical assumption that  $f$  was either integrable or a Wigner function. If we wish to make the additional technical assumption that not only is  $f$  (and hence  $\hat{f}$ ) in  $L^2$  but, for some  $\epsilon > 0$  (however small)  $|\hat{f}|^2 |\mathbf{k}|^\epsilon$  is also integrable over  $\mathbb{R}^d$ , then we can show the following:  $\hat{f}_n$  converges not merely in strong  $L^2$  but also in strong  $L^1$ , and consequently  $f_n$  converges uniformly to a Gaussian. Thus

$$\sup_{\mathbf{x}} |f_n(\mathbf{x}) - g(\mathbf{x})| \rightarrow 0 \text{ as } n \rightarrow \infty,$$

which rules out a number of physically uninteresting pathological cases, such as  $f_n(\mathbf{x})$  converging to the Gaussian  $g(\mathbf{x})$  for all  $\mathbf{x}$  except for a set of measure zero.

**Lemma (C6):** If  $W$  is a Wigner function, then  $\hat{W}(\mathbf{z}) \leq \hat{W}(\mathbf{0}) = 1$ , where the inequality is strict if  $\mathbf{z} \neq \mathbf{0}$ .

*Proof:* For a normalized wave function  $\psi$  in  $n$  dimensions (a pure state), the Fourier transform of the Wigner function is

$$\begin{aligned} \hat{W}(\mathbf{k}, \mathbf{x}) &= \int e^{-i(\mathbf{k}\cdot\mathbf{q} + \mathbf{x}\cdot\mathbf{p})} W(\mathbf{q}, \mathbf{p}) d^n q d^n p \\ &= \int e^{-i\mathbf{k}\cdot\mathbf{q}} [\psi(\mathbf{q} - \mathbf{x}/2)]^* \psi(\mathbf{q} + \mathbf{x}/2) d^n q, \end{aligned}$$

where the integral is to be taken over all space. Thus

$$|\hat{W}(\mathbf{k}, \mathbf{x})| \leq \int |\psi(\mathbf{q} - \mathbf{x}/2)| |\psi(\mathbf{q} + \mathbf{x}/2)| d^n q.$$

Using the trivial inequality  $AB \leq (A^2 + B^2)/2$  (with strict inequality unless  $A = B$ ), we obtain

$$\begin{aligned} |\hat{W}(\mathbf{k}, \mathbf{x})| &\leq \frac{1}{2} \int |\psi(\mathbf{q} - \mathbf{x}/2)|^2 d^n q + \frac{1}{2} \int |\psi(\mathbf{q} + \mathbf{x}/2)|^2 d^n q \\ &= \int |\psi(\mathbf{q})|^2 d^n q = 1. \end{aligned}$$

The case of a mixed state, where the Wigner function is a weighted average of Wigner functions of pure states, follows directly from superposition.

That we have strict inequality for  $\mathbf{z} \neq \mathbf{0}$  is seen as follows: The second inequality above is an equality only if  $\psi(\mathbf{q} - \mathbf{x}/2) = \psi(\mathbf{q} + \mathbf{x}/2)$  almost everywhere, i.e., if  $\psi$  has

period  $\mathbf{x}$ . But since  $\int \psi(\mathbf{q}) d^n q = 1$ ,  $\psi$  cannot be periodic, and the only possibility is  $\mathbf{x} = \mathbf{0}$ . Thus setting  $\mathbf{x} = \mathbf{0}$  in the first inequality and subtracting unity from both sides shows that we have equality only if

$$\int [1 - \cos(\mathbf{k} \cdot \mathbf{q})] |\psi(\mathbf{q})|^2 d^n q = 0.$$

Since the integrand is non-negative, it must vanish identically. Since  $\|\psi\|_2 = 1 > 0$ ,  $\psi$  cannot vanish almost everywhere, and the only possibility is  $\mathbf{k} = \mathbf{0}$ .

*Theorem III:* The Gaussian result in Theorem (II) is always obtained if  $f$  is a Wigner function with finite first and second moments.

*Proof:* Let  $f(\mathbf{z}) = W(\mathbf{z}) = W(\mathbf{p}, \mathbf{q})$  be a Wigner function in  $n$  dimensions, i.e., take  $d = 2n$ . We only need to check that all the conditions of Theorem (II) hold, i.e., that all such Wigner functions are indeed proper quasiprobability densities. All Wigner functions integrate to unity and are square integrable (indeed  $\|W\|_2 \leq (2\pi\hbar)^{-n/2}$ , with equality only for pure states [14]). Lemma (C6) showed that all Wigner functions are indeed proper. Thus, all that remains to be shown is that the covariance matrix  $V = C$  is strictly positive definite. This is a well-known fact, basically a corollary to the multidimensional uncertainty principle, but we give a brief proof here for completeness.

Without loss of generality, we may assume that the Wigner function corresponds to a pure state, since the covariance matrix of a mixed state is simply the weighted average of the covariance matrices in the mixture, and the weighted average of positive definite matrices is always positive definite. That  $V$  is positive semidefinite fol-

lows immediately from the fact that Wigner functions are proper:

$$|\hat{W}(\mathbf{z})|^2 = 1 - V_{jk} z_j z_k + O(|\mathbf{z}|^3),$$

where  $j$  and  $k$  are to be summed over from 1 to  $d$ , so if  $V$  would have a negative eigenvalue, then there must exist a point near the origin where  $|\hat{W}| > 1$ , a contradiction. The multidimensional uncertainty relationship [30,31] states that

$$\det V \geq (\hbar/2)^{2n}$$

for all Wigner functions, with equality only for Gaussian pure states, so none of the eigenvalues of  $V$  can vanish, and  $V$  must be positive definite. Thus all Wigner functions are proper quasiprobability densities, and the proof is complete.

Note that in contrast to the case of positive densities, second moments can vanish not merely in pathological cases, but also for well-behaved functions. Such an example is

$$\hat{W}(\mathbf{z}) = e^{-z^4},$$

for which  $V = 0$ . Also note that the requirement that the first and second moments be finite is necessary for the classical CLT as well. Finite first and second moments with respect to momentum is equivalent to the kinetic energy being finite. Finite first and second spatial  $\mathbf{q}$  moments can be interpreted as the system being spatially localized. Indeed, if the Hamiltonian is quadratic and positive definite (as it was in all cases treated in this paper), then all first and second moments must be finite if the total energy of the system is finite.

- 
- [1] W. H. Zurek, S. Habib, and J. P. Paz, *Phys. Rev. Lett.* **70**, 1187 (1993).  
 [2] O. Kübler and H. D. Zeh, *Ann. Phys. (Paris)* **76**, 405 (1973).  
 [3] W. H. Zurek, *Phys. Today* **44** (10), 36 (1991).  
 [4] E. Joos and H. D. Zeh, *Z. Phys. B* **59**, 223 (1985).  
 [5] M. R. Gallis and G. N. Fleming, *Phys. Rev. A* **42**, 38 (1989).  
 [6] M. Tegmark, *Found. Phys. Lett.* **6**, 571 (1993).  
 [7] L. Diósi, in *Proceedings of the Workshop in Trani on Waves and Particles in Light and Matter*, edited by A. Garuccio and A. vd der Merwe (Kluwer, City, 1993).  
 [8] S. W. Hawking, *Commun. Math. Phys.* **87**, 395 (1982).  
 [9] J. Ellis, S. Mohanty, and D. V. Nanopoulos, *Phys. Lett. B* **221**, 113 (1989).  
 [10] M. A. Huerta and H. S. Robertson, *J. Stat. Phys.* **1**, 393 (1969).  
 [11] M. Tegmark and L. Yeh, *Physica A* **202**, 342 (1994).  
 [12] G. W. Ford, J. T. Lewis, and R. F. O'Connell, *J. Stat. Phys.* **53**, 439 (1988).  
 [13] E. P. Wigner, *Phys. Rev.* **40**, 749 (1932).  
 [14] M. Hillery, R. H. O'Connell, M. O. Scully, and E. P. Wigner, *Phys. Rep.* **106**, 121 (1984).  
 [15] Y. S. Kim and M. E. Noz, *Phase Space Picture of Quantum Mechanics: Group Theoretical Approach* (World Scientific, Singapore, 1991).  
 [16] E. Schrödinger, *Naturwissenschaften* **14**, 664 (1926).  
 [17] R. J. Glauber, *Phys. Rev.* **131**, 2766 (1963).  
 [18] P. J. Davis, *Circulant Matrices* (Wiley, New York, 1979).  
 [19] G. W. Ford, M. Kac, and P. Mazur, *J. Math. Phys.* **6**, 504 (1965).  
 [20] A. Rényi, *Foundations of Probability* (Holden-Day, San Francisco, 1970).  
 [21] W. Hoeffding and H. Robbins, *Duke Math. J.* **15**, 773 (1948).  
 [22] Z. A. Lomnicki and S. K. Zaremba, *Math. Z.* **66**, 490 (1957).  
 [23] W. Feller, *An Introduction to Probability Theory and its Applications*, 3rd ed. (Wiley, New York, 1968).  
 [24] Y. V. Prohorov and Y. A. Rozanov, *Probability Theory* (Springer, Berlin, 1969).  
 [25] E. Schrödinger, *Ann. Phys. (Leipzig)* **44**, 916 (1914).  
 [26] P. Mazur and E. Montroll, *J. Math. Phys.* **1**, 70 (1960).  
 [27] R. P. Feynman, *Statistical Mechanics* (Benjamin, Reading, 1972).  
 [28] A. Zygmund, *Trigonometric Series* (Cambridge University Press, Cambridge, 1968), p. 198.  
 [29] E. Hewitt and K. Stromberg, *Real and Abstract Analysis* (Springer, Berlin, 1965).  
 [30] R. Simon, E. C. G. Sudarshan, and N. Mukunda, *Phys. Lett.* **124A**, 223 (1987).  
 [31] L. Yeh, Ph.D. thesis, U.C. Berkeley, 1993.