

BRIEF REPORTS

Brief Reports are accounts of completed research which do not warrant regular articles or the priority handling given to Rapid Communications; however, the same standards of scientific quality apply. (Addenda are included in Brief Reports.) A Brief Report may be no longer than four printed pages and must be accompanied by an abstract. The same publication schedule as for regular articles is followed, and page proofs are sent to authors.

Power law distributions of burst duration and interburst interval in the solar wind: Turbulence or dissipative self-organized criticality?

M. P. Freeman, N. W. Watkins, and D. J. Riley*

British Antarctic Survey, High Cross, Madingley Road, Cambridge, CB3 0ET, United Kingdom

(Received 25 February 2000; revised manuscript received 27 June 2000)

We calculate the probability density functions P of burst energy e , duration T , and interburst interval τ for a known turbulent system in nature. Bursts in the Earth-Sun component of the Poynting flux at 1 AU in the solar wind were measured using the MFI and SWE experiments on the NASA WIND spacecraft. We find $P(e)$ and $P(T)$ to be power laws, consistent with self-organized criticality (SOC). We find also a power-law form for $P(\tau)$ that distinguishes this turbulent cascade from the exponential $P(\tau)$ of ideal SOC, but not from some other SOC-like sandpile models. We discuss the implications for the relation between SOC and turbulence.

PACS number(s): 05.65.+b, 47.27.Sd, 47.65.+a, 96.50.Bh

In their seminal papers [1,2], Bak, Tang, and Wiesenfeld (BTW) demonstrated that a discrete cellular automaton model of an artificial sandpile had a spatial response to slow fuelling that was characterized by a scale-free distribution of energy release events or ‘‘avalanches’’ (see also Ref. [3]). Scale invariance was shown by a power-law probability density function (PDF) P of avalanche area A , $P(A) = CA^{-\alpha}$. This scale-invariant spatial structure led BTW [1] to propose the sandpile as a toy model of turbulence because, in Kolmogorov turbulence [4], long-wavelength, injection-range perturbations cause a scale-free forward cascade of energy transport until the dissipation scale is reached and therefore one might expect the PDFs of burst quantities in turbulent systems to be power laws too. These have recently been shown in burst area A for a generic inverse cascade model [5], in burst energy e and duration T for both a shell model [6], and reduced 2D MHD turbulence simulations [7], and in peak burst power for 1D MHD turbulence [8].

Boffetta *et al.* [6] (hereafter B99) have also shown that the PDF P of interburst intervals τ in a shell model of turbulence is a power law too but that this is not so for the BTW sandpile in which $P(\tau)$ is exponential. B99 postulated that the power law $P(\tau)$ found for solar flares [9] was consistent with a shell model of turbulence rather than the BTW sandpile. Here we demonstrate that the predicted avalanche phenomenology [power laws in $P(e)$, $P(T)$, and $P(\tau)$] of a shell model of turbulence is observed within a natural system—the solar wind—for which there is direct independent evidence of turbulence [10].

The solar wind is a near-radial supersonic plasma outflow from the solar corona which carries with it solar magnetic flux into interplanetary space by virtue of the plasma’s very

high electrical conductivity. In this ideal magnetohydrodynamic (MHD) approximation, the electric field \mathbf{E}' in the rest frame of the moving plasma is given by $\mathbf{E}' = \mathbf{E} + \mathbf{v} \times \mathbf{B} = \mathbf{0}$ from Ohm’s law. The electromagnetic energy (Poynting) flux $\mathbf{E} \times \mathbf{H}$ along the Sun-Earth line x can be approximated by $v(B_y^2 + B_z^2)/\mu_0$ assuming a radial solar wind. This quantity was calculated from ‘‘key parameter’’ measurements of \mathbf{B} and $|\mathbf{v}|$ from the MFI and SWE experiments, respectively, on the WIND spacecraft [11] between January 1995 and December 1998 inclusive. The typically 80–100 s averaged measurements of $|\mathbf{v}|$ were interpolated on to the 46 s time samples of \mathbf{B} .

In the resulting time series, bursts were identified, by the method used in Ref. [12], as intervals when the Poynting flux exceeded a given fixed threshold. Thresholds were set at the 10, 20, . . . 90 percentiles of the cumulative probability distribution of the Poynting flux. For each threshold, the PDF of the burst energy e , burst lifetime T , and interburst interval τ was calculated, where the burst energy is the sum of the Poynting flux samples over the burst lifetime T . The PDFs are shown in Fig. 1. The burst energy PDF (top panel) can be seen to have a power-law region over about 4 orders of magnitude between about 10^{-5} and 10^{-1} J m $^{-2}$. The burst lifetime PDF (second panel) also exhibits a power-law region and can be fitted by a power law with exponential cutoff similar to that found previously for the solar wind ε function [13]. In these respects, the solar wind Poynting flux has the avalanche phenomenology common to both the BTW sandpile and turbulence.

The interburst interval PDF has been plotted on both a log-log scale (third panel) and a log-linear scale (bottom panel). It is readily seen that this PDF is a power law rather than an exponential. A power law with an exponent of 1.67 is shown by the thick dashed curve in the third panel. This

*Present address: University of Cambridge, Cambridge, UK.




FIG. 1. Probability density functions of burst measures for the solar wind Poynting flux. From top to bottom, the measures are burst energy e , duration T , and interburst interval τ . The PDFs of all measures have power-law regions.

power-law form distinguishes the solar wind from a system having the properties of the BTW sandpile and instead shows it to be consistent with the shell model of turbulence used by B99. This is the same result they found for solar flares, for which there was not the direct independent evidence of turbulence that there is for the solar wind.

It is possible that the solar wind avalanche phenomenology is simply dominated by the advection of an already tur-

bulent fluid from the sun rather than by an energy cascade within the solar wind itself [Chapman (personal communication)]. We can expect the solar wind outflow from the Sun to be strongly influenced by energy dissipation events in the solar corona such as nanoflares [14] because these events can change the thermal pressure gradient that drives the solar wind [15] and/or allow reconfigurations of the solar magnetic field that aid or inhibit plasma outflow from the Sun. These observations are also topical in magnetospheric physics because we have previously shown [13] a similarity between the avalanche phenomenology present in geomagnetic perturbations [12,13] (which measure dissipative currents in the Earth's ionosphere) and that in the energy delivered by the solar wind to the Earth's magnetic and plasma environment. Independently, an analysis of the R/S Hurst exponents of solar wind variables and magnetospheric indices [16] has drawn similar conclusions.

So what does the observation of avalanche phenomenology in a natural system tell us about its physics? BTW postulated [1,2] that the appearance of “avalanche phenomenology” (power-law burst PDFs) in Nature was due to an underlying “critical” fixed point in the dynamics which was attractive (“self-organized”)—self-organized criticality (SOC). Renormalization group studies [17] have demonstrated that the Abelian BTW model indeed exhibits such an attractive fixed point. However, although BTW argued that SOC implied avalanche phenomenology, the converse is not true; and, in particular, the observation of avalanche phenomenology in natural systems [18] does not by itself prove that such systems are SOC.

There are many examples of systems that are either not self-organized or not critical, or both, that nevertheless present avalanche phenomenology. Avalanche phenomenology has been seen in the forest fire model [19] controlled by a repulsive rather than an attractive fixed point; it thus has to be tuned to exhibit scaling [20]. Some other models [18] exhibit power-law distributions without finite size scaling and so are not *bona fide* critical. Avalanche phenomenology can also be produced by coherent noise driving [21] or by “sweeping of an instability” [22]. In addition, the fixed-threshold method of estimating burst sizes that was used in Ref. [12] and the present work may generally result in scale-free PDFs if applied to certain types of time series. The action of slicing through a fractional Brownian motion (fBM) time series at a fixed level generates a set of crossing times known as an isoset, for which the PDF of the time interval between two subsequent crossings has a power law form [23]. Hence the burst duration and interburst interval statistics drawn from such an fBM time series by the fixed threshold method would also be expected to be power laws.

Clearly it is not sensible to apply the SOC label generally to systems exhibiting avalanche *phenomenology* [24]. Instead we should follow B99 in using a restricted definition of SOC, implicit in BTW's choice of name, as being the *mechanism* of self-organization to a critical state. From this point of view, in order to show the presence of SOC, one has to demonstrate those properties of self-organization and criticality that are unique to the process of SOC rather than simply observing the avalanche phenomena that SOC was designed to account for.

In consequence, the important question remains [7] as to the generality of B99's identification of an exponential $P(\tau)$ with the SOC mechanism. Exponential $P(\tau)$ implies that energy release episodes are uncorrelated in time because of the standard result that Poisson-distributed random numbers have an exponential distribution of waiting times. This will give rise to a $1/f^2$ power spectrum [18] for frequencies higher than those corresponding to the longest correlation time. In the BTW model, this is the time for the longest avalanche and is set by the system length. Jensen *et al.* [25] found that the BTW system had a $1/f^2$ high frequency power spectrum in energy flow down the sandpile, rather than the $1/f$ spectrum indicative of long-time correlation.

Whilst exponential $P(\tau)$ certainly holds for the BTW sandpile [26,6], this is not true for some other sandpile models. For example, let us consider the nearest-neighbor OFC model [27,18]. The conservative form of this model has been shown to be critical [28] and to evolve to a steady state [18]. In this case, $P(\tau)$ is found to be exponential [27]. However, there is also a nonconservative form of the nearest-neighbor OFC model [27,18] in which dissipation is introduced. This was recently shown to cease to be critical [28] and, in this dissipative case, $P(\tau)$ is found to differ from an exponential [27,29]. This supports the identification of exponential $P(\tau)$ with SOC.

Three classes of sandpile model, all of which modify aspects of BTW SOC, exhibit time correlation between bursts—variously reported as a nonexponential $P(\tau)$ in the dissipative OFC model [27,29] and as a “ $1/f$ ” power spectrum in both running [30] and continuous (e.g., Ref. [31])

sandpiles. However, it has yet to be shown that any of these systems are still SOC in the sense of both possessing an attractive fixed point and showing finite size scaling.

If B99 are correct in identifying time correlation of bursts as a diagnostic for the absence of SOC, then there should then be no instance of a model that has an attractive fixed point and finite size scaling (self-organized and critical) and which also has time correlated bursts of energy flow [specifically a $1/f$ spectrum or nonexponential $P(\tau)$]. That is, the time correlation in dissipative, running and continuous sandpiles is actually the signature of the breakdown of self-organized criticality. The apparent paradox of the observation of scale-free burst PDFs in such models is resolved when one recognizes that scaling may survive away from the fixed point, and can thus coexist with time correlation [28]. Scaling in both space and time can thus be a robust “generic” property of such “near-SOC” systems even if exact criticality is not. The test of Boffetta *et al.* can then test for the presence of SOC but cannot distinguish any of the modified sandpiles from turbulence models, and hence such “near-SOC” models remain possible descriptions of turbulence.

We are grateful to R. P. Lepping and K. W. Ogilvie for solar wind data from the NASA WIND spacecraft. We acknowledge valuable discussions with Sandra Chapman, Iain Coleman, Tim Horbury, Sean Oughton, Carmen do Prado, Dave Tetreault, and Tom Chang. We appreciate the provision of preprints by Giuseppe Consolini, Channon Price, Jouni Takalo, and Donald Turcotte. N.W.W. acknowledges the generous hospitality of the people at MIT.

-
- [1] P. Bak, C. Tang, and K. Wiesenfeld, *Phys. Rev. Lett.* **59**, 381 (1987).
- [2] P. Bak, C. Tang, and K. Wiesenfeld, *Phys. Rev. A* **38**, 364 (1988).
- [3] J.I. Katz, *J. Geophys. Res.* **91**, 10 412 (1986).
- [4] U. Frisch, *Turbulence: The Legacy of A. N. Kolmogorov* (Cambridge University Press, Cambridge, UK, 1995).
- [5] D.L. Turcotte, *Rep. Prog. Phys.* **62**, 1 (1999); A. Gabriellov, W.I. Newman, and D.L. Turcotte, *Phys. Rev. E* **60**, 5293 (1999).
- [6] G. Boffetta *et al.*, *Phys. Rev. Lett.* **83**, 4662 (1999); P. Giuliani *et al.*, *Physica A* **280**, 75 (2000).
- [7] G. Einaudi and M. Velli, *Phys. Plasmas* **6**, 4146 (1999).
- [8] S. Galtier and A. Pouquet, *Sol. Phys.* **179**, 141 (1998); S. Galtier, astro-ph/0003420 (unpublished).
- [9] G. Pearce, A.K. Rowe, and J. Yeung, *Astrophys. Space Sci.* **208**, 99 (1993).
- [10] W.H. Matthaeus and M.L. Goldstein, *J. Geophys. Res.* **87**, 6011 (1982); **87**, 10 347 (1982); D.A. Roberts, L.W. Klein, M.L. Goldstein, and W.H. Matthaeus, *ibid.* **92**, 11 021 (1987); D.A. Roberts, M.L. Goldstein, L.W. Klein, and W.H. Matthaeus, *ibid.* **92**, 12 023 (1987); For reviews see, e.g., A. Mangeney, R. Grappin, and M. Velli, in *Advances in Solar System Magnetohydrodynamics*, edited by E.R. Priest and A.W. Hood (Cambridge University Press, Cambridge, 1991); C.Y. Tu, and E. Marsch, *Space Sci. Rev.* **73**, 1 (1995).
- [11] M.H. Acuna *et al.*, *Space Sci. Rev.* **71**, 5 (1995). For MFI and SWE instrument descriptions see R.P. Lepping *et al.*, *ibid.* **71**, 207 (1995); K.W. Ogilvie *et al.*, *ibid.* **71**, 55 (1995).
- [12] J. Takalo, Ph.Lic. thesis, Department of Physics, University of Jyväskylä, Finland, 1993; J. Takalo *et al.*, *Geophys. Res. Lett.* **26**, 1813 (1999); *J. Atmos. Sol-Terr. Phys.* (to be published); G. Consolini, in *Cosmic Physics in the Year 2000*, proceedings of the 8th GIFCO Conference, Como, Italy, edited by S. Aiello *et al.* (Societa Italiana di Fisica, Bologna, 1997), p. 123; G. Consolini and P. de Michelis, *J. Atmos. Sol-Terr. Phys.* (to be published).
- [13] M.P. Freeman, N.W. Watkins, and D.J. Riley, *Geophys. Res. Lett.* **27**, 1087 (2000).
- [14] E.N. Parker, *Astrophys. J.* **330**, 474 (1988).
- [15] E.N. Parker, *Astrophys. J.* **128**, 664 (1958).
- [16] C.P. Price and D.E. Newman, *J. Atmos. Sol-Terr. Phys.* (to be published).
- [17] L. Pietronero, A. Vespignani, and S. Zapperi, *Phys. Rev. Lett.* **72**, 1690 (1994); A. Vespignani, S. Zapperi, and L. Pietronero, *Phys. Rev. E* **51**, 1711 (1995); A. Erzan, L. Pietronero, and A. Vespignani, *Rev. Mod. Phys.* **67**, 545 (1995).
- [18] H.J. Jensen, *Self-Organized Criticality: Emergent Complex Behaviour in Physical and Biological Systems* (Cambridge University Press, Cambridge, 1998).
- [19] V. Loreto, L. Pietronero, A. Vespignani, and S. Zapperi, *Phys. Rev. Lett.* **75**, 465 (1995); V. Loreto, A. Vespignani, and S. Zapperi, *J. Phys. A* **29**, 2981 (1996).
- [20] T.S. Chang, *Phys. Plasmas* **6**, 4137 (1999).
- [21] M.E.J. Newman and K. Sneppen, *Phys. Rev. E* **54**, 6226 (1996).

- [22] D. Sornette, *J. Phys. I* **4**, 209 (1994).
- [23] P.S. Addison, *Fractals and Chaos: An Illustrated Guide* (Institute of Physics, Bristol, 1997).
- [24] P. Bak, *How Nature Works: The Science of Self-Organised Criticality* (Oxford University Press, Oxford, 1997).
- [25] H.J. Jensen, K. Christensen, and H.C. Fogedby, *Phys. Rev. B* **40**, 7425 (1989).
- [26] M.S. Wheatland, P.A. Sturrock, and J.M. McTiernan, *Astrophys. J.* **509**, 448 (1998).
- [27] K. Christensen and Z. Olami, *Geophys. Res. Lett.* **97**, 8729 (1992).
- [28] J.X. de Carvalho and C.P.C. Prado, *Phys. Rev. Lett.* **84**, 4006 (2000).
- [29] Z. Olami and K. Christensen, *Phys. Rev. A* **46**, R1720 (1992); F. Angulo-Brown and A. Muñoz-Diosdado, *Geophys. J. Int.* **139**, 410 (1999).
- [30] T. Hwa and M. Kardar, *Phys. Rev. A* **45**, 7002 (1992).
- [31] H.F. Chau and K.S. Cheng, *Phys. Rev. A* **46**, R2981 (1992).