

Simulated dark-matter halos as a test of nonextensive statistical mechanics

Chloé Féron* and Jens Hjorth

Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark

(Received 14 June 2007; revised manuscript received 4 January 2008; published 29 February 2008)

In the framework of nonextensive statistical mechanics, the equilibrium structures of astrophysical self-gravitating systems are stellar polytropes, parametrized by the polytropic index n . By careful comparison to the structures of simulated dark-matter halos we find that the density profiles, as well as other fundamental properties, of stellar polytropes are inconsistent with simulations for any value of n . This result suggests the need to reconsider the applicability of nonextensive statistical mechanics (in its simplest form) to equilibrium self-gravitating systems.

DOI: 10.1103/PhysRevE.77.022106

PACS number(s): 05.40.-a, 04.40.-b, 05.90.+m, 95.35.+d

I. INTRODUCTION

Nonextensive statistical mechanics is a generalization of thermodynamics and statistical mechanics proposed by Tsallis in 1988 [1], aiming to overcome the limitations of Boltzmann entropy in its conventional applications [2–5]. The theory has considerably widened the fields of application of statistical physics, allowing a description of systems affected by nonlocal effects, such as long-range forces and memory effects.

The study of astrophysical self-gravitating systems was one of the first applications of the theory [6]. This approach has recently witnessed a renewed interest in the hope that it may provide a theoretical basis for the description of the universal structure of dark-matter (DM) halos [7–10]. Nonextensive statistical mechanics predicts their equilibrium states to be stellar polytropes (SPs) [6,11], which have been claimed to fit DM halos as well as the usual Navarro-Frenk-White model [9,10] (NFW model [12]). Moreover, SPs have the distinct advantage of being analytically derived from nonextensive statistical mechanics, while most models describing DM halos are empirical fits to N -body simulations. In this context, nonextensive statistical mechanics appears as an attractive framework for providing a theoretical understanding of the structure of DM halos and self-gravitating systems in general.

In this paper we compare, in a parameter-independent way, the equilibrium configuration of astrophysical self-gravitating systems predicted by nonextensive statistical mechanics to simulated DM halos. We clarify the issue of which central boundary conditions to use for comparing SPs to simulated halos. On this basis, we establish that simulated DM halos do not corroborate the predictions of nonextensive statistical mechanics. This result calls into question the direct applicability of nonextensive statistical mechanics to equilibrium astrophysical self-gravitating systems.

II. NONEXTENSIVE STATISTICAL MECHANICS AND STELLAR POLYTOPES

Nonextensive statistical mechanics is based on the formulation of a generalized entropy

$$S_q = k_B \frac{1 - \sum_{i=1}^W p_i^q}{q-1} \quad (q \in \mathbb{R}), \quad (1)$$

which reduces in the limit $q \rightarrow 1$ to the Boltzmann-Gibbs entropy $S_{BG} = -k_B \sum_{i=1}^W p_i \ln p_i$ (p_i is the probability of finding the system in the microstate i , W is the number of microstates i for a given macrostate, and k_B is the Boltzmann constant). When the index $q \neq 1$, the entropy of the system is nonextensive, i.e.; it is not possible to sum the entropies $S_q(A)$ and $S_q(B)$ of two subsystems A and B , but the generalized entropy S_q is conveniently pseudoadditive: $S_q(A+B) = S_q(A) + S_q(B) + (1-q)S_q(A)S_q(B)$. The third term accounts for coupling between the subsystems, due to nonlocal effects.

In this framework, astrophysical self-gravitating systems at equilibrium are described by SPs. This solution arises from extremizing the entropy S_q under the constraints of fixed total mass and energy as derived in [6,11] (see Ref. [13] about the choice of constraints), leading to a spherically symmetric isotropic system. The resulting distribution function depends on the parameter q and, via the identification $n = 1/2 + 1/(1-q)$,¹ has the same form as the SP distribution function $f(\epsilon) \propto \epsilon^{n-3/2}$ for $\epsilon > 0$ and $f(\epsilon) = 0$ for $\epsilon < 0$, where n is the polytropic index and ϵ the relative energy per unit mass [14].

In order to compare the predictions of nonextensive statistical mechanics to the results of N -body simulations, we compute the radial profiles of fundamental quantities of astrophysical self-gravitating systems: the matter density $\rho(r)$, the logarithmic density slope $d \log_{10}(\rho)/d \log_{10}(r)$, the integrated mass $M(r)$, and the circular velocity $V_c(r)$ (see Fig. 1). SPs are solutions of the Lane-Emden equation [14], depending on three free parameters: the polytropic index n , the central density ρ_0 , and the central velocity dispersion σ_0 . The value of n (and q) is an intrinsic property of each system, reflecting its degree of nonextensivity, but has not been determined from first principles and is therefore a free param-

¹An earlier version of the theory used $n = 3/2 + \frac{1}{q-1}$, where the relation between n and q depends on the choice of the constraints when extremizing the generalized entropy S_q [11,13]. A third relation $n = \frac{5-3q}{2(1-q)}$ has been proposed in [30].

*chloe@dark-cosmology.dk




FIG. 1. (Color online) From top to bottom are presented the density, the logarithmic density slope, the integrated mass, and the circular velocity. The profiles are scaled to r_{-2} , the radius at which the slope equals -2 . Stellar polytropes predicted by nonextensive statistical mechanics are represented by the solid curves with the shades corresponding to different polytropic index n . Simulated dark-matter halos are represented by the NFW model (dashed curve), the 3D Sérsic model (dotted curve, $\alpha=0.17$), and the Hernquist model (dash-dotted curve, $a=0.45$).

eter. SPs with $n > 3/2$ are stable due to Antonov's stability criterion—i.e., $df(\epsilon)/d\epsilon < 0$ [14]. Other values of n are unrealistic: $n = 3/2$ corresponds to a distribution function independent of ϵ and $n < 3/2$ to a distribution function diverging

at the escape energy $\epsilon=0$. Therefore, we compute numerical solutions of the Lane-Emden equation for $n > 3/2$, choosing values of n ranging from 2 to ∞ . We verified the consistency of our profiles using analytical solutions of the Lane-Emden equation, as well as the isothermal sphere, which is the asymptotic solution when $n \rightarrow \infty$.²

Nonextensive statistical mechanics does not specify the boundary conditions for SPs, which are needed to solve numerically the Lane-Emden equation. Simulated halos necessarily have a finite gravitational potential at the center; therefore, we look for SPs fulfilling this condition. For SPs the density ρ is linked to the gravitational potential ψ by $\rho = cst \times \psi^n$ [14], so we require solutions of the Lane-Emden equation with finite density at the center. Chandrasekhar proved that such solutions have $d \log_{10}(\rho)/d \log_{10}(r) = 0$ at $r=0$ [15]. Therefore, the class of SPs we can compare consistently to simulated halos has the natural boundary conditions at the center: $\rho = \rho_0$ and $d \log_{10}(\rho)/d \log_{10}(r) = 0$. We present the corresponding profiles in Fig. 1. In [9,16], the authors arbitrarily fix a nonzero logarithmic density slope near the center, chosen to match the NFW density profile [32]. This class of solutions must be treated with caution, as it does not ensure physical boundary conditions at the center.

III. COMPARISON TO SIMULATED DM HALOS

Cold dark-matter (CDM) halos resulting from N -body simulations provide an excellent way of testing the predictions of nonextensive statistical mechanics. They are collisionless systems whose particles interact via long-range gravitational interaction only, allowing a direct comparison to theory. DM structures in N -body simulations evolve from primordial fluctuations via hierarchical clustering to form halos of universal shape, well described by the NFW model [12]. These halos formed through accretion and mergers are sufficiently similar to those obtained by monolithic collapse—i.e., isolated halos—so that we do not take into account in this study the influence of formation history and cosmological environment on the final state of simulated halos [17].

Collisionless systems have very long collisional relaxation time scales to reach “true” equilibrium, but access a stable quasiequilibrium state faster (dynamical time scales are short compared to cosmological time scales—i.e., the age of the Universe) through the processes of violent relaxation and phase mixing [14]. The quasiequilibria predicted by numerical cosmological simulations and by nonextensive statistical mechanics, respectively, may in either case be considered the most probable state the system reaches.

Many empirical models have been proposed as fitting the universal profile of CDM halos. We use in our study three of them to represent simulated halos. The NFW model is the more commonly used [12]. The Sérsic model has been proposed recently as fitting more accurately the inner part of

²Analytical solutions exist only for $n=0,1,5$. We approximate $n \rightarrow \infty$ by $n=500$, as large values of n tend asymptotically towards the isothermal sphere.

high-resolution halos [18,19]. The Hernquist model provides a simple description of self-gravitating systems, for which all the profiles we use have analytical expressions [20]. As seen in Fig. 1, these profiles form a group with similar shapes and we do not need to use actual simulations in our study.

To compare simulated DM halos to SPs, we select the radial range for which recent N -body simulations are robustly resolved. Based on the high-resolution simulations published in Navarro *et al.* [18], this corresponds to $0.1r_{-2} < r < 10r_{-2}$, where r_{-2} is the radius at which the logarithmic density slope equals -2 . To compare SPs of all values of n with simulated halos, we need a scaling suitable for finite- and infinite-mass halos, circular velocity profiles with and without a maximum, and density profiles with very different steepness: r_{-2} provides such a universal scaling.³ Moreover, it allows us to take equally into account the inner and outer parts of the halo, respectively defined as having a slope shallower and steeper than -2 [12]. Finally, our study is independent of scaling parameters: SP profiles are scaled depending on the choice of ρ_0 and σ_0 , but the shape depends only on n . Therefore, scaling to r_{-2} makes identical any profiles of the same n but different ρ_0 and σ_0 , keeping only information about the shape. All the profiles we present, as scaled to r_{-2} , depend only on structural parameters: the range of n for SPs, $\alpha=0.17$ for the three-dimensional Sérsic profile [18], $a=0.45$ for the Hernquist profile [20], and r_s for the NFW profile.⁴ In Fig. 1 we compare SPs with simulated CDM halos, scaling to r_{-2} .

In the outer parts, where $r > r_{-2}$, SPs have very different shapes depending on the polytropic index n . Profiles with $n < 5$ correspond to finite-mass halos, while $n \geq 5$ polytropes have infinite mass, tending to the isothermal sphere when $n \rightarrow \infty$. While finite-mass polytropes may appear more attractive from a physical point of view, they also provide worse fits to simulated DM halos, with an outer slope as steep as $d \log_{10}(\rho) / d \log_{10}(r) < -5$ at $10r_{-2}$. For infinite-mass SP with larger values of the polytropic index, $n=16.5$ (as found in [9]) agrees with NFW and Sérsic models, $n=10$ with the Hernquist model. Therefore, *some infinite-mass SPs can provide a good description of the outer parts of simulated halos.*

In the inner parts, where $r < r_{-2}$, SPs share a similar property: they have a large core, the shape and the extent of which depend little on the value of n . This feature is in striking disagreement with the predictions of N -body simulations, which show steeper inner slopes. While this core structure itself has been advanced as an advantage of SPs over the NFW profile [10], solving the well-known cusp problem between simulations and observations [21], it appears in Fig. 1 that this is not the case: the difference between SPs and simulated halos in the density, density slope, and mass profiles is as high as one order of magnitude at $0.1r_{-2}$. Therefore, *for any polytropic index n , SPs do not properly describe the inner parts of simulated halos.*

³The scaling radius r_{-2} is determined uniquely in the case of a monotonically decreasing slope. However, SPs of $n \geq 5$ have a logarithmic density slope oscillating around -2 [14,31]; we therefore define r_{-2} as the smallest radius at which the slope is -2 .

⁴In the NFW model, the scale radius r_s is equal to r_{-2} .

IV. DISCUSSION

We have established here that the predictions of nonextensive statistical mechanics are not corroborated by simulations of DM halos. Our results are based on readily observable quantities—i.e., the matter density itself and profiles derived from it. These findings are in direct contrast to previous works which found reasonable agreement because they either considered only the outer parts of simulated halos [8], used a nonzero density slope as initial condition near the center [9], or used values of $n < 5$ that led to too steep density profiles in the outer parts [10]. Support for our conclusions, however, comes from Barnes *et al.* [16] who, even though fixing a nonzero density slope near the center to match the NFW density profile, found inconsistency between SP and DM halo velocity dispersion profiles. While we are cautious about the physical relevance of such profiles, it is of interest to emphasize that neither cored nor cuspy polytropes can describe properly simulated DM halos.

These results imply a fundamental difference in the matter distribution of astrophysical self-gravitating systems predicted by nonextensive statistical mechanics and by N -body simulations. Such a discrepancy raises three main questions. Is our comparison with simulated DM halos valid? Is nonextensive statistical mechanics the proper theory to describe collisionless long-range interaction systems at equilibrium? And how does this study relate to observed DM halos? We briefly address these issues below.

(1a) We study idealized systems which do not take into account complex effects in DM halos such as velocity anisotropy [22] or triaxiality [23]. However, the error we make by using isotropic and spherically averaged models to represent DM halos is small compared to the discrepancy we find between SPs and simulated halos. Moreover, we do not address here general problems of statistical mechanics of self-gravitating systems like, e.g., infinite mass [24], so as to focus on the issues specific to nonextensive statistical mechanics.

(1b) N -body simulations depend on various parameters, such as the choice of the softening length and the number of particles, which can introduce numerical effects in the resulting halos. The softening of the gravitational force on a small scale, introduced to avoid two-body interactions, creates a core at the center of the halo, approximately the size of the softening length. The number of particles fixes the maximum phase-space density resolved at redshift zero, and if not large enough, a core appears due to the lack of particles in the center. Both effects lead to a shallower density profile in the center of halos if the simulation is of low resolution [25]. Therefore, the discrepancy we observe between simulated profiles and SPs is genuine and increases with the resolution of numerical simulations.

(1c) We compare theoretical statistical predictions to cosmological simulations, but the latter depend on initial conditions and cosmological parameters. Initial conditions are fixed by the shape of the power spectrum of initial fluctuations $P(k) \propto k^{n_s}$ —i.e., by the choice of the index n_s . From CMB observations, we have an indication that $n_s = 0.958 \pm 0.016$ [26], but tests on cosmological simulations proved that they depend very little on the choice of n_s and on

the cosmological model used [27]. Therefore, the universal profiles of simulated halos can be compared to statistical theories as a general prediction.

(1d) The choice of a scaling is necessary to compare DM halos models, but can bring a visualization bias. We checked if the use of r_{-2} leads us to overestimate the disagreement between nonextensive statistical mechanics and N -body simulations. However, adjusting the fit in the outer parts of the halo—e.g., scaling to the virial radius—increases even more the discrepancy in the inner parts, while scaling profiles to fit well at smaller radii makes the discrepancy appear in the outer parts of the halo.

(2) Nonextensive statistical mechanics relies on three assumptions: (a) the generalized entropy S_q is the right form to describe long-range interaction systems, (b) the system is at equilibrium, and its most probable state is given by the maximum entropy principle, and (c) S_q is maximized at fixed energy, leading to a system whose distribution function depends only on the energy per unit mass, and has isotropic velocity dispersion. Though simulated halos show evidence of velocity anisotropy, the relation between density slope and velocity anisotropy [22] shows that it is isotropic in the center, where violent relaxation [28] (the process by which strong potential fluctuations efficiently drives the system towards equilibrium) is most effective. Therefore, assumptions (b) and (c) hold true in the center. However, it is in the inner parts that the disagreement between SP and simulated halos is the strongest. Hence, we suggest that assumption (a)—i.e.,

the choice of the generalized entropy S_q motivated by non-extensive statistical mechanics—cannot be used to predict the equilibrium structures of simulated DM halos.

(3) Simulated DM halos are in good agreement with observations [29], except for the inner core found from spiral galaxy rotation curves [21]. Stellar polytropes, interestingly, have an inner core too, but the discrepancy we observe between SPs and simulated DM halos is too large to explain the core of observed DM halos. Adding adiabatic contraction in simulations results in DM halos with steeper inner parts, which would not change our conclusions.

In summary, we have established that nonextensive statistical mechanics [1,6,11], a theory generalizing classical statistical mechanics and thermodynamics, does not describe the equilibrium state of astrophysical self-gravitating systems, as represented by DM halos formed in N -body simulations.

ACKNOWLEDGMENTS

It is a pleasure to thank S. H. Hansen, E. van Kampen, and E. I. Barnes for valuable discussions and comments, and to acknowledge the referees for helping us to improve this paper. The Dark Cosmology Centre is funded by the Danish National Research Foundation. This work was supported in part by the European Community's Sixth Framework Marie Curie research Training Network Programme, Contract No. MRTN-CT-2004-505183 "ANGLES."

-
- [1] C. Tsallis, *J. Stat. Phys.* **52**, 479 (1988).
 [2] I. V. Karlin, M. Grmela, and A. N. Gorban, *Phys. Rev. E* **65**, 036128 (2002).
 [3] A. Taruya and M. A. Sakagami, *Phys. Rev. Lett.* **90**, 181101 (2003).
 [4] P. H. Chavanis and C. Sire, *Physica A* **356**, 419 (2005).
 [5] D. H. E. Gross, *Physica E (Amsterdam)* **29**, 251 (2005).
 [6] A. R. Plastino and A. Plastino, *Phys. Lett. A* **174**, 384 (1993).
 [7] S. H. Hansen, D. Egli, L. Hollenstein, and C. Salzmann, *New Astron.* **10**, 379 (2005).
 [8] M. P. Leubner, *Astrophys. J. Lett.* **632**, 1 (2005).
 [9] T. Kronberger, M. P. Leubner, and E. van Kampen, *Astron. Astrophys.* **453**, 21 (2006).
 [10] J. Zavala, D. Núñez, R. A. Sussman, L. G. Cabral-Rosetti, and T. Matos, *J. Cosmol. Astropart. Phys.* **06** (2006), 008.
 [11] A. Taruya and M. Sakagami, *Physica A* **322**, 285 (2003).
 [12] J. F. Navarro, C. S. Frenk, and S. D. M. White, *Astrophys. J.* **462**, 563 (1996).
 [13] C. Tsallis, R. S. Mendes, and A. R. Plastino, *Physica A* **261**, 534 (1998).
 [14] J. Binney and S. Tremaine, *Galactic Dynamics* (Princeton University Press, Princeton, 1987).
 [15] S. Chandrasekhar, *An introduction to the study of stellar structure* (The University of Chicago Press, Chicago, 1939).
 [16] E. I. Barnes, L. L. R. Williams, A. Babul, and J. J. Dalcanton, *Astrophys. J.* **655**, 847 (2007).
 [17] A. Huss, B. Jain, and M. Steinmetz, *Astrophys. J.* **517**, 64 (1999).
 [18] J. F. Navarro, E. Hayashi, C. Power, A. R. Jenkins, C. S. Frenk, S. D. M. White, V. Springel, J. Stadel, and T. R. Quinn, *Mon. Not. R. Astron. Soc.* **349**, 1039 (2004).
 [19] A. W. Graham, D. Merrit, B. Moore, J. Diemand, and B. Terzić, *Astron. J.* **132**, 2701 (2006).
 [20] L. Hernquist, *Astrophys. J.* **356**, 359 (1990).
 [21] G. Gentile, P. Salucci, U. Klein, D. Vergani, and P. Kalberla, *Mon. Not. R. Astron. Soc.* **351**, 903 (2004).
 [22] S. H. Hansen and B. Moore, *New Astron.* **11**, 333 (2006).
 [23] E. Hayashi, J. F. Navarro, and V. Springel, *Mon. Not. R. Astron. Soc.* **377**, 50 (2007).
 [24] J. Hjorth and J. Madsen, *Mon. Not. R. Astron. Soc.* **253**, 703 (1991).
 [25] B. Moore, F. Governato, T. Quinn, J. Stadel, and G. Lake, *Astrophys. J. Lett.* **499**, 5 (1998).
 [26] D. N. Spergel, R. Bean, O. Doré, M. R. Nolta, C. L. Bennett, J. Dunkley, G. Hinshaw, N. Jarosik, E. Komatsu, L. Page *et al.*, *Astrophys. J., Suppl. Ser.* **170**, 377 (2007).
 [27] J. F. Navarro, E. C. S. Frenk, and S. D. M. White, *Astrophys. J.* **490**, 493 (1997).
 [28] D. Lynden-Bell, *Mon. Not. R. Astron. Soc.* **136**, 101 (1967).
 [29] E. Pointecouteau, M. Arnaud, and G. W. Pratt, *Astron. Astrophys.* **435**, 1 (2005).
 [30] J. A. S. Lima and R. E. de Souza, *Physica A* **350**, 303 (2005).
 [31] M. V. Medvedev and G. Rybicki, *Astrophys. J.* **555**, 863 (2001).
 [32] E. I. Barnes (private communication); E. van Kampen (private communication).