

**Anomalous negative differential thermal resistance in a momentum-conserving lattice**Wei-Rong Zhong,<sup>1</sup> Mao-Ping Zhang,<sup>1</sup> Bao-Quan Ai,<sup>2,\*</sup> and Bambi Hu<sup>3</sup><sup>1</sup>*Department of Physics and Siyuan Laboratory, College of Science and Engineering, Jinan University, Guangzhou 510632, China*<sup>2</sup>*Laboratory of Quantum Information Technology, ICMP and SPTE, South China Normal University, Guangzhou 510006, China*<sup>3</sup>*Department of Physics, University of Houston, Houston, Texas 77204-5005, USA*

(Received 16 May 2011; revised manuscript received 31 August 2011; published 28 September 2011)

A two-segment Fermi-Pasta-Ulam lattice has been investigated by using nonequilibrium molecular dynamics. Here we present an anomalous negative differential thermal resistance (NDTR) that has not been reported in Frenkel-Kontorova and  $\phi_4$  lattices up to the present. The NDTR disappears in the low-temperature region. The region of NDTR shifts from the large to the small temperature difference region as the system size increases. The anomalous dependence of NDTR on the temperature can be explained as the negative effect induced by the nonlinear coupling. The explanation can also cover the phenomenon of NDTR in momentum-nonconserved lattices.

DOI: [10.1103/PhysRevE.84.031130](https://doi.org/10.1103/PhysRevE.84.031130)

PACS number(s): 05.60.-k, 44.10.+i, 66.10.cd

**I. INTRODUCTION**

Negative differential thermal resistance (NDTR), which may imply an intrinsic physical mechanism for some thermal devices, has attracted a lot of attention in recent years. NDTR is first found in asymmetric Frenkel-Kontorova (FK) lattices [1], and then in pure anharmonic systems [2]. Analytical and numerical studies have been carried to reveal the original property of NDTR [1–7]. Up to now NDTR has been found in many low-dimensional systems, such as the anharmonic lattice with gradient mass [8], two-segment FK lattices with weak links [1,5], the pure  $\phi_4$  and FK lattice [2,6], the double-stranded systems [9], and so on. It has been reported that NDTR depends on the nonlinear external potential, the finite size, and the temperature of the system.

The main explanations about the phenomenon of NDTR include several points: (1) the mismatch of the phonon spectra of the two particles [1,8]; (2) the competition between the temperature difference, which acts as an external field, and the temperature-dependent thermal boundary conductance [2]; (3) the effective phonon-band shifts [7]; (4) a ballistic transport that induce the competition between the molecular occupation factor and the temperature difference [5]; and (5) the phonon-lattice scattering that becomes so significant that NDTR occurs [2,6]. All these explanations suggest a common view that the NDTR cannot occur when the system size increases, or in the system without nonlinear external potential, or in the system as the temperature is increasing. In other words, NDTR only appears at low temperature in a small system with nonlinear external potential. For the sake of comparison, here we call this kind of NDTR the normal NDTR. According to the normal NDTR, it is proposed that the momentum-conserved system without nonlinear external potential, such as a Fermi-Pasta-Ulam (FPU) lattice, behaves with no NDTR phenomenon. Since nanomaterials such as carbon nanotubes are a momentum-conserved system [10,11], this view about no NDTR in the momentum-conserved system once placed the studies of the NDTR in

carbon nanotubes in a difficult position. Actually, it is not the case.

In this paper, a two-segment FPU lattice with a weak link is studied by molecular dynamics simulations. We will present that the NDTR can also occur in a momentum-conserved system. It is surprising that NDTR in the FPU lattice shows an anomalous dependence on the temperature and the system size, which is different from the NDTR found in the system with nonlinear external potential. Here we introduce a competition between the linearity and the nonlinearity to interpret the origin of NDTR. Our explanation can cover some of the phenomena of NDTR that have been reported in FK models.

**II. MOMENTUM-CONSERVING AND MOMENTUM-NONCONSERVING LATTICES**

*Momentum-conserving lattice* [12]. A chain of  $N$  coupled atoms, in which only nearest-neighbor interactions will be considered for simplicity. The first class of models we wish to consider are defined by a Hamiltonian of the form ( $p_l = m_l \dot{x}_l$ )

$$\mathcal{H} = \sum_{l=1}^{N_M} \left[ \frac{p_l^2}{2m_l} + V(x_{l+1} - x_l) \right]. \quad (1)$$

Boundary conditions need also to be specified by defining  $x_0$  and  $x_{N+1}$ . Typical choices are periodic, fixed, or free boundaries. As only internal forces, which depend on relative positions, are present, the total momentum is conserved and thus a zero mode exists. The important examples are the well-known Lennard-Jones potential and Fermi-Pasta-Ulam (FPU) potential.

*Momentum-nonconserving lattice* [12]. At the simplest level of modelization, this can be described by adding an external, on-site potential to Eq. (1). For instance, neglecting the transverse motion leads to one-dimensional models of the form

$$\mathcal{H} = \sum_{l=1}^{N_M} \left[ \frac{p_l^2}{2m_l} + U(x_l) + V(x_{l+1} - x_l) \right]. \quad (2)$$

\* aibq@scnu.edu.cn

The substrate potential  $U(x_l)$  breaks the invariance  $x_l \rightarrow x_l + \text{const.}$  of Eq. (2) and the total momentum is no longer a constant of the motion. An important example is the well-known Frenkel-Kontorova (FK) potential.

### III. MODELS AND METHODS

The nonlinear lattices that we use in this paper consist of two segments, a left segment (L) and a right segment (R). Each segment is a FPU lattice. Segments L and R are coupled via a spring of constant  $k_{\text{int}}$ . The total Hamiltonian of the model is

$$H = H_L + H_R + H_{\text{int}}, \quad (3)$$

and the Hamiltonian of each segment can be written as

$$H_M = \sum_{i=1}^{N_M} \left[ \frac{p_{M,i}^2}{2m_M} + \frac{k_M}{2} (x_{M,i+1} - x_{M,i})^2 + \frac{\beta_M}{4} (x_{M,i+1} - x_{M,i})^4 \right], \quad (4)$$

where  $x_{M,i}$  and  $p_{M,i}$  denote the displacement from the equilibrium position and the conjugate momentum of the  $i$ th particle in segment  $M$ , where  $M$  stands for L or R. The parameters  $k_M$  and  $\beta_M$  are the harmonic and anharmonic spring constant of the FPU lattice, respectively. We couple the last particle of segments L and R via a harmonic spring. Thus,  $H_{\text{int}} = \frac{k_{\text{int}}}{2} (x_{L,N} - x_{R,1})^2$ . For the sake of simplicity, we set the mass of particle  $m = 1$  and the Boltzmann constant  $k_B = 1$ .

In our simulations we use fixed boundary conditions and the chain is connected to two heat baths at temperatures  $T_L$  and  $T_R$ . It is reported that the Nosé-Hoover thermostat may induce an inauthentic NDTR in the system without nonlinear external potential [6,13]. To avoid the adverse effect of the Nosé-Hoover thermostat we use the Langevin heat baths and integrate the equations of motion by using the Verlet frog-jumping algorithm [14,15]. The local temperature is defined as  $T_i = \langle p_i^2 \rangle$ . The local heat flux is defined as  $j_i = k_M \langle p_i(x_i - x_{i-1}) \rangle + \beta_M \langle p_i(x_i - x_{i-1})^3 \rangle$ , and the total heat flux is  $J = Nj$ . The simulations are performed long enough to allow the system to reach a steady state in which the local heat flux is constant along the chain. For the sake of comparison, we define a heat current ratio,  $J_R = J/J_{\text{max}}$ , in which  $J_{\text{max}}$  is the maximum heat current under a fixed temperature  $T_R$  of the right heat bath. The transport coefficient is an important quantity for characterizing the transport mode of a thermal transport process [16,17]. The thermal conductance evaluated as  $\sigma = Nj/\Delta T$  represents an effective transport coefficient that includes both boundary and bulk resistances [12]. Here we fix the temperature of the hot heat baths and change the temperature of the cold heat baths. The temperature difference increases with the decrease of the temperature of cold heat baths.

### IV. RESULTS AND DISCUSSION

Figure 1 shows the temperature dependence of NDTR in the two-segment FPU lattice. In the lattices with nonlinear external potential, which are also called momentum-nonconserved systems, many studies have shown that the NDTR disappears




FIG. 1. (Color online) Heat current as a function of the temperature difference for various fixed high temperatures  $T_L = 0.05, 0.1, 0.35, 0.60, 1.0$ . The linear and nonlinear couplings are respectively  $k = 1.0$  and  $\beta = 0.5$ , the weak-link coupling  $k_{\text{int}} = 0.01$ , and the system size is 64.

with the increasing temperature of the systems [2,5–7]. However, here it is presented that the NDTR occurs with the increasing temperature of the hot heat baths. Interestingly, the results for momentum-conserved systems are contrary to those for momentum-nonconserved systems.

This anomalous NDTR can be understood from the competition between linear and nonlinear interactions. First we would like to discuss the role of linear and nonlinear interaction on the thermal conductivity. It is well known that positive effects of the linear interaction on the thermal conductivity of the system have been confirmed by many studies, which include not only numerical but also analytical results [4,12,18]. This can be described as a function  $K \sim \lambda^\alpha$ , where  $K$  and  $\lambda$  are respectively thermal conductivity and the linear coupling, with the exponent  $\alpha > 0$ . Correspondingly, it is also shown that the nonlinear external potential has a negative influence on the thermal conductivity [4,12,18]. The relationship is written as  $K \sim V^\beta$ , in which  $V$  is the strength of the nonlinear coupling or the external potential, with the exponent  $\beta < 0$ . The NDTR depends exactly on the competition between these negative and the positive effects. When the negative effects surpass the positive ones, the NDTR occurs. Otherwise, the NDTR disappears.

For the FPU lattices, Fig. 2 shows that the heat current decreases with the nonlinear coupling constant, which implies a nonlinear inhibition on the thermal conductivity. In the absence of nonlinear coupling, i.e., in the case of a harmonic lattice, as shown in Fig. 2, when the nonlinear coupling  $\beta$  equals zero, the heat current increases linearly with the temperature difference and no NDTR occurs. When  $\beta$  increases to a value larger than 0.3, due to the enhancement of the nonlinear effect the NDTR appears at the region of the large temperature difference.

In order to give more details for this explanation, we plot Fig. 3 to show the dependence of the linear and nonlinear forces on the displacement. As mentioned above, in FPU lattices the NDTR disappears for low temperature and occurs for high temperature. For the system with low temperature,




FIG. 2. (Color online) Heat current as a function of temperature difference for various nonlinear couplings. The temperature  $T_L = 0.35$ , and the remaining parameters are the same as those for Fig. 1.

the particles oscillate near the equilibrium position and the displacement between two nearest-neighbor particles is small, i.e.,  $\Delta x_i \sim 0$ . As shown in Fig. 3(a), when  $\Delta x_i \sim 0$ , the force from linear coupling ( $-k\Delta x_i$ ) is larger than that from nonlinear coupling ( $-\beta\Delta x_i^3$ ). Therefore, the linear coupling determines the thermal transport and no NDTR occurs. As the temperature of the system increases,  $|\Delta x_i|$  goes to a value larger than  $|\Delta x_C|$  and the force from  $-\beta\Delta x_i^3$  is larger than  $-k\Delta x_i$ . The nonlinear coupling as the negative effect determines the thermal transport and then NDTR occurs. However, in the FK lattices an opposite temperature dependence of NDTR is observed. The Hamiltonian of the FK model is  $H_{FK} = \sum_{i=1}^N \left\{ \frac{p_i^2}{2m} + \frac{k_0}{2} (x_{M,i+1} - x_{M,i})^2 + \frac{V}{2\pi} [1 - \cos(\frac{2\pi}{a} x_i)] \right\}$ , where  $x_i$  and  $p_i$  denote the displacement from equilibrium position and the conjugate momentum of the  $i$ th particle. The parameters  $k_0$  and  $V$  are the harmonic spring constant and the strength of the external potential, respectively. Here we can also give




FIG. 3. (Color online) Force depends on the displacement of the particle. (a) Linear and nonlinear coupling force in FPU model. (b) Linear coupling and nonlinear on-site force in FK model.




FIG. 4. (Color online) Heat current as a function of temperature difference for different weak-link couplings  $k_{int}$ . Here the remaining parameters are  $k = 1.0$ ,  $\beta = 0.5$ ,  $T_L = 0.35$ , and  $N = 64$ .

an explanation from the relationship between the force and the displacement. As shown in Fig. 3(b), for the system with low temperature, the force from linear coupling ( $-k_0\Delta x_i$ ,  $k_0 = 0.8$ ) is smaller than that from nonlinear coupling ( $-V \sin x_i$ ,  $V = 0.6$ ) and the nonlinear coupling determines the thermal transport and then NDTR occurs. On the contrary, for the FK lattices with high temperature, the superior linear coupling induces the disappearance of the NDTR. If one sets  $k_0 = 3.0$ , the linear force is always larger than the nonlinear force, then the NDTR disappears, which has been confirmed in Refs. [2,6].

Figure 4 shows how the coupling constant  $k_{int}$  affects the temperature dependence of the heat flux  $J/J_{max}$ . The region of NDTR diminishes for increasing  $k_{int}$  until it vanishes at a critical coupling constant. The weak link  $k_{int}$  is a linear coupling, as also depicted in Fig. 3(a); when  $k_{int}$  is small, the nonlinear effect can easily get above the linear one at  $\Delta x_i \sim 0$  in the interface of the two FPU lattices and then NDTR occurs. However, the NDTR vanishes for strong linear link  $k_{int}$ .

In addition, the system-size dependence of NDTR in FPU lattices is also different from that found in FK and  $\phi_4$  lattices. In FK and  $\phi_4$  lattices [2,6], with the increasing of the system size, the NDTR region shifts from the small to the large temperature difference region and vanishes in the end. On the contrary, as shown in Fig. 5, in FPU lattices the NDTR disappears for the small system. When the system size increases, the region of NDTR (the regime with a frame) shifts from the large to the small temperature difference region.

## V. COMPARISONS

Here we would like to compare our explanations about NDTR with those that have been presented. As mentioned above, the NDTR have been explained as the phonon-lattice scattering, the crossover of the transport mode, the competition between the temperature difference and the boundary conductance, the mismatch of phonon spectra, and so on. As has been reported, the mismatch of phonon spectra and the effective phonon-band shifts can give a good interpretation




FIG. 5. (Color online) NDTR depends on the system size. Here the remaining parameters are  $k = 1.0$ ,  $\beta = 0.5$ ,  $T_L = 0.35$ , and  $k_{\text{int}} = 0.05$ .

for the NDTR in two-segment lattices, but fail to interpret the NDTR in pure single lattices [2,6,7]. Moreover, the competition between the temperature difference and the temperature-dependent thermal boundary conductance can only explain the small systems with strong boundary effect [2,6], but will be invalid for large systems mentioned here. Finally, the phonon-lattice scattering and the crossover of the transport mode approach the main reason of NDTR at low temperature [2,6], but not an appropriate origin of NDTR at high temperature. Actually, all these explanations are based upon a negative effect, which maybe imply a fundamental physical mechanism of NDTR. And here this negative effect comes from the nonlinearity that includes both the nonlinear coupling and the nonlinear external potential.

A comparison of NDTR in the two cases is also presented from three aspects:

(1) *The temperature dependence of NDTR.* As shown in Fig. 2 (two-segment FK model) of Ref. [5], a temperature dependence of NDTR is reported. When the temperature of hot heat baths  $T_R$  is small, there exists NDTR. As  $T_R$  increases, the NDTR disappears. However, here Fig. 1 presents that the NDTR occurs with increasing temperature of the hot heat baths.

(2) *The system-size dependence of NDTR.* In the previous works (FK and  $\Phi_4$  model), as shown in Fig. 6 (deformable FK model) of Ref. [6], Figs. 3 (FK model) and 6 ( $\Phi_4$  model) of Ref. [2], and Fig. 4 (two-segment FK model) of Ref. [7], NDTR becomes weak and disappears with the increasing of the system size. The most important property is that the NDTR region shifts from the small to the large temperature difference region and vanishes in the end. In our current works, on the

contrary, as shown in Fig. 5, in two-segment FPU lattices the NDTR disappears for the small system. When the system size increases, the region of NDTR (the regime with a frame) shifts from the large to the small temperature difference region.

(3) *The nonlinear potential.* As shown in Eqs. (1) and (2), the Hamiltonian of the momentum-conserving lattice is much different from that of the momentum-nonconserving lattice. Many reports have shown that the momentum-conserving lattice has an anomalous thermal conductivity [19]. Then it is natural to regard the NDTR of the momentum-conserving lattice as an “anomalous” one. Additionally, the NDTR in this paper shows very different or even opposite behavior from the NDTR that has been reported in the previous papers; thus we have to call it “anomalous NDTR.”

## VI. CONCLUSIONS AND APPLICATIONS

In summary, we have reported an anomalous negative differential thermal resistance in two-segment FPU chains through molecular dynamics simulations. The NDTR occurs in the system when the system size as well as the temperature increases. This anomalous NDTR effect can be understood from the negative effect induced by the nonlinear force, which can give a more fundamental explanation about NDTR. Since the FPU lattice is a momentum-conserved system, our results provide a new view about the NDTR in the system without external potential, which is regarded as a necessary condition for NDTR up to the present. Furthermore, our results have also suggested that the NDTR can be achieved in two-segment nanoscale materials with a weak linear link, which may be exciting information for fabricating a nanoscale thermal device.

The experimental motivation of the model refers to the carbon nanotubes and their extended models. The Lennard-Jones potential or Brenner-Tersoff potential, which describes the momentum-conserving lattice, is usually used to study carbon nanotubes [10,11]. An advanced structure of the nanotubes is more similar to our model. For example, one can connect two carbon nanotubes with silicon nanotubes. Here silicon nanotubes are weak connections and carbon nanotubes are momentum-conserving lattices.

## ACKNOWLEDGMENTS

We would like to thank the Siyuan clusters for running part of our programs. This work was supported in part by the National Natural Science Foundation of China (Grant Nos. 11004082 and 11175067), the National Natural Science Foundation of Guangdong Province, China (Grant Nos. 10451063201005249 and S2011010003323), and the Fundamental Research Funds for the Central Universities, JNU (Grant No. 21609305).

[1] B. Li *et al.*, *Appl. Phys. Lett.* **88**, 143501 (2006); L. Wang and B. Li, *Phys. Rev. Lett.* **99**, 177208 (2007).  
 [2] D. He, S. Buyukdagli, and B. Hu, *Phys. Rev. B* **80**, 104302 (2009); D. He, B. Q. Ai, H. K. Chan, and B. Hu, *Phys. Rev. E* **81**, 041131 (2010).

[3] D. Segal and A. Nitzan, *Phys. Rev. Lett.* **94**, 034301 (2005); L.-A. Wu and D. Segal, *ibid.* **102**, 095503 (2009); D. Segal, *Phys. Rev. E* **79**, 012103 (2009).  
 [4] E. Pereira, *Phys. Rev. E* **82**, 040101(R) (2010); E. Pereira and H. C. F. Lemos, *ibid.* **78**, 031108 (2008).

- [5] W. R. Zhong, P. Yang, B. Q. Ai, Z. G. Shao, and B. Hu, *Phys. Rev. E* **79**, 050103(R) (2009).
- [6] B. Q. Ai and B. Hu, *Phys. Rev. E* **83**, 011131 (2011); B. Q. Ai, W. R. Zhong, and B. Hu, *ibid.* **83**, 052102 (2011).
- [7] Z. G. Shao, L. Yang, H. K. Chan, and B. Hu, *Phys. Rev. E* **79**, 061119 (2009).
- [8] N. Yang, N. Li, L. Wang, and B. Li, *Phys. Rev. B* **76**, 020301(R) (2007).
- [9] W. R. Zhong, *Phys. Rev. E* **81**, 061131 (2010).
- [10] G. Wu and B. Li, *Phys. Rev. B* **76**, 085424 (2007).
- [11] H. Rafii-Tabar, *Computational Physics of Carbon Nanotubes* (Cambridge University Press, New York, 2008).
- [12] S. Lepri, R. Livi, and A. Politi, *Phys. Rep.* **377**, 1 (2003).
- [13] S. Nose, *J. Chem. Phys.* **81**, 511 (1984); W. G. Hoover, *Phys. Rev. A* **31**, 1695 (1985).
- [14] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes* (Cambridge University Press, Cambridge, 1992).
- [15] D. C. Rapaport, *The Art of Molecular Dynamics Simulation* (Cambridge University Press, New York, 2004).
- [16] D. Li, Y. Wu, P. Kim, L. Shi, P. Yang, and A. Majumdera, *Appl. Phys. Lett.* **83**, 2934 (2003).
- [17] J. S. Wang, *Phys. Rev. Lett.* **99**, 160601 (2007); Y. Xu, J. S. Wang, W. Duan, B. L. Gu, and B. Li, *Phys. Rev. B* **78**, 224303 (2008).
- [18] F. Bonetto, J. L. Lebowitz, J. Lukkarinen, and S. Olla, *J. Stat. Phys.* **134**, 1097 (2009); F. Bonetto, J. L. Lebowitz, and J. Lukkarinen, *ibid.* **116**, 783 (2004); D. Roy, *Phys. Rev. E* **77**, 062102 (2008).
- [19] O. Narayan and S. Ramaswamy, *Phys. Rev. Lett.* **89**, 200601 (2002); B. Li and J. Wang, *ibid.* **91**, 044301 (2003).