

Correlations and critical behavior of the q model

Alexander St. John¹ and Harsh Mathur²

¹*Department of Physics, University of Wisconsin Parkside, 900 Wood Road, Kenosha, Wisconsin 53144, USA*

²*Department of Physics, Case Western Reserve University, 10900 Euclid Avenue, Cleveland, Ohio 44106, USA*

(Received 14 August 2011; revised manuscript received 16 October 2011; published 9 November 2011)

The q model is a random walk model used to describe the flow of stress in a stationary granular medium. Here we derive the exact horizontal and vertical correlation functions for the q model in two dimensions. We show that close to a critical point identified in earlier work these correlation functions have a universal scaling form reminiscent of thermodynamic critical phenomena. We determine the form of the universal scaling function and the associated critical exponents ν and z .

DOI: [10.1103/PhysRevE.84.051303](https://doi.org/10.1103/PhysRevE.84.051303)

PACS number(s): 81.05.Rm, 02.50.Ey, 73.43.-f

I. INTRODUCTION

The q model was introduced by Coppersmith *et al.* [1] to describe the flow of stress in a static granular medium motivated by the experiments of Ref. [2]. Since then there have been significant advances in experiments and theory (see, for example, [3] and references therein). Although the q model has been superseded, it remains of some interest as a simple tractable model of a granular medium. Moreover, the q model is closely related to models that describe the process of aggregation in statistical mechanics [4], the transport of electrons on the surface of an integer quantum Hall multilayer [5], passive scalar turbulence [6], and the branching of river networks [7], among others. In earlier work by Lewandowska *et al.* [8] it was shown that although the q model describes physics far from equilibrium, nonetheless, for a particular value of its parameters, the q model behaves in a manner reminiscent of the critical point in an equilibrium thermodynamic phase transition. It was found that the q model is remarkably soluble and a number of exact critical exponents and universal scaling functions near to the critical point were obtained. The purpose of the present work is to strengthen the analogy to thermodynamic critical phenomena by calculating the two-point correlation functions for the q model.

In experiments that the q model is used to describe, a pack of beads is loaded from above with a uniform stress. The distribution of load at the bottom of the bead pack as well as the propagation of stress through the pack have been measured experimentally [2]. In the q model it is assumed that the beads lie at the sites of a regular lattice. The beads in each layer are supported by their nearest neighbors in the layer below. For simplicity let us consider the q model in two spatial dimensions. Let us suppose that the beads are arranged in a square lattice as shown in the Fig. 1. For each bead a fraction f of its load is supported by its neighbor to the left in the layer below and a fraction $1 - f$ by its neighbor to the right. These fractions are assumed to vary randomly from bead to bead. We denote the depth of the layer by t and the position of a bead within a layer by n . The load on bead n in layer $t + 1$ is then determined by the recursion relation

$$w_n(t + 1) = f_{n+1,t} w_{n+1}(t) + (1 - f_{n-1,t}) w_{n-1,t} + I_n(t + 1). \quad (1)$$

Given the random fractions f one can use Eq. (1) to propagate an applied uniform load in the top layer downward. The goal is to obtain a statistical description of the load as a function of depth by analytic or numerical solution of Eq. (1). In the original q model an extra term corresponding to the weight of the beads was included. Here we have neglected that term, as is appropriate for the experiments of Ref. [2].

In the q model it is generally assumed that the fractions are independent identically distributed random variables with a distribution $P(f)$ that satisfies the symmetry requirement $P(f) = P(1 - f)$. The simplest choice is to assume that each fraction is either zero or one with equal probability; this is called the singular distribution, and it was shown in Ref. [8] that the behavior of the q model for this distribution is reminiscent of a critical point in equilibrium thermodynamics. Other distributions can be characterized by a parameter δ (defined precisely below) that measures how far the distribution is from the critical point.

In this paper we calculate the horizontal and vertical two-point correlations of the load on the beads by generalizing the methods of Ref. [8]. The horizontal load correlation

$$c(m, t) = \langle w_n(t) w_{n+2m}(t) \rangle \quad (2)$$

is the covariance of the load on two beads that are in the same layer at depth t and have a horizontal separation $2m$. The brackets $\langle \dots \rangle$ denote an average over the random fractions. By translational invariance the covariance depends only on the separation of the two beads ($2m$) and not on the absolute location of either bead (n and $n + 2m$, respectively). By analogy to thermodynamic critical phenomena we would expect the horizontal correlation length to diverge as $\xi_{\text{horizontal}} \propto 1/\delta^\nu$ and the vertical correlation length to diverge as $\xi_{\text{vertical}} \propto 1/\delta^{\nu z}$ close to the critical point. Furthermore, we would expect the horizontal correlation function to have the universal scaling form

$$c_h(m, t) - 1 = \frac{1}{\delta^\alpha} \mathcal{G}(m\delta^\nu, t\delta^{\nu z}). \quad (3)$$

We find that in two dimensions the correlation function does indeed have a universal scaling form for large depth t and small δ that we determine. Moreover, we find the correlation length exponent $\nu = 1$, the dynamical exponent $z = 2$, and the amplitude exponent $\alpha = 0$.




FIG. 1. Upper panel: The q model in two dimensions (reproduced from Ref. [8]). The beads are assumed to be arranged in a square lattice. Each bead is assumed to be supported by its nearest neighbors in the layer below. A random fraction f of the load is transferred to the neighbor at left, the remainder $(1 - f)$ to the neighbor to the right. The fractions vary from bead to bead and are assumed to be independent identically distributed random variables. t specifies the horizontal layer occupied by a bead. The horizontal layers are numbered by consecutive integers. The position of a bead within a horizontal layer is given by n . n is even in a layer with even t and odd, for odd t . Thus the horizontal separation of two beads in the same layer must be an even integer. Similarly the horizontal separation for two beads in layers with an even vertical separation must also be an even integer. Lower panel: The distribution of load in layer $t = 100$ computed numerically for a single realization of the q model with the random fractions f drawn from the singular distribution. A uniform load is applied to the top layer $t = 1$. It is assumed that there are 100 beads in each layer and periodic boundary conditions have been applied. The gray bar shows the horizontal correlation length appropriate to the singular q model at this depth derived in Sec. II B.

The vertical two-point correlation

$$c_v(m, t, \tau) = \langle w_n(t) w_{n+2m}(t + 2\tau) \rangle \quad (4)$$

is the covariance of the load on two beads that are located in horizontal layers separated by a depth of 2τ and that have a horizontal separation of $2m$. Since the q model is only translationally invariant in the horizontal direction, the vertical two-point correlation depends on both the separation of the two layers (2τ) as well as the absolute location of the first layer (t). Again, by analogy to thermodynamic critical phenomena

we would expect the vertical correlation to have the universal scaling form

$$c_v(m, t, \tau) - 1 = \frac{1}{\delta^\beta} \mathcal{H}(m\delta^\nu, t\delta^{\nu z}, \tau\delta^{\nu z}). \quad (5)$$

We find that in two dimensions the correlation function does indeed have a universal scaling form for large depth t and small δ that we determine. Moreover, we find the correlation length exponent $\nu = 1$ and the dynamical exponent $z = 2$, in agreement with the values obtained from the horizontal correlation calculation and the amplitude exponent $\beta = 1$.

The relationship of the q model to random walk models can be understood as follows. Suppose that instead of a uniform load, a unit load is applied to a small number of beads in the top layer. In the singular case the subsequent propagation of the load follows the trajectories of random walkers that coalesce upon contact and move together thereafter. The nonsingular case can be interpreted as walkers that have a certain probability to fission. By contrast, in common random walk models the walkers either pass through each other in the noninteracting case or bounce off each other or annihilate each other upon contact in interacting models [9]. It is seen below that the horizontal correlation functions of the q model decay as Gaussians rather than exponentials, reflecting their close kinship to random walks.

In the prior work of Lewandowska *et al.* [8] the critical point was characterized by computing the covariance of the load on a single bead. Reference [8] derived the universal scaling function that describes the evolution of this quantity and computed a number of critical exponents including the product $\nu z = 2$ (but not the separate values of ν and z). The present work adds to this body of knowledge the universal scaling forms for the two-point correlations and the important critical exponents ν and z . The present work is limited to two dimensions where the critical behavior is most interesting. Reference [8] also studied higher dimensions and thereby discovered that the upper critical dimension for the q model is three. Moreover, Ref. [8] considered the effect of an “injection term” that takes into account the weight of the beads, which has been neglected in comparison to the applied load in Eq. (1). In other related work Rajesh and Majumdar [10] computed the two-point correlations for the q model but at large depth and far from the critical regime. Finally, Snoeijer and van Leeuwen [11,12] have studied the distribution of load at asymptotically large depths, as discussed further in the Conclusion.

II. HORIZONTAL CORRELATIONS

A. Exact solution

Our purpose in this subsection is to evaluate the evolution of the correlations Eq. (2) with depth by use of Eq. (1). We begin by noting that the random fractions that appear in Eq. (1) are assumed to be independent identically distributed random variables with a symmetric distribution $P(f)$ that satisfies the requirement $P(f) = P(1 - f)$. The distribution of fractions may be characterized by a parameter ϵ defined via

$$\left\langle \left(f - \frac{1}{2} \right)^2 \right\rangle = \frac{\epsilon}{4}. \quad (6)$$

It is easy to verify that $\epsilon = 1$ for the singular distribution and $\epsilon = \frac{1}{3}$ for the uniform distribution. Thus δ defined as

$$\delta = 1 - \epsilon \quad (7)$$

is a measure of the distance of a distribution from the singular distribution.

We assume that a uniform load is applied to the top layer. Hence the correlation in the top layer $c_h(m,0) = 1$. As shown in Sec. II A of Ref. [8], the subsequent evolution of the correlation is governed by

$$\begin{aligned} c_h(m,t+1) &= \sum_n H_{mn} c(n,t) \\ &= \left(\frac{1}{2} + \frac{\epsilon}{2} \delta_{m,0} \right) c_h(m,t) \\ &\quad + \left(\frac{1}{4} - \frac{\epsilon}{4} \delta_{m,1} \right) c_h(m-1,t) \\ &\quad + \left(\frac{1}{4} - \frac{\epsilon}{4} \delta_{m,-1} \right) c_h(m+1,t). \end{aligned} \quad (8)$$

The evolution matrix H may be interpreted as the Hamiltonian of a quantum particle on a lattice with a non-Hermitian barrier at the origin. It was shown in Sec. II B of Ref. [8] that one can obtain an integral expression for the correlation at depth t by making a biorthogonal expansion in the left and right eigenstates of the non-Hermitian Hamiltonian H . An important subtlety in this procedure is that one has to verify that the eigenstates of H are complete, since there is in general no guarantee that a non-Hermitian operator will have a complete set of eigenstates. Making use of the biorthogonal expansion and the expressions for the left and right eigenvectors of H derived in Ref. [8] we obtain¹

$$\begin{aligned} c_h(m,t) &= 1 + \int_{-\pi}^{\pi} \frac{dk}{2\pi} \cos^{2t} \left(\frac{k}{2} \right) e^{ikm} \\ &\quad \times \frac{[-i\epsilon^2 \sin k - \epsilon(1-\epsilon)(1+\cos k)]}{[(1-2\epsilon+2\epsilon^2) + (1-2\epsilon)\cos k]}. \end{aligned} \quad (9)$$

Equation (9) is an exact expression for the horizontal correlation function and is the main result of this subsection. We note that Eq. (9) applies for all m except $m = 0$. For $m = 0$ the horizontal correlation function $c_h(0,t)$ coincides with the variance calculated by Ref. [8] and the expression for $c_h(0,t)$ is given by their Eq. (41).

B. Critical exponents and scaling limit

To obtain the scaling limit of the correlation function we now simplify Eq. (9), assuming $t \gg 1$ and $\delta \ll 1$, but without making any assumptions about the relative size of t , δ , and m . Specifically, we first approximate $\cos^{2t}(k/2) \approx \exp(-k^2 t/4)$. This form is justified for small k and leads to negligible

¹To be specific, we start from the formal expression for $c_h(m,t)$ given in Eq. (38) of Ref. [8]. Expressions for $a^{(\pm)}(k)$ are given by Eqs. (35) and (40) of Ref. [8]; for $\phi^{(\pm)}(k)$ by Eqs. (28) and (30) of Ref. [8]; and for $\lambda(k)$ by Eq. (29) of Ref. [8]. Substituting these expressions in Eq. (38) of Ref. [8] and simplifying yields Eq. (9) of the present manuscript.

error for large k , since both the exact expression and the approximation are negligible for large k compared to $1/\sqrt{t}$. For the same reason we may extend the range of integration in Eq. (9) to infinity with negligible error. Finally, since the integrand has negligible weight for large k , we may expand both numerator and denominator in the second line of Eq. (9) to leading order in k and δ . After making these approximations and a rescaling of the integration variable we find

$$c_h(m,t) - 1 = -f(\theta,\mu) - \frac{\partial}{\partial \mu} f(\theta,\mu), \quad (10)$$

where $\theta = \delta\sqrt{t}$ is a scaled measure of depth and $\mu = 2m\delta$ is a scaled measure of horizontal separation and the function

$$f(\theta,\mu) = \int_{-\infty}^{\infty} \frac{du}{\pi} \frac{e^{-u^2\theta^2} e^{-i\mu u}}{1+u^2}. \quad (11)$$

It is evident from Eqs. (10) and (11) that the horizontal correlations have the expected scaling form Eq. (3). Furthermore, from the form of the scaled variables θ and μ we infer that the horizontal correlation length exponents $\nu = 1$ and the dynamical exponent $z = 2$.

To derive the asymptotic behavior of the correlation function in the scaling regime it is convenient to rewrite the correlation function as

$$\begin{aligned} c_h(m,t) - 1 &= -\frac{1}{\sqrt{\pi}} \int_{\mu/\theta}^{\infty} ds \\ &\quad \times \exp \left[-\theta \left(s - \frac{\mu}{\theta} \right) \right] \exp \left(-\frac{1}{4} s^2 \right). \end{aligned} \quad (12)$$

This form is obtained by noting that $f(\theta,\mu)$ is an integral over a product of a Gaussian and a Lorentzian. Using Parseval's theorem it may therefore be written as an integral over a product of the Fourier transforms of the Gaussian and Lorentzian factors. Using this transformed representation for $f(\theta,\mu)$ leads from Eqs. (10) to (12).

Now let us consider the regime of small depth, $\theta \ll 1$, or equivalently, $t\delta^2 \ll 1$ (the "critical regime"). In this regime we would expect the correlation function to be indistinguishable from the correlation function at the critical point (since, for fixed t , the case that $t\delta^2 \ll 1$ corresponds to the limit that $\delta \rightarrow 0$). Indeed, we find that for small θ Eq. (12) simplifies to

$$c_h(m,t) - 1 = -\frac{1}{\sqrt{\pi}} \int_{\mu/\theta}^{\infty} ds \exp \left(-\frac{1}{4} s^2 \right). \quad (13)$$

Thus the correlation function depends only on μ/θ or equivalently, m/\sqrt{t} and is independent of δ . Equation (13) reveals that in the critical regime the loads (or to be precise, the deviations in load from the mean) on neighboring beads are strongly anticorrelated. For small distances (m/\sqrt{t}) Eq. (13) simplifies to

$$c_h(m,t) - 1 \approx -1 + \frac{2m}{\sqrt{\pi t}} + \dots \quad (14)$$

At large distances ($m/\sqrt{t} \gg 1$) Eq. (13) simplifies to

$$c_h(m,t) - 1 \approx -\sqrt{\frac{t}{\pi m^2}} \exp \left(-\frac{m^2}{t} \right); \quad (15)$$

in other words, the anticorrelations decay as a Gaussian. In summary, we find that in the critical regime there are strong

horizontal anticorrelations whose range grows with the square root of the depth.

Next let us consider the regime of large depth, $\theta \gg 1$, or equivalently, $t\delta^2 \gg 1$ (the ‘‘saturated regime’’). In this regime Eq. (12) simplifies to

$$c_h(m, t) = -\frac{1}{\sqrt{\pi t \delta^2}} \exp\left(-\frac{m^2}{t}\right). \quad (16)$$

Thus in the saturated regime there are weak anticorrelations. The amplitude of these anticorrelations falls off inversely with depth while their range continues to grow as the square root of the depth.

III. VERTICAL CORRELATIONS

A. Exact solution

There is a remarkably simple exact relationship between the horizontal and vertical correlations for the q model [10]. In order to establish this relationship it is instructive to first consider the correlations between beads in two consecutive layers, t and $t + 1$, with a horizontal separation $2m + 1$. By use of the evolution Eq. (1) we may write

$$\begin{aligned} \langle w_n(t)w_{n+2m+1}(t+1) \rangle &= \langle f_{n+2m+2,t}w_n(t)w_{n+2m+2}(t) \rangle \\ &+ \langle (1 - f_{n+2m,t})w_n(t)w_{n+2m}(t) \rangle. \end{aligned} \quad (17)$$

Note that the load on the beads in layer t is independent of the fractions that appear on the right-hand side of Eq. (17); these fractions determine that the subsequent propagation of load from layer t propagates to layer $t + 1$. Hence the averages on the right-hand side of Eq. (17) factorize. Recalling that by virtue of the symmetry condition on the distribution of the fractions, $\langle f \rangle = \langle (1 - f) \rangle = 1/2$, and making use of the definition of the horizontal correlation Eq. (2) we obtain

$$\langle w_n(t)w_{n+2m+1}(t+1) \rangle = \frac{1}{2}c_h(2m+2, t) + \frac{1}{2}c_h(2m, t). \quad (18)$$

Similarly one can relate the correlation between beads separated by two layers, $c_v(m, t, \tau \rightarrow 1)$, to horizontal correlations by twice using the evolution Eq. (1) to obtain

$$c_v(m, t, 1) = \frac{1}{4}c_h(m+2, t) + \frac{1}{2}c_h(m, t) + \frac{1}{4}c_h(m-2, t). \quad (19)$$

By now the astute reader will have noted the appearance of binomial coefficients in Eqs. (18) and (19) and indeed, it is not difficult to show by induction that

$$c_v(m, t, \tau) = \frac{1}{2^{2\tau}} \sum_{k=-\tau}^{+\tau} \binom{2\tau}{\tau+k} c_h(m-k, t). \quad (20)$$

Equation (20) is the exact relationship between vertical and horizontal correlations and is the main result of this subsection.

B. Critical exponents and scaling limit

In this section we derive the universal scaling behavior of the vertical correlation function. There are three circumstances to consider. It is clear from Fig. 1 that each bead sits at the bottom of an inverted cone of beads that are partially supported by it. The correlation between two beads will evidently be strong if the upper bead of the pair lies inside the support

cone of the lower bead. Under this circumstance, which we call the ‘‘direct case,’’ $|m| \leq \tau$. On the other hand, if the two beads are so far apart that their support cones do not intersect at all the correlation between their loads is rigorously zero. The ‘‘uncorrelated case’’ corresponds to the condition $|m| > 2t + \tau$. Finally there is the intermediate ‘‘indirect case’’ in which the two support cones do intersect but the upper bead does not lie inside the support cone of the lower bead.

We first consider the scaling limit for the direct case. The sum in Eq. (20) can be separated into two parts. The first part corresponds to the single term with $k = m$. This term determines the contribution of the covariance in load of the upper bead to the vertical correlation with the lower bead. It will emerge that this covariance contribution leads to positive correlations and tends to dominate the direct vertical correlations. The second part corresponds to the remaining terms in Eq. (20). It will emerge that this contribution leads to anticorrelations and is generally subdominant.

We now compute the covariance contribution to the direct vertical correlation. Retaining only the $k = m$ term in Eq. (20) we obtain

$$c_v(m, t, \tau) = \frac{1}{2^{2\tau}} \binom{2\tau}{\tau+m} c_h(0, t). \quad (21)$$

Assuming that $\tau \gg 1$ and $|m| \gg 1$, we may approximate the binomial coefficient as a Gaussian. Finally, using the scaling limit of $c_h(0, t)$ given in Eq. (48) of Ref. [8] we obtain²

$$\begin{aligned} c_v(m, t, \tau) - 1 &= \frac{1}{\sqrt{\pi \tau \delta^2}} \exp\left[-\frac{(m\delta)^2}{\tau \delta^2}\right] \\ &\times \left[1 - \frac{1}{\sqrt{\pi}} \int_0^\infty ds \exp(-s\sqrt{t\delta^2})e^{-s^2/4}\right]. \end{aligned} \quad (22)$$

Equation (22) gives the covariance contribution to the direct vertical correlations in the scaling limit. It agrees with the conjectured universal scaling form in Eq. (5) and we find that the critical exponents are $\beta = 0, \nu = 1$, and $z = 2$.

To gain further insight into the covariance contribution to the direct vertical correlations, it is useful to consider the limiting behavior of Eq. (22) in the critical regime ($t\delta^2 \ll 1$) and the saturated regime ($t\delta^2 \gg 1$). In the former limit the dependence on δ cancels and we find

$$c_v(m, t, \tau) - 1 = \frac{2}{\pi} \sqrt{\frac{t}{\tau}} \exp\left[-\frac{m^2}{\tau}\right]. \quad (23)$$

Thus in the critical regime there are strong vertical correlations (not anticorrelations) and these correlations grow with the square root of the depth t . The horizontal range of the correlations grows as the square root of the vertical separation between the two beads, τ . If we imagine increasing the horizontal separation of the beads, keeping the vertical separation fixed, then the correlations die away well before

²The alert reader will note that we have also transformed Eq. (48) of Ref. [8], which features an integral of the product of a Gaussian and a Lorentzian into an integral over the corresponding Fourier transforms instead by use of Parseval’s theorem.

we reach the boundary of the support cone. In the saturated regime we find

$$c_v(m, t, \tau) = \frac{1}{\delta} \frac{1}{\sqrt{\pi\tau}} \exp\left(-\frac{m^2}{\tau}\right). \quad (24)$$

Here, too, there are strong vertical correlations (not anticorrelations), but these correlations have saturated and are no longer growing with depth t . Their horizontal range grows as the square root of the vertical separation between the beads, τ . The correlations die away well before we reach the boundary of the support cone, just as they do in the critical regime.

We turn now to the second contribution to the vertical correlations in the direct case, namely, the sum over terms with $k \neq m$ in Eq. (20). In the scaling regime we may approximate the binomial coefficients in Eq. (20) as a Gaussian, convert the sum into an integral, and extend the range of integration to infinity to obtain

$$c_v(m, t, \tau) = \int_{-\infty}^{\infty} dk \frac{1}{\sqrt{\pi\tau}} \exp\left(-\frac{k^2}{\tau}\right) c_h(m - k, t). \quad (25)$$

Using Eq. (12) for c_h we may perform the integral over k explicitly to obtain

$$c_v(m, t, \tau) - 1 = -\frac{1}{\sqrt{\pi}} \exp\left[-\frac{(m\delta)^2}{(t + \tau)\delta^2}\right] \int_0^{\infty} ds e^{-s^2/4} \times \exp\left[-s\left(\sqrt{(t + \tau)\delta^2} - \frac{m\delta}{\sqrt{(t + \tau)\delta^2}}\right)\right]. \quad (26)$$

Note that this contribution to the vertical correlation function is also of the conjectured form of Eq. (5), with the same critical exponents $\beta = 0$, $\nu = 1$, and $z = 2$. It is noteworthy that this contribution is a function only of $(t + \tau)\delta^2$ and the ratio $m\delta/\sqrt{(t + \tau)\delta^2}$; it is a more restricted function of its arguments than required by the scaling form Eq. (5). The full vertical correlation in the direct case is the sum of Eq. (22) and Eq. (26); however, as noted above and shown below, the contribution of Eq. (26) is generally dominated by the contribution of Eq. (22).

To gain further insight into the second contribution to the direct vertical correlations, it is useful to consider the limiting behavior of Eq. (26) in the critical regime ($t\delta^2 \ll 1$) and the saturated regime ($t\delta^2 \gg 1$). In the former limit the dependence on δ cancels and we find

$$c_v(m, t, \tau) - 1 = -\frac{1}{\pi} \exp\left(-\frac{m^2}{t + \tau}\right) \int_0^{\infty} ds e^{-s^2/4} \times \exp\left(\frac{ms}{\sqrt{t + \tau}}\right). \quad (27)$$

Equation (27) is still a formidable expression but it simplifies to $c_v(m, t, \tau) - 1 \approx -1$ for $m/\sqrt{t + \tau} \ll 1$ and to $c_v(m, t, \tau) - 1 \approx -2$ for $m/\sqrt{t + \tau} \gg 1$. Thus the second contribution to the direct vertical correlation in the critical regime corresponds to a weak anticorrelation that is negligible compared to the strong positive correlation implied by the first contribution, Eq. (23). Next let us consider the saturated regime ($t\delta^2 \gg 1$).

In this regime Eq. (26) simplifies to

$$c_v(m, t, \tau) - 1 = -\frac{1}{\delta} \frac{1}{\sqrt{\pi(t + \tau)}} \exp\left(-\frac{m^2}{t + \tau}\right). \quad (28)$$

Equation (28) should be compared to the first contribution, Eq. (24). Thus in the saturated regime, too, the second contribution to the direct vertical correlations is smaller than the first contribution.

In summary, in the direct case the vertical correlation is the sum of two contributions, both of which have the conjectured universal scaling form, Eq. (5), with exponents $\beta = 0$, $\nu = 1$, and $z = 2$. The first contribution, Eq. (22), corresponds to a positive correlation and generally dominates. The second contribution, Eq. (26), corresponds to an anticorrelation and is generally subdominant. It is illuminating to look at the limiting behavior of the two components in the critical regime ($t\delta^2 \ll 1$) and the saturated regime. In the critical regime the first contribution is given by Eq. (23) and the second contribution, which lies between -1 and -2 , is given by Eq. (27). In the saturated regime the first contribution is given by Eq. (24) and the second contribution by Eq. (28).

Finally, we now turn to the vertical correlation in the indirect case, $|m| > \tau$. In this case the first contribution is absent, since the term $k = m$ lies outside the range of the sum in Eq. (20). Thus in the indirect case there is only a weak anticorrelation between beads given by Eq. (26). This concludes our analysis of the vertical correlations.

IV. CONCLUSION

In this paper we add to the existing body of known results about the q model by obtaining results on the exact horizontal and vertical correlation functions. We show that the correlation functions have a universal scaling form close to the critical point identified in previous work [8] and we determine the scaling form and associated critical exponents. It is intriguing that the q model, which describes physics far from equilibrium, nonetheless shows behavior reminiscent of equilibrium thermodynamic critical phenomena. Among the many questions that our work leaves open, we here mention two. First we note that it may be possible to calculate three-point and higher correlation functions by means of a non-Hermitian generalization of the Bethe ansatz [13]. Second, it is desirable to determine the dynamics of the entire distribution of load on a single bead. Reference [8] obtained the dynamics of the entire distribution right at the critical point, and Snoeijer and van Leeuwen [12] studied the large depth asymptotic dynamics of the load distribution in a case far from the critical point. Close to the critical point Ref. [8] made a scaling hypothesis about the form of this distribution and it remains of interest to verify this conjecture via numerics or exact solution.

ACKNOWLEDGMENTS

A.S. acknowledges support from the REU program at Case Western Reserve University via NSF Grant No. DMR-0850037. H.M. acknowledges support by the DOE. We also acknowledge valuable past discussions with Onuttom Narayan that motivated the present work and wish to thank Philip Taylor for discussion of this work and comments on the manuscript.

- [1] S. N. Coppersmith, C.-H. Liu, S. N. Majumdar, O. Narayan, and T. A. Witten, *Phys. Rev. E* **53**, 4673 (1996).
- [2] C.-H. Liu, S. R. Nagel, D. A. Schecter, S. N. Coppersmith, S. N. Majumdar, O. Narayan, and T. A. Witten, *Science* **269**, 513 (1995); D. M. Mueth, H. M. Jaeger, and S. R. Nagel, *Phys. Rev. E* **57**, 3164 (1998).
- [3] C. Goldenberg and I. Goldhirsch, *Nature* **435**, 188 (2005); T. S. Majumdar and R. P. Behringer, *ibid.* **435**, 1079 (2005); E. I. Corwin, H. M. Jaeger, and S. R. Nagel, *ibid.* **435**, 1075 (2005); A. R. T. van Eerd, W. G. Ellenbroek, M. van Hecke, J. H. Snoeijer, and T. J. H. Vlugt, *Phys. Rev. E* **75**, 060302(R) (2007).
- [4] H. Takayasu, *Phys. Rev. Lett.* **63**, 2563 (1989); H. Takayasu, I. Nishikawa, and H. Tasaki, *Phys. Rev. A* **37**, 3110 (1988).
- [5] J. T. Chalker and A. Dohmen, *Phys. Rev. Lett.* **75**, 4496 (1995).
- [6] For a review, see B. I. Shraiman and E. D. Siggia, *Nature* **405**, 639 (2000).
- [7] A. E. Scheidegger, *Bull. Intl. Assoc. Sci. Hydrol.* **12**, 15 (1967); P. S. Dodds and D. H. Rothman, *Phys. Rev. E* **59**, 4865 (1999).
- [8] M. Lewandowska, H. Mathur, and Y. K. Yu, *Phys. Rev. E* **64**, 026107 (2001).
- [9] M. E. Fisher, *J. Stat. Phys.* **34**, 667 (1984).
- [10] R. Rajesh and S. N. Majumdar, *Phys. Rev. E* **62**, 3186 (2000).
- [11] J. H. Snoeijer and J. M. J. van Leeuwen, *Phys. Rev. E* **65**, 051306 (2002).
- [12] J. H. Snoeijer and J. M. J. van Leeuwen, *J. Stat. Phys.* **109**, 449 (2002).
- [13] H. Mathur (unpublished).