

From closed to open one-dimensional Anderson model: Transport versus spectral statistics

S. Sorathia,¹ F. M. Izrailev,^{1,2,*} V. G. Zelevinsky,² and G. L. Celardo³

¹*Instituto de Física, Universidad Autónoma de Puebla, Apartado Postal J-48, Puebla, Puebla 72570, Mexico*

²*NSCL and Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824-1321, USA*

³*Dipartimento di Matematica e Fisica, Università Cattolica, via Musei 41, 25121, Brescia, and INFN, Sezione di Pavia, Italy*

(Received 30 April 2012; revised manuscript received 11 July 2012; published 31 July 2012)

Using the phenomenological expression for the level spacing distribution with only one parameter $0 \leq \beta \leq \infty$ covering all regimes of chaos and complexity in a quantum system, we show that transport properties of the one-dimensional Anderson model of finite size can be expressed in terms of this parameter. Specifically, we demonstrate a strictly linear relation between β and the normalized localization length for the whole transition from strongly localized to extended states. This result allows one to describe all transport properties in the open system entirely in terms of the parameter β and the strength of the coupling to the continuum. For nonperfect coupling, our data show a quite unusual interplay between the degree of internal chaos defined by β and the degree of openness of the model. The results can be experimentally tested in single-mode waveguides with either bulk or surface disorder.

DOI: [10.1103/PhysRevE.86.011142](https://doi.org/10.1103/PhysRevE.86.011142)

PACS number(s): 05.60.Gg, 46.65.+g, 71.23.An, 72.15.Qm

I. INTRODUCTION

In spite of remarkable progress in understanding the statistical properties of quantum systems, either deterministic or disordered, one of the important problems still awaits detailed analysis. The specific question is as follows: To what extent can we predict the scattering properties of a complex *open* system, if we know the global properties of eigenstates and energy spectra of the corresponding *closed* system? This problem was solved for a specific case of closed systems with maximal chaotic properties described by fully random matrices. A proper mathematical tool in these studies is based on effective non-Hermitian Hamiltonians of a certain structure [1], and many analytical results have been obtained; see, for example, [2–4]. The key point in this method is that the scattering matrix of an open system is expressed in terms of eigenvalues and eigenfunctions of the related closed system along with their decay amplitudes.

A much more difficult problem emerges if the closed system is not fully chaotic, being characterized by additional parameters related to the degree of chaos. Recently, this problem was analyzed numerically in Refs. [5–7] where the global characteristics of scattering or signal transmission through a system have been studied in dependence on two control parameters, the degree of internal chaos and the strength of coupling to the continuum. In particular, it was found that, independently of the degree of chaos, increasing continuum coupling leads the system from the quasistable regime of isolated narrow resonances to the “super-radiant” regime of overlapping resonances coexisting with long-lived compound states. However, the specific features of this evolution may critically depend on the degree of chaos and therefore the observation of the signal transmission provides important information on the regular or chaotic character of the intrinsic dynamics.

The above studies were performed with the use of random matrices (canonical Gaussian ensembles or two-body random

interactions). Some statistical assumptions may be questionable in application to realistic physical systems. Below we study the one-dimensional (1D) Anderson model, paying most attention to the relation between the scattering properties of an open model and those of the eigenstates and spectral statistics of the closed model. We discover unexpected effects that give additional insight into the problem of scattering through finite disordered systems.

II. THE MODEL

The tight-binding Anderson model is often used to describe electron propagation in random media. In the 1D case the corresponding Hamiltonian with diagonal disorder takes the standard form,

$$H_{mn} = \epsilon_n \delta_{mn} - \nu(\delta_{m,n+1} + \delta_{m,n-1}). \quad (1)$$

Here ν is the hopping amplitude connecting the nearest sites (in what follows we set $\nu = 1$); the site energies ϵ_n are assumed to be uniformly distributed in the interval $[-W/2, W/2]$, giving rise to the disorder variance $\sigma^2 = W^2/12$. Our interest is in the transmission properties through samples of *finite size* N with arbitrary *coupling amplitudes* $\sqrt{\gamma^L}$ and $\sqrt{\gamma^R}$ connecting the left and right edges with attached semi-infinite *ideal* leads in which $\epsilon_n = 0$; see Fig. 1. For zero disorder, open tight-binding models were studied in [8,9]. For nonzero disorder, so far, the main interest has been related to the statistics and distributions of resonances for one open channel [10,11]; the relation of the resonances to the transport properties was studied in [12].

Without disorder, the spectrum of the closed chain consists of Bloch waves with the nodes at the ends and energies inside the band $|E| \leq 2$. In the limit $N \rightarrow \infty$ and for weak disorder, $\sigma^2 \ll 1$, all eigenstates are exponentially localized with the characteristic length $l_\infty(E)$ given by the Thouless relation [13]

$$l_\infty^{-1}(E) = \frac{\sigma^2}{8(1 - E^2/4)}. \quad (2)$$

This expression is valid everywhere apart from the vicinity of the band edges, $|E| = 2$, and band center, $E = 0$.

*felix.izrailev@gmail.com




FIG. 1. Disordered 1D lattice consisting of N sites connected at both ends to ideal semi-infinite tight-binding leads.

III. LEVEL REPULSION IN A CLOSED MODEL

To quantify the degree of chaos in finite samples with no continuum couplings, $\gamma^L = \gamma^R = 0$, we employ the well-known results of the random matrix theory (RMT). In deterministic quantum models with chaotic behavior in the classical counterpart, chaos is usually characterized by the Wigner-Dyson (WD) distribution $P(s)$ of normalized spacings s between the nearest energy levels. In the opposite case of integrable classical counterparts, $P(s)$ is typically close to the Poisson distribution. Therefore, one can take the distribution $P(s)$ as a measure of chaos in a closed model.

For zero disorder, $\epsilon_n = 0$, the energy spectrum near the band center is equidistant, so that $P(s) \rightarrow \delta(s - 1)$, and the eigenstates are extended and regular (standing waves). In the limit of strong disorder, all eigenstates are effectively localized on the scale of the sample size $N \gg 1$; thus the form of $P(s)$ should be close to Poissonian. In between these limits, for $l_\infty \propto N$, it is natural to expect a kind of “intermediate statistics” which can be compared with the WD distribution. However, the latter is known to emerge when the eigenstates are chaotic with random fluctuations of their components, as happens in ensembles of full random matrices [14]. It is also known that chaotic eigenstates appear in quasi-one-dimensional models described by band random matrices, when the localization length is larger than the sample size. Such a situation physically corresponds to the diffusion of wave packets [15]. In contrast, in our model described by tridiagonal matrices [see Eq. (1)], the diffusion scale is absent since the localization length is proportional to the mean free path. For this reason, in the theory of disordered solids an emergence of the WD distribution for the one-dimensional Anderson model of finite size is of special interest [16].

In order to describe the entire evolution of $P(s)$ as a function of the strength of disorder, we use the phenomenological expression suggested in Ref. [17],

$$P_\beta(s) = B_1 z^\beta (1 + B_2 \beta z)^{f(\beta)} \exp \left[-\frac{1}{4} \beta z^2 - \left(1 - \frac{\beta}{2}\right) z \right], \quad (3)$$

where

$$f(\beta) = \beta^{-1} 2^\beta \left(1 - \frac{\beta}{2}\right) - 0.16874. \quad (4)$$

Here $z = \pi s/2$ and the parameters B_1 and B_2 are determined by the normalization conditions

$$\int_0^\infty P_\beta(s) ds = \int_0^\infty s P_\beta(s) ds = 1. \quad (5)$$

The above formula (3) was suggested in Refs. [17,18] by using the analytical expressions derived by Dyson [19] for a classical gas of two-dimensional charged particles moving on a ring at temperature $1/\beta$. This model was found to be very effective since for the values $\beta = 1, 2, 4$ the partition

function giving the probability of finding the particles at specific positions coincides with the partition function for eigenvalues in canonical Gaussian ensembles.

One of the original motivations for the expression (3) was to have a unique formula for $P(s)$ that would provide correct results in particular cases of conventional random matrix ensembles. Specifically, the function $f(\beta)$ is constructed in such a way that for the values $\beta = 1, 2, 4$ corresponding to Gaussian ensembles of random matrices of a given symmetry (orthogonal, unitary, and symplectic, respectively) it is close to the expressions for $P(s)$ obtained in the RMT for those ensembles [14]. As shown in Ref. [18], for these values of β the dependence (3) is more accurate (in the whole range of s) than the WD distribution typically used in the literature.

Since for $\beta = 0$ Eq. (3) coincides with the Poisson distribution, and for $\beta = \infty$ it reproduces the δ function, this interpolation formula is a perfect candidate for the description of the intermediate statistics in the closed Anderson model (1). An expression similar to Eq. (3) [20] has already been used to describe $P(s)$ in finite one- and two-dimensional Anderson models [21,22], resulting in values of β from 0.05 to 19.2 depending on the disorder strength.

It should be stressed that in application to our model the parameter β in Eq. (3) should be considered as a parameter which *globally* describes the distribution $P(s)$, rather than a parameter determining the repulsion of energy levels for very small spacings $s \ll 1$. To date, there are indications that the repulsion between neighboring energy levels in the finite Anderson model cannot be rigorously associated with the symmetry of the matrix H_{mn} [see Eq. (1)], as happens in the case of full random matrices. Indeed, in Refs. [23,24] it was analytically shown that for 3×3 real symmetric matrices with matrix elements $a_{13} = a_{31} = 0$ and other elements random, the repulsion for small $s \ll 1$ is nonlinear, $P(s) \sim s \ln(1/s)$. For 4×4 tridiagonal random matrices the repulsion appears to be $P(s) \sim s \ln^2(1/s)$ [24]. The extensive numerical study performed in Ref. [24] allows one to predict the general dependence $P(s) \sim s \ln^{N-2}(1/s)$ for any size N of tridiagonal matrix with all nonzero elements random. For a slightly different type of random tridiagonal matrix, the rigorous result obtained in Ref. [25] gives $P(s) \sim s \ln^{N-3}(1/s)$. All these results indicate that the level repulsion (defined in the limit $s \rightarrow 0$) in disordered tridiagonal matrices strongly depends on the specific properties of the system.

With Eq. (3) we performed an extensive numerical study of $P(s)$ by changing the degree of disorder for a closed chain in a large range of the control parameter $x = l_\infty/N$ with l_∞ defined by Eq. (2); see the examples in Fig. 2. Our data demonstrate that Eq. (3) gives an amazingly good correspondence (supported by the χ^2 criteria) with the numerical data in a very large range of x . Therefore, one can make the unexpected conclusion that the distribution $P(s)$ for the Anderson model on a finite scale N can be described by the Dyson Coulomb gas model, where s is the distance between the nearest particles on a ring.




FIG. 2. (Color online) Examples of $P(s)$ for $|E| < 0.2$ (excluding the energies very close to 0), with $N = 1000$. The data are obtained for 120 disorder realizations with the χ^2 fit to Eq. (3) for $r = 40$ bins. The reduced χ^2 statistic is shown with the corresponding confidence levels given in parentheses.

Another result is that for the specific degree of disorder, namely, for $x = l_\infty/N \approx 0.435$ [see Eq. (6) below], the distribution $P(s)$ is indistinguishable from the WD distribution. This is in correspondence with the numerical results earlier obtained in Ref. [21]. It is known that in the RMT an emergence of the WD distribution is directly related to the random character of eigenstates [14]. The numerical data [21], indeed, show that, in the situation when $P(s)$ is close to the WD distribution, some of the global statistical characteristics of the eigenstates are in good correspondence with the assumption of their randomness. However, more detailed studies of the eigenstates are needed in order to claim that they have the same degree of randomness as in the conventional ensembles of random matrices. Our preliminary results show that, in spite of a very good correspondence of $P(s)$ with the WD distribution, the more small-scale characteristics of eigenstates demonstrate small but clear deviations from the predictions of the RMT.

IV. LOCALIZATION LENGTH VERSUS REPULSION

Now we can establish the relation between x and β . It is qualitatively clear that they express the same phenomenon of gradual transformation of standing waves into localized states, but the exact relation between them was unknown. All the data are summarized in Fig. 3. We see a precise *linear dependence* between x and β in the whole range of x values independently of the chosen energy range. The fit of the data as $\beta = Ax + C$ gives the slope $A = 2.3 \pm 0.1$ with C essentially zero. This result, obtained carefully with the use of χ^2 statistical criteria, shows that the repulsion parameter β is just the normalized localization length l_∞ ,

$$\beta \approx 2.3 \frac{l_\infty}{N}. \quad (6)$$

The factor 2.3 in Eq. (6) can be attributed to the fluctuations of components of the eigenstates.




FIG. 3. (Color online) Repulsion parameter β versus $x = l_\infty/N$ for $E \approx 0$ (circles) and $E \approx -1$ (squares); see Fig. 2.

The localization length l_∞ of a given state with the site components ψ_n can be defined through the Shannon entropy,

$$S = - \sum_{n=1}^N w_n \ln w_n \quad (7)$$

with $w_n = \psi_n^2$. For random states with a Gaussian distribution of ψ_n one gets $S = \ln(N/2.07)$. We define the normalized *entropic localization length* $d_N = 2.07 \exp(S)$, where $\langle \dots \rangle$ represents an ensemble average. With this definition we have $d_N = N$ for fully chaotic eigenfunctions occupying N sites. The data show that the onset of strong chaos occurs when $d_N \approx 2.1l_\infty$ [26] which is equivalent to $\beta \approx d_N/N \approx 1$. Therefore, one arrives at the same result through an analysis of the spectrum as through the eigenstates.

According to this important result, the repulsion parameter β is nothing but the properly rescaled localization length. This was observed in numerical studies of the kicked rotor [27] and Wigner banded random matrices [28]. Here, this result emerges for the standard Anderson model, thus indicating a generic effect. Very recently, a linear relation between the repulsion parameter β and localization length was found in an experimental study of disordered elastic rods [29].

V. OPEN MODEL: NON-HERMITIAN HAMILTONIAN

The scattering properties of *open* systems can be formulated [1] with the effective non-Hermitian Hamiltonian [8,10,12,26]

$$\mathcal{H}_{mn}(E) = H_{mn} + \mathcal{D}(E)(\gamma^L \delta_{n,1} \delta_{m,1} + \gamma^R \delta_{n,N} \delta_{m,N}), \quad (8)$$

where

$$\mathcal{D}(E) = \frac{E}{2} - \frac{i}{2} \sqrt{4 - E^2}. \quad (9)$$

This expression is valid for any disorder ϵ_n , continuum coupling γ^L, γ^R , and energy E . Near the center of the band (the general case, $-2 < E < 2$, is studied in Ref. [26]) Eq. (8) reduces to the canonical form,

$$\mathcal{H}_{mn}(E) = H_{mn} - \frac{i}{2} W_{mn} \quad (10)$$

with

$$W_{mn} = 2\pi \sum_{c=L,R} A_m^c(E) A_n^c(E), \quad (11)$$

where $W_{mn}(E)$ is defined by the coupling amplitudes

$$A_n^{L,R}(E) = \sqrt{\gamma^{L,R}/\pi} \left[1 - \frac{E^2}{4} \right]^{1/4} (\delta_{n,1}^{(L)} + \delta_{n,N}^{(R)}). \quad (12)$$

Here H_{mn} in Eq. (1) describes the internal dynamics, while the non-Hermitian part $W(E)$ is factorized in terms of the coupling amplitudes $A_n^c(E)$ between the internal states $|n\rangle$ and open decay channels, $c = L, R$, where L and R stand for left and right, respectively.

The non-Hermitian Hamiltonian (10) allows us to construct the scattering matrix S in the space of channels,

$$S = \frac{1 - i\pi K}{1 + i\pi K}, \quad (13)$$

where the reaction matrix K is defined as

$$K^{ab}(E) = \sum_j \frac{\tilde{A}_j^a \tilde{A}_j^b}{E - E_j}, \quad \tilde{A}_j^c = \sum_m A_m^c \psi_m^{(j)}, \quad (14)$$

and $\psi_m^{(j)}$ is the m component of the eigenstate $|j\rangle$ of the closed Hermitian Hamiltonian (1).

For weak disorder, $\sigma^2 \ll 1$, the strength of coupling to the leads is characterized by the coupling parameter

$$\kappa^c = \frac{2\pi\gamma^c}{ND}. \quad (15)$$

Here D is the mean level spacing at the center of the energy band in a closed chain. By this definition, the channel transmission coefficient is maximal for *perfect coupling* when $\kappa^c = 1$. Below we consider symmetric coupling $\gamma^c \equiv \gamma$, $\kappa^c \equiv \kappa$. In all numerical simulations we take $N = 1000$ sites and combine an ensemble average over a large number of realizations with a spectral average over 1000 energies across an energy window $|E| < 0.2$.

VI. TRANSPORT CHARACTERISTICS

For *continuous* weak random potentials the problem of scattering through finite one-dimensional samples is rigorously solved by various analytical approaches (see, for example, [30,31]). It was shown that the distribution function for the transmission coefficient T (conductance) depends on the ratio of the localization length l_∞ to the length N of a sample. Therefore, the knowledge of the localization length (defined in the limit $N \rightarrow \infty$) gives a full description of scattering (for perfect coupling with the leads).

In contrast with continuous models, for the *tight-binding model* (1) there are no rigorous results, either for the distribution of T or for the moments $\langle T^q \rangle$, that would be valid everywhere inside the energy spectra $|E| < 2$, even for weak disorder. The reason is the existence of the so-called resonances for specific energies $E_r = \pm 2 \cos(\pi r/s)$ with r, s integer, for which standard perturbation theory fails. The most studied case is the band center where the Thouless expression (2) has to be corrected due to an anomaly precisely at $E = 0$ for which $l_\infty \simeq 105.2/W^2$ instead of $96/W^2$ (for details and references, see [31] and [32]). On the other hand, one can expect that Eqs. (16) and (17) can also be used if the energy is not too close to the band center or band edges, with l_∞ defined by Eq. (2). We have checked this conjecture for the two lowest moments of the transmission coefficient, defined




FIG. 4. (Color online) Upper panel: $\langle T \rangle$ versus $\beta = 2.3x$ for different coupling strengths κ ; the smooth curve shows the analytical expression Eqs. (16) and (17). Lower panel: Dependence $\langle T \rangle$ versus κ for different values of β . For $\beta \rightarrow \infty$, the average is found to tend towards $2\kappa/(1 + \kappa^2)$, depicted as the smooth curve.

as $T = |S^{LR}|^2$ [30,31],

$$\langle T^q \rangle = \sqrt{\frac{2x^3}{\pi}} \exp\left(-\frac{1}{2x}\right) \int_0^\infty f_q(z) \exp\left(-\frac{z^2 x}{2}\right) dz, \quad (16)$$

where $x = l_\infty/N$. Here $\langle \dots \rangle$ stands for the ensemble average, and the functions f_q for $q = 1, 2$ are

$$f_1(z) = \frac{z^2}{\cosh z}, \quad f_2(z) = \frac{2z^2 + z \sinh 2z}{4 \cosh^3 z}. \quad (17)$$

Our numerical data, indeed, manifest the correspondence with the theoretical expression for the mean transmission with $\kappa = 1$; see the upper panel of Fig. 4. Here and in the following we express the degree of localization in terms of β using Eq. (6).

In this figure (upper panel) we also present the transmission coefficient $\langle T \rangle$ for the case of nonperfect coupling, for which the analytical results are absent. The maximal value of $\langle T \rangle$ occurs for perfect coupling, $\kappa = 1$. As for nonperfect coupling, the lower panel of Fig. 4 shows that the transmission coefficient is symmetric with respect to the change $\kappa \rightarrow 1/\kappa$, which is known for models described by full random matrices in place of H_{mn} in Eq. (10) (see, e.g., [3]). However, our results manifest that even in the presence of strongly localized, $\beta \ll 1$, or regular extended eigenstates, $\beta \gg 1$, in a closed model, this symmetry is preserved when the system is open.

A very good correspondence was also observed for $\langle T^2 \rangle$. This gives the possibility for a discussion of the variance of the transmission, $\text{Var}(T) \equiv \langle T^2 \rangle - \langle T \rangle^2$, characterizing *mesoscopic fluctuations* of the transmission. In the upper panel of Fig. 5, we plot $\text{Var}(T)$ versus β for different fixed values of κ . Once more, an excellent agreement between the analytical expressions given by Eqs. (16) and (17) and the numerical data based on the discrete model (10) can be seen when $\kappa = 1$. For this case, the variance reaches a maximum for $\beta \approx 4.0$ which is much larger than that corresponding to maximal internal chaos ($\beta = 1$), as one may naively expect. For $\beta \approx 4$, the eigenstates are extended; however, they are neither strongly chaotic nor




FIG. 5. (Color online) Upper panel: $\text{Var}(T)$ versus β for different values of κ . The solid curve corresponds to the analytical value for $\text{Var}(T)$ obtained from Eqs. (16) and (17). Lower panel: $\text{Var}(T)$ versus κ for different values of β .

regular. Our data demonstrate a nonmonotonic dependence of $\text{Var}(T)$ on both sides of $\kappa = 1$ for some critical value $\beta_{cr} \sim 4$ and above. Below this critical value the single maximum of $\text{Var}(T)$ occurs at perfect coupling whereas above β_{cr} there are two maxima on both sides.

Moving away from nonperfect coupling, a clear symmetry is seen for the mutually inverse values $\kappa = 10$ and 0.1 ; this symmetry is also observed in the lower panel of the same figure where we plot the variance of T versus κ for different values of β . Such a symmetry has been repeatedly mentioned in the literature; see, for example, [5]. It follows from the physics of many intrinsic states coupled to the same decay channels; one of the first derivations follows from the Moldauer-Simonius relation [33]. With κ growing beyond the super-radiant transition at $\kappa \sim 1$, the broad state becomes a part of the background while the narrow resonances return to the nonoverlap regime.

The analytical estimates (16) and (17) for $\langle T \rangle$ are valid for perfect coupling, $\kappa = 1$, only. On the other hand, one can obtain an analytical expression for $\langle \ln T \rangle$ which is valid for any value of κ and again reveals the symmetry $\kappa \leftrightarrow 1/\kappa$,

$$\langle \ln T \rangle = -\frac{2N}{l_\infty} + 2 \ln \left[\frac{4\kappa}{(1+\kappa)^2} \right]. \quad (18)$$

It can be derived via the product of $N + 2$ transfer matrices, by tracing out two (nonrandom) matrices describing the coupling to the leads. This expression can be used for the definition of the localization length l_∞ in the presence of nonperfect coupling. Our numerical data for $\langle \ln T \rangle$ manifest excellent correspondence with Eq. (18), at the band center (see Fig. 6), as well as for different values of β (not shown).

VII. CORRELATIONS

The knowledge of the scattering matrix S defined by Eq. (13) allows one to study many details of scattering. One problem of both theoretical and experimental interest is to understand how the correlations between different cross




FIG. 6. (Color online) $\langle \ln T \rangle - f(\kappa)$ vs N for different values of κ . Here $f(\kappa) = 2 \ln [4\kappa/(1+\kappa)^2]$ [see Eq. (18)] and $W = 1$; $l_\infty = 105.2$ for $E = 0$.

sections depend on the degree of localization for different coupling strengths. This problem was recently addressed in Refs. [5–7] where the non-Hermitian Hamiltonians contained, apart from a diagonal part, bandlike random symmetric matrices. Such models are typically studied in application to open complex systems, like heavy nuclei and quantum dots. There, the amplitudes A_m^c connecting intrinsic states $|m\rangle$ with many open channels $c = 1, \dots, M \gg 1$ were assumed to be completely random and independent of internal dynamics. In contrast, in our case the derivation of the Hamiltonian (10), for specified internal disorder $\{\epsilon_n\}$, does not involve any additional assumption of randomness for the coupling to continuum. Our main interest here is to understand how the properties of cross section correlations depend on whether the internal eigenstates are localized or extended, and how the correlations depend on the coupling to the continuum.

The average cross section can be divided into two parts corresponding to *direct* and *fluctuating* processes, respectively (see, e.g., [5]). The direct processes correspond to a very fast passage of an incoming particle (or wave) through the scattering region, thus resulting in broad short-lived states of an open system. In contrast, the fluctuating parts of cross sections describe narrow long-lived states (resonances) known in nuclear physics as “compound” states. In our model with $M = 2$ channels, the transmission and reflection coefficients are defined as follows:

$$T = |S^{LR}|^2 = |S_{fi}^{LR}|^2, \quad R = |S^{LL}|^2 = |(S^{LL}) + S_{fi}^{LL}|^2, \quad (19)$$

where “ fl ” stands for the fluctuating parts. With the usual definition of the (partial) cross section $\sigma^{ab} = |\delta_{ab} - S^{ab}|^2$, we define the fluctuating cross sections as $\sigma^{LR} = |S_{fi}^{LR}|^2$ and $\sigma^{LL} = |S_{fi}^{LL}|^2$. Therefore, the average fluctuating part of the two cross sections can be expressed in terms of the average transmission T and reflection R coefficients:

$$\langle \sigma^{LR} \rangle = \langle T \rangle, \quad \langle \sigma^{LL} \rangle = \langle R \rangle - \langle S^{LL} \rangle^2. \quad (20)$$

The average elements of the scattering matrix can be written as [compare with Eq. (13)]

$$\langle S^{ab} \rangle = \frac{1-\kappa}{1+\kappa} \delta_{ab}. \quad (21)$$

This expression is well known in the RMT (see, for example, [3]). One can show that it is also valid for the considered




FIG. 7. (Color online) Correlations C_1 (upper) and C_2 (lower) versus β for $\kappa = 1, 0.5, 0.2$. The solid curves are the analytical expressions (16) and (17) for $-\text{Var}(T)$ and $\text{Var}(T)$ shown in the upper and lower panels, respectively.

Anderson model [26]. Note that for perfect coupling, $\kappa = 1$, the mean values $\langle \sigma^{LL} \rangle$ and $\langle \sigma^{LR} \rangle$ of the cross sections are nothing but the average transmission and reflection coefficients, respectively.

Two types of correlations of our interest are defined as

$$C_1 = \langle \sigma^{LL} \sigma^{LR} \rangle - \langle \sigma^{LL} \rangle \langle \sigma^{LR} \rangle, \quad (22)$$

$$C_2 = \langle \sigma^{LL} \sigma^{RR} \rangle - \langle \sigma^{LL} \rangle \langle \sigma^{RR} \rangle. \quad (23)$$

In the literature they are referred to as the *covariances*, widely studied in connection with transmission through waveguides with bulk and surface scattering (see, for example, [34] and references therein). The result of computation of the correlations (22)–(23) is presented in Fig. 7 as a function of the repulsion parameter β for different values of κ . For perfect coupling, the data are well described by the analytical

expressions, since in this case we simply have

$$C_1 = -\text{Var}(T), \quad C_2 = \text{Var}(T). \quad (24)$$

Referring to Fig. 8 for nonperfect coupling, one has to stress the following. First, both for C_1 and C_2 there is a critical value of coupling, for which the sign of the correlation changes for any value of β . Second, the symmetry $\kappa \leftrightarrow 1/\kappa$ is seen for any value of the repulsion parameter β , and therefore for any degree of localization in the closed model. As a whole, the dependence of the correlations C_1 and C_2 on the coupling κ and on the degree of internal chaos (in our case defined by the repulsion parameter β) qualitatively agrees with that found in Refs. [7] for random matrix models.

VIII. CONCLUSION

We studied the transport properties of the 1D Anderson model in dependence on the degree of internal chaos and strength of the coupling to the continuum. We found that the level spacing distribution $P(s)$ for a closed model of finite size N is well described by the phenomenological expression (3) where the repulsion parameter β changes from $\beta = 0$ to $\beta = \infty$. This expression originates from the two-dimensional Coulomb gas model with the temperature $1/\beta$, and gives the distribution of spacings between the nearest charged particles moving on a ring. This fact may be used for further analytical studies of the spectrum statistics in the finite Anderson model.

In the closed model we established an important linear relation between the parameter β of spectral statistics and the normalized localization length l_∞/N of the eigenfunctions. This result still awaits a rigorous analysis. The important point is that the parameter β can be used to describe the transformation of extended standing waves into localized states, when the degree of disorder is increased. In passing from extended to localized states, our data clearly manifest the Wigner-Dyson distribution occurring in a quite narrow region of the disorder strength, for $\beta \approx 1$.

Opening the system at the ends, we used the effective non-Hermitian Hamiltonian to study the transport properties of the model. For perfect coupling we demonstrated that both the transmission coefficient (conductance) and its variance can be analytically described by the theoretical expressions developed for disordered models with continuous potentials. This fact allows one to fully predict how the mesoscopic fluctuations depend on the degree of internal chaos quantified by the spectral parameter β , or, which is the same, by the normalized localization length. For nonperfect couplings, we have developed an expression for the mean logarithm of the conductance, which is confirmed by our numerical data. Our extensive numerical study of nonperfect coupling reveals specific properties of transport characteristics in dependence on *both* the internal chaos and the coupling strength to the continuum.

Our special interest was in the study of correlations between two cross sections related to the transmission and reflection of scattering waves. The data show that the dependence of these correlations on the degree of disorder, and on the coupling




FIG. 8. (Color online) Correlations C_1 (upper) and C_2 (lower) versus κ for different values of β as indicated.

strength with the continuum, is qualitatively of the same type as found in Ref. [7] for models described by random non-Hermitian matrices. Since our original model (1) is adequate for one-mode waveguides with inserted scatterers [35], one can suggest that the properties of scattering revealed in our study can be experimentally observed.

ACKNOWLEDGMENTS

V.Z. acknowledges the NSF Grants No. PHY-0758099 and No. PHY-1068217, F.M.I. acknowledges support from CONAcYT Grant No. N-161665, and S.S. acknowledges support from the Leverhulme Trust.

-
- [1] C. Mahaux and H. A. Weidenmüller, *Shell Model Approach to Nuclear Reactions* (North-Holland, Amsterdam, 1969).
- [2] J. J. M. Verbaarschot, H. A. Weidenmüller, and M. R. Zirnbauer, *Phys. Rep.* **129**, 367 (1985).
- [3] V. V. Sokolov and V. G. Zelevinsky, *Phys. Lett. B* **202**, 10 (1988); *Nucl. Phys. A* **504**, 562 (1989).
- [4] Y. V. Fyodorov and B. A. Khoruzhenko, *Phys. Rev. Lett.* **83**, 65 (1999); Y. V. Fyodorov and H.-J. Sommers, *J. Phys. A* **36**, 3303 (2003).
- [5] G. L. Celardo, F. M. Izrailev, V. G. Zelevinsky, and G. P. Berman, *Phys. Rev. E* **76**, 031119 (2007); *Phys. Lett. B* **659**, 170 (2008).
- [6] G. L. Celardo, F. M. Izrailev, S. Sorathia, V. G. Zelevinsky, and G. P. Berman, in *Nuclei and Mesoscopic Physics*, edited by P. Danielewicz, P. Piecuch, and V. Zelevinsky, AIP Conf. Proc. No. 995 (AIP, Melville, NY, 2008), p. 75.
- [7] S. Sorathia, F. M. Izrailev, G. L. Celardo, V. G. Zelevinsky, and G. P. Berman, *Europhys. Lett.* **88**, 27003 (2009).
- [8] V. V. Sokolov and V. G. Zelevinsky, *Ann. Phys. (N.Y.)* **216**, 323 (1992).
- [9] A. F. Sadreev and I. Rotter, *J. Phys. A* **36**, 11413 (2003).
- [10] A. Volya and V. Zelevinsky, in *Nuclei and Mesoscopic Physics*, edited by V. Zelevinsky, AIP Conf. Proc. No. 777 (AIP, Melville, NY, 2005), p. 229.
- [11] M. Titov and Y. V. Fyodorov, *Phys. Rev. B* **61**, R2444 (2000); M. Terraneo and I. Guarneri, *Eur. Phys. J. B* **18**, 303 (2000); H. Kunz and B. Shapiro, *J. Phys. A* **39**, 10155 (2006); *Phys. Rev. B* **77**, 054203 (2008).
- [12] G. L. Celardo and L. Kaplan, *Phys. Rev. B* **79**, 155108 (2009); G. L. Celardo, A. M. Smith, S. Sorathia, V. G. Zelevinsky, R. A. Sen'kov, and L. Kaplan, *ibid.* **82**, 165437 (2010).
- [13] D. J. Thouless, in *Ill-Condensed Matter*, edited by R. Balian, R. Maynard, and G. Toulouse (North-Holland, Amsterdam, 1979).
- [14] T. A. Brody, J. Flores, J. B. French, P. A. Mello, A. Pandey, and S. S. M. Wong, *Rev. Mod. Phys.* **53**, 385 (1981).
- [15] C. W. J. Beenakker, *Rev. Mod. Phys.* **69**, 731 (1997); F. M. Izrailev, T. Kottos, A. Politi, and G. P. Tsironis, *Phys. Rev. E* **55**, 4951 (1997); D. Cohen, F. M. Izrailev, and T. Kottos, *Phys. Rev. Lett.* **84**, 2052 (2000).
- [16] Y. V. Fyodorov and A. D. Mirlin, *Int. J. Mod. Phys. B* **8**, 3795 (1994).
- [17] G. Casati, F. Izrailev, and L. Molinari, *J. Phys. A* **24**, 4755 (1991).
- [18] F. M. Izrailev, in *Quantum Chaos*, Proceedings of the International School of Physics "Enrico Fermi," Course CXIX, Varenna, 1991, edited by G. Casati, I. Guarneri, and U. Smilansky (North-Holland, Amsterdam, 1993) p. 265.
- [19] F. J. Dyson, *J. Math. Phys.* **3**, 140 (1962); **3**, 157 (1962); **3**, 166 (1962).
- [20] F. M. Izrailev, *Phys. Lett. A* **134**, 13 (1988).
- [21] B. Wischmann and E. Müller-Hartmann, *Z. Phys. B* **79**, 91 (1990).
- [22] M. P. Sørensen and T. Schneider, *Z. Phys. B* **82**, 115 (1991).
- [23] L. Molinari and V. V. Sokolov, *J. Phys. A* **22**, L999 (1989).
- [24] B. Grammaticos, A. Ramani, and E. Cauries, *J. Phys. A* **23**, 5855 (1990).
- [25] A. V. Vagov and O. K. Vorov, *Phys. Lett. A* **232**, 91 (1997).
- [26] S. Sorathia, Ph.D. thesis, Instituto de Física, BUAP, Puebla, Mexico, 2010.
- [27] F. M. Izrailev, *J. Phys. A* **22**, 865 (1989); *Phys. Rep.* **196**, 299 (1990).
- [28] G. Casati, B. V. Chirikov, I. Guarneri, and F. M. Izrailev, *Phys. Rev. E* **48**, R1613 (1993).
- [29] J. Flores, G. Monsivais, P. Mora, A. Morales, R. A. Méndez-Sánchez, A. Díaz-de-Anda, and L. Gutiérrez, in *Symmetries in Nature: Symposium in Memoriam Marcos Moshinsky*, edited by K. B. Wolf, L. Benet, J. M. Torres, and P. O. Hess, AIP Conf. Proc. No. 1323 (AIP, Melville, NY, 2010), p. 62; J. Flores, L. Gutiérrez, R. A. Méndez-Sánchez, G. Monsivais, P. Mora, and A. Morales, [arXiv:1203.4241v1](https://arxiv.org/abs/1203.4241v1).
- [30] I. M. Lifshits, S. A. Gredeskul, and L. A. Pastur, *Introduction to the Theory of Disordered Systems* (Wiley, New York, 1988).
- [31] F. M. Izrailev, A. A. Krokhin, and N. M. Makarov, *Phys. Rep.* **512**, 125 (2012).
- [32] V. E. Kravtsov and V. I. Yudson, *Ann. Phys. (N.Y.)* **326**, 1672 (2011).
- [33] M. Simonius, *Phys. Lett. B* **52**, 279 (1974).
- [34] A. García-Martín, F. Scheffold, M. Nieto-Vesperinas, and J. J. Sáenz, *Phys. Rev. Lett.* **88**, 143901 (2002).
- [35] U. Kuhl, F. M. Izrailev, A. A. Krokhin, and H.-J. Stöckmann, *Appl. Phys. Lett.* **77**, 633 (2000); A. Krokhin, F. Izrailev, U. Kuhl, H.-J. Stöckmann, and S. E. Ulloa, *Physica E* **13**, 695 (2002); U. Kuhl, F. M. Izrailev, and A. A. Krokhin, *Phys. Rev. Lett.* **100**, 126402 (2008).