

Magnetic moment nonconservation in magnetohydrodynamic turbulence models

S. Dalena,^{1,2} A. Greco,¹ A. F. Rappazzo,² R. L. Mace,³ and W. H. Matthaeus²

¹*Dipartimento di Fisica, Università della Calabria, I-87036 Cosenza, Italy*

²*Bartol Research Institute, Department of Physics and Astronomy, University of Delaware, Newark, Delaware 19716, USA*

³*School of Chemistry and Physics, University of KwaZulu-Natal, Westville Campus, South Africa*

(Received 3 April 2012; published 11 July 2012)

The fundamental assumptions of the adiabatic theory do not apply in the presence of sharp field gradients or in the presence of well-developed magnetohydrodynamic turbulence. For this reason, in such conditions the magnetic moment μ is no longer expected to be constant. This can influence particle acceleration and have considerable implications in many astrophysical problems. Starting with the resonant interaction between ions and a single parallel propagating electromagnetic wave, we derive expressions for the magnetic moment trapping width $\Delta\mu$ (defined as the half peak-to-peak difference in the particle magnetic moments) and the bounce frequency ω_b . We perform test-particle simulations to investigate magnetic moment behavior when resonance overlapping occurs and during the interaction of a ring-beam particle distribution with a broadband slab spectrum. We find that the changes of magnetic moment and changes of pitch angle are related when the level of magnetic fluctuations is low, $\delta B/B_0 = (10^{-3}, 10^{-2})$, where B_0 is the constant and uniform background magnetic field. Stochasticity arises for intermediate fluctuation values and its effect on pitch angle is the isotropization of the distribution function $f(\alpha)$. This is a transient regime during which magnetic moment distribution $f(\mu)$ exhibits a characteristic one-sided long tail and starts to be influenced by the onset of spatial parallel diffusion, i.e., the variance $\langle(\Delta z)^2\rangle$ grows linearly in time as in normal diffusion. With strong fluctuations $f(\alpha)$ becomes completely isotropic, spatial diffusion sets in, and the $f(\mu)$ behavior is closely related to the sampling of the varying magnetic field associated with that spatial diffusion.

DOI: [10.1103/PhysRevE.86.016402](https://doi.org/10.1103/PhysRevE.86.016402)

PACS number(s): 94.05.Pt, 95.30.Qd, 96.50.Ci

I. INTRODUCTION

Magnetic moment μ conservation is an important topic in plasma physics. Many commonly used theories that describe particle motion in perturbed magnetic fields assume that particle magnetic moment is on average constant. When μ is not conserved, the associated effects, such as particle energization, can have a bearing on astrophysical phenomena such as coronal heating, cosmic ray transport, temperature anisotropies in the solar wind [1], and energy release by magnetic reconnection [2]. Furthermore μ conservation is strictly related to particle confinement in plasma machines and dynamically chaotic systems [3]. The conditions that guarantee conservation of μ are well understood in ideal circumstances (e.g., Ref. [4]). However, the various conditions that can lead to significant levels of nonconservation remain the subject of ongoing discussion, especially in turbulent systems, with potentially major impact on understanding plasma dissipation and related topics. Recent studies have clarified some aspects of this problem [2,5].

To provide additional clarification of this issue, in this paper we study magnetic moment conservation for charged particles in the presence of a single electromagnetic wave as well as in the presence of turbulent magnetic fields having one-dimensional spectra comparable to those measured in the solar wind. The goal is to establish the validity range of the adiabatic invariance and the key mechanisms that regulate magnetic moment nonconservation.

The guiding center approximation [4] splits charged particle motion into guiding center motion and the gyromotion around it. When analyzing motion in nonuniform electromagnetic fields, it is desirable to neglect the rapid and relatively uninteresting gyromotion, focusing instead on the far slower

motion of the guiding center. Averaging the particle equation of motion over the gyrophase, we obtain a reduced equation that describes the guiding center motion. In the nonrelativistic case the equation of motion of the guiding center in the direction parallel to the magnetic field reads

$$\frac{dp_{\parallel}}{dt} = -\mu\nabla_{\parallel}B + qE_{\parallel}, \quad (1)$$

where the particle magnetic moment is defined as $\mu = v_{\perp}^2/B$ and $\nabla_{\parallel} = (\hat{\mathbf{B}} \cdot \nabla)$ is the spatial derivative along the field direction. In the perpendicular direction the guiding center drifts with the velocity

$$\mathbf{v}_D = \frac{\mathbf{F} \times \mathbf{B}}{qB^2}, \quad (2)$$

where $\mathbf{F} = [q\mathbf{E} - \mu\nabla B - (mv_{\parallel}^2)\nabla_{\parallel}\mathbf{B}]$ is the total force acting on the guiding center, averaged over a gyroperiod, in the (non-inertial) frame comoving with the guiding center. Therefore, as long as a particle moves through slowly varying electric and magnetic fields, its guiding center behaves like a particle with a magnetic moment μ , which is conserved.

This approximation is valid when the smallest length scales of the electromagnetic fields are much larger than the particle's Larmor radius. This corresponds to the well-known Born-Oppenheimer approximation in quantum mechanics. This description of particle motion in a nonuniform magnetic field is also useful for numerical simulations. Indeed direct simulations of kinetic equations (Vlasov, Boltzmann) with a large magnetic field require the resolution of small spatial and time scales associated with gyration about the magnetic field. This difficulty is alleviated by approaches such as the guiding center approximation, or gyrokinetics, that make

use of μ conservation, to provide approximate models for particle motion in a strong magnetic field. However, in the presence of turbulence the assumption of slow variation of the magnetic field over the particle Larmor radius can break down. Turbulent magnetic fluctuations are observed in space plasmas in practically all environments and at all scales. Furthermore collisionless wave-particle interactions can introduce dissipation processes that act to change the magnetic moment. In such a case the validity of the guiding center theory might be questioned.

When the amplitude of the magnetic fluctuations is lower than that of the mean magnetic field (averaged over the fluctuation time scale), a perturbation approach called the *quasilinear approximation* is applicable [6–8]. In this case the *resonant* fluctuations make the dominant contribution to particle scattering. The resonance condition for wave-particle interaction is given by

$$\omega - k_{\parallel} v_{\parallel} = n\Omega, \tag{3}$$

where ω is the wave frequency, k_{\parallel} and v_{\parallel} are respectively the wave vector and the particle velocity components along the mean magnetic field \mathbf{B}_0 , and $\Omega = qB/m$ is the particle gyrofrequency. Landau resonance [9] is found at $n = 0$, while $n = \pm 1, \pm 2, \dots$ are the cyclotron resonances. In linear theory these resonances are represented by δ functions. In the presence of well-developed magnetohydrodynamic turbulence we expect the discrete resonances to be significantly broadened due to the rapid decorrelation of the wave’s phases in strong turbulence [10–12].

The particle reaction to the perturbation is always periodic except when condition (3) is satisfied. In this case the perpendicular electric force due to the wave remains in phase with the particle cyclotron motion and the particle reaction is *secular* or resonant and, over short times, nonoscillatory. The secular electric force acting on a given particle is constant over a particle gyroperiod, so that the magnetic moment is no longer conserved.

Charged particles are scattered by their interaction with the waves and undergo pitch angle diffusion. The pitch angle $\theta = \arctan(v_{\perp}/v_{\parallel})$ is the angle between the direction of the magnetic field and the particle’s helical trajectory. Scattering from magnetic fluctuations causes the distribution of the pitch angle cosine $\alpha = v_{\parallel}/|v|$ to become isotropic. The familiar formulation of the pitch angle scattering involves computation of a Fokker-Planck coefficient, which under suitable conditions satisfies an adiabatic approximation [6,13,14]. For cosmic ray diffusion in isotropic magnetostatic turbulence [15], the limits of validity of this approximation have been explored.

The magnetic moment μ is formally related to the time averages of the cosine of pitch angle by

$$\mu \sim \frac{v_{\perp}^2}{|\mathbf{B}|} = \frac{v^2}{|\mathbf{B}|}(1 - \alpha^2). \tag{4}$$

We therefore expect the behavior of the magnetic moment to be strongly related to the pitch angle behavior. A comment on the historical connection between magnetic moment changes and pitch angle scattering is in order. The authors of Ref. [16] showed that the Landau resonance [in Eq. (3)] is connected with violation of the first adiabatic invariant. In fact, the general existence of a Dirac δ function in pitch angle scattering

corresponds to the prediction of the quasilinear approximation for the contribution of mirroring to pitch angle scattering. The fact that mirroring exhibits itself as a δ function is a clear indication that this particular pitch angle scattering mechanism is misordered within the quasilinear theory, which assumed the effects of the random field on the motion of particles to be of second order in the random field strength.

Recently, in Ref. [5], plasma heating was studied by examining charged test particles in three-dimensional numerical simulations of weakly compressible magnetohydrodynamic turbulence. This study found systematic changes in the standard deviation of the magnetic moment distribution function for both high-gyrofrequency ($\omega \ll \Omega$) and low-gyrofrequency ($\omega \sim \Omega$) particles. This was attributed to resonance broadening in the nonlinear turbulence. The important issue of parallel vs perpendicular heating was a focus in Ref. [5].

In the present paper the emphasis is not on heating or realistic three-dimensional turbulence spectra, but rather on the relationship between velocity space pitch angle scattering and magnetic moment changes. We are especially interested in how these two distinct (but related) quantities separately approach a diffusive limit. Pitch angle scattering is known to be dominated by parallel resonances, and is therefore sensitive to the bandwidth of the turbulence [17]. To address this issue we therefore adopted more simplified models, but with realistically large bandwidth, to gain a basic understanding of the conditions for the onset of magnetic moment nonconservation.

II. STOCHASTIC MOTION, TRAPPING WIDTH, AND RESONANCE OVERLAPPING

Wave-particle interactions usually involve multiple resonances. Particle motion is substantially different depending on whether these resonances overlap or not. Numerical simulations show a complex behavior that cannot be approached analytically, e.g., it is not possible to write an equation for the evolution of particle distributions when two resonances overlap [18]. Such motions in the presence of overlapping resonances are commonly labeled *stochastic*.

It is important to distinguish between two different kinds of stochasticity. Wave-particle interaction in the presence of uncorrelated small-amplitude electromagnetic waves or plasma turbulence is called *extrinsically diffusive* [19]. In this case the regular phase space structure for a charged particle interacting resonantly with an electromagnetic wave is perturbed by neighboring uncorrelated waves. This leads to *extrinsic stochasticity* and diffusive behavior. On the other hand, nonlinear systems, such as a particle interacting resonantly with a large-amplitude, obliquely propagating (with respect to \mathbf{B}_0) electromagnetic plasma wave, can exhibit *intrinsic stochasticity*. Indeed, when the wave amplitude is sufficiently large, the resonances at the gyrofrequency harmonics are sufficiently broadened that they overlap with adjacent primary resonances. Therefore particles interacting even with a single monochromatic wave may exhibit *intrinsically stochastic* and

diffusive behavior [20]. This is the regime of nonlinear diffusion and irreversible chaotic mixing of orbits.

Because one of the main hypotheses of quasilinear theory is that particles dynamics is adequately modeled by their unperturbed trajectories, the quasilinear time scale τ_c must be much smaller than the time scale for the onset of nonlinear orbit effects τ_{nl} [cf. [21–23]]:

$$\tau_c \ll \tau_{nl} \sim \frac{1}{\omega_b}, \quad (5)$$

where ω_b is the bounce frequency. This means that the turbulent spectrum should be broad enough so that the typical time scale for a charged particle to interact with a resonant wave packet would be much less than its typical bounce time $\tau_b = 2\pi/\omega_b$ in a monochromatic wave at the characteristic wave number and frequency of the wave packet. The bounce time τ_b for a particle in resonance with an electromagnetic wave is proportional to its oscillation period in the pseudopotential well governing the resonant wave-particle interaction [20]. This interaction can be approximated by a pendulum Hamiltonian in the vicinity of the resonance point.

Particles in resonance with a single finite-amplitude fluctuation undergo finite-amplitude and long-period oscillations, which can lead to a variety of interesting nonlinear effects (see, e.g., Refs. [24,25]). This is the so-called trapping width Δv_{\parallel} , given by the half peak-to-peak difference in the particle velocity's parallel component. The trapping width and the bounce frequency for a nonrelativistic particle interacting resonantly with an electromagnetic wave are given by Eqs. (5a)–(5c) of Ref. [25]. These approximate expressions for Δv_{\parallel} and ω_b yield considerable physical insight into the diffusion process [23] when used in conjunction with the quasilinear diffusion coefficient.

III. MAGNETIC MOMENT TRAPPING WIDTH

From the trapping width Δv_{\parallel} and bounce frequency ω_b computed in Ref. [23] for the case of a circularly polarized electromagnetic wave (see the Appendix), it is possible to derive the pitch angle trapping half-width as

$$\Delta\alpha = \frac{\Delta v_{\parallel}}{v} = 2 \left[(1 - \alpha^2)^{1/2} |\alpha| \frac{\delta B}{B_0} \right]^{1/2}. \quad (6)$$

As the magnetic moment μ is related to α by Eq. (4), we can write the trapping width for the magnetic moment as

$$\Delta\mu = 2\alpha\Delta\alpha = 4\alpha \left[(1 - \alpha^2)^{1/2} |\alpha| \frac{\delta B}{B_0} \right]^{1/2}. \quad (7)$$

These expressions apply to a circularly polarized wave. From Eq. (7) we expect that μ continues to be a good adiabatic invariant when resonances are not present or when a particle interacts with extremely small-amplitude waves.

IV. MODEL AND GOVERNING EQUATIONS

We investigate magnetic moment behavior first during the resonant interaction between one ion and a circularly polarized magnetic wave, then when resonance overlapping occurs, and

finally during the interaction between a distribution of particles and a broadband turbulent spectrum. Because some of our normalization quantities are expressed in terms of typical time and length scales of the turbulence slab model [6,17], we first give a general summary of the slab model.

For the general one-dimensional slab description, turbulence is made up of a sum of right- and left-handed circularly polarized nondispersive plane Alfvén waves propagating in the parallel direction. The magnetic field fluctuations are perpendicular to both the wave vector and the mean field. The fields are assumed to be magnetostatic. This amounts to the auxiliary assumption that the average particle speed is well in excess of the phase speed of the underlying linear wave mode. We ignore nonlinear wave-wave couplings in the spirit of quasilinear theory (see, e.g., Ref. [26–28]).

Considering Alfvén waves propagating with $\omega/k = \omega/k_{\parallel} \simeq \pm v_A$, the magnetostatic approximation implies $|\mathbf{v}| \gg v_A$ (strictly $|v_{\parallel}| \gg v_A$). Since particle energy is conserved in a frame moving at the parallel component of the phase velocity of the wave (ω/k_{\parallel}), quasilinear theory [26] implies that

$$(v_{\parallel} - \omega/k_{\parallel})^2 + v_{\perp}^2 = \text{const.}$$

Because of the magnetostatic assumption, particle energy is conserved, i.e., energy diffusion is forbidden and in velocity space the resonant interaction diffuses the pitch angle and gyrophase only. Finally, we ignore all interparticle correlations resulting from their mutual interaction through their microfields (e.g., Coulomb collisions, Debye shielding, and polarization). Furthermore the feedback of the particles on the macroscopic fields is ignored, i.e., we consider only test particles in prescribed macroscopic magnetostatic fields. By virtue of the inequality $|v_{\parallel}| \gg v_A$, the turbulent electric field of the order $(\delta B/B_0)v_A B_0$ is negligible compared to the motional electric field of the particle, $v_{\parallel} B_0$.

The dispersionless hypothesis rules out phase mixing and, hence, phase decorrelation due to this process. Consequently the only way for a particle to experience phase decorrelation of a “wave packet” is to traverse an autocorrelation length of the turbulence [29]. The autocorrelation time in this case is given by [23]

$$\tau_c = \frac{1}{|\Delta(\omega - k_{\parallel}v_{\parallel})|} = \frac{1}{|v_{\parallel}\Delta k_{\parallel}|} \simeq \frac{\lambda_c}{|v_{\parallel}|}, \quad (8)$$

where λ_c is the turbulence correlation length.

The behavior of a test particle is described by its time-dependent position $\mathbf{r}(t)$ and three-dimensional velocity $\mathbf{v}(t)$, which are advanced according to $d\mathbf{r}/dt = \mathbf{v}$ and the Lorentz force equation

$$m \frac{d\mathbf{v}}{dt} = q \left[\mathbf{E} + \frac{\mathbf{v}}{c} \times \mathbf{B} \right]. \quad (9)$$

In order to render the equations nondimensional, we normalize the lengths to l_z , the turbulence coherence length related to the turbulence correlation length λ_c ($\lambda_c = 0.747 l_z$ for our particular slab configuration [30]), velocities to the Alfvén speed v_a , times to the Alfvén crossing time $\tau_A = l_z/v_a$, the magnetic and the electric field to $B_0 = \sqrt{4\pi\rho} v_a$ and $E_n = (v_a/c)B_0 = v_a^2 \sqrt{4\pi\rho}/c$, respectively. For the static case the light speed may also be used as a characteristic quantity [31]. The introduction of an Alfvén speed in our test particle

model, where the waves are treated as static, may appear rather artificial. However, the magnetostatic assumption is valid here provided that $|v_{\parallel}| \gg v_A$ and we introduce v_A in anticipation of future work where we will drop the magnetostatic hypothesis.

With our choice for the characteristic quantities the dimensionless equations of motion of our charged test particles are given by

$$\frac{d\mathbf{r}}{dt} = \mathbf{v}, \quad (10)$$

$$\frac{d\mathbf{v}}{dt} = \beta(\mathbf{E} + \mathbf{v} \times \mathbf{B}). \quad (11)$$

Here $\beta = \Omega\tau_A$ (cf. the α parameter in Ref. [32]) couples particle and field spatial and temporal scales and provides a particularly useful means to relate our numerical experiments to space and astrophysical plasmas. In general, in a turbulent collisionless plasma the bandwidth of the inertial range fluctuations may extend from large fluctuations at the correlation scale, λ_c , to small fluctuations at the ion inertial scale. In this case $\beta \gg 1$ and the turbulent time scales are much slower than the typical particle gyroradius [33].

The resonant condition for the static case in terms of β is given by

$$k_{\text{res}}\lambda = \frac{n\beta}{\alpha(v/v_A)} = \frac{n\beta}{(v_{\parallel}/v_A)}. \quad (12)$$

Time is advanced through a fourth-order Runge-Kutta integration method with an adaptive time step (pp. 708–716 of Ref. [34]).

V. NUMERICAL SIMULATIONS

Particles are loaded randomly in space at $t = 0$ throughout a one-dimensional simulation box of length L . The fields are described in the following sections. In spherical coordinates, with the polar axis along the z direction parallel to the mean magnetic field of strength B_0 , the particle velocity components are

$$v_x = v \sin \theta \cos \phi, \quad v_y = v \sin \theta \sin \phi, \quad v_z = v \cos \theta. \quad (13)$$

The particles' initial velocities are randomly distributed in the gyrophase ϕ in the range $[0, 2\pi]$, while the velocity magnitude v and pitch angle θ are determined by the particular numerical experiment.

Typical particle velocities used in our simulations are $10v_A$ and $100v_A$, satisfying the magnetostatic constraint. In our analysis magnetic moments are expressed in units of the characteristic quantity $\mu_n = v^2/B_0$. We also define $\delta b = \delta B/B_0$.

The statistic analysis of the particle magnetic moment involves averaging trajectories over the particle gyroperiod $\tau_g = 2\pi/\Omega$. For each simulation we compute the effective number of gyroperiods N_{τ_g} that particles complete in a given magnetic field configuration as

$$N_{\tau_g} = \int_0^t \frac{dt}{2\pi} \frac{eB(t)}{mc}, \quad (14)$$

where $B(t)$ is the intensity of the total magnetic field. When $\delta b \ll 1$, $B(t) \simeq B_0$; however, as δb increases toward unity the

wave's contribution to the strength of the total magnetic field $B(t)$ is not negligible.

A. Single wave

We start studying the ion motion in the presence of a constant magnetic field \mathbf{B}_0 and a perpendicular left-handed circularly polarized wave with

$$\mathbf{B} = \delta B_x \cos(k_0 z) \hat{\mathbf{e}}_x - \delta B_y \sin(k_0 z) \hat{\mathbf{e}}_y + B_0 \hat{\mathbf{e}}_z, \quad (15)$$

where δB_x and δB_y are the amplitudes of the wave and k_0 is the wave vector. We assume $\delta B_x = \delta B_y = \delta B$ for the rms average values. In these simulations $\beta = 10^3$, $v = 100v_A$, and $\alpha = 0.125$ ($\theta = 82^\circ$).

We follow the test particles until they complete $N_{\tau_g} = 100$ gyroperiods. For the resonance condition, Eq. (12), we set $k_0 = 80/\lambda$. Particles injected with a pitch angle cosine different from $\alpha = 0.125$ will not be in resonance with this wave, exhibiting a different behavior. For a direct comparison we also inject nonresonant particles, i.e., with $\alpha = 0.5$ ($\theta = 60^\circ$).

Figure 1 shows the time evolution of the cosine of the pitch angle α , the particle magnetic moment μ , and the parallel component of the induced electric field E_z , for a resonant particle ($\alpha = 0.125$). Different columns corresponds to different values of the wave amplitude: $\delta b = 0.001$ (first column), $\delta b = 0.01$ (second column), $\delta b = 0.1$ (third column), and $\delta b = 1.0$ (fourth column).

When the parallel component of the induced electric field is almost constant and equal to $E_z \sim -v_{\perp} \delta b$, the resonant interaction produces variations that are secular over a gyroperiod. However, an oscillation occurs over a longer time, the bounce period $\tau_b = 2\pi/\omega_b$ (where ω_b is the bounce frequency discussed in Sec. II). This is the typical time scale over which the velocity, and hence the particle trajectory, exhibits significant deviations from the linear $v_{\parallel} = \text{const}$ and $v_{\perp} = \text{const}$ case.

In Sec. III we derived the analytical expression for the half trapping width of the magnetic moment for a particle interacting with a left- or right-handed circularly polarized wave [see Eq. (7)]. We now compute the values of the half peak-to-peak difference in α and μ , $\Delta\alpha = (\alpha_{\text{max}} - \alpha_{\text{min}})/2$ and $\Delta\mu = (\mu_{\text{max}} - \mu_{\text{min}})/2$, for the resonant interaction simulations. These values and those obtained from the theoretical expressions (6) and (7) are listed in Table I and are in good agreement, confirming the validity of Eqs. (6) and (7) and reinforcing the intuitive idea that the magnetic moment and pitch angle behaviors are strictly related.

To compare resonant and nonresonant dynamics, we show in Fig. 2 the time evolution of the cosine of the pitch angle α (first row) and the magnetic moment μ (third row), and their distribution functions $f(\alpha)$ (second row) and $f(\mu)$ (fourth row) at the end of the simulation, for a *resonant particle* with $\alpha = 0.125$ (left column), and a *nonresonant* one with $\alpha = 0.5$ (right column). In contrast with the resonant case in which α and μ exhibit well-known secular variations with typical period equal to τ_b , the α and μ profiles for a nonresonant particle show a regular oscillating behavior, a distinctive signature of regular particle motion. The values of the half peak-to-peak difference in α and μ obtained from the simulation are $\Delta\alpha_{\text{sim}} = 0.0025$ and $\Delta\mu_{\text{sim}} = 0.003$. These




FIG. 1. Gyroresonant interaction between a circularly polarized wave and a particle with $v = 100v_A$ and $\alpha = 1/8$: cosine of pitch angle α (top row), particle magnetic moment μ (middle row), and parallel component of the induced electric field (bottom row). Different columns correspond to different wave amplitudes: $\delta b = 0.001$ (first column), $\delta b = 0.01$ (second column), $\delta b = 0.1$ (third column), and $\delta b = 1.0$ (fourth column).

are smaller than the theoretical values computed from Eqs. (6) and (7) with $\delta b = 0.01$ and $\alpha = 0.5$, for which we obtain $\Delta\alpha_{\text{th}} = 0.1316$ and $\Delta\mu_{\text{th}} = 0.1316$.

The distribution functions $f(\alpha)$ and $f(\mu)$ (Fig. 2) for a resonant particle are more spread in α and μ and are centered around their initial values $\alpha = 0.125$ and $\mu = 0.98$. In the nonresonant case, $f(\mu)$ remains peaked at its initial value, i.e., its magnetic moment is constant during particle motion. The spread in α of its distribution is $\sim 10\%$, small compared to the resonant case spreading of $\sim 40\%$.

Figure 3 shows the distribution functions $f(\alpha)$ and $f(\mu)$ at the end of the simulation for 1000 resonant and nonresonant particles injected in the simulation box with random positions and phases. For nonresonant particles (right column) the distributions remain peaked around their initial values $\alpha = 0.5$ and $\mu/\mu_n = 0.75$ with very little spreading. For the resonant particles (left column) $f(\alpha)$ acquires a Gaussian shape centered around its initial value $\alpha = 0.125$. Furthermore it spreads by ~ 0.1 , comparable to the trapping width for the

single particle $2\Delta\alpha = 0.014$ (Fig. 2). The magnetic moment distribution for the resonant case has a characteristic shape found for μ in the parameter range in which the pitch angle exhibits a Gaussian distribution and the density distribution function is still isotropic (the particle free-streaming regime). As for the pitch angle, the spread in the magnetic moment distribution of ~ 0.03 is comparable to the trapping width for the single particle $2\Delta\mu = 0.00352$ [see Eq. (7) and Fig. 2].

B. Overlapping resonances

In order to understand the effect of overlapping resonances on particle magnetic moment, we perform a numerical experiment with four different particles in the simulation box with random initial positions, and the same initial velocity $v = 100v_A$, but different values for the pitch angle cosine: $\alpha_1 = 1/2$, $\alpha_2 = 1/4$, $\alpha_3 = 1/8$, $\alpha_4 = 1/32$. For $\beta = 10^3$, making use of the resonance condition for the static case [Eq. (12)], the cyclotron resonances $n = 1$ for the different values of α are expected for $k_1\lambda = 20$, $k_2\lambda = 40$, $k_3\lambda = 80$, and $k_4\lambda = 320$.

The total magnetic field is given by

$$\mathbf{B} = B_0\hat{\mathbf{e}}_z + \sum_{i=1}^4 \delta b \cos[k_i z + \phi_i]\hat{\mathbf{e}}_x - \sum_{i=1}^4 \delta b \sin[k_i z + \phi_i]\hat{\mathbf{e}}_y, \quad (16)$$

where the ϕ_i are random phases. Taking into account resonance broadening effects, all particles with parallel velocities in the range

$$v_{\parallel} - \Delta v_{\parallel} < v_{\parallel} < v_{\parallel} + \Delta v_{\parallel} \quad (17)$$

TABLE I. Trapping width values for α and μ : comparison between theoretical (subscript th) and numerical (subscript sim) values.

δb	$\Delta\alpha_{\text{th}}$	$\Delta\alpha_{\text{sim}}$	$\Delta\mu_{\text{th}}$	$\Delta\mu_{\text{sim}}$
0.001	0.022	0.02	0.0056	0.0055
0.01	0.07	0.075	0.0176	0.02
0.1	0.2227	0.2	0.0556	0.055
1.0	0.704	0.6	0.176	0.175




FIG. 2. (Color online) Time evolution of the cosine of the pitch angle α (first row) and its distribution function $f(\alpha)$ (second row), and time evolution of the magnetic moment μ (third row) and its distribution function $f(\mu)$ (fourth row) of a resonant ($\alpha = 0.125$, left column) and a nonresonant particle ($\alpha = 0.5$, right column). $v = 100 v_A$.

can potentially resonate with a wave, whose wave number is $k_{\parallel} = \Omega/v_{\parallel}$. As found in Ref. [35], the direct evidence




FIG. 3. (Color online) $f(\alpha)$ (first row) and $f(\mu)$ (second row) at the end of the simulation for an initial distribution of 1000 resonant (left column) and nonresonant (right column) particles randomly distributed in the simulation box. $\delta b = 0.01$.

TABLE II. Values of Δv_{\parallel} for $\alpha = 1/2$, $1/4$, $1/8$, and $1/32$ resonances at different δb .

δb	$\alpha = 1/2$	$\alpha = 1/4$	$\alpha = 1/8$	$\alpha = 1/32$
0.001	4.16	3.1	2.227	1.3
0.01	13.1	9.85	7.042	3.583
0.1	41	31.1	22.27	11.33
1.0	131	98.3	70.42	35.83

of resonances overlapping is the disappearance of constants of motion, i.e., the onset of stochasticity in the Hamiltonian formalism. We undertook simulations with four different wave amplitudes $\delta b = 0.001$, 0.01 , 0.1 , and 1.0 . The values of the trapping half-widths Δv_{\parallel} computed for the different pitch angles with Eq. (A8) are listed in Table II for the different δb considered.

Figure 4 shows time histories of the pitch angle cosine α (left column) and magnetic moment μ (right column) profiles for various δb . Again similar behavior is seen for α and μ . For the smallest wave amplitude, $\delta b = 0.001$ (first row), it is possible to recognize very well the four different resonances in the profiles of α and μ . For $\delta b = 0.01$ (second row) the resonance at $\alpha_3 = 1/8$ is overlapping with the resonance at $\alpha_4 = 1/32$. Indeed, the initial parallel velocity of the particle injected at the smallest pitch angle, $v_{\parallel,4} = 3.125 v_A$, lies in the range of velocities [see Eq. (17)] in possible resonance with




FIG. 4. (Color online) Transition from nonoverlapping to overlapping resonances: α (left column) and μ (right column) profiles as the wave amplitude is varied: $\delta b = 0.001$ (first row), $\delta b = 0.01$ (second row), $\delta b = 0.1$ (third row), and $\delta b = 1.0$ (fourth row).




FIG. 5. (Color online) Transition from nonoverlapping to overlapping resonances with varying wave amplitude: $\delta b = 0.001$ (first row), $\delta b = 0.01$ (second row), $\delta b = 0.1$ (third row), and $\delta b = 1.0$ (fourth row). The distribution functions $f(\alpha)$ (left column), $f(\mu)$ (central column), and $f(\delta z)$ (right column) are averaged over time.

$k_{\parallel} = k_3$. For higher wave amplitudes, $\delta b = 0.1$ (third row) and $\delta b = 1.0$ (fourth row), the condition (17) is satisfied by all particle velocities. Stochasticity arises and the different resonances are indistinguishable.

The distribution functions (Fig. 5) $f(\alpha)$ (left column), $f(\mu)$ (central column), and $f(\delta z)$ (right column) (where $\delta z = z - z_0$ is the displacement along z relative to the particle initial position z_0) after $100\tau_g$ exhibit similar characteristics.

For $\delta b = 0.001$ (first row), $f(\alpha)$ and $f(\mu)$ are peaked in correspondence with their four initial values because of the good resonance separation. $f(\delta z)$ shows that the particles are simply free streaming in the parallel direction and, depending on their initial parallel velocity, they cover shorter or longer distances along z .

For $\delta b = 0.01$ (second row), $f(\alpha)$ spreads around its initial four peaks because the particles interact resonantly with waves of larger amplitude, and the resonances overlap for $\alpha < 1/4$, as discussed previously. Similar effects are shown also by $f(\mu)$, confirming that for small δb the resonant interaction affects the magnetic moment and pitch angle in similar ways.

While for $\delta b = 0.01$ particles continue to free stream in the z direction, different profiles for $f(\delta z)$ appear for $\delta b = 0.1$ (third row). The pitch angle distribution begins to isotropize and the magnetic moment exhibits a one-sided long tail distribution extending toward smaller μ . This behavior is similar to the regime found previously in the single-wave experiment when $f(\alpha)$ is nearly isotropic, $f(\delta z)$ still indicates particles free streaming, and the magnetic moment distribution displays a long tail.

For $\delta b = 1.0$ (fourth row), by $t = 100\tau_g$ the pitch angle cosine distribution $f(\alpha)$ has become completely isotropic, while $f(\delta z)$ approaches a Gaussian distribution indicative




FIG. 6. (Color online) Power spectrum of the turbulent magnetic field. k is normalized to the coherence length l_z .

of spatial diffusion. In this regime $f(\mu)$ loses its long tail and starts to acquire a Gaussian shape. In that way we have identified three distinct regimes of statistical magnetic moment behavior with increasing degree of turbulence.

VI. SLAB SPECTRUM

In this section we present the results of our numerical simulations of test particles in the presence of a broadband slab spectrum [see Eq. (19) and Fig. 6]. We have performed simulations for different particle velocities and amplitudes of the magnetic field fluctuations.

The simulations use a unidimensional computational box of length $L = 10000 l_z$ ($l_z = 1$ is the coherence scale for the slab spectrum) with $N_z = 2^{28} = 268\,435\,456$ grid points [36]. The magnetic field in physical space is generated from a spectrum $P(k)$ in Fourier space, via inverse fast Fourier transform. The turbulent magnetic field is given by

$$\mathbf{B}(z) = B_0 \mathbf{e}_z + \delta \mathbf{B}(z), \quad (18)$$

with $\delta \mathbf{B}(z) = \delta B_x(z) \hat{\mathbf{e}}_x + \delta B_y(z) \hat{\mathbf{e}}_y$, and the solenoidality condition is identically satisfied.

The modes of the magnetic field components in k space are given by

$$\begin{aligned} \delta B_x(k_n) &= [P(k_n)]^{1/2} e^{i\Phi_n}, \\ \delta B_y(k_n) &= [P(k_n)]^{1/2} e^{i\Psi_n}, \end{aligned}$$

where $k_n = 2\pi n/L$ and Φ_n and Ψ_n are random phases. The slab spectrum $P(k)$ is given by

$$P(k_n) = \begin{cases} C_{\text{slab}} [1 + (k_n l_z)^2]^{-5/6} & \text{for } k_n < k_{\text{diss}}, \\ C_{\text{diss}} \left(\frac{k_n}{k_{\text{diss}}}\right)^{-7/3} & \text{for } k_n \geq k_{\text{diss}}, \end{cases} \quad (19)$$

where $C_{\text{slab}} = 2\lambda_c \delta b_{x,\text{slab}}^2$ is a constant specific to this form of the slab model, $\delta b_{x,\text{slab}}^2$ is the mean square fluctuation, k_{diss} is the dissipation range wave number, and $C_{\text{diss}} = C_{\text{slab}} [1 + (k_{\text{diss}} l_z)^2]^{-5/6}$ is the constant for the dissipation range (set by the continuity of the spectrum $P(k)$ at k_{diss}). The vectors of Fourier coefficients are zero padded for $N_{\text{max}} + 1 \leq n \leq N_z$ providing an extra level of smoothness to the fields by an effective trigonometric interpolation [30]. In all the simulations we use $N_{\text{max}} = 6.7 \times 10^7$ and a simple

TABLE III. Characteristic scales in the spectrum.

Wave number index	Wave number value
$N_{k_{\min}} = 1$	$k_{\min} = 6.28 \times 10^{-4}$
$N_{kl_z} = 10^4$	$k_{l_z} = 2\pi$
$N_{k_{\text{diss}}} = 1.6 \times 10^6$	$k_{\text{diss}} = 10^3$
$N_{k_{\text{max}}} = 6.7 \times 10^7$	$k_{\text{max}} = 4.2 \times 10^4$
$N_{k_{\text{max}1}} = 1.3 \times 10^8$	$k_{\text{max}1} = 8.4 \times 10^4$

linear interpolation to compute the fields at the test particle position.

The resulting spectrum is shown in Fig. 6. Several important scales are present in the system. They are labeled as k_{\min} , k_{l_z} , k_{diss} , k_{max} , and k_{N_z} . The discrete wave numbers are obtained through $k_n = 2\pi n/L$ as

$$N_k = \frac{L}{2\pi} k \sim 1600k. \quad (20)$$

We summarize the values for k and N_k used in our simulations in Table III, where:

(a) $k_{\min} = 2\pi/L$ is the minimum wave vector of the spectrum, corresponding to $N_k = N_{k_{\min}} = 1$.

(b) $k_{l_z} = 2\pi/l_z = 2\pi$ is the wave vector that marks the beginning of the *inertial range*. Three decades of *energy-containing range* from k_{\min} to k_{l_z} ensure turbulence homogeneity. l_z or $\lambda_c = 0.747l_z$ (see Ref. [30]) correspond to the typical length scales over which the particles attain diffusive behavior of the pitch angle. Three decades of inertial range with $P(k) \propto k^{-5/3}$ well represent solar wind conditions.

(c) k_{diss} is the wave vector corresponding to the beginning of the *dissipation range*. In our model, the spectrum extends beyond k_{diss} with $P(k) \propto k^{-7/3}$.

(d) At two decades higher wave number, $k_{\text{max}} = \sqrt{m_i/m_e} k_{\text{diss}}$ determines the end of the dissipation range.

(e) Extending for two decades beyond $k_{\text{max}} = 4.2 \times 10^4$, the spectrum includes zero padding up to $k_{\text{max}1} = 8.4 \times 10^4$.

(f) Another important scale, not labeled in Fig. 6 because it depends on test particle velocity, is the wave vector corresponding to $z_{\text{max}} = vT_{\text{tot}}$, the distance covered by a charged test particle moving at speed v in the simulation running time T_{tot} . To avoid periodicity effects it is important that the box length L is large enough so that particle trajectories are limited to a small fraction of the full length, i.e., $L \gg z_{\text{max}}$ or $k_{\min} \ll 1/z_{\text{max}}$. Periodicity might indeed give rise to artificial field line diffusion.

We fix the value of the β parameter [see Eq. (10)] equal to 10^4 . This corresponds approximately to observed solar wind turbulence properties at 1 AU, as follows:

$$\beta = \Omega\tau_A = \left(\frac{q}{m}\right) \frac{\lambda_c \sqrt{4\pi\rho}}{c} = \frac{\omega_{pi}\lambda_c}{c} = \frac{\lambda_c}{\lambda_{ii}}, \quad (21)$$

where λ_c is the turbulence correlation length, $\omega_{pi} = (4\pi n_{oi} q_i^2 / m_i)^{1/2}$ is the ion plasma frequency (q_i and m_i are respectively the ion charge and mass), and $\lambda_{ii} = c/\omega_{pi} = (c^2 m_i / 4\pi n_i e^2)^{1/2}$ is the ion inertial length. For $n_i = n_e$ then $\lambda_{ii} = (m_i/m_e)^{1/2} \rho_{ie}$. Because the solar wind density at 1 AU is

TABLE IV. Typical values used in the simulations.

V (units of v_A)	r_L (units of l_z)	k_{res} (units of $1/l_z$)	ϵ	t_c (units of τ_A)
10	10^{-3}	8×10^3	1.33×10^{-3}	0.0747
100	10^{-2}	8×10^2	1.33×10^{-2}	0.00747

approximately $n \sim (1, 10) \text{ cm}^{-3}$ on average $\lambda_{ii} \sim 1000 \text{ km}$. At the same distance the turbulence correlation length λ_c is approximately 10^6 km [37] and $\beta \simeq 10^4$.

Typically 1000 particles are injected in the simulation with random initial positions. Particles are loaded from a cold ring-beam [see Eqs. (13)] distribution with constant velocity magnitude, $\sin\theta$ is set equal to $(1 - \alpha_0^2)^{1/2}$, where α_0 is the initial pitch angle cosine with respect to the background field B_0 . The initial gyrophase ϕ is chosen randomly. For all the simulations $\alpha_0 = 0.125$ ($\theta \simeq 82^\circ$).

From the previous section, we know that the behavior of the magnetic moment is related to the pitch angle behavior for a low level of magnetic fluctuation ($\delta b = 0.001, 0.01$). The pitch angle and magnetic moment exhibit Gaussian distribution functions typical of normal diffusion processes. With increase in the turbulence level, the pitch angle distribution approaches isotropy and a transient regime is observed with the magnetic moment starting to be influenced by the onset of spatial parallel diffusion. When $f(\alpha)$ becomes completely isotropic, spatial diffusion sets in and the $f(\mu)$ behavior is closely related to the sampling of the varying magnetic field strength associated with that spatial diffusion.

From quasilinear theory we know that velocity and real space diffusion occur at two different time scales. Typically, velocity space diffusion takes place with the time scale $\tau_c = \lambda_c/v$ shorter than the typical time scale at which parallel diffusion occurs $\tau_{\parallel} = \lambda_{\parallel}/v$, where $\lambda_{\parallel} = 3D_{\parallel}/v$ is the parallel mean free path. For this reason we follow test particles in the simulation box for a time $T > \tau_c$, typically with $T = 20\tau_c$. Particle parameters used in the simulations are listed in Table IV.

An important parameter in the description of energetic test particles is $\epsilon = r_L/\lambda_c$, which is sometimes called the dimensionless particle rigidity. It can be related to the bendover wave number of the turbulence, $k_{bo} = 1/\lambda_c$, and the minimum resonant wave number, $k_{\min}^r = 1/r_L$, as $\epsilon = k_{bo}/k_{\min}^r$. For example, when $r_L \gg \lambda_c$ the particles experience all possible k modes in few gyroperiods resonating with the energy-containing scale ($k_{\min}^r \ll k_{bo}$). For lower energies the test particles resonate in the inertial range. Those with $v = 10v_A$ will resonate at the end of the inertial range ($1/r_1$ in Fig. 6), while those with $v = 100v_A$ at the middle of the inertial range ($1/r_2$ in Fig. 6). Furthermore, as explained previously, the condition $k_{\min} \ll z_{\text{max}}$ is necessary to avoid artificial effects in particle transport associated with periodicity of the magnetic field.

Figure 7 shows $f(\alpha)$ (left column), $f(\mu)$ (central column), and $f(\delta z)$ (right column) for a distribution of particles moving with an initial velocity $v = 10v_A$ in the presence of the slab spectrum [Eq. (19), Fig. 6]. Different rows correspond to different values of δb (0.001 first row, 0.01 second row, 0.1




FIG. 7. (Color online) Distribution functions of the cosine of the pitch angle $f(\alpha)$ (left column), the magnetic moment $f(\mu)$ (central column), and the particle displacements relative to the initial position $f(\delta z)$ (right column) at $20\tau_c$ for different values of δb : $\delta b = 0.001$ (first row), $\delta b = 0.01$ (second row), $\delta b = 0.1$ (third row), and $\delta b = 1.0$ (fourth row). The blue and the green (light gray) lines indicate the initial value and the mean value of each distribution. Particle parameters at injection: $v = 10 v_A$ and $\alpha_0 = 0.125$.

third row, and 1.0 fourth row). All the distribution functions are computed at the end of the simulation, i.e., after $20\tau_c$. The blue line and the green (light gray) line indicate the initial value and the mean value of each distribution. As particles are injected at different positions, it is convenient to define the quantity $\delta z = z(j) - z(0)$ (j is a temporal index). In this way it is possible to take out from the distribution function $f(\delta z)$ both the drift effect ($v_D t$) and the particle diffusion relative to their own positions (Δz_i). The general expression for the z position of the i th particle is given by

$$z_i = z_i(0) + v_D t + \Delta z_i = z_i(0) + \delta z_i. \quad (22)$$

The primary diagnostic for studying particle diffusion is the variance $\sigma^2(t) \propto t^s$ of the particle's cosine of pitch angle, magnetic moment, and position parallel to the mean

field direction. Figure 8 illustrates the time evolution of the variances $\langle(\Delta\alpha)^2\rangle$ [Fig. 8(a)], $\langle(\Delta\mu)^2\rangle$ [Fig. 8(b)], and $\langle(\Delta z)^2\rangle$ [Fig. 8(c)] for a particle distribution moving with initial velocity equal to $10v_A$. Different colors correspond to different δb values: $\delta b = 0.001$ black line (1), $\delta b = 0.01$ purple line (2), $\delta b = 0.1$ red line (3), and $\delta b = 1.0$ blue line (4). The variances are fitted with the gray lines: the dotted line is used for $s = 2$, the dot-dashed line for $s = 1$, the dashed line for $s = 0.8$, and the three-dot-dashed line for $s = 0.7$.

For $\delta b = 0.001$, α and μ display Gaussian distributions while particles free stream in the z direction. Particles that cover greater distance in z are more scattered in pitch angle and consequently in μ . Figure 8 shows superdiffusive behavior (black line, $s = 2$) with particles free streaming along z , and later, variance characteristic of normal diffusion with $\langle(\Delta\alpha)^2\rangle$ and $\langle(\Delta\mu)^2\rangle$ scaling $\propto t$.

For $\delta b = 0.01$, the particles cover only one side of the α hemisphere, continuing to travel along z [purple line (2) in Fig. 8]. This is the transient regime already observed in Fig. 5 when $f(\mu)$ exhibits a one-sided long tail distribution toward smaller μ .

For $\delta b = 0.1$, the pitch angle distribution becomes completely isotropic and spatial diffusion sets in, as shown by the slope $s = 1$ of $\langle(\Delta z)^2\rangle$ in Fig. 8 [red line (3)] at the end of the simulation. The deviation from purely free-streaming or ballistic behavior means that, while the system has not become fully diffusive along the mean field direction, there are signs that diffusive processes in velocity space are beginning to diminish the free streaming. Although $f(\mu)$ still exhibits a long tail, the influence of spatial diffusion starts to appear. The well-pronounced peak observed in $f(\mu)$ for $\delta b = 0.01$ is substantially reduced and the mean value of the magnetic moment decreases. Moreover μ displays subdiffusive behavior up to $0.02\tau_c$. After this time particles diffuse in space and $\langle(\Delta\mu)^2\rangle$ attains a plateau. The Gaussian shape is not reached yet, probably because spatial diffusion is just beginning.

For $\delta b = 0.5$ (see Fig. 9) and $\delta b = 1.0$, $f(\alpha)$ is isotropic, particle motion is completely diffusive in real space, as the slope $s = 1$ in $\langle(\Delta z)^2\rangle$ in Fig. 8 [blue line (4)] shows, and the $f(\mu)$ behavior is closely related to the sampling of varying




FIG. 8. (Color online) Statistics for $v = 10v_A$. Variances for the cosine of the pitch angle α (a), magnetic moment μ (b), and particle displacements relative to the initial position δz (c) at different values of δb : $\delta b = 0.001$ [black (1) line], $\delta b = 0.01$ [purple (2) line], $\delta b = 0.1$ [red (3) line], and $\delta b = 1.0$ [blue (4) line]. The s values indicate the different slopes for the scalings $\langle(\Delta\alpha)^2\rangle$, $\langle(\Delta\mu)^2\rangle$, and $\langle(\Delta z)^2\rangle \propto t^s$. The variances are fitted with the gray lines: the dotted line is $s = 2$, dot-dashed line $s = 1$, dashed line $s = 0.8$, and three-dot-dashed line $s = 0.7$.




FIG. 9. (Color online) Statistics for $v = 10v_A$ and $\delta b = 0.5$. Magnetic moment variance $\langle(\Delta\mu)^2\rangle$ (first plot) and mean square displacement $\langle(\Delta z)^2\rangle$ (second plot) versus time, and magnetic moment distribution function $f(\mu)$ after $0.01\tau_c$ (third plot) and at the end of the simulation (fourth plot). The s values indicate the different slopes for the scaling of the variances $\langle(\Delta\mu)^2\rangle$ and $\langle(\Delta z)^2\rangle \propto t^s$. The variances are fitted with the red dot-dashed lines and the blue dotted lines.

magnetic field strength associated with that spatial diffusion, displaying a Gaussian distribution centered at the middle of μ space.

From a more detailed analysis of the case $\delta b = 0.5$ (Fig. 9) we notice that the magnetic moment variance (first figure) scales according to $\langle(\Delta\mu)^2\rangle \propto t^{0.17}$ (red dot-dashed line) up to $0.002\tau_c$, and after $0.005\tau_c$ a plateau is attained (blue dotted line). Instead, the particle motion [see the time evolution of $\langle(\Delta z)^2\rangle$, second plot] becomes fully diffusive (blue dotted line) only after $0.007\tau_c$. The magnetic moment distribution $f(\mu)$ in the first part of the evolution (third plot) is in the transient regime characterized by the long tail. In contrast, when the particles diffuse in real space (fourth plot), $f(\mu)$ reacquires the Gaussian profile. Thus the final stage of the magnetic moment variance evolution, i.e., the formation of the plateau, can be considered as a precursor for the onset of the parallel diffusion of particles in space. Of course this effect is present in the pitch angle variance too, but in addition in the μ behavior we have a direct signature of the onset of the spatial diffusion, that is, the reappearance of the Gaussian shape in the distribution function, while the pitch angle distribution remains completely isotropic.

Thus these transitions in magnetic moment behavior are related not just to the variation of the turbulence level, but also to the different time scale at which magnetic moment conservation is studied.

The magnetic moment distribution functions $f(\mu)$ and variances $\langle(\Delta\mu)^2\rangle$ in the case $v = 100v_A$ (not shown) exhibit the same features observed for $v = 10v_A$. However, with increasing particle speed the total number of gyroperiods, N_{τ_g} , performed by each particle decreases; as a consequence,

faster particles sample less variation in magnetic field strength. This leads to a slower spatial diffusion, i.e., for $100v_A$ spatial diffusion occurs on a time scale longer than $20\tau_c$.

For $v = 100v_A$ we show in Fig. 10 the magnetic moment standard deviation σ_μ/μ_{\min} (blue triangle, left plot) and the changes in its mean $\Delta\mu/\mu_{in} = (\bar{\mu} - \mu_{in})/\mu_{in}$ (red circle, right plot) versus δb after $20\tau_c$. As δb increases toward unity the changes in magnetic moment distribution start to increase more quickly.

VII. CONCLUSIONS

In this paper we have investigated the conservation of a charged particle magnetic moment in the presence of turbulent magnetic fields. For slow spatial and temporal variations of the magnetic field with respect to the particle gyroradius and gyroperiod, the magnetic moment μ is an adiabatic invariant of the particle motion. Nonconservation of the magnetic moment can influence the particle acceleration and has considerable implications in many astrophysical problems such as coronal heating, cosmic ray transport, and temperature anisotropies in the solar wind. These applications motivate the present basic study of the degree to which magnetic moments are conserved in increasingly complex models of one-dimensional spectra. While all the models considered here have been very oversimplified relative to the spectra observed for example in the solar wind [17,38] or in simulations of magnetohydrodynamic turbulence [5,39], the present study is intended to contribute to the basic understanding of the conditions for the onset of magnetic moment nonconservation. We point the interested reader also to a recent study in Ref. [5] that addresses this issue from a somewhat different perspective.

In order to reproduce and extend some of the results obtained in Ref. [25], we started to study the *resonant interaction between ions and a single parallel propagating electromagnetic wave* (see Sec. V A). Using the specialized expression for the trapping width Δv_{\parallel} found by Ref. [23] in the case of a single circularly polarized wave, we have been able to write a similar expression for magnetic moment [see Eq. (7)]. In the presence of a single finite-amplitude fluctuation the magnetic moment of a resonant particle undergoes a finite-amplitude nonlinear oscillation too. We have performed several simulations, changing both the particle velocity and the amplitude of the wave. For each of them we compare the values of $\Delta\mu$ and Δv_{\parallel} with those




FIG. 10. (Color online) Standard deviation σ_μ/μ_{\min} (blue triangle, left plot) and variation in magnetic moment $\Delta\mu/\mu_{in}$ (red circle, right plot) versus δb at $20\tau_c$ for $v = 100v_A$ and $\alpha = 0.125$.

obtained using our specialized expression, and they are in good agreement.

We designed a particular experiment to study the *effects of resonances overlapping* (see Sec. VB). From the analysis of the distribution functions of the particle pitch angle, $f(\alpha)$, magnetic moment, $f(\mu)$, and z position, $f(z)$, we distinguish three different regimes. First, for a low level of magnetic fluctuation, i.e., $\delta B/B_0 = 0.001, 0.01$, the magnetic moment distribution half-width is directly related to the pitch angle distribution. Second, for $\delta B/B_0 = 0.1$ stochasticity arises as a consequence of overlapping resonances and its effect on the pitch angle is isotropization of the distribution function. This is a transient regime during which the magnetic moment exhibits a one-sided long tail distribution and starts to be influenced by the onset of spatial parallel diffusion. Finally, when $f(\alpha)$ becomes completely isotropic, spatial diffusion sets in ($\delta B/B_0 = 0.1$), the $f(\mu)$ behavior is closely related to the sampling of varying magnetic field strength associated with that spatial diffusion.

Other studies regarding particle interaction with two electromagnetic waves as well as a flat turbulent spectrum (not shown) were also conducted and they confirmed this general picture.

Motivated by these results we studied the *behavior of many particles interacting with a broadband slab spectrum*, generated in order to mimic some of the major features of the solar wind (see Fig. 6): (a) three decades of the energy-containing scale ensure turbulence homogeneity, (b) three decades of inertial range well reproduce the observations, and (c) two decades of dissipation range enable us to cross the “ α_{\min} barrier” related to the “resonance gap” predicted by quasilinear theory [29,40]. After that there are almost two other decades of zero padding, important for the trigonometric interpolations and for the smoothness of the field. This is implemented using a numerical grid with $N_z = 2^{28} = 268\,435\,456$ points, corresponding to $134\,217\,728$ wave vectors for the spectrum. Apart from the obvious limitation that this spectrum is purely one dimensional, it is constructed to correspond roughly to features of solar wind spectra observed by single spacecraft, where the fully three-dimensional spectrum is in effect reduced to a one-dimensional form. Information is lost in the process (see, e.g., Ref. [38]).

In order to gain insight into magnetic moment conservation, we have performed simulations changing both particle velocity $v = (10, 100)v_A$ and the amplitude of the magnetic field fluctuations $\delta B/B_0 = (0.001, 0.01, 0.1, 0.5, 1.0)$. Particles injected at different velocities start to resonate at different points of the spectrum. We analyzed the distribution functions (see Fig. 7) and the variance (see Fig. 8) of the pitch angle cosine α , magnetic moment μ , and parallel position z .

From the experiment with resonances overlapping, we know that the three different regimes of statistical behavior of μ are related to other two effects: diffusion in velocity space and spatial parallel diffusion. These take place at different characteristic times τ_c and τ_{\parallel} , respectively. In order to investigate the effects of both processes on magnetic moment distributions, we followed test particles in the simulation box for times $T > \tau_c$.

For a low level of magnetic fluctuations particles free stream in the z direction while α and μ exhibit Gaussian

distributions around their initial values. For $\delta B/B_0 = 0.01$, particles cover completely one side of the α hemisphere, continuing to stream freely along z . This is the transient regime during which $f(\mu)$ exhibits a one-sided long tail distribution in the direction of smaller μ that appears to be a typical feature of magnetic moment distribution. During this transient regime the distribution of particles nearly conserves its magnetic moment. With increase in the value of $\delta B/B_0$, spatial diffusion starts to take place, and $f(\mu)$ recovers the typical Gaussian shape centered in the middle of μ space. These different regimes of magnetic moment statistical behavior are related not just to the variation of the turbulence level $\delta B/B_0$, but also to the different time scale at which magnetic moment conservation is studied (see Fig. 9).

In spite of the limitations of the present approach, the results presented here provide a basic view of how magnetic moments are modified in simplified models, and in particular how magnetic moment changes are related to pitch angle changes and sampling of magnetic variations due to spatial diffusion. It is clear that additional study is required to understand more fully the influences of turbulence on magnetic moment statistics. For example, realistic three-dimensional models of the magnetic field turbulence, as well as incorporation of electric field fluctuation effects, are expected to have significant effects. It is also possible that non-Gaussian features of magnetic field fluctuations, such as are associated with intermittency effects, may also influence magnetic moment changes, much as they influence spatial transport due to trapping and related influences [41,42]. In this regard the present results, along with those of Ref. [5], may be considered as baseline or minimal quantification of nonconservation of magnetic moments of a distribution of test particles in turbulence. Planned future studies will investigate quantitatively how additional realism in the modeling might produce even more significant departures from magnetic moment conservation.

ACKNOWLEDGMENTS

This research was supported in part by the NASA Heliophysics Theory Program No. NNX11AJ44G, and by the NSF Solar Terrestrial and SHINE programs (Grants No. AGS-1063439 and No. AGS-1156094), by the NASA MMS and Solar probe PLUS Projects, and by Marie Curie Project No. FP7 PIRSES-2010-269297 (Turboplasmas).

APPENDIX: DERIVATION OF TRAPPING HALF-WIDTH FOR A CIRCULARLY POLARIZED WAVE

Using Eqs. (5a) and (5b) of Ref. [25] it is possible to derive a simplified expression for the trapping half-width and the bounce frequency in the case of an Alfvén static wave [23]. For this particular case $k_{\perp} = 0$ and $\phi = 0$. We can rewrite Eq. (5c) of Ref. [25] as

$$Z_n = mc^2 \left\{ \frac{v_{\perp}}{2c} \left[\left(\epsilon_2 - \frac{k_{\parallel}}{k} \sigma \epsilon_1 \right) J_{n-1}(k_{\perp} \rho) - \left(\epsilon_2 + \frac{k_{\parallel}}{k} \sigma \epsilon_1 \right) J_{n+1}(k_{\perp} \rho) \right] + \sigma \left(\frac{v_{\parallel} k_{\perp}}{ck} \epsilon_1 + \epsilon_3 \right) J_n(k_{\perp} \rho) \right\}, \quad (\text{A1})$$

with $\cos \alpha = 1$ and $\sin \alpha = 0$. Because $\mathbf{k} \parallel \mathbf{B}_0$ we can choose $\hat{\mathbf{e}}_z = \mathbf{B}_0/|\mathbf{B}_0|$, $\hat{\mathbf{e}}_y$ is any arbitrary direction perpendicular to $\hat{\mathbf{e}}_z$, and $\hat{\mathbf{e}}_x = \hat{\mathbf{e}}_y \times \hat{\mathbf{e}}_z$. The vector potential can be obtained from the magnetic field, $\nabla \times \mathbf{B}_\perp = B_x \hat{\mathbf{e}}_x + B_y \hat{\mathbf{e}}_y$. In Fourier space $\nabla \rightarrow ik_z \hat{\mathbf{e}}_z$, so we have

$$A_x = -\frac{i}{k_\parallel} B_y, \quad A_y = \frac{i}{k_\parallel} B_x. \quad (\text{A2})$$

Considering only a single circularly polarized wave in space, for the two different possible helicities we can write

$$\mathbf{B}_\pm = (B_\pm \hat{\mathbf{e}}_\pm) \exp[i(k_\parallel z)], \quad (\text{A3})$$

where

$$B_\pm = \frac{1}{\sqrt{2}}(B_x \mp i B_y) \quad \text{and} \quad \hat{\mathbf{e}}_\pm = \frac{1}{\sqrt{2}}(\hat{\mathbf{e}}_x \mp i \hat{\mathbf{e}}_y) \quad (\text{A4})$$

are respectively the complex amplitudes and the orthogonal polarization unit vectors. The $+$ ($-$) polarization state is the positive (negative) helicity, i.e., the vector \mathbf{B} is rotating counterclockwise (clockwise). First, let us consider only the left-handed polarized wave \mathbf{B}_+ . Assuming $B_+ = \sqrt{2} \delta B e^{-i\pi/2}$ we can write the x and y components of the wave magnetic field as

$$\begin{aligned} B_x &= \delta B \exp[i(k_\parallel z - \pi/2)], \\ B_y &= \delta B \exp(ik_\parallel z). \end{aligned} \quad (\text{A5})$$

Inserting these two expressions into Eq. (A2) we obtain

$$\begin{aligned} A_x &= \frac{\delta B}{k_\parallel} \exp[i(k_\parallel z - \pi/2)], \\ A_y &= \frac{\delta B}{k_\parallel} \exp(ik_\parallel z). \end{aligned}$$

Comparing the real parts of these equations with Eq. (1) b) of Ref. [25] we obtain an expression for the coefficients A_1 and A_2 and for the normalized components of the wave polarization vector ϵ_1 , ϵ_2 , and ϵ_3 :

$$A_1 = \eta \frac{\delta B}{k_\parallel}, \quad A_2 = \frac{\delta B}{k_\parallel} \quad \text{where} \quad \eta = \frac{k_\parallel}{|k_\parallel|}, \quad (\text{A6})$$

$$\epsilon_1 = \frac{|q|\eta\delta B}{mc^2 k_\parallel}, \quad \epsilon_2 = \frac{|q|\delta B}{mc^2 k_\parallel}, \quad \epsilon_3 = 0. \quad (\text{A7})$$

TABLE V. Wave polarization and resonance contribution to trapping width.

Polarization	η	resonance n
B_+ left handed	1 parallel	-1
B_- right handed	-1 antiparallel	1
B_+ left handed	-1 parallel	1
B_- right handed	1 parallel	-1

Similarly, for a right-handed circularly polarized wave \mathbf{B}_- we have

$$\begin{aligned} A_1 &= -\eta \frac{\delta B}{k_\parallel}, \quad A_2 = \frac{\delta B}{k_\parallel} \quad \text{where} \quad \eta = \frac{k_\parallel}{|k_\parallel|}, \\ \epsilon_1 &= -\frac{|q|\eta\delta B}{mc^2 k_\parallel}, \quad \epsilon_2 = \frac{|q|\delta B}{mc^2 k_\parallel}, \quad \epsilon_3 = 0. \end{aligned}$$

In the case of a single circularly polarized wave propagating parallel (or antiparallel) to the magnetic field there is only one resonance present and the particle motion is integrable [25]: indeed $J_n(0) = 0$ unless $n = 0$. Therefore depending on the polarization of the wave and on its direction of propagation η only $l = 1$ or $l = -1$ resonances contribute to the trapping width, as shown in Table V. Thus, considering Eqs. (5a) and (5b) of Ref. [25], and Eqs. (A1), (A6), and (A7) with $J_0(0) = 1$, the authors of Ref. [23] find a specialized formula for the trapping half-width and bounce frequency applied to the case of a circularly polarized wave propagating parallel, $k_\parallel > 0$ and $n = -1$, or antiparallel, $k_\parallel > 0$ and $n = 1$, to \mathbf{B}_0 :

$$\begin{aligned} \Delta v_\parallel^{(-1)} &= 2v \left[(1 - \alpha^2)^{1/2} |\alpha| \frac{\delta B}{B_0} \right]^{1/2}, \\ \omega_b^{(-1)} &= \Omega_0 \left[\frac{(1 - \alpha^2)^{1/2} \delta B}{|\alpha| B_0} \right]^{1/2}, \end{aligned} \quad (\text{A8})$$

if $k_\parallel v_\parallel > 0$ and zero otherwise, in which $\alpha = \cos \theta$ is the cosine of the pitch angle. Exactly the same set of equations holds for $\Delta v_\parallel^{(+1)}$ and $\omega_b^{(+1)}$. However, the condition for their being nonzero is reversed, i.e., $k_\parallel v_\parallel > 0$. We omit the superscripts (± 1) because of this degeneracy.

[1] E. Marsch, in *Physics of the Inner Heliosphere, Vol. 2: Particles, Waves and Turbulence*, edited by E. Marsch (Springer, Berlin, 1991), pp. 45–133.
[2] K. Knizhnik, M. Swisdak, and J. F. Drake, *Astrophys. J. Lett.* **743**, L35 (2011).
[3] B. V. Chirikov, *Proc. R. Soc. London, Ser. A* **413**, 145 (1987).
[4] B. Rossi and S. Olbert, *Introduction to the Physics of Space* (McGraw-Hill, New York, 1970).
[5] R. Lehe, I. J. Parrish, and E. Quataert, *Astrophys. J.* **707**, 409 (2009).
[6] J. R. Jokipii, *Astrophys. J.* **146**, 480 (1966).
[7] I. H. Uch, *Astrophys. Space Sci.* **46**, 389 (1977).
[8] F. C. Jones, J. R. Jokipii, and M. G. Baring, *Astrophys. J.* **509**, 238 (1998).

[9] L. D. Landau, *J. Phys. (Moscow)* **10**, 25 (1946).
[10] A. J. Owens, *Astrophys. J.* **191**, 235 (1974).
[11] M. L. Goldstein, *Astrophys. J.* **204**, 900 (1976).
[12] B. D. G. Chandran, *Phys. Rev. Lett.* **85**, 4656 (2000).
[13] A. J. Klimas and G. Sandri, *Astrophys. J.* **169**, 41 (1971).
[14] A. J. Klimas and G. Sandri, *Astrophys. J.* **180**, 937 (1973).
[15] A. J. Klimas and G. Sandri, *Astrophys. J.* **184**, 955 (1973).
[16] M. L. Goldstein, A. J. Klimas, and G. Sandri, *Astrophys. J.* **195**, 787 (1975).
[17] J. W. Bieber, W. H. Matthaeus, C. W. Smith, W. Wanner, M.-B. Kallenrode, and G. Wibberenz, *Astrophys. J.* **420**, 294 (1994).
[18] G. R. Smith and A. N. Kaufman, *Phys. Fluids* **21**, 2230 (1978).
[19] A. J. Lichtenberg and B. P. Wood, *Phys. Rev. Lett.* **62**, 2213 (1989).

- [20] H. Karimabadi, K. Akimoto, N. Omid, and C. R. Menyuk, *Phys. Fluids B* **2**, 606 (1990).
- [21] J. Weinstock, *Phys. Fluids* **12**, 1045 (1969).
- [22] R. C. Davidson, *Methods in a Nonlinear Plasma Theory* (Academic Press, New York, 1972), p. 356.
- [23] R. L. Mace, S. Dalena, and W. H. Matthaeus, *Phys. Plasmas*, **19**, 032309 (2012).
- [24] P. J. Palmadesso, *Phys. Fluids* **15**, 2006 (1972).
- [25] H. Karimabadi, D. Krauss-Varban, and T. Terasawa, *J. Geophys. Res.* **97**, 13853 (1992).
- [26] C. F. Kennel and H. E. Petschek, *J. Geophys. Res.* **71**, 1 (1966).
- [27] D. G. Swanson, *Plasma Waves* (Academic Press, Boston, 1989).
- [28] *Waves in Plasmas*, edited by T. H. Stix (American Institute of Physics, New York, 1992).
- [29] T. B. Kaiser, T. J. Birmingham, and F. C. Jones, *Phys. Fluids* **21**, 361 (1978).
- [30] R. L. Mace, W. H. Matthaeus, and J. W. Bieber, *Astrophys. J.* **538**, 192 (2000).
- [31] J. Minnie, R. A. Burger, Parhi S., W. H. Matthaeus, and J. W. Bieber, *Adv. Space Res.* **35**, 543 (2005).
- [32] J. Ambrosiano, W. H. Matthaeus, M. L. Goldstein, and D. Plante, *J. Geophys. Res.* **93**, 14383 (1988).
- [33] M. L. Goldstein, W. H. Matthaeus, and J. J. Ambrosiano, *Geophys. Res. Lett.* **13**, 205 (1986).
- [34] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes*, 2nd ed. (Cambridge University Press, New York, 1992).
- [35] B. V. Chirikov, *Fiz. Plazmy* **4**, 521 (1978).
- [36] S. Dalena, P. Chuychai, R. L. Mace, A. Greco, G. Qin, and W. H. Matthaeus, *Comput. Phys. Commun.* **183**, 1974 (2012).
- [37] W. H. Matthaeus, M. L. Goldstein, and J. H. King, *J. Geophys. Res.* **91**, 59 (1986).
- [38] J. W. Bieber, W. Wanner, and W. H. Matthaeus, *J. Geophys. Res., A* **101**, 2511 (1996).
- [39] S. Oughton, E. R. Priest, and W. H. Matthaeus, *J. Fluid Mech.* **280**, 65 (1994).
- [40] T. B. Kaiser, F. C. Jones, and T. J. Birmingham, *Astrophys. J.* **180**, 239 (1973).
- [41] D. Ruffolo, W. H. Matthaeus, and P. Chuychai, *Astrophys. J.* **597**, 169 (2003).
- [42] P. Chuychai, D. Ruffolo, W. H. Matthaeus, and J. Meechai, *Astrophys. J.* **659**, 1761 (2007).