

Scaling laws and accurate small-amplitude stationary solution for the motion of a planar vortex filament in the Cartesian form of the local induction approximation

Robert A. Van Gorder*

Department of Mathematics, University of Central Florida, Orlando, Florida 32816-1364, USA

(Received 18 January 2013; revised manuscript received 15 March 2013; published 23 April 2013)

We provide a formulation of the local induction approximation (LIA) for the motion of a vortex filament in the Cartesian reference frame (the extrinsic coordinate system) which allows for scaling of the reference coordinate. For general monotone scalings of the reference coordinate, we derive an equation for the planar solution to the derivative nonlinear Schrödinger equation governing the LIA. We proceed to solve this equation perturbatively in small amplitude through an application of multiple-scales analysis, which allows for accurate computation of the period of the planar vortex filament. The perturbation result is shown to agree strongly with numerical simulations, and we also relate this solution back to the solution obtained in the arclength reference frame (the intrinsic coordinate system). Finally, we discuss nonmonotone coordinate scalings and their application for finding self-intersections of vortex filaments. These self-intersecting vortex filaments are likely unstable and collapse into other structures or dissipate completely.

DOI: [10.1103/PhysRevE.87.043203](https://doi.org/10.1103/PhysRevE.87.043203)

PACS number(s): 47.32.C–

I. INTRODUCTION

The self-induced velocity of a vortex filament has been described by the local induction approximation (LIA) $\mathbf{v} = \gamma \kappa \mathbf{t} \times \mathbf{n}$ [1–3], where \mathbf{t} and \mathbf{n} are unit tangent and unit normal vectors to the vortex filament, respectively, κ is the curvature, and γ is the strength of the vortex filament. A number of methods have been employed to study the LIA. Exact stationary solutions to the LIA in extrinsic coordinate space have been found by Kida [4] in the case of torus knots, and these solutions were given in terms of elliptic integrals. By rewriting the LIA in cylindrical-polar coordinates, Ricca also obtained torus knot solutions—which were asymptotically equivalent to Kida’s solutions—in explicit analytic form and derived a stability criterion [5]. Static solutions to the LIA have also been found by Lipniacki [6]. Physical invariants obtained under LIA are discussed by Ricca in Ref. [7].

The fully nonlinear Schrödinger (NLS) equation governing the self-induced motion of a vortex filament in the LIA was previously derived by Van Gorder [8,9] in the Cartesian coordinate space. Dmitriyev [10] considered a linear approximation of the LIA, while Shivamoggi and van Heijst [11] considered a more sophisticated approximation, obtaining a cubic derivative NLS equation. The full nonlinear equation was obtained in Ref. [8]. Some existence results for space-periodic planar vortex filaments will be given in a forthcoming work. In particular, it has been shown that there exists a maximal amplitude beyond which space-periodic planar vortex filament solutions do not exist. We should remark that the aforementioned studies have considered the Cartesian form of the LIA. Umeki [12] obtained an alternate formulation, applying an arclength-based coordinate system as opposed to a Cartesian coordinate system. While the Cartesian and arclength formulations are obtained through different derivations, both formulations are equivalent to the LIA. Van Gorder [13] obtained exact stationary solutions for this model in terms of elliptic functions.

In the present paper, we provide a formulation of the LIA for the motion of a vortex filament in the Cartesian reference frame (the extrinsic coordinate system) which allows for scaling of the reference coordinate. For general monotone scalings of the reference coordinate, we derive an equation for the planar solution to the derivative NLS equation governing the LIA. We proceed to solve this equation perturbatively in small amplitude through an application of multiple-scales analysis, which allows for accurate computation of the period of the planar vortex filament. The perturbation result is shown to agree strongly with numerical simulations, and we also relate this solution back to the solution obtained in the arclength reference frame (the intrinsic coordinate system). Finally, we discuss nonmonotone coordinate scalings and their application for finding self-intersections of vortex filaments. These self-intersecting vortex filaments are unstable and collapse into other structures or dissipate completely.

II. FORMULATION AND SCALING OF THE LIA

Alternate scaling of the LIA can be useful both for physical analysis and for computational reasons. For instance, the infinite domain due to $x \in \mathbb{R}$ can be mapped into a closed and bounded interval, which can assist with analytical and numerical analyses. We first determine the influence of such transforms. Let us consider the scaled position vector

$$\mathbf{r} = f(x)\mathbf{i}_x + y(x,t)\mathbf{i}_y + z(x,t)\mathbf{i}_z, \quad (1)$$

where $f(x)$ denotes a general scaling of the x coordinate. This is one of two possible equivalent such scalings, with the other being

$$\mathbf{r} = x\mathbf{i}_x + y(f^{-1}(x),t)\mathbf{i}_y + z[f^{-1}(x),t]\mathbf{i}_z \quad (2)$$

provided f^{-1} , the inverse map of f , exists. For this reason, we will often be interested in monotone scalings f so that the inversion f^{-1} is well-defined. We chose to work with Eq. (1) as opposed to Eq. (2) since it gives more computationally

*rav@knights.ucf.edu

tractable results. From Eq. (1), we compute

$$\mathbf{t} = \frac{d\mathbf{r}}{ds} = \frac{d\mathbf{r}}{dx} \frac{dx}{ds} = (f', y_x, z_x) \frac{dx}{ds}$$

and $\mathbf{v} = (0, y_t, z_t)$, where

$$\frac{dx}{ds} = \frac{1}{\sqrt{f'^2 + y_x^2 + z_x^2}}.$$

We then have $\kappa \mathbf{n} = \frac{d\mathbf{t}}{ds} = \frac{d\mathbf{t}}{dx} \frac{dx}{ds}$, giving

$$\begin{aligned} \kappa \mathbf{n} &= [f''(y_x^2 + z_x^2) - f'(y_x y_{xx} + z_x z_{xx})] \frac{dx}{ds} \mathbf{i}_x \\ &+ [y_{xx} z_x - y_x z_{xx} + y_{xx} f'^2 - f' f'' y_x] \frac{dx}{ds} \mathbf{i}_y \\ &+ [z_{xx} y_x^2 - z_x y_x y_{xx} + z_{xx} f'^2 - f' f'' z_x] \frac{dx}{ds} \mathbf{i}_z, \end{aligned}$$

so that $\mathbf{v} = \gamma \kappa \mathbf{t} \times \mathbf{n} = \gamma \mathbf{t} \times (\kappa \mathbf{n})$ becomes

$$\begin{aligned} \mathbf{v} &= \gamma(y_x z_{xx} - z_x y_{xx}) \left(\frac{dx}{ds}\right)^3 \mathbf{i}_x \\ &- \gamma(f' z_{xx} - f'' z_x) \left(\frac{dx}{ds}\right)^3 \mathbf{i}_y \\ &+ \gamma(f' y_{xx} - f'' y_x) \left(\frac{dx}{ds}\right)^3 \mathbf{i}_z. \end{aligned}$$

Matching the two representations of \mathbf{v} , we obtain the constraint $y_x z_{xx} - z_x y_{xx} = 0$ and the real-valued system

$$\begin{aligned} y_t &= -\gamma(f' z_{xx} - f'' z_x) \left(\frac{dx}{ds}\right)^3, \\ z_t &= \gamma(f' y_{xx} - f'' y_x) \left(\frac{dx}{ds}\right)^3. \end{aligned}$$

Introducing the complex potential function

$$\Phi(x, t) = y(x, t) + iz(x, t),$$

the PDE system reduces to

$$i\Phi_t + \gamma(f'\Phi_{xx} - f''\Phi_x)(f'^2 + |\Phi_x|^2)^{-3/2} = 0. \quad (3)$$

Note that Eq. (3) is a complicated NLS equation with variable coefficients (f' and f'' in general depend on x). However, with the scaling $\Phi(x, t) = \Psi(\mu, t)$, where $\mu = f(x)$, we can reduce Eq. (3) to

$$i\Psi_t + \gamma(1 + |\Psi_\mu|^2)^{-3/2} \Psi_{\mu\mu} = 0, \quad (4)$$

for nondegenerate f . Hence, Eq. (3) yields solutions of the LIA. The transformed equation (4) matches exactly that studied in Refs. [8,9].

The form of Eq. (3) [and hence Eq. (4)] is $U(1)$ invariant, just like many of its derivative NLS relatives; hence it makes sense to consider stationary solutions of the form $\Phi(x, t) = e^{-i\gamma t} \phi(x)$ to Eq. (3) [and $\Psi(\mu, t) = e^{-i\gamma t} \psi(\mu)$ to Eq. (4)].

To summarize, the permitted scalings are those for which

- (1) the LIA is invariant under monotone scalings of the x coordinate and
 - (2) the LIA is invariant under scalings of the form $e^{-i\gamma t}$.
- Together, these conditions guarantee the existence of planar vortex filaments described by $\Psi(x, t) = e^{-i\gamma t} \psi(\mu(x))$. In the




FIG. 1. Plot of the spatial geometry. The curve represents the planar vortex filament described by $\Phi(x, t) = e^{-i\gamma t} \phi(\mu(x))$ for periodic $\psi(\mu(x))$. As time increases, the structure rotates about the x axis.

extrinsic three-dimensional Cartesian frame, the position of the planar vortex filament at any time t is then given by

$$\mathbf{r} = \mu(x)\mathbf{i}_x + \cos(\gamma t)\psi(\mu(x))\mathbf{i}_y - \sin(\gamma t)\psi(\mu(x))\mathbf{i}_z. \quad (5)$$

To better visualize such vortex filaments, see Fig. 1, where we consider a periodic function, $\psi(\mu(x))$. The vortex filament rotates about the x axis as time increases. So, by determining $\psi(\mu(x))$, we determine the spatial structure of the planar vortex filament completely, the inclusion of the factor $e^{-i\gamma t}$ providing the motion of such a filament in time.

III. ACCURATE PERTURBATION APPROACH FOR THE STATIONARY SOLUTION

Let us consider the stationary solution $\Psi(\mu, t) = Ae^{-i\gamma t} \psi(\mu)$ to the scaled equation (4), where we let the parameter $A > 0$ hold the amplitude and normalize $\max \psi = 1$. Then, we obtain the ordinary differential equation

$$\psi + (1 + A^2 \psi'^2)^{-3/2} \psi'' = 0. \quad (6)$$

The simplest nonlinear approximation of the equation takes the form

$$\psi + \left(1 - \frac{3}{2} A^2 \psi'^2\right) \psi'' = 0. \quad (7)$$

As discussed in a forthcoming work, Eq. (7) has periodic real-valued solutions for $A < 1/\sqrt{3} \approx 0.577$. For small A , Eq. (7) is a good approximation of Eq. (6). It then makes sense to consider a perturbation solution, in terms of small parameter A^2 . However, standard perturbation will yield inaccurate solutions which fall out of resonance with the true solution due to the appearance of secular terms. Hence, we are interested in applying the method of multiple scales to Eq. (7). To proceed, assume there exists the parameter $\delta(A^2)$ such that $d/d\mu = \delta(A^2)(d/d\eta)$, where $\eta = \delta(A^2)\mu$. Then, we consider the perturbation solution $\psi(\mu) = \hat{\psi}(\eta; A^2) = \psi_0(\eta; A^2) + A^2 \psi_1(\eta; A^2) + O(A^4)$, $\delta(A^2) = \delta_0 + A^2 \delta_1 + O(A^4)$.

Equation (7) becomes

$$\hat{\psi} + \delta^2 \left(1 - \frac{3}{2} A^2 \delta^2 \hat{\psi}_\eta^2\right) \hat{\psi}_{\eta\eta} = 0, \quad (8)$$

giving

$$\delta_0^2 \psi_{0,\eta\eta} + \psi_0 = 0, \quad \psi_0(0) = 1, \quad \psi_{0,\eta}(0) = 0, \quad (9)$$

$$\begin{aligned} \delta_0^2 \psi_{1,\eta\eta} + \psi_1 &= \frac{3}{2} \delta_0^4 \psi_{0,\eta}^2 (\psi_0)_{\eta\eta} - 2\delta_0 \delta_1 \psi_{0,\eta\eta}, \\ \psi_1(0) &= 0 = (\psi_1)_\eta(0). \end{aligned} \quad (10)$$

The quantities at $\eta = 0$ follow from the fact that we desire space-periodic ψ with amplitude A . As we assume the solution $\Psi = A\psi \exp(-i\gamma t)$, it follows that the amplitude of ψ must be 1 (then the amplitude of Ψ is A). Without loss of generality, we take $\eta = 0$ to correspond to a peak (this can be translated by $\eta \rightarrow \eta' + \eta_0$ if need be). Thus, $\psi_\eta(x) = 0$. Assuming $\psi = \psi_0 + A^2 \psi_1 + \dots$, it follows that $\psi_0(0) = 1$, $\psi_{0,\eta}(0) = 0$, $\psi_1(0) = 1$, and $\psi_{1,\eta}(0) = 0$.

Normalizing to get 2π -periodic solutions, we pick $\delta_0 = 1$, obtaining $\psi_0(\eta) = \cos(\eta)$. From here, we have

$$(\psi_1)_{\eta\eta} + \psi_1 = \left(2\delta_1 - \frac{3}{8}\right) \cos(\eta) + \frac{3}{8} \cos(3\eta), \quad (11)$$

so picking $\delta_1 = 3/16$ prevents any secular terms. We then obtain

$$\psi_1(\eta) = \frac{3}{64} [\cos(\eta) - \cos(3\eta)] = \frac{3}{16} \sin^2(\eta) \cos(\eta). \quad (12)$$




FIG. 2. (Color online) Plot of the perturbation solutions (13) for $\psi(x)$ obtained through the method of multiple scales against numerical solutions obtained via the Runge-Kutta-Fehlberg method (RKF45) [14]. The valid region for the approximation (7) is $A < 1/\sqrt{3} \approx 0.577$, and in this region the results agree nicely. For larger A , the agreement breaks down as the solutions fall out of resonance with the true solutions.




FIG. 3. (Color online) Plot of the absolute error between the perturbation solutions (13) for $\psi(x)$ obtained through the method of multiple scales and the numerical solutions obtained via the Runge-Kutta-Fehlberg method (RKF45) [14]. The agreement is strong for small-amplitude solutions, while the agreement gradually breaks down for larger amplitudes.

Therefore, we have obtained the perturbation solution

$$\begin{aligned} \psi(\mu) &= \cos\left(\left[1 + \frac{3}{16} A^2\right] \mu\right) \\ &+ \frac{3}{16} A^2 \sin^2\left(\left[1 + \frac{3}{16} A^2\right] \mu\right) \\ &\times \cos\left(\left[1 + \frac{3}{16} A^2\right] \mu\right). \end{aligned} \quad (13)$$

Consider the standard case $\mu(x) = x$. From Eq. (13), we see that the approximate period of small-amplitude solutions satisfies

$$T(A) \approx 2\pi \left[1 + \frac{3}{16} A^2\right]^{-1} \approx 2\pi - \frac{3\pi}{8} A^2 + \frac{9\pi}{128} A^4. \quad (14)$$

In order to demonstrate the agreement between the solution (13) and the true solution, we plot the numerical solution along with the perturbation solution in Fig. 2. Since the perturbation and numerical results agree so nicely, the difference between the two is not easily ascertainable, so we plot their errors separately, in Fig. 3.

IV. CONNECTION WITH ARCLENGTH SOLUTION AND IMPLICIT SOLUTION

In this section, take $\mu(x) = x$, so that $\Phi(x, t) = \Psi(\mu, t)$. In Van Gorder [13], an exact stationary solution for the arclength formulation of the LIA was given by

$$v(s, t) = e^{-it} q(s) = B e^{-it} \operatorname{sn}\left(\frac{s - \hat{s}}{\sqrt{1 - B^2}}, Bi\right), \quad (15)$$

where B is the amplitude (in the arclength frame), \hat{s} is a constant, and s is the arclength element. It was shown in Ref. [12] that the Cartesian quantity $\Phi(x, t)$ and the arclength

quantity $v(s,t)$ are related by

$$|\Phi_x|^2 = \frac{2|v|^2}{(1-|v|^2)^2}, \quad \frac{dx}{ds} = \frac{1-|v|^2}{1+|v|^2}. \quad (16)$$

Noting that $|\Phi_x| = \phi'(x)$ and $|v| = q(s)$, we have that $\phi'^2 = 2q^2(1-q^2)^{-2}$. Separating variables, and using the form of dx/ds given in Eq. (16), we obtain

$$\phi(x) = \sqrt{2} \int_{\delta}^{s(x)} \frac{B \operatorname{sn}\left(\frac{s-\delta}{\sqrt{1-B^2}}, Bi\right)}{1+B^2 \operatorname{sn}^2\left(\frac{s-\delta}{\sqrt{1-B^2}}, Bi\right)} ds. \quad (17)$$

Performing the integration exactly is not possible (in closed form). And then, one must still contend with the arclength variable $s(x)$. So, while this formula offers a connection between the exact arclength solution to the planar vortex filament problem and that of the Cartesian problem, it is not very practical. We can compare this formula to the direct solution for $\phi(x)$. A first integral of Eq. (6) [when $\mu = x$ and hence $\psi(\mu) = \phi(x)$] is

$$\phi^2 - \frac{2}{A^2 \sqrt{1+A^2 \phi'^2}} = -\mathcal{E}. \quad (18)$$

If $\phi(0) = 1$ and $\phi'(0) = 0$, then $\mathcal{E} = (2-A^2)/A^2 > 0$ since $|A| < \sqrt{2}$ for any periodic solution. Solving Eq. (18) for ϕ' and separating variables as needed,

$$x = \pm \int_{\phi}^1 \frac{A^3(\zeta^2 + \mathcal{E})}{\sqrt{4-A^4(\zeta^2 + \mathcal{E})^2}} d\zeta. \quad (19)$$

Changing variables to $\xi = \zeta^2 + \mathcal{E}$,

$$x = \pm \frac{A^3}{2} \int_{\phi^2+\mathcal{E}}^{2/A^2} \frac{\xi d\xi}{\sqrt{(\xi-\mathcal{E})(2-A^2\xi)(2+A^2\xi)}}. \quad (20)$$

Equation (20) is an implicit solution which is not easily inverted. However, we may still extract information out of this relation more easily than is the case when dealing with Eq. (17). In the previous section, we approximated the period of a space-periodic planar vortex filament using perturbation. We shall now be interested in comparing that approximation with a true exact relation between the period T and the amplitude A for a space-periodic solution to the vortex filament problem.

If we consider the phase portrait, a quarter-period $T(A)/4$ occurs when ϕ goes from $\phi = 0$ to $\phi = 1$, so from Eq. (20) we obtain the exact yet implicit relation

$$T(A) = 2A^3 \int_{\mathcal{E}}^{2/A^2} \frac{\xi d\xi}{\sqrt{(\xi-\mathcal{E})(2-A^2\xi)(2+A^2\xi)}}. \quad (21)$$

Now, in the valid region $0 < A < \sqrt{2}$, the definite integral (21) can be evaluated in terms of elliptic integrals to give the relation

$$T(A) = 8E(A/2) - 4K(A/2), \quad (22)$$

where K is the complete elliptic integral of the first kind and E is the complete elliptic integral of the second kind. Recall that the period of the solutions in the arclength representation [13], the period of the space-periodic solution, was a bit simpler, involving only the elliptic integral K .




FIG. 4. (Color online) Plot of the x -period $T(A)$ for the stationary solution x -dependence function $\phi(x)$. In addition to the exact value (22), we plot two approximate quantities, namely, the approximation found through multiple scales (14) and the asymptotic approximation (25) of the true result (22). We consider $A \in [0, \sqrt{2}]$.

In order to extract more information from Eq. (22), we turn to the small- θ asymptotics

$$K(\theta) = \frac{\pi}{2} \left(1 + \frac{1}{4} \frac{\theta^2}{1-\theta^2} - \frac{1}{8} \frac{\theta^4}{1-\theta^2} \right), \quad (23)$$

$$E(\theta) = \frac{\pi}{2} \left(1 - \frac{1}{4} \theta^2 - \frac{3}{64} \theta^4 \right). \quad (24)$$

Using Eqs. (23) and (24) in Eq. (22), and approximating where needed,

$$T(A) \approx 2\pi - \frac{3\pi}{8} A^2 - \frac{7\pi}{256} A^4. \quad (25)$$

Note that the approximation (25) of the period $T(A)$ obtained through the fully nonlinear relation (20) for ϕ is in extremely good agreement with the approximation obtained through the method of multiple scales (14) for the period $T(A)$. In Fig. 4, we plot the exact period $T(A)$ found in Eq. (22), along with the approximations shown in Eqs. (14) and (25). In Fig. 5, we plot the relative error between the approximations and the exact values.

V. NONMONOTONE SPACE SCALES AND NUMERICAL SELF-INTERSECTION OF FILAMENTS

Up to this point we have considered only monotone scalings $f(x)$ in Eq. (1), since these permit well-behaved solutions to Eq. (4). As we have shown, such solutions can be studied analytically, and in some cases exactly. However, in situations where $f(x)$ is nonmonotone, we may still assume a stationary solution of the form $\Phi(x,t) = e^{-i\gamma t} \phi(x)$. While $\psi(\mu)$ from Eq. (6) was defined on the real μ axis in the case of monotone $\mu = f(x)$, for nonmonotone $f(x)$ it is possible that the domain of $\phi(x)$ will be restricted. Assuming the solution




FIG. 5. (Color online) We demonstrate the relative error between the approximations of the period $T(A)$ and the true solution (22). Both are extremely accurate for small A , and gradually lose accuracy for larger A , though the asymptotic approximation (25) outperforms the multiple-scale approximation (14) nicely. That said, in its region of validity ($A < 1/\sqrt{3}$), the multiple-scale approximation (14) is rather accurate for only a first-order perturbation result.

$\Phi(x, t) = e^{-i\gamma t} \phi(x)$, Eq. (3) reduces to

$$\phi + \frac{f'\phi'' - f''\phi'}{(f'^2 + \phi'^2)^{3/2}} = 0. \quad (26)$$

The ordinary differential equation (26) is degenerate when f is not strictly monotone, i.e., if there exists a point $x = a$ at which $f'(a) = 0$. In order for a planar vortex filament to have self-intersections, there should exist points $x_* < x^*$ such that $f(x_*) = f(x^*)$ and $\phi(x_*) = \phi(x^*)$, but for $x_* < x_1 < x_2 < x^*$, $f(x_1) = f(x_2)$ and $\phi(x_1) = \phi(x_2)$ cannot hold simultaneously. If such x_1 and x_2 exist, then there can be a loop (if not, then we just have a constant valued function). Then from Eq. (1) we must have $\mathbf{r}(x_*, t) = \mathbf{r}(x^*, t)$ for all $t \geq 0$. If we have such points $x_* < x_1 < x_2 < x^*$, there is at least one loop formed. This loop is parametrized by $\theta \in [x_*, x^*]$ as

$$\mathbf{r}(\theta, t) = f(\theta)\mathbf{i}_x + \cos(\gamma t)\phi(\theta)\mathbf{i}_y - \sin(\gamma t)\phi(\theta)\mathbf{i}_z, \quad (27)$$

with the loop closing since $\mathbf{r}(x_*, t) = \mathbf{r}(x^*, t)$. In Fig. 6 we provide a schematic of the planar loop vortex filament. Now that we have some conditions on parametrized crossings and loop structures on a vortex filament, we provide some examples to show that these structures can actually occur as solutions to the equation governing a vortex filament of planar type.

A. Single-loop case

As our first numerical case, we consider an example of a parametrized single loop on a vortex filament. Let us consider




FIG. 6. (Color online) Schematic of a self-intersection for the planar vortex filament governed by a solution $\phi(x)$ to Eq. (26). Self-intersection occurs at a solution x^* where the parametrization x attains the value x^* such that $f(x_*) = f(x^*)$ and $\phi(x_*) = \phi(x^*)$. It is necessary for $\phi(x_1) \neq \phi(x_2)$ for all $x_* < x_1 < x_2 < x^*$ in order to have a single loop. For multiple loops, similar yet more complicated conditions must hold.

the scaling $f(x) = x^2/2$. We then have

$$\phi + \frac{x\phi'' - \phi'}{(x^2 + \phi'^2)^{3/2}} = 0. \quad (28)$$

Unlike in the simpler case of monotone f , here we cannot easily solve the differential equation (28) analytically. So, we resort to numerical solutions. It is useful to assign a specific x_N as a numerical initial point. Picking $x_N = 0$ is problematic, since Eq. (28) is degenerate at that point. So, we shall take x_N to be small yet positive. We find that loops are not obtained for many parameter values. However, they can occur for our choice of f . Taking, for instance, $x_N = 0.1$, $\phi(x_N) = 0.6$, and $\phi'(x_N) = -0.1$, we find that $\phi(2.059) = \phi(-2.059)$ while $f(x) = f(-x)$ by the form of f selected, so we pick $x_* = -2.059$ and $x^* = 2.059$. To make sure the loop is closed, the derivatives should differ at each point. We calculate $\phi'(x_*) = 3.589$ while $\phi'(x^*) = 0.857$, so the loop does close. So, in the prescribed geometry, we have found a closed filament loop. As mentioned above, the loop must remain closed for all $t \geq 0$. The resulting single-loop planar vortex filament is displayed in Fig. 7.

B. Double-loop case

Let us now consider a double-loop structure on a vortex filament. Let us take the scaling $f(x) = \cos(x)$. We then have

$$\phi + \frac{-\sin(x)\phi'' + \cos(x)\phi'}{(\sin^2(x) + \phi'^2)^{3/2}} = 0. \quad (29)$$

Taking $x_N = 0.1$, $\phi(x_N) = 0.5$, and $\phi'(x_N) = -0.095$, we numerically solve Eq. (29). Defining $-x_*^{[1]} = 2.35 = x_*^{[1]}$, $x_*^{[2]} = -3.89$, and $x_*^{[2]} = 2.395$, we have that $\phi(x_*^{[1]}) = \phi(x_*^{[1]})$ and $\phi(x_*^{[2]}) = \phi(x_*^{[2]})$. Yet, since $f(x) = \cos(x)$, we have $f(x_*^{[1]}) = f(x_*^{[1]})$ and $f(x_*^{[2]}) = f(x_*^{[2]})$. So, the conditions for crossing are satisfied at spatial coordinates $\cos(x_*^{[1]}) = -0.70$ and $\cos(x_*^{[2]}) = -0.73$. We verify that the




FIG. 7. (Color online) Plot of the numerical solution for a single-loop vortex filament described by $\phi(x)$ when $\phi(x)$ satisfies Eq. (28), $\phi(0.1) = 0.6$, and $\phi'(0.1) = -0.1$. The x scaling is $f(x) = x^2/2$. The space coordinate is parametrized by $x \in [-2.12, 3.00]$.

derivatives differ at each point, so the loop structures close off at the required points. (If the derivatives do not differ, then the filament may become tangent to itself, and therefore not close to form a loop, at the required point.) Hence, we have obtained a double-loop structure on a vortex filament. The resulting double-loop planar vortex filament is displayed in Fig. 8.

One may continue with multiloop structures, but these get progressively harder to construct, since one must guess an appropriate transform of the space variable $f(x)$ and deduce values of the crossings. Further, since this is done numerically




FIG. 8. (Color online) Plot of the numerical solution for a double-loop vortex filament described by $\phi(x)$ when $\phi(x)$ satisfies Eq. (29), $\phi(0.1) = 0.5$, and $\phi'(0.1) = -0.095$. The x scaling is $f(x) = \cos(x)$. The space coordinate is parametrized by $x \in [-4.0, 2.5]$.

(such analytical constructions are very challenging), there is a bit of guess work involved in the initial conditions which permit solutions $\phi(x)$ which allow for the crossings.

While these loop structures have been shown to exist numerically for appropriate scales f and planar components ϕ , in practice one would not expect these structures to persist. A more physically relevant situation would be for a vortex filament to cross (or come close to crossing, since physically the vortex core has a nontrivial diameter), a loop structure is momentarily formed, and then the filament is disrupted. Depending on the ambient fluid, one could have that (i) the vortex filament sheds the loop and realigns as a noncrossing well-defined curve; (ii) the loop collapses, with the filament stretching laterally to realign as a nonintersecting curve; or (iii) the loop dominates, with the “tails” decaying, resulting in a vortex ring. While the present results point toward any of those outcomes, note that one would need to retain more structure than the LIA permits in order to fully model the dynamics of such vortex filament crossings. In order to study such complicated dynamics, the full integral form of the Biot-Savart law would be required. At best, the LIA provides a sort of first-order approximation of such behavior, though it fails to pick up on the complicated dynamics of these situations, which would lead from a vortex filament crossing to one of the possible outcomes listed. That the LIA can pick up on the occurrence of such interactions, given its simplicity relative to the full Biot-Savart law, is still beneficial.

C. Analytical calculation

While numerical results are easiest for the case of self-intersections, we remark that analytical approximations can be obtained, at a cost. Indeed, when $f'(x_0) = 0$ for some x_0 , then Eq. (26) degenerates (the coefficient of ϕ'' vanishes, decreasing the order of the equation). To counter this, we must have two solution branches, which we match at x_0 . However, while the matching preserves continuity, it cannot preserve continuity of the first derivative (on each side of x_0 , that is, $x < x_0$ and $x > x_0$, the slope of the branches must differ). Without loss of generality, take $x_0 = 0$. Then, in order to match a positive branch and a negative branch, we consider the following piecewise defined solution:

$$\phi(x) = \begin{cases} -\psi(f(x)), & x_* < x < 0, \\ 0, & x = 0, \\ \psi(f(x)), & 0 < x < x^*, \end{cases} \quad (30)$$

where $\psi(\mu)$ is a solution as was found in the monotone case and $x_* < 0 < x^*$ such that $f(x_*) = f(x^*) = T/2$, where T is the period of ψ . From the form of Eq. (26), if ψ is a solution, then so is $-\psi$. Hence, each branch is a solution (when $f' \neq 0$). This representation is not unique, as we could have reversed the signs in Eq. (30). To get both functions to match at $x = 0$, we use a modified form of Eq. (13) where $\psi(0) = 0$ and $\psi'(0) = 1$ (which gives a sine, as opposed to cosine, representation). This is equivalent to translation of the solution in Eq. (13) by $-\pi/2$ on the x axis. So, to lowest order (one can add higher-order corrections, but we suppress them for brevity)

Eq. (30) becomes

$$\phi(x) = \begin{cases} -\sin\left\{\left[1 + \frac{3}{16}A^2\right]f(x)\right\}, & x_* < x < 0, \\ 0, & x = 0, \\ \sin\left\{\left[1 + \frac{3}{16}A^2\right]f(x)\right\}, & 0 < x < x^*. \end{cases} \quad (31)$$

Note that $\phi(x_*) = -\psi(f(x_*)) = -\psi(T/2) = 0 = \psi(T/2) = \psi(f(x^*)) = \phi(x^*)$ by construction, so $\phi(x_*) = \phi(x^*)$.

For example, consider again the $f(x) = x^2/2$ case. We pick

$$x_*(A) = -\sqrt{2\pi\left[1 + \frac{3}{16}A^2\right]^{-1}},$$

$$x^*(A) = \sqrt{2\pi\left[1 + \frac{3}{16}A^2\right]^{-1}}.$$

We then get (to lowest order)

$$\phi(x) = \begin{cases} -\sin\left(\left[1 + \frac{3}{16}A^2\right]\frac{x^2}{2}\right), & x_*(A) < x < 0, \\ 0, & x = 0, \\ \sin\left(\left[1 + \frac{3}{16}A^2\right]\frac{x^2}{2}\right), & 0 < x < x^*(A). \end{cases} \quad (32)$$

Note that $\phi(x_*(A)) = -\sin(\pi) = 0 = \sin(\pi) = \phi(x^*(A))$. Furthermore, let

$$x_1(A) = -\sqrt{\pi\left[1 + \frac{3}{16}A^2\right]^{-1}},$$

$$x_2(A) = \sqrt{\pi\left[1 + \frac{3}{16}A^2\right]^{-1}}.$$

Then $\phi(x_1(A)) = -\sin(\pi/2) = -1 \neq 1 = \sin(\pi/2) = \phi(x_2(A))$. So, there exist $x_1(A)$ and $x_2(A)$ such that $x_*(A) < x_1(A) < x_2(A) < x^*(A)$, $\phi(x_*(A)) = \phi(x^*(A))$, and $\phi(x_1(A)) \neq \phi(x_2(A))$; so a loop is indeed formed. The solution (32) is shown in Fig. 9, in the case of $A = 0.25$. The single-loop structure is prominent.




FIG. 9. (Color online) Plot of the analytical solution for a single-loop vortex filament described by $\phi(x)$ when $\phi(x)$ satisfies Eq. (32). The x scaling is $f(x) = x^2/2$, while the amplitude of the solution is taken to be $A = 0.25$. The space coordinate is parametrized by $x \in [x_*(A), x^*(A)]$ while on the loop.

We remark that since ϕ is continuous on $x \in [-\sqrt{\pi}, \sqrt{\pi}]$, yet ϕ' has a discontinuity at a single point $x = 0$, the matched solution is a class of “weak” solution.

VI. CONCLUSIONS

We have derived the fully nonlinear form of the LIA governing the motion of a vortex filament. Permitting a scaling of the free coordinate along which the vortex is aligned (x , in our case) permits us to have greater flexibility in computing solutions, both analytically and numerically. Such vortex solutions are a variation on the theme of planar vortex filaments and take the form

$$\mathbf{r} = [f(x), \cos(\gamma t)\phi(x), -\sin(\gamma t)\phi(x)].$$

The main analytical benefit is that such a solution form can capture a greater range of physical behaviors (particularly when the scale is nonmonotone), while numerical simulations can be made easier by taking the scale $f: \mathbb{R} \rightarrow \mathcal{I}$, where \mathcal{I} is a compact interval (numerical integration on such a compact interval can often be simpler than on an unbounded domain such as the real line).

In the case of monotone scalings $f(x) = \mu$, we have a very elegant way to determine the planar contribution $\phi(x) = \psi(\mu)$ to the vortex filament structure, obtaining a nonlinear ordinary differential equation (ODE) governing ψ [see Eq. (6)]. For monotone scalings, we therefore find that ψ is a strict function of μ and therefore the ODE (6) has only constant coefficients, making its solution possible. The planar solution is equivalent to a stationary solution of the form $\Phi = e^{-\gamma t}\psi(\mu)$. The main stationary solution of interest is periodic for small amplitudes A , so this is the solution we focus on in Sec. III. While numerical solutions can be obtained, we compute a perturbation solution, scaling both the function and the variable by the amplitude A of solutions through a multiple-scales approach. We compare the perturbation solution to numerical solutions, finding that the perturbation solution accurately captures the structure of the planar vortex filament (in particular, the spacial period of oscillation for such solutions). We find that the spatial period $T(A)$ is given by the approximation

$$T(A) \approx 2\pi\left(1 + \frac{3}{16}A^2\right)^{-1},$$

for small A .

Properties of the planar vortex filament in the arclength system (the intrinsic coordinate frame) are considered in Ref. [13], and in Sec. IV we have compared the two formulations. The primary benefit of the arclength frame is that it allows for exact solutions, in terms of elliptic sn functions. The Cartesian framework, however, gives us a clearer view of exactly what is going on with the structure of the vortex filament. While there is no exact solution, the perturbation result does work nicely for small-amplitude periodic solutions. Despite the fact that there is no exact closed-form solution for $\psi(\mu)$, we are able to derive an exact relation for the period $T(A)$ in terms of elliptic integrals, obtaining

$$T(A) = 8E(A/2) - 4K(A/2),$$

which agrees nicely with the approximation found through perturbation for small A (see Fig. 4). This is also reminiscent

of the period for the arclength representation of the planar solution discussed in Ref. [13]. Note that there is a bound $A < \sqrt{2}$ on the amplitude A of the space-periodic function $\psi(\mu)$, which will be discussed in a forthcoming work. As such, the maximal period occurs with amplitude $A = 0$ and is $T(0) = 2\pi$, while the minimal period solution occurs with amplitude $A = \sqrt{2}$ and is $T(\sqrt{2}) = 3.3886$. Hence, the period $T(A)$ of a space-periodic solution is related in an inverse manner to the amplitude A of such a solution.

For monotone scalings f we were able to obtain the nice analytical results discussed above. We also discuss nonmonotone coordinate scalings f and their application for finding self-intersections of vortex filaments. An equation governing the stationary solution $\phi(x)$ is given in Eq. (26). This equation becomes singular at points where monotonicity of f fails, yielding solutions which, in some cases, permit self-intersection of the curve \mathbf{r} given in Eq. (1). Such a self-intersection results in a vortex filament loop. While such a situation is not tractable analytically, we provide numerical simulations to demonstrate that such results are at the very least mathematically possible. We also outline some general criteria which would permit a loop filament

structure. These self-intersecting vortex filaments essentially “break” the LIA formulation, meaning that, once intersection occurs, the LIA is not sufficient to study the dynamics of the loop solutions. Such solutions are likely unstable and collapse into other structures or dissipate completely. These types of dynamics are quite interesting and certainly merit future work. Analytical results, under weaker conditions than monotonicity, yet stronger conditions than just arbitrary nonmonotone transforms, could be possible, maybe in the case of the specific examples considered here. Some analytical results were given for the nonmonotone scalings, and it was shown that such solutions may be constructed in a piecewise manner. These analytical solutions are continuous, yet fail to have a continuous derivative. In this sense, we may view such solutions as weak solutions. Nevertheless, these analytical results agree qualitatively with the numerical simulations.

ACKNOWLEDGMENTS

R.A.V. was supported in part by NSF Grant No. 1144246. The author appreciates the comments of the reviewers, which led to the improvement of the paper.

-
- [1] L. S. Da Rios, *Rend. Circ. Mat. Palermo* **22**, 117 (1906).
 - [2] R. J. Arms and F. R. Hama, *Phys. Fluids* **8**, 553 (1965).
 - [3] R. L. Ricca, *Nature* **352**, 561 (1991).
 - [4] S. Kida, *J. Fluid Mech.* **112**, 397 (1981).
 - [5] R. L. Ricca, *Chaos* **3**, 83 (1993); R. L. Ricca, D. C. Samuels, and C. F. Barenghi, *J. Fluid Mech.* **391**, 29 (1999); F. Maggioni, S. Z. Alamri, C. F. Barenghi, and R. L. Ricca, *Il Nuovo Cimento C* **32**, 133 (2009); *Phys. Rev. E* **82**, 026309 (2010).
 - [6] T. Lipniacki, *J. Fluid Mech.* **477**, 321 (2003).
 - [7] R. L. Ricca, *Phys Fluids A* **4**, 938 (1992).
 - [8] R. A. Van Gorder, *Theor. Comput. Fluid Dyn.* **26**, 161 (2012).
 - [9] R. A. Van Gorder, *Theor. Comput. Fluid Dyn.* **26**, 591 (2012).
 - [10] V. P. Dmitriyev, *Am. J. Phys.* **73**, 563 (2005).
 - [11] B. K. Shivamoggi and G. J. F. van Heijst, *Phys. Lett. A* **374**, 1742 (2010).
 - [12] M. Umeki, *Theor. Comput. Fluid Dyn.* **24**, 383 (2010).
 - [13] R. A. Van Gorder, *Phys. Rev. E* **86**, 057301 (2012).
 - [14] E. Fehlberg, *Computing (Arch. Elektron. Rechnen)* **6**, 61 (1970).