

Contact time periods in immunological synapse

Daniel R. Bush and Amit K. Chattopadhyay*

Non-linearity and Complexity Research Group - Aston University, Aston Triangle, Birmingham B4 7ET, United Kingdom

(Received 5 February 2014; revised manuscript received 25 June 2014; published 8 October 2014)

This paper resolves the long standing debate as to the proper time scale (τ) of the onset of the immunological synapse bond, the noncovalent chemical bond defining the immune pathways involving T cells and antigen presenting cells. Results from our model calculations show $\langle\tau\rangle$ to be of the order of seconds instead of minutes. Close to the linearly stable regime, we show that in between the two critical spatial thresholds defined by the integrin:ligand pair ($\Delta_2 \sim 40\text{--}45$ nm) and the T-cell receptor TCR:peptide-major-histocompatibility-complex pMHC bond ($\Delta_1 \sim 14\text{--}15$ nm), $\langle\tau\rangle$ grows monotonically with increasing coreceptor bond length separation δ ($= \Delta_2 - \Delta_1 \sim 26\text{--}30$ nm) while $\langle\tau\rangle$ decays with Δ_1 for fixed Δ_2 . The nonuniversal δ -dependent power-law structure of the probability density function further explains why only the TCR:pMHC bond is a likely candidate to form a stable synapse.

DOI: [10.1103/PhysRevE.90.042706](https://doi.org/10.1103/PhysRevE.90.042706)

PACS number(s): 87.16.dj, 05.40.-a, 87.18.Tt

I. INTRODUCTION

Cell to cell contacts define key chemical pathways that articulate immune response signaling through cellular signal transduction [1]. Signals are transported in the event of attached integrin-ligand pairs, whereupon they are carried through intracellular signaling pathways. Such biomechanical signaling, mediated by surrounding coreceptor molecules (CD4, CD8, CD45) proliferates an immune response in the body through cellular level interactions [2], often resulting in the formation of the immunological synapse (IS) bond between the immune cells (T and B cells) and the antigen presenting cells (APC). In T cells, some of these signaling pathways are directed toward the nucleus, where conformational changes lead to cell proliferation and triggering of the immune effector functions.

The T-cell:APC bond, a “close contact” patch between the membranes, is known to have a diameter ~ 10 μm and contains a large number of membrane bound molecules [3]. Important cell surface molecules on the T cell include the T-cell receptor (TCR) and the leukocyte function associated-1 adhesion molecule (LFA-1). The T-cell receptor (TCR) binds with peptide bound major histocompatibility complexes (pMHC) on the APC. Similarly, the integrin LFA-1 molecule has a natural ligand in the intercellular adhesion molecule-1 (ICAM-1) on the APC surface.

Nascent TCR:pMHC and LFA-1:ICAM-1 bonds begin to form following the initial cellular attachment. Fluorescent tagging reveals heterogeneous segregation and aggregation of the surface molecules throughout the interface [3,4], a result that is attributed to the presence of multiple length scales ($\sim 14\text{--}15$ nm for TCR:pMHC and $\sim 40\text{--}45$ nm for LFA-1:ICAM-1 [5,6]) leading to the dynamic reorganization of the cell surface molecules as a pathway towards IS formation [7–9]. Initially, the longer bonds (LFA-1:ICAM-1) localize at the center of the contact zone, with small patches of shorter bonds (TCR:pMHC) toward the edge of the contact zone. Such nonlinear patterning have been studied in details [10,11] with theoretical models convincingly proving that the dynamical

reorganization is the result of the presence of multiple length scales in the problem along with descriptions of conditions in which such changes in receptor configurations [3] occur. In what follows, we will use this fact as an input but otherwise focus on a different aspect of quantifying the strength of the immunological synapse bond that was beyond the scope of either of these papers.

From the binding assay measurements, we know that the affinity of the TCR-pMHC complex may vary drastically depending on the proliferated peptide, TCR sequences, and the MHC allele. What triggers the sequential patterning process leading primarily to the aggregation-segregation mechanism, followed by binding of the IS bond, are defined by the binding energies of the respective of the emergent synapse. While such a binding affinity is not exactly proportional to the separation length of the individual cell membranes, it has now been proved [12] that the TCR:pMHC interaction has to be long enough to complete proximal signaling while the dissociation rates have to be sufficiently short to allow multiple TCRs engaging with the same pMHC. As explained in [12], following an initial clustering of the TCRs on the cell surface, affinity rates are affected by the average “dwell times” with affinity increasing monotonically as dwell times up to a threshold followed by a saturation regime; so while it would be ideal to model the TCR:pMHC interaction with affinity as the switch, defining the average dwell time as our regulation “order parameter” allows for more direct theoretical modeling of the IS dynamics and in turn the patterning process. At this minimalist level, we are considering only one-no-one TCR-pMHC interactions which is why TCR intensity resulting from the aggregation on the cell membranes will only linearly affect the IS kinetics.

The early patterning signals cause polarization of the microtubule organizing center, effectively stopping T-cell migration and orienting the internal machinery toward the contact area. At a time scale of up to 5 min, the pattern inverts such that the TCR:pMHC bonds migrate to the center of the initial close-contact patch and the LFA-1:ICAM-1 bonds form a tight adhesion ring around the periphery [3]. Such a *pattern inversion* is strictly a “nonuniversal” feature, since dynamic interactions (kinapses) between the cells are also shown to initiate immune effector functions [13,14].

*a.k.chattopadhyay@aston.ac.uk

It has been largely recognized that the mature synapse is required to trigger the immune effector function [9]. A primary objective of this study is to focus on the kinetic behavior at the start of this dynamical bond formation process. When the TCR is attached to an agonist pMHC, intracellular signaling molecules can phosphorylate the cytoplasmic portion of the TCR and signal transduction occurs. However, it is not well understood if signaling continues after the TCR has disengaged from the pMHC, although some studies indicate this may be possible [7]. Recent studies have also identified TCR microclusters, small patches of membrane enriched in TCR and signaling molecules, continuously forming at the periphery of the synapse throughout the contact duration [15]. Signaling from these microclusters peaks while in the periphery and diminishes as they migrate toward the center of the synapse. The time dynamics of these signals are in the order of seconds, consistent with upregulation of Ca^{2+} levels, before the synapse matures minutes later [13].

The strength and start time of the IS bond patterning is defined by the average time the two randomly forced (due to thermal fluctuations) fluctuating membranes (T cell and APC) remain in contact with each other above a minimum threshold Δ that is defined by the bond lengths of the participating molecules, a model that was successfully implemented previously [16] in estimating the average lengths scale of the interacting T Cell:pMHC patch sizes. In line with the Chattopadhyay-Burroughs' model [16], here we analyze the average time of contact of these close-contact patches, at the start of patterning, based on an analogous one-membrane–two-threshold model (theoretical architecture follows [16]). This is a linearized (around the linearly stable fixed point) version of the nonlinear reaction-diffusion model due to Qi *et al.* [10,11,17,18] in line with our previous work in this sequel [16]. Numerical analysis of the nonlinear models [10,17] have been shown to be in near quantitative agreement with the early images of the synapse [3,4]. Our linearized model portrays the nonstationary state dynamics of a fluctuating membrane $\phi(\mathbf{x},t)$ close to the linearly stable point and across a range of mean separation distances defined by the bond lengths of relevant coreceptor molecules (15–45 nm). As shown in [10,19], incorporation of the first nonlinear (cubic) perturbation in the linearized (stochastic) model predicts a Hopf-bifurcation point below which the linear regime dominates and above which nonlinear patterning [17] takes over. Our focus here is to study the crossover from the linear to the nonlinear regime.

II. THE TCR:APC MEMBRANE FLUCTUATION MODEL

The dynamics that we are studying here stems from the interaction of two stochastically driven membranes. With a membrane separation distance designated by $\phi(\mathbf{x})$ where \mathbf{x} is any point on the membrane observed at time t , our linearized stochastic continuum model can be written as

$$M\dot{\phi} = -B\nabla^4\phi + \gamma\nabla^2\phi - \lambda\phi + \eta(\mathbf{x},t). \quad (1)$$

Here B is the coefficient of the membrane rigidity, γ is the surface tension, λ quantifies the linearized relaxation kinetics close to equilibrium, and M is the membrane damping constant. As in a standard membrane dynamics, the membrane

rigidity term and the surface relaxation terms create a force balance by working against each other while the contribution from the surrounding coreceptor molecules is encapsulated in the linear $-\lambda\phi$ term. The thermal noise $\eta(\mathbf{x},t)$ is assumed to be a spatio-temporally independent Gaussian white noise defined through fluctuation-dissipation kinetics [16]:

$$\langle \eta(\mathbf{x},t) \rangle = 0, \quad (2a)$$

$$\langle \eta(\mathbf{x},t)\eta(\mathbf{x}',t') \rangle = 2k_B T M \delta(\mathbf{x} - \mathbf{x}')\delta(t - t'). \quad (2b)$$

Although broadly argued from a thermodynamic perspective, the above stochastic (model) has its origin in a more detailed nonlinear dynamical system architecture as propounded in [10,11]. The linearized model focuses on the contact events (TCR:APC synapse) and arises from a linear stability analysis of the nonlinear model defined in these references close to the equilibrium point. Detailed descriptions of the linear stability criteria and its application are available in [11] and have been previously employed in [16]. It must be mentioned that the range of validity of this linear model is limited to the start of the immunological synapse patterning and can not account for the eventual self-organized criticality leading to the mature synapse formation.

A. The two threshold model

Our starting model is a generalized version of the model described in [16]. In this earlier work, we defined a one membrane model fluctuating across a threshold as an analog of our physical system. In that model, this single threshold cloned the TCR:APC and other small coreceptor bond length range but considered all larger length bonds (e.g., integrin:ligand) as a constant. This present improved model considers two thresholds Δ_1 (TCR:pMHC) and Δ_2 (integrin:ligand) in acknowledgment of the presence of two different length scales, one large and other small, in a reminder of [17]. Two opposing points on the membranes are said to be within close contact if the separation distance is less than $-\Delta_i$ nm ($i = 1,2$), that, in the one membrane model, translates to a configuration of a fluctuating membrane staying above a critical threshold $-\Delta_i$ nm through a certain average distance $\langle X_+ \rangle$ [16] and an average time $\langle \tau^+ \rangle$ (result from this article). Figure 1 explains the time persistence behavior of the fluctuating membrane where $t^{\pm}_{\Delta_i}$ ($i = 1,2$) gives the “bottom-up/up-bottom” crossover times across the lines $\phi = \Delta_i$, where the duration of time for close contact is given by $t^-_{\Delta_i} = t_2 - t_1$. In accordance with the reflection symmetry of this model, $t^+_{\Delta_i} = t_2 - t_1$ represents the close-contact duration under the conditions

$$\phi(t_1) = \phi(t_2) = \Delta_i, \quad (3a)$$

$$\phi(t) \geq \Delta_i, \quad t_1 < t < t_2. \quad (3b)$$

We shall use the latter definition for notational purposes, as detailed in the Fig. 1 caption.

B. Sec. II A for a two threshold model

As in [16], we assume each successive crossing of the $\phi = \Delta_i$ line to be statistically independent {*independent interval approximation* (Sec. II A) [21,22]}. The time persistence characteristics between two thresholds Δ_1 and




FIG. 1. t_{Δ}^+ and t_{Δ}^- regions: The time evolution of the separation distance for a point in the membrane interface, $\phi(\mathbf{x}, t)$. The $t_{\Delta_i}^+$ regions are periods of time where close contact exists between the membrane surfaces at distances Δ_i ($i = 1, 2$) and during the $t_{\Delta_i}^-$ period the membrane separation distance is not favorable for ligand-receptor bond formation. The shaded regions indicate the time persistence [20] between two thresholds, Δ_1 and Δ_2 .

Δ_2 ($\Delta_2 > \Delta_1$) that are separated by a distance $\delta = \Delta_2 - \Delta_1$ represents a set of four different events where a signal ϕ persists between the two thresholds (Fig. 1): (a) the fluctuation enters the δ region from below the lower threshold, persists within the δ region, and returns below the lower threshold t_{11} ; (b) the fluctuation enters the δ region from below the lower threshold, persists, and becomes larger than the upper threshold t_{12} ; (c) the fluctuation enters the δ region from above the upper threshold, persists, and exits below the lower threshold t_{21} ; and finally, (d) the fluctuation enters from above the upper threshold, persists, and exits above the upper threshold t_{22} . A consummate representation of the average time of persistence considering all four scenarios together can then be defined as

$$\langle t_{\delta} \rangle = \sum_{i,j=1}^2 w_{ij}(\delta) \langle t_{ij}(\delta) \rangle, \quad (4)$$

where $w_{ij}(\delta)$ is the δ -dependent probability of occurrence of the event t_{ij} , with $\sum_{i,j=1}^2 w_{ij}(\delta) = 1$. Detailed quantitative depictions of statistics in each of these zones will be detailed in the following section. In the numerical simulation of the model, the above normalization condition was always adhered to.

III. ANALYSIS

The theoretical routine focuses both on the analytical as well as on the numerical aspects. For the former, the target is to recast the model solutions within a Gaussian stationary process (GSP) framework. For the latter, Euler integration of the stochastic model is followed by the estimation of the probably density function of the persistent crossings. The following subsections detail these separately.

A. On to GSP

In the Fourier transformed space, Eq. (1) admits of a stochastic solution

$$\phi(\mathbf{x}, t) = \frac{1}{2\pi M} \int d\mathbf{k} \int_0^t e^{-\alpha(\mathbf{k})(t-t') + i\mathbf{k}\cdot\mathbf{x}} \tilde{\eta}(\mathbf{k}, t') dt' + \tilde{\phi}_0, \quad (5)$$

where $\alpha(\mathbf{k}) = \frac{Bk^4 + \gamma k^2 + \lambda}{M}$ under initial conditions $\tilde{\phi}_0$. $\phi(\mathbf{x}, t)$ is a *Gaussian process* with zero mean, whereupon it can be entirely characterized by the two point correlation function (vide [16] for detailed reference; notations are in line with this parent article)

$$\begin{aligned} c_{12}(t_1, t_2) &= \langle \phi(\mathbf{x}, t_1) \phi(\mathbf{x}, t_2) \rangle \\ &= \frac{k_B T}{(2\pi)^2 M} \int d\mathbf{k} \frac{e^{-\alpha(\mathbf{k})(t_2 - t_1)}}{\alpha(\mathbf{k})} \\ &\quad - \frac{k_B T}{(2\pi)^2 M} \int d\mathbf{k} \frac{e^{-\alpha(\mathbf{k})(t_1 + t_2)}}{\alpha(\mathbf{k})}, \end{aligned} \quad (6)$$

where $0 < t_1 < t_2$. In the large time limit, where any of t_1 or t_2 is large with $|t_2 - t_1|$ still being finite, the second integral term tends to zero and we have a form that is solely dependent on $\tau = |t_2 - t_1|$, which is a *Gaussian stationary process* (GSP), with the following correlation functions:

$$c_{12}(\tau) = \frac{k_B T}{(2\pi)^2 M} \int d\mathbf{k} \frac{e^{-\alpha(\mathbf{k})\tau}}{\alpha(\mathbf{k})}, \quad (7)$$

$$c_{12}(0) = c_{11} = \frac{k_B T}{(2\pi)^2 M} \int d\mathbf{k} \frac{1}{\alpha(\mathbf{k})}. \quad (8)$$

The above is the representation of our linearized nonequilibrium model in the Gaussian stationary state limit. Since we are only interested in close-contact times, we introduce the formalism of a conditional correlator for an arbitrary time t during cell-to-cell contact by using the variable $\sigma = \text{sgn}(\phi(\mathbf{x}, t) - \Delta_i)$, that changes sign about $\phi(\mathbf{x}, t) = \Delta_i$ [21]. The conditional correlator $A_+(\phi_1, \phi_2) = \langle \text{sgn}(\phi(\mathbf{x}, t_1) - \Delta_i) \text{sgn}(\phi(\mathbf{x}, t_2) - \Delta_i) \rangle$ for states where $\phi > \Delta_i$ can be expressed as

$$A_+(\phi_1, \phi_2) = N \int_{\Delta_i}^{\infty} d\phi_1 e^{-\frac{\text{detc}}{2c_{11}} \phi_1^2} \int_{\Delta_i}^{\infty} d\phi_2 e^{-\frac{c_{11}}{2} (\phi_2 + \frac{c_{12}}{c_{11}} \phi_1)^2}, \quad (9)$$

with $\text{detc} = c_{11}^2 - c_{12}^2$ and the normalization factor

$$N = \frac{\sqrt{\text{detc}}}{\text{acot}\left(\frac{c_{12}}{\sqrt{\text{detc}}}\right)}.$$

As before, $\phi_j = \phi(\mathbf{x}, t_j)$ ($j = 1, 2$) and the curly brackets “ $\langle \rangle$ ” refer to ensemble averages over all noise realizations. In the limit of Sec. II A, the probability that the field ϕ has crossed the line $\phi = \Delta_i$ in a small enough time interval τ will be given by $\frac{\tau}{\langle \tau^+ \rangle}$, leading to the relation $A_+ = 1 - \frac{\tau}{\langle \tau^+ \rangle}$ [21,22]. This leads to

$$\langle \tau^+ \rangle = -\frac{1}{A'_+}, \quad (10)$$

where A'_+ is the partial derivative of A_+ with respect to τ . In practice, due to the lack of a closed form solution, $A'_+ = \frac{\partial A_+}{\partial c_{12}} \cdot \frac{\partial c_{12}}{\partial \tau}$ is being evaluated numerically.

B. Numerical scheme

Continuum Eq. (1) is discretized and simulated using a first-order forward difference Euler scheme

$$\phi_{i,j}(t + \Delta t) = \phi_{i,j}(t) + \Delta t \dot{\phi}_{i,j}(t) \quad (11)$$

on a two-dimensional mesh with $\Delta x = \Delta y = 1$, using periodic boundary conditions. We use the experimental parameter settings in accord with cellular membranes [16]: $M = 4.7 \times 10^6 k_B T s \mu m^{-4}$, $B = 11.8 k_B T$, $\gamma = 5650 k_B T \mu m^{-2}$, and $\lambda = 6.0 \times 10^5 k_B T \mu m^{-4}$. For each term to contribute to the dynamics, simple dimensional analysis will show that $B \nabla^4[\phi] \sim \gamma \nabla^2[\phi] \sim \lambda[\phi]$. The numbers here are all in energy units that help us to nondimensionalize the eventual outcomes. The noise in this case is perturbative whose strength is just enough to stimulate dynamical fluctuations. In what follows, we compare results obtained from the approximate analytical solutions (Sec. II A based) with the numerical evaluation of the starting model equation.

IV. RESULTS

Figure 2 shows the ensemble average for the time persistence above any of the (Δ_1 or Δ_2) thresholds keeping the other fixed, for both the numerical and the scaled analytical solution. The parameters used give rise to persistent close-contact patches in the order of magnitude required for the biological problem, that is, 10–50 nm. The numerical results (in dots: Fig. 2) are consistent with the analytical solutions (continuous line: Fig. 2), where the close-contact time decreases as the separation distance increases, suggesting that the TCR:pMHC bond persists longer than the LFA-1:ICAM-1.

The result shown in Fig. 2 can be qualitatively understood from simple probabilistic considerations. As the threshold value increases, it becomes more difficult for the randomized (Gaussian) fluctuations to cross this threshold, resulting in reduced average time spent above the threshold value. More nontrivial, though, is the functional nature of the decay in




FIG. 2. (Color online) Time persistence for different thresholds, $\langle \tau^+ \rangle$ vs Δ_i : The ensemble average for the time persistence above the Δ_i threshold. The dots show the result obtained by numerically solving Eq. (1) based on the scheme detailed in Eq. (11) while the solid line indicates the analytical result obtained from a solution of Eq. (10). The results have been linearly scaled for comparison.




FIG. 3. (Color online) Time persistence, $\langle \tau_\delta \rangle$ vs δ : The ensemble average time persistence between two thresholds, where the distance between the thresholds is given by δ .

the $\langle \tau^+ \rangle$ value against Δ . As opposed to a simplistic (and incorrect) visual impression, the decay profile here is not exponential, rather it is defined through an intricate balance between power-law scaled fluctuations against the statistics of deterministically decaying membrane fluctuation modes.

A vital part of this model study is the analysis of the dynamics of the randomly driven membrane in between the two given thresholds. In line with the parlance used previously as well as in [16], this can be represented by an estimation of the time persistence between two threshold values, Δ_1 and Δ_2 , where Δ_1 is the T-cell analog of the TCR:pMHC bond length (~ 15 nm) while Δ_2 symbolizes the ICAM-1:LFA-1 bond separation length (~ 45 nm).

The result for the variation of the average time between the two thresholds as a function of the distance δ between the thresholds is shown in Fig. 3. The calculations were done by starting with $\Delta_1 = 1$ nm and then varying Δ_2 between 1 and 50 nm. The results shown in Fig. 2 are the average over multiple such initial choices of Δ_1 and then varying Δ_2 accordingly to generate the appropriate range for δ .

As previously, the dotted points refer to the numerical simulation results while the continuous line represents the interpolation of the same to maintain continuity. The time persistence initially increases as δ increases, then it saturates and asymptotically approaches the $\langle \tau^+ \rangle_{\Delta_1}$ value. As $\delta \rightarrow 0$ the average time persistence tends toward the smallest time length scale used, for our case $dt = 0.1$. Once again, a quantitative understanding of Fig. 3 can be had from the fact that an increase in the δ value can be wrought about in either of the two possible ways—an increase in Δ_2 for fixed Δ_1 , or else a decrease in Δ_1 for a fixed Δ_2 . For the first case when Δ_2 increases at a fixed Δ_1 , it is easier for a fluctuation mode to remain within the upper limit Δ_2 than to cross it while for the second case, with a fixed value of the upper threshold Δ_2 , a lowering of Δ_1 increases the probability of a fluctuation mode crossing this line resulting in an increased value of $\langle \tau_\delta \rangle$.

Figures 2 and 3, respectively, express the variation of the average “persistent times” against bond lengths above a critical threshold and that in between two thresholds (Δ_1 and Δ_2). These results are subsets of a bigger ensemble defined by a membrane that fluctuates across two thresholds Δ_1




FIG. 4. (Color online) Variation of the probability density function $p(\tau)$ with τ for $\delta = 15$ nm between two thresholds for the τ_{11} , τ_{21} , τ_{21} , and τ_{22} cases.

and Δ_2 , or else that of two membranes whose fluctuations are measured across a single threshold Δ , that, as already explained earlier, are equivalent analytical descriptions. In line with our model of a single membrane fluctuating across two thresholds detailed in the previous section and as depicted in Fig. 1, the statistics can be classified in to four broad zones—“11”, “12”, “21”, and “22”. While 11 defines the fluctuation regime for a crossing from a region in $\phi < \Delta_1$ across the line $\phi = \Delta_1$ but for $\phi < \Delta_2$, 22 encapsulates the complementary regime for a crossing across $\phi = \Delta_2$ from a point $\phi > \Delta_2$. 12 and 21 represent statistics when crossings are restricted within $\Delta_1 < \phi < \Delta_2$ as explained in Fig. 1. Representing the corresponding average persistent time scales by τ_{ij} ($i, j = 1, 2$), we find that due to reasons of reflection symmetry in the time correlators [$A_+(t_+, \Delta_1) = -A_-(t_-, \Delta_2)$] and [$A_-(t_+, \Delta_1) = -A_+(t_-, \Delta_2)$], τ_{11} and τ_{22} are identical, as are τ_{12} and τ_{21} . Comparing with the notations used previously, $\tau_{12}(=\tau_{21})$ may be identified with τ_δ while $\tau_{11}(=\tau_{22})$ may be




FIG. 5. (Color online) Variation of the probability density function $p(\tau_{12})$ with time τ_{12} between thresholds Δ_1 and Δ_2 for a range of δ values. As expected, the smallest δ ($=15$ nm) turns up the highest peak (represented by crosses), followed by $\delta = 20$ nm (squares), with $\delta = 25$ nm showing the lowest peak with the maximum spread (solid squares).

identified with $\langle \tau^+ \rangle$ under the constraint $\phi < \Delta_2$. For the same reason (reflection symmetry), as shown in Fig. 4, the respective density distributions too conform to these symmetry lines.

The result presented in Fig. 4 is not a special case, as shown in Fig. 5, where the τ_{11} and τ_{12} scenarios are plotted for three values of δ .

As δ increases, the probability density for the τ_{12} case shows an increase in the probability for longer time persistence, a result that matches with the observation presented earlier through Fig. 3. On the other hand, the number of τ_{12} events become less frequent as δ is increased. The τ_{11} densities also change slightly, again showing an increase in the probability for longer time persistence.

V. CONCLUSIONS

The analysis presented here has two major immunological implications. First, Figs. 2 and 3 clearly prove that the onset of patterning at the immature kinapse level, when the central LFA-1:ICAM-1 bond gives way to the smaller TCR:pMHC bond, occurs at the time scale of seconds. This discovery is expected to confirm the start time of mature synapse formation. Admittedly, though, the parameter values used make the result a subjective case in that the time scale predicted for a different membrane:membrane dynamics may as well be in minutes or hours, instead of in seconds. Our calculation pins down the time scale to within 2–4 s that is a further improvement on the <12 s window as suggested in [12]. Second, the nonuniversal character of the time correlation of the IS bond as evident in the dependence on the separation length δ and separation times τ , ensuring that the probability density $p(t)$ is a function of the system parameters along with being functions of δ (Fig. 3) and τ_{12} (Fig. 5), confirms a widely acknowledged belief in the community that the TCR:pMHC bond is non-self-organizing in nature and hence is the only stable bond at this spatio-temporal regime. An immediate impact of this can be seen in the projected time scale for the crossover from the linear to the nonlinear phase that, as is shown in Fig. 3, spans 2–4 s (peak time of the PDF profile). The implication of this analysis is that of a time scale difference of an order of magnitude related to the start time of the “immature” IS bond formation, a time scale that is also associated with the transition from the linear to the nonlinear regime (and hence our emphasis on a study of the linear stability regime of an otherwise nonlinear dynamics). As to how such crossover is affected by kinase-phosphatase pathways (these pathways act as signal transduction inhibitors and thereby control the rate of the IS bond formation) and what modification this may bring about in the prediction of the time scale of a mature IS bond are some of the exciting topics that we are presently working on. Results are shortly to be communicated on the quantitative nature of the extremal value statistics of these fluctuations and how such nonuniversal exponents affect the life time and strength of IS bonds.

ACKNOWLEDGMENT

AKC acknowledges insightful discussions with Darren Flower.

- [1] K. Murphy, P. Travers, and M. Walport, *Janeway's Immunobiology* (Garland Science, Taylor & Francis, New York, 2006).
- [2] K. Choudhuri, J. Llodra, E. W. Roth, J. Tsai, S. Gordo, K. W. Wucherpfennig, L. C. Kam, D. L. Stokes, and M. L. Dustin, *Nature (London)* **507**, 118 (2014).
- [3] A. Grakoui, S. K. Bromley, C. Sumen, M. M. Davis, A. S. Shaw, P. M. Allen, and M. L. Dustin, *Science* **285**, 221 (1999).
- [4] C. R. F. Monks, B. A. Frieberg, H. Kupfer, N. Sciaky, and A. Kupfer, *Nature (London)* **395**, 82 (1998).
- [5] A. N. Barclay, M. H. Brown, S. K. A. Law, A. J. McKnight, M. G. Tomlinson, and P. A. van der Merwe, *The Leucocyte Antigen Factsbook*, 2nd ed., Factsbook Series (Academic Press Ltd., London, 1997).
- [6] T. A. Springer, *Nature (London)* **346**, 425 (1990).
- [7] S. C. Bunnell, in *Immunological Synapse*, edited by B. Saito (Springer, London, 2010).
- [8] T. Yokosuka and T. Saito, in *Immunological Synapse*, edited by T. Saito and F. D. Batista (Springer, London, 2010).
- [9] S. J. Davis and P. A. van der Merwe, *Nat. Immunol.* **7**, 803 (2006).
- [10] S. Y. Qi, J. T. Groves, and A. Chakraborty, *Proc. Natl. Acad. Sci. (USA)* **98**, 6548 (2001).
- [11] N. J. Burroughs and C. Wülfing, *Biophys. J.* **83**, 1784 (2002).
- [12] J. D. Stone, A. S. Chervin, and D. M. Kranz, *Immunology* **126**, 165 (2009).
- [13] M. L. Dustin, *Immunological Rev.* **221**, 77 (2008).
- [14] M. Gunzer, A. Schäfer, S. Borgmann, S. Grabbe, K. S. Zänker, E.-B. Bröcker, and P. Friedl, *Immunity* **13**, 323 (2000).
- [15] T. Yokosuka, K. Sakata-Sogawa, W. Kobayashi, M. Hiroshima, A. Hashimoto-Tane, M. Tokunaga, M. L. Dustin, and T. Saito, *Nat. Immunol.* **6**, 1253 (2005).
- [16] A. K. Chattopadhyay and N. J. Burroughs, *Europhys. Lett.* **77**, 48003 (2007).
- [17] N. J. Burroughs, K. Köhler, V. Miloserdov, M. L. Dustin, P. A. van der Merwe, and D. M. Davis, *PLoS Comput. Biol.* **7**, e1002076 (2011).
- [18] S. Raychaudhuri, A. K. Chakraborty, and M. Kardar, *Phys. Rev. Lett.* **91**, 208101 (2003).
- [19] A. Košmrlj, M. Kardar, and A. K. Chakraborty, *Annu. Rev. Condens. Matter Phys.* **4**, 339 (2013).
- [20] C. Sire, *Phys. Rev. Lett.* **93**, 130602 (2004).
- [21] S. N. Majumdar and A. J. Bray, *Phys. Rev. Lett.* **81**, 2626 (1998).
- [22] B. Derrida, V. Hakim, and R. Zeitak, *Phys. Rev. Lett.* **77**, 2871 (1996).