

Ice nucleation on carbon surface supports the classical theory for heterogeneous nucleation

Raffaella Cabriolu and Tianshu Li*

Department of Civil and Environmental Engineering, George Washington University, Washington, DC 20052, USA

(Received 13 December 2014; published 13 May 2015)

The prevalence of heterogeneous nucleation in nature was explained qualitatively by the classical theory for heterogeneous nucleation established over more than 60 years ago, but the quantitative validity and the key conclusions of the theory have remained unconfirmed. Employing the forward flux sampling method and the coarse-grained water model (mW), we explicitly computed the heterogeneous ice nucleation rates in the supercooled water on a graphitic surface at various temperatures. The independently calculated ice nucleation rates were found to fit well according to the classical theory for heterogeneous nucleation. The fitting procedure further yields the estimate of the potency factor, which measures the ratio of the heterogeneous nucleation barrier to the homogeneous nucleation barrier. Remarkably, the estimated potency factor agrees quantitatively with the volumetric ratio of the critical nuclei between the heterogeneous and homogeneous nucleation. Our numerical study thus provides a strong support to the quantitative power of the theory and allows understanding ice nucleation behaviors under the most relevant freezing conditions.

DOI: [10.1103/PhysRevE.91.052402](https://doi.org/10.1103/PhysRevE.91.052402)

PACS number(s): 64.60.Q–, 64.70.D–, 82.60.Nh

I. INTRODUCTION

The freezing of water nearly all proceeds with assistance from foreign substances, a process known as heterogeneous nucleation. In clouds, the dominant candidates for heterogeneous ice nucleation are bacteria, pollen grains, mineral dusts, soot particles, and high-molecular-weight organic compounds [1,2]. Despite its ubiquity, the microscopic picture behind such prevailing process still remains elusive because of the complex and stochastic nature of the heterogeneous nucleation event. In particular, the mechanisms controlling heterogeneous ice nucleation are not well understood.

Although a molecular understanding is still missing, the thermodynamic rationale behind the heterogeneous nucleation was already provided in 1950s by the classical theory for heterogeneous nucleation [3,4], an extension to the classical nucleation theory (CNT) [5] for homogeneous nucleation, on the basis of macroscopic arguments. According to CNT, the formation of a critical nucleus needs to overcome a free energy barrier ΔG_{hom}^* through spontaneous fluctuations. In the case of a spherical solid nucleus forming from the supercooled liquid, the free energy barrier can be expressed as

$$\Delta G_{\text{hom}}^* = \frac{16\pi\gamma_{ls}^3}{3(\rho\Delta\mu_{ls})^2}, \quad (1)$$

where γ_{ls} is the solid-liquid interface free energy, $\Delta\mu_{ls}$ is the chemical potential difference between liquid and solid, and ρ is the density of liquid. The homogeneous nucleation rate R_{hom} varies with the nucleation temperature T following the Arrhenius equation [6]:

$$R_{\text{hom}} = A_{\text{hom}} \exp(-\Delta G_{\text{hom}}^*/k_B T), \quad (2)$$

where A_{hom} is the kinetic prefactor. For homogeneous ice nucleation, both experiments [7–11] and simulations [12–17] suggest that the temperature dependence of the homogeneous ice nucleation rate may be quantitatively described by CNT,

with parametrization being refined by the controlled experiments [18].

When a foreign flat wall (W) is present, the solid nucleus can preferentially form at the interface between the liquid and the wall [Fig. 1(a)]. At its critical size, the solid embryo is under the unstable equilibrium with respect to the dissolution and the growth, which also indicates a mechanical equilibrium. At the liquid-solid-wall triple junction, solving the equation of equilibrium yields the Young's equation

$$\gamma_{lw} = \gamma_{sw} + \gamma_{ls} \cos \theta_c, \quad (3)$$

where γ_{lw} and γ_{sw} are the surface tensions for liquid-wall and solid-wall interfaces. θ_c defines the contact angle of solid embryo on the flat wall, with $\theta_c = 0$ and $\theta_c = 180^\circ$ indicating the complete wetting of the wall by solid and liquid, respectively. If the solid nucleus is further assumed to be part of the sphere, i.e., a spherical cap, its volume V_{cap} can be expressed as $V_{\text{cap}} = f(\theta_c)V_{\text{sphere}}$, where

$$f(\theta_c) = (1 - \cos \theta_c)^2(2 + \cos \theta_c)/4, \quad (4)$$

and V_{sphere} is the volume of the sphere containing the cap. Remarkably, under the framework of classical theory for heterogeneous nucleation, the factor $f(\theta_c)$ coincides with the ratio of the free energy barriers between the heterogeneous and homogeneous nucleation, i.e., $f(\theta_c) = \Delta G_{\text{het}}^*/\Delta G_{\text{hom}}^*$. It thus follows that

$$f(\theta_c) = \frac{\Delta G_{\text{het}}^*}{\Delta G_{\text{hom}}^*} = \frac{V_{\text{cap}}}{V_{\text{sphere}}}. \quad (5)$$

Equation (5) provides a simple but robust explanation for the preference of the heterogeneous nucleation over the homogeneous nucleation. Instead of forming a spherical nucleus from spontaneous thermal fluctuations, only part of the sphere V_{cap} needs to be nucleated when a foreign surface is present. Accordingly, the free energy barrier is reduced by the same factor $f(\theta_c)$ by which the volume of critical nucleus is reduced. Since $f(\theta_c)$ measures the degree of the free energy reduction, it is also known as the potency factor. According to Eq. (4), the potency factor $f(\theta_c)$ for a foreign wall

*Corresponding author: tsli@gwu.edu




FIG. 1. (Color online) (a) A solid critical nucleus forms at the interface between liquid and a foreign flat wall, with a solid contact angle θ_c . The contact angle θ_c can be determined by the surface tensions through Eq. (3). The solid nucleus is assumed to be part of a sphere (dashed line) with a radius r^* , i.e., a spherical cap with the volume $V_{\text{cap}} = f(\theta_c)V_{\text{sphere}}$. (b) The potency factor $f(\theta_c)$ increases from 0 to 1 (homogeneous), as the solid contact angle θ_c varies from 0 to 180°.

is determined by the solid contact angle θ_c and varies between 0 and 1, as shown in Fig. 1(b). A wall with a lower solid contact angle yields a lower potency factor $f(\theta_c)$, thus further enhancing heterogeneous nucleation. Then the heterogeneous nucleation rate can be expressed by

$$R_{\text{het}} = A_{\text{het}} \exp[-f(\theta_c)\Delta G_{\text{hom}}^*/k_B T]. \quad (6)$$

Although the classical theory for heterogeneous nucleation offers a qualitative explanation to the prevalence of heterogeneous nucleation, its quantitative validity remains unconfirmed. Auer and Frenkel [19] employed the umbrella sampling method to calculate the nucleation barrier of the hard-sphere crystal that completely wets the smooth walls, and found that the computed barrier height is substantially higher than that predicted by the CNT. The disagreement was attributed to the omission in the CNT of the line tension at the liquid-solid-wall triple junction, which may become non-negligible when crystal completely wets the wall. In a recent study by Winter *et al.* [20], the total surface energies of the liquid nuclei (of both spherical and spherical cap shape) were directly obtained by Monte Carlo simulation for the Ising lattice gas model. It was found that the obtained surface energies could be comparable with the capillary approximation

employed by CNT, if the line tension effects at the triple junction are considered.

In this work, we show that the heterogeneous ice nucleation on a graphitic surface indeed supports the quantitative power of the theory. In particular, the validity of Eqs. (5) and (6) is strongly supported from the ice nucleation rates computed explicitly using the forward flux sampling method over a wide temperature range. Our work thus provides the first validation of heterogeneous CNT in the partially wetting regime where the potency factor is far enough from zero.

II. SIMULATION DETAILS

Our molecular dynamics (MD) simulations were carried out using the coarse-grained model of water (mW) [21]. The intermolecular interaction between water and carbon was adopted from a recent parametrization of the two-body term of the mW model, so that the strength of the water-carbon interaction reproduces the experimental contact angle (86°) of water on graphite [22]. The model was recently employed in direct MD simulations to study the heterogeneous ice nucleation on carbon surfaces [22–24], where the nonequilibrium freezing temperature of ice was found to increase due to the preferential nucleation of ice on a carbon surface. Here we employ the forward flux sampling (FFS) method [25,26] to systematically and explicitly compute the heterogeneous ice nucleation rates at various temperatures where spontaneous ice nucleation becomes too slow to occur in direct simulation. The details of the rate constant calculations can be found in Appendix A. Our MD simulation includes 4096 water molecules and 1008 carbon atoms, in a nearly cubic cell with a periodic boundary condition. The isobaric-isothermal canonical ensemble (NPT) with a Noe-Hoover thermostat was employed, with a relaxation time of 1 and 15 ps for temperature and pressure, respectively. A time step of 5 fs was used. It should be noted that while the homogeneous nucleation rate is measured by the nucleation frequency per unit volume, the heterogeneous nucleation rate should be characterized by the nucleation frequency per unit area. However, because the simulation volume of liquid is small and ice nucleation on the carbon surface is strongly preferred, it is convenient to describe the heterogeneous nucleation rate R_{het} on the basis of volume in order to facilitate a direct comparison with R_{hom} .

III. NUMERICAL VERIFICATION OF EQ (5)

Figure 2 shows the computed heterogeneous ice nucleation rates (in logarithm) as a function of nucleation temperature, in the range of 220–240 K. To quantitatively explore the catalytic activity of the graphitic surface, we compare the obtained heterogeneous ice nucleation rates with the reported homogeneous ice nucleation rates from the previous work [12] using the FFS method and the mW water model, as shown in Fig. 2. It is clear that a graphitic surface yields the significantly enhanced ice nucleation rates under all temperatures studied. Our results thus support the finding by Lupi *et al.* [22] and confirm the enhanced ice nucleation capacity of the carbon surface.

The calculated heterogeneous ice nucleation rates at various temperatures allow assessing the quantitative validity of




FIG. 2. (Color online) Temperature dependence of ice nucleation rate (logarithm) in the mW water model for both homogeneous ice nucleation and the heterogeneous nucleation on a graphitic surface. The solid red dots and blue squares represent the calculated ice nucleation rates by using the FFS method. The dashed red and blue lines indicate the fitting on the basis of the CNT for homogeneous nucleation [Eq. (7)] and the extension for heterogeneous nucleation [Eq. (8)], respectively. The data of homogeneous nucleation were extracted from Ref. [12].

Eq. (6). To do this, we fit the obtained heterogeneous ice nucleation rates at various temperatures according to the theory of nucleation, using the procedure employed by Li *et al.* in analyzing the homogeneous ice nucleation [12]. In this procedure, the chemical potential difference $\Delta\mu$ is approximated as a linear function of temperature, i.e., $\Delta\mu = H(T - T_m)/T_m$, where T_m is the equilibrium melting temperature (274.6 K) of ice in the mW model and H is a constant. The liquid-solid interface energy γ_{ls} is assumed to be temperature independent. It is noted that both assumptions have been verified for the mW water model in different simulation studies [17,27]. For homogeneous ice nucleation, it was shown that the independently calculated homogeneous ice nucleation rate can be fitted according to the following expression:

$$\ln(R_{\text{hom}}) = \ln(A_{\text{hom}}) + \frac{C_{\text{hom}}}{(T - T_m)^2 T}, \quad (7)$$

where $\ln(A_{\text{hom}}) = 114.07 \pm 1.86$ and $C_{\text{hom}} = -16\pi\gamma_{ls}^3 T_m^2 / (3k_B\rho^2 H^2) = -3.72 \pm 0.08 \times 10^7 \text{ K}^3$ are the fitting constants [12]. We note that the nucleation barrier $\Delta G_{\text{hom}}^* = k_B C_{\text{hom}} / (T - T_m)^2$. The fitting yielded an estimate of $\gamma_{ls} = 31.01 \pm 0.21 \text{ mJ m}^{-2}$, which agrees well with the surface tensions computed through other approaches for the mW water model [17,28].

Remarkably, the obtained heterogeneous ice nucleation rates are also found to fit well the classical theory for heterogeneous nucleation, as shown in Fig. 2. Specifically, the calculated ice nucleation rates R_{het} can be fitted according to

$$\ln(R_{\text{het}}) = \ln(A_{\text{het}}) + \frac{C_{\text{het}}}{(T - T_m)^2 T}. \quad (8)$$

The fitting yields the estimate of the kinetic prefactor for heterogeneous nucleation $\ln(A_{\text{het}}) = 102 \pm 7.70$, which

is consistent with that for the homogeneous nucleation $\ln(A_{\text{hom}}) = 114.07 \pm 1.86$ [12]. More importantly, the other fitting constant $C_{\text{het}} = -1.70 \pm 0.07 \times 10^7 \text{ K}^3$ allows estimating the reduction of the nucleation barrier, as $C_{\text{het}}/C_{\text{hom}} = \Delta G_{\text{het}}^* / \Delta G_{\text{hom}}^* \equiv f_b(\theta_c)$. By comparing the fitting constants C from the heterogeneous and homogeneous ice nucleation, we obtain the potency factor for the graphitic surface $f_b(\theta_c) = 0.456 \pm 0.019$. This corresponds to a solid contact angle of $\theta_c \sim 86.6^\circ$. It is noted here that θ_c is the contact angle between ice and graphene and should not be confused with the water-graphene contact angle 86° , although their magnitudes coincide here.

It is then of interest to further test the validity of Eq. (5), namely, the potency factor $f_b(\theta_c)$ can be also quantitatively related to the volumetric ratio $f_v(\theta_c)$ of the critical nucleus of the heterogeneous nucleation and the homogeneous nucleation. We note that such verification becomes possible in our study because the size of the critical nucleus λ^* can be independently estimated from the ensemble of nucleation trajectories obtained in the FFS calculation. Using the definition that the critical nucleus has the equal probabilities of dissolving and growing completely, i.e., with a committer probability $p_{\mathbf{B}} = 0.5$ [29], we obtained the estimate of the critical nucleus size λ (number of water molecules contained in the critical ice nucleus) at various nucleation temperatures, as shown in Fig. 3. According to CNT, the critical size of the spherical nucleus in homogeneous nucleation is expressed by $\lambda_{\text{hom}}^* = 32\pi\gamma^3 / [3\rho^2(\Delta\mu)^3]$. For the mW water model, since γ is nearly temperature independent and $\Delta\mu = H(T - T_m)/T_m$ [17], the critical size λ_{hom}^* exhibits the following temperature dependence:

$$\lambda_{\text{hom}}^*(T) = \frac{32\pi\gamma^3}{3\rho^2 H^3} \frac{1}{\left(\frac{T_m}{T} - 1\right)^3} = \frac{B_{\text{hom}}}{\left(\frac{T_m}{T} - 1\right)^3}, \quad (9)$$

where B_{hom} is the temperature-independent constant. The obtained critical size λ_{hom}^* at various temperatures are found to fit well according to Eq. (9), as shown in Fig. 3(a). The good fit is not unexpected because previous studies [13,15] have shown the critical ice nucleus from homogeneous nucleation is nearly spherical. The fitting procedure yields the constant $B_{\text{hom}} = 1.752 \pm 0.027$. For the heterogeneous ice nucleation, the same fitting procedure was found to equivalently apply to the calculated critical nucleus size λ_{het}^* , through

$$\lambda_{\text{het}}^*(T) = \frac{B_{\text{het}}}{\left(\frac{T_m}{T} - 1\right)^3}, \quad (10)$$

which yields the constant $B_{\text{het}} = 0.840 \pm 0.014$. By comparing the two fitting constants B , one obtains the volumetric ratio $f_v(\theta_c) \equiv \lambda_{\text{het}}^* / \lambda_{\text{hom}}^* = B_{\text{het}} / B_{\text{hom}}$. Remarkably, the obtained volumetric ratio $f_v(\theta_c) = 0.480 \pm 0.011$ agrees *quantitatively* with the potency factor $f_b(\theta_c) = 0.456 \pm 0.019$ estimated from the nucleation barriers. The quantitative validity of Eq. (5), an important conclusion from CNT and its extension, is thus strongly supported through the molecular simulation results based on the mW water model.




FIG. 3. (Color online) (a) Variation of the critical ice nucleus size with temperature for both homogeneous and heterogeneous nucleation. The critical size is estimated based on the committor probability analysis, which yields λ_{het}^* : 145 ± 5 at 225 K, 205 ± 10 at 230 K, 288 ± 7 at 235 K, and 410 ± 10 at 240 K, respectively. (See Appendixes A and B for more details.) The critical size for homogeneous nucleation was obtained in Ref. [12]. The simulation data and the fitted curves are represented by data points and solid lines, respectively. (b) A snapshot of the critical ice nucleus forming from homogeneous ice nucleation at 240 K. (c) A snapshot of the critical ice nucleus forming on graphene surface at 240 K.

IV. DISCUSSION

The verified quantitative validity of the CNT (and its extension) then allows predicting ice nucleation behavior in the presence of a heterogeneous nucleation center. The nucleation efficacy of the foreign surface can be generally described based on its potency factor f . Using the fitted kinetic prefactor A_{hom} ($\approx A_{\text{het}}$) and C_{hom} , and Eq. (8), one obtains both the heterogeneous ice nucleation rate and the corresponding critical nucleus size, as a function of the nucleation temperature, for different potency factors f . As shown in Fig. 4, the predicted ice nucleation rates clearly indicate the preference and the relevance of the heterogeneous nucleation at the moderate and low supercooling. For example, at 250 K, a nucleation center with the potency factor of $f = 0.1$ (equivalent to a solid contact angle $\theta_c = 52.5^\circ$) yields an ice nucleation rate about 100 orders of magnitude higher than that of homogeneous nucleation at the same temperature. Intriguingly, the sizes of the critical nuclei from these relevant nucleation events fall within the range of a few hundred to a few thousand water molecules. This implies that the most relevant ice nucleation events mediated by an effective nucleation center under a low supercooling, *i.e.*, $-10^\circ\text{C} \sim -20^\circ\text{C}$, can




FIG. 4. (Color online) The predicted temperature variation of (a) ice nucleation rate $R(T)$ and (b) critical size of ice nucleus $\lambda^*(T)$ in the mW water model due to the presence of a heterogeneous nucleation center with different potency factor f . A small potency factor f indicates a strong ice nucleation efficacy, and $f = 1$ corresponds to homogeneous ice nucleation. (c) The predicted variation of the critical size λ^* for different potency factor f , subject to a fixed nucleation rate.

be possibly modeled by molecular simulations through using a reasonable number ($10^3 \sim 10^4$) of water molecules.

Interestingly, when the *ice nucleation rate* R is fixed, the theory of nucleation predicts that the critical size λ^* decreases

with the potency factor f [Fig. 4(c)], and a homogeneous nucleation yields the minimum critical nucleus. This may appear surprising, but the prediction can be understood by the fact that the heterogeneous nucleation producing the same ice nucleation rate occurs at a much elevated temperature [Fig. 4(a)]. For example, an ice nucleation rate of $10^{20} \text{ m}^{-3} \text{ s}^{-1}$ would require a homogeneous nucleation temperature $T_{\text{hom}} \approx 225 \text{ K}$ but a heterogeneous nucleation temperature $T_{\text{het}} \approx 260 \text{ K}$ for the nucleation center with a potency factor $f = 0.1$. The prediction [Fig. 4(c)] shows the critical nucleus for such heterogeneous nucleation at 260 K contains about 1220 water molecules (i.e., 1/10 of the critical size of the homogeneous nucleation at 260 K), larger than the critical size (310) of the homogeneous nucleation at 225 K. As the solid contact angle θ_c decreases with the nucleation efficacy (Fig. 1), a strong ice nucleation center yields a more “flat” ice nucleus that appears increasingly two-dimension like. It is noted that in such scenarios the possible effect from the line tension at the triple junction can be non-negligible [19,20]. However, its quantitative effect on the ice nucleation rate is unclear. Using the density of ice, one can estimate the radius of the spherical segment (i.e., the frustum of the spherical cap) to be of the order of a few nanometers, implying that the dimension of an effective ice nucleation site is typically of the order of 10^1 nm . This estimate may be used as an important parameter in experiments for potentially observing ice nucleation *in situ* and designing an effective strategy for controlling ice nucleation.

ACKNOWLEDGMENTS

The authors thank V. Molinero for valuable discussion. The authors acknowledge support from the American Chemical Society Petroleum Research Fund through Grant No. 52101-DNI6, the NSF through Grant No. CBET-1264438, and the Sloan Foundation through the Deep Carbon Observatory.

APPENDIX A: CALCULATION OF ICE NUCLEATION RATES

We employed the forward flux sampling (FFS) method [26] to compute the ice nucleation rate on graphitic surface. The method has been applied to successfully study the homogeneous ice nucleation [12] using the mW water model. In FFS, the rate constant for the transition from the basin A to the basin B is obtained through the “effective positive flux” expression [30]: $R_{AB} = \dot{\Phi}_{\lambda_0} P(\lambda_B|\lambda_0)$, where $\dot{\Phi}_{\lambda_0}$ and $P(\lambda_B|\lambda_0)$ are the initial flux rate crossing the first interface λ_0 from basin A , and the probability for a trajectory starting from λ_0 to reach basin B , respectively. The interfaces are




FIG. 5. (Color online) Calculated growth probability $P(\lambda|\lambda_0)$ as a function of ice nucleus size λ for heterogeneous ice nucleation on a graphitic surface at 225 K, 230 K, 235 K, and 240 K. The interfaces λ_i used in FFS simulation correspond to the abscissa of the data points.

defined based on an order parameter λ , i.e., number of water molecules contained in the largest ice cluster. The ice-like and liquid-like water molecules are identified based on the local order parameter q_6 [12], and an ice-like water molecule is defined as the one with $q_6 > 0.5$. More details of computing q_6 can be found in Ref. [12]. The initial flux rate can be computed by direct molecular dynamics simulation, through $\dot{\Phi}_{\lambda_0} = N_0/(t_0 V)$, where N_0 is the number of successful crossings to the interface λ_0 from basin A , t_0 is the total time of the initial sampling, and V is the simulation volume. As our recent work [31] has demonstrated the importance of the initial sampling on the reliability of the final nucleation rate, particularly in a heterogeneous environment, a sufficiently large sampling time ($t_0 > 300 \text{ ns}$) is used for computing $\dot{\Phi}_{\lambda_0}$ to ensure the convergence. The growth probability $P(\lambda_B|\lambda_0)$ is obtained through $P(\lambda_B|\lambda_0) = \prod_{i=1}^n P(\lambda_i|\lambda_{i-1})$ and the individual crossing probability $P(\lambda_i|\lambda_{i-1}) = N_i/M_{i-1}$, where N_i is the number of successful crossings (~ 110) to the interface λ_i , and M_{i-1} is the number of trial shootings (typically of 10^3) from the interface λ_{i-1} . More details for computing $P(\lambda_B|\lambda_0)$ are explained in Refs. [32] and [12]. The statistical uncertainty of nucleation rate is mainly attributed to the error in the calculated $P(\lambda_i|\lambda_{i-1})$, which includes the variance of the binomial distribution of N_i and the landscape variance of the configurations collected at the previous interface λ_{i-1} [26]. The calculated initial flux rate $\dot{\Phi}_{\lambda_0}$, growth probability $P(\lambda_B|\lambda_0)$, and the final nucleation

TABLE I. Calculated initial flux rate $\dot{\Phi}_{\lambda_0}$, growth probability $P(\lambda_B|\lambda_0)$, and nucleation rate R for heterogeneous ice nucleation on a graphitic surface using the mW water model and the FFS method. The critical nucleus size λ^* is determined by the committor probability analysis on the basis of the computed growth probability.

T/K	$\dot{\Phi}_{\lambda_0} (\text{m}^{-3} \text{ s}^{-1})$	$P(\lambda_B \lambda_0)$	Nucleation rate $R (\text{m}^{-3} \text{ s}^{-1})$	Critical size λ^*
225	1.69×10^{34}	$7.64 \pm 2.16 \times 10^{-4}$	$1.29 \pm 0.36 \times 10^{31}$	145 ± 5
230	1.12×10^{34}	$4.83 \pm 1.83 \times 10^{-6}$	$5.41 \pm 2.05 \times 10^{28}$	205 ± 10
235	8.90×10^{33}	$1.51 \pm 0.74 \times 10^{-10}$	$1.34 \pm 0.66 \times 10^{24}$	288 ± 7
240	4.19×10^{33}	$2.23 \pm 1.49 \times 10^{-15}$	$9.34 \pm 6.24 \times 10^{18}$	410 ± 10




FIG. 6. (Color online) Calculated committor p_B as a function of the nucleus size λ for heterogeneous ice nucleation on a graphitic surface at various nucleation temperatures. The dotted lines indicate the upper and lower boundaries of the calculated p_B on the basis of the estimated error bar of p_B . The horizontal dash line corresponds to a $p_B = 0.5$ and intersects the both boundaries of the calculated p_B , which yields the estimate of the uncertainty of λ^* .

rates R are listed in Table I for the heterogeneous ice nucleation at 225, 230, 235, and 240 K. The computed growth probability $P(\lambda|\lambda_0)$, as a function of nucleus size λ , is also shown in Fig. 5.

At 220 K, ice nucleation becomes accessible in direct molecular dynamics simulation. Three out of four trajectories lead to spontaneous ice crystallization, with a total simulation time around 30 ns, which yields an estimate of the nucleation rate of the order of $10^{32} \text{ m}^{-3} \text{ s}^{-1}$.

The size of the critical ice nucleus λ^* can be numerically determined by the committor probability p_B [29]. A critical crystalline nucleus is defined as the one with 50% probability of growing completely into crystal, i.e., $p_B = 0.5$. The committor p_B as a function of nucleus size λ is obtained on the basis of the computed growth probability $P(\lambda_i|\lambda_{i-1})$, and the




FIG. 7. (Color online) The computed histogram of committor p_B for heterogeneous ice nucleation at 230 K at the interface $\lambda = 210$. The Gaussian distribution function with the same intrinsic mean and standard deviation is shown as the red line.

critical sizes λ^* under different temperatures are determined (Fig. 6) and listed in Table I.

APPENDIX B: p_B HISTOGRAM ANALYSIS

To understand whether the size of the nucleus λ is a good approximation to the actual reaction coordinate, we carried out the committor distribution analysis near the critical size $\lambda^* \approx 205$ for the heterogeneous nucleation at 230 K. There were 110 configurations selected from the interface $\lambda = 210$ and each one received 20 shootings with Boltzmann distributed velocities. The obtained committor distribution is shown in Fig. 7, along with the Gaussian distribution with the intrinsic mean μ and standard deviation σ [33]. The calculated distribution is peaked at $\mu = 0.521$, with an overall agreement with the intrinsic committor distribution, indicating that λ fairly accurately describes the actual reaction coordinate and is thus a good order parameter.

-
- [1] W. Cantrell and A. Heymsfield, *Bull. Am. Meteorol. Soc.* **86**, 795 (2005).
- [2] B. J. Murray, D. O'Sullivan, J. D. Atkinson, and M. E. Webb, *Chem. Soc. Rev.* **41**, 6519 (2012).
- [3] D. Turnbull, *J. Chem. Phys.* **18**, 198 (1950).
- [4] D. Turnbull, *J. Appl. Phys.* **21**, 1022 (1950).
- [5] M. Volmer and A. Weber, *Z. Phys. Chem. (Leipzig)* **119**, 277 (1926).
- [6] K. Kelton, in *Solid State Physics—Advances in Research and Applications*, edited by H. Ehrenreich and D. Turnbull (Academic Press, New York, 1991), Vol. 45, p. 75.
- [7] B. Kramer, O. Hubner, H. Vortisch, L. Woste, T. Leisner, M. Schwell, E. Ruhl, and H. Baumgartel, *J. Chem. Phys.* **111**, 6521 (1999).
- [8] T. Koop, B. Luo, A. Tsias, and T. Peter, *Nature (London)* **406**, 611 (2000).
- [9] P. Stockel, I. Weidinger, H. Baumgartel, and T. Leisner, *J. Phys. Chem. A* **109**, 2540 (2005).
- [10] P. Taborek, *Phys. Rev. B* **32**, 5902 (1985).
- [11] B. J. Murray, S. L. Broadley, T. W. Wilson, S. J. Bull, R. H. Wills, H. K. Christenson, and E. J. Murray, *Phys. Chem. Chem. Phys.* **12**, 10380 (2010).
- [12] T. Li, D. Donadio, G. Russo, and G. Galli, *Phys. Chem. Chem. Phys.* **13**, 19807 (2011).
- [13] T. Li, D. Donadio, and G. Galli, *Nat. Commun.* **4**, 1887 (2013).
- [14] E. B. Moore and V. Molinero, *Nature (London)* **479**, 506 (2011).
- [15] A. Reinhardt and J. P. K. Doye, *J. Chem. Phys.* **136**, 054501 (2012).
- [16] E. Sanz, C. Vega, J. R. Espinosa, R. Caballero-Bernal, J. L. F. Abascal, and C. Valeriani, *J. Am. Chem. Soc.* **135**, 15008 (2013).

- [17] J. R. Espinosa, E. Sanz, C. Valeriani, and C. Vega, *J. Chem. Phys.* **141**, 18C529 (2014).
- [18] T. Koop and B. Zobrist, *Phys. Chem. Chem. Phys.* **11**, 10839 (2009).
- [19] S. Auer and D. Frenkel, *Phys. Rev. Lett.* **91**, 015703 (2003).
- [20] D. Winter, P. Virnau, and K. Binder, *Phys. Rev. Lett.* **103**, 225703 (2009).
- [21] V. Molinero and E. B. Moore, *J. Phys. Chem. B* **113**, 4008 (2009).
- [22] L. Lupi, A. Hudait, and V. Molinero, *J. Am. Chem. Soc.* **136**, 3156 (2014).
- [23] L. Lupi and V. Molinero, *J. Phys. Chem. A* **118**, 7330 (2014).
- [24] A. Reinhardt and J. P. K. Doye, *J. Chem. Phys.* **141**, 084501 (2014).
- [25] R. J. Allen, D. Frenkel, and P. R. T. Wolde, *J. Chem. Phys.* **124**, 024102 (2006).
- [26] R. J. Allen, D. Frenkel, and P. R. T. Wolde, *J. Chem. Phys.* **124**, 194111 (2006).
- [27] L. C. Jacobson, W. Hujo, and V. Molinero, *J. Phys. Chem. B* **113**, 10298 (2009).
- [28] D. T. Limmer and D. Chandler, *J. Chem. Phys.* **137**, 044509 (2012).
- [29] P. G. Bolhuis, D. Chandler, C. Dellago, and P. L. Geissler, *Annu. Rev. Phys. Chem.* **53**, 291 (2002).
- [30] T. van Erp, D. Moroni, and P. Bolhuis, *J. Chem. Phys.* **118**, 7762 (2003).
- [31] Y. Bi and T. Li, *J. Phys. Chem. B* **118**, 13324 (2014).
- [32] T. Li, D. Donadio, and G. Galli, *J. Chem. Phys.* **131**, 224519 (2009).
- [33] B. Peters, *J. Chem. Phys.* **125**, 241101 (2006).