

**Accuracy of rate coding: When shorter time window and higher spontaneous activity help**Marie Levakova,<sup>1,\*</sup> Massimiliano Tamborrino,<sup>2</sup> Lubomir Kostal,<sup>1</sup> and Petr Lansky<sup>1</sup><sup>1</sup>*Department of Computational Neuroscience, Institute of Physiology of the Czech Academy of Sciences, Videnska 1083, 14220 Prague 4, Czech Republic*<sup>2</sup>*Institute for Stochastics, Johannes Kepler University Linz, Altenbergerstraße 69, 4040 Linz, Austria*

(Received 22 November 2016; published 22 February 2017)

It is widely accepted that neuronal firing rates contain a significant amount of information about the stimulus intensity. Nevertheless, theoretical studies on the coding accuracy inferred from the exact spike counting distributions are rare. We present an analysis based on the number of observed spikes assuming the stochastic perfect integrate-and-fire model with a change point, representing the stimulus onset, for which we calculate the corresponding Fisher information to investigate the accuracy of rate coding. We analyze the effect of changing the duration of the time window and the influence of several parameters of the model, in particular the level of the presynaptic spontaneous activity and the level of random fluctuation of the membrane potential, which can be interpreted as noise of the system. The results show that the Fisher information is nonmonotonic with respect to the length of the observation period. This counterintuitive result is caused by the discrete nature of the count of spikes. We observe also that the signal can be enhanced by noise, since the Fisher information is nonmonotonic with respect to the level of spontaneous activity and, in some cases, also with respect to the level of fluctuation of the membrane potential.

DOI: [10.1103/PhysRevE.95.022310](https://doi.org/10.1103/PhysRevE.95.022310)**I. INTRODUCTION**

The question of how the information is encoded in the brain activity remains one of the major unresolved problems in neuroscience. A pioneering work in this direction was done by Adrian [1], who did experiments on the stretch receptor in a muscle spindle and demonstrated that the number of spikes emitted by the receptor neuron increases with the force applied to the muscle. Since then, the neuronal firing rate has been traditionally thought to contain relevant information about external stimuli. The firing rate is usually determined by counting the number of spikes that occur within a predefined time window [2]. The irregularities encountered in real spike trains make the determination of the firing rate complicated and thus a sufficiently long time window is necessary to achieve a satisfactory precision.

It is clear that an approach based only on the firing rate neglects all the information possibly contained in the exact timing of the spikes [3–5], leading to the concept of temporal coding [6–10]. One of the common objections against reducing the complexity of neural coding only to firing rates is based on behavioral experiments, which suggest that reaction times are often rather short [11] and thus are against the common understanding that the reliability of rate coding increases proportionally to the applied time window.

In the past two decades, the performance of both temporal and rate coding was investigated for various settings, ranging from single neurons to neuronal networks (see, for example, [12–18]). The problem of quantifying the code performance is often approached by examining the accuracy with which the stimulus can be ultimately decoded from the observed response and the best decoding precision is evaluated using the Cramér-Rao lower bound on the mean-square error (see, e.g., [19–24]). Since the probability distributions of the number

of spikes are rarely known for advanced neuronal models, two approaches can be mostly encountered in the literature. Either processes based on a statistical description of the neuronal activity and enabling a straightforward way to express the count of spikes are used, e.g., Bernoulli or Poisson processes [25–27], or the rate is inferred from interspike intervals [28].

In this work the probability distribution of the number of spikes in a fixed time window for the stochastic perfect integrate-and-fire model is derived. We then use Fisher information to infer the rate coding accuracy and investigate how it is influenced by the key parameters of the model. In particular, our first goal is to study the role of the observation time window. Throughout the paper, the time window is always the window used by the nervous system and not a window determined by an external observer. Our second goal is to investigate how the amount of noise in the form of either presynaptic spontaneous activity or the random fluctuation of the membrane potential impacts the decoding accuracy and if there is an optimal level of noise enhancing the signal, like, for example, in stochastic resonance and other similar phenomena, which were found in many settings (see, e.g., [29–36]). As we show, counting spikes in a longer time window does not always improve the decoding accuracy. Moreover, the loss in decoding precision caused by a change in the length of the time window might be, at least partially, compensated by increasing the level of spontaneous activity of presynaptic neurons. Finally, we identify several stochastic-resonance-like phenomena related to the level of spontaneous activity and the level of fluctuation of the membrane potential.

**II. METHODS****A. Fisher information**

A common approach to analyze the decoding performance is to ask how well an optimal decoder can estimate the true value of the presented stimulus  $s$  based on a stochastic neuronal response  $R$  [26,31,37,38]. To quantify the estimation accuracy,

\*marie.levakova@fgu.cas.cz

the mean-square error is used,

$$\text{MSE}(\hat{s}) = \mathbb{E}[(\hat{s} - s)^2], \quad (1)$$

where  $\hat{s}$  is the estimator of  $s$  from  $R$ . If the estimator  $\hat{s}$  is unbiased, i.e.,  $\mathbb{E}(\hat{s}) = s$ , the mean-square error is equal to the variance of the estimator  $\text{MSE}(\hat{s}) = \text{var}(\hat{s})$ . According to the Cramér-Rao theorem [39,40], every unbiased and regular estimator  $\hat{s}$  satisfies the inequality

$$\text{var}(\hat{s}) \geq \frac{1}{J(s)}, \quad (2)$$

where  $J(s)$  denotes the Fisher information about  $s$  carried by  $R$ . For the sake of simplicity and analytical tractability, many theoretical studies on neural coding use the inverse of Fisher information as an approximation of the mean-square error, which implies that a higher Fisher information is assumed to reflect a higher decoding accuracy [29,31].

From the rate coding perspective, the response  $R$  is the number of spikes after the stimulus onset. Hence,  $R$  is a discrete quantity, with probability mass function  $p_R(r; s)$ , and the Fisher information is given by

$$J(s) = \sum_r \frac{1}{p_R(r; s)} \left[ \frac{\partial}{\partial s} p_R(r; s) \right]^2. \quad (3)$$

Note that the stimulus level  $s$  plays the role of a parameter of the distribution of  $R$ .

## B. Neuronal model

Throughout the study, we describe the neuronal activity by the stochastic perfect integrate-and-fire model introduced by Gerstein and Mandelbrot [41]. The idea of the model is to approximate the membrane potential dynamics by a Wiener process  $X(t)$ , which is given as the solution to a stochastic differential equation

$$dX(t) = \mu dt + \sigma dW(t), \quad X(0) = 0,$$

where  $W(t)$  is the standard (driftless) Wiener process,  $\mu > 0$  is the drift, and  $\sigma > 0$  is the diffusion parameter. The membrane potential is thus driven by a deterministic force  $\mu$  accompanied by white noise. Once  $X(t)$  reaches a constant threshold  $B > 0$ , a spike is elicited,  $X(t)$  is immediately reset to its starting value 0, and the accumulation of the membrane potential starts anew. The resulting spike train is then a renewal point process, where interspike intervals are independent and identically distributed as  $T \sim IG(B/\mu, B^2/\sigma^2)$ , an inverse Gaussian distribution with mean  $\mathbb{E}(T) = B/\mu$ , and variance  $\text{var}(T) = B\sigma^2/\mu^3$ .

Denote by  $X_0$  the value of the membrane potential at a randomly chosen time. The distribution of  $X_0$  can be derived from the known transition probability density of the Wiener process under the absorbing boundary  $B$ , which gives the probability density that the process starting from zero at time zero attains the value  $x$  at time  $t$  while the boundary  $B$  has not been crossed during that time [42,43],

$$f_X^B(x, t) = \frac{1}{\sqrt{2\pi\sigma^2 t}} \left\{ \exp \left[ -\frac{(x - \mu t)^2}{2\sigma^2 t} \right] - \exp \left[ \frac{2\mu B}{\sigma^2} - \frac{(x - 2B - \mu t)^2}{2\sigma^2 t} \right] \right\}. \quad (4)$$

The conditional density of  $X_0$  given that the time elapsed since the last spike is  $t$  can be written as

$$f_{X_0|T>t}(x|t) = \frac{f_X^B(x, t)}{1 - F_T(t)}, \quad (5)$$

where  $F_T(t)$  is the cumulative distribution function (CDF) of the interspike interval  $T$ . The time interval between the randomly chosen time point and the last preceding spike is the backward recurrence time  $S$ , which has the probability density function (PDF)  $f_S(t) = [1 - F_T(t)]/\mathbb{E}(T)$  [44]. Multiplying the conditional density  $f_{X_0|T>t}(x|t)$  by  $f_S(t)$  and integrating over all possible  $t$  yields the unconditional distribution of  $X_0$  as follows:

$$\begin{aligned} f_{X_0}(x) &= \int_0^\infty f_{X_0|T>t}(x|t) f_S(t) dt = \int_0^\infty \frac{f_X^B(x, t)}{\mathbb{E}(T)} dt \\ &= \frac{1}{B} \left[ \exp \left( \frac{\mu(x - |x|)}{\sigma^2} \right) - \exp \left( \frac{2\mu(x - B)}{\sigma^2} \right) \right]. \end{aligned} \quad (6)$$

We consider a situation where a stimulus of level  $s$  is presented at time  $t_0$ . The spontaneous activity before  $t_0$  results only from the spontaneous activity of presynaptic neurons, while the evoked activity after  $t_0$  is affected by the stimulation. Before  $t_0$ , the parameters of the Wiener process are  $\mu = \mu_0$  and  $\sigma^2 = \sigma_0^2$ . The presentation of the stimulus of intensity  $s$  at  $t_0$  changes the parameters to  $\mu = \mu(s)$  and  $\sigma^2 = \sigma^2(s)$  (Fig. 1).

Throughout the paper, we assume that the evoked drift  $\mu(s)$  is the sum of the spontaneous drift  $\mu_0$  and the stimulus-driven increment  $\Delta\mu(s)$ . The specific functional form of  $\mu(s)$  is derived from the Hill function [45], which is frequently employed in both theoretical and experimental studies [46–50]. For a stimulus level  $s$  expressed on a logarithmic




FIG. 1. Schematic illustration of the studied model. Shown on top is a sample path of the membrane potential  $X(t)$ , described by a Wiener process. When  $X(t)$  exceeds a constant threshold  $B > 0$ , an action potential is generated. Then  $X(t)$  is reset to 0 and its evolution starts anew. At time  $t_0$ , a stimulus is applied and the parameters of the process change from  $\mu_0$  and  $\sigma_0^2$  to  $\mu(s)$  and  $\sigma^2(s)$ . The times from the stimulus onset to the first, second, third, etc.,  $n$ th evoked spike are denoted by  $T_1, T_2, T_3, \dots, T_n$ . Shown on the bottom is the corresponding counting process of evoked spikes  $N(t^*)$ , which gives the number of spikes in a time window  $[t_0, t_0 + t^*]$ .

scale, it can be written as

$$\mu(s) = \mu_0 + \frac{A}{1 + e^{-b(s-s_0)}}, \quad s \in (-\infty, \infty). \quad (7)$$

The parameter  $A$  is the maximum possible increment in the drift  $\mu(s)$ ,  $b$  controls the steepness of the function, and  $s_0$  is the inflection point.

For the diffusion parameter  $\sigma^2(s)$ , we assume a linear dependence on the drift  $\mu(s)$ , that is,

$$\sigma^2(s) = k\mu(s) + m, \quad (8)$$

where  $k, m \geq 0$ . The linear relationship  $\sigma^2(s) = k\mu(s) + m$  can be derived for the membrane potential described by a randomized random walk with Poissonian input, where the amount of excitatory input is proportional to the stimulus intensity and the inhibitory input is constant [51]. By letting either  $k$  or  $m$  be equal to zero, we obtain the following two special cases:  $\sigma^2(s) = k\mu(s)$ , which corresponds to balanced excitatory and inhibitory input [52,53], and  $\sigma^2(s) = m$ , which, together with the requirement  $\lim_{s \rightarrow -\infty} \sigma^2(s) = \sigma_0^2$ , gives a diffusion parameter independent of the stimulation, that is,  $\sigma^2(s) = \sigma_0^2$ . In the following we study the accuracy with which the stimulus intensity  $s$  can be decoded from  $N(t^*)$ , the number of spikes observed in a time window of duration  $t^*$ .

### III. RESULTS

#### A. Fisher information

The reported Fisher information  $J(s)$  about  $s$  based on the observation of  $N(t^*)$  was calculated numerically using the formulas for the distribution of  $N(t^*)$  given in the Appendix. Throughout the paper, we consider two different scenarios corresponding to two possible beginnings of the observation time window. (i) The observation time window starts with an evoked spike. In this scenario the initial value of the membrane potential is known and  $X_0 = 0$  [Figs. 2 and 4(a)]. (ii) The observation time window starts at  $t_0$ . Here the initial value of the membrane potential  $X_0$  is random and has the PDF (6), with  $\mu = \mu_0$  and  $\sigma^2 = \sigma_0^2$  [Figs. 3 and 4(b)]. We focus on the dependence of the Fisher information on the duration of the time window  $t^*$ , but we also take into account the level of spontaneous activity  $\mu_0$  and the parameters  $k$  and  $m$  governing the diffusion parameter  $\sigma^2(s)$ . Without loss of generality, the threshold  $B$  is set to  $B = 1$  and, as a consequence, drift  $\mu$  represents also the firing rate. Moreover, we set  $A = 50$ ,  $b = 1$ , and  $s_0 = 0$ .

##### 1. Fisher information with respect to the length of the time window $t^*$

In general, Fisher information roughly increases as the time window gets longer [see Figs. 2(a) and 2(b), where the observation time window starts with a spike, i.e.,  $X_0 = 0$ , and Figs. 3(a) and 3(b), where the time window starts at  $t_0$ , i.e.,  $X_0$  is random]. In both cases, however, the Fisher information is nonmonotonic and has many local maxima and minima. The local extremes are more pronounced when  $X_0 = 0$ , but they are also clearly visible when  $X_0$  is random. Therefore, observing the process for a longer time may not necessarily lead to a higher decoding accuracy.

##### 2. Fisher information with respect to the level of spontaneous activity $\mu_0$

When the response starts with a spike, the Fisher information has again a nonmonotonic behavior with respect to  $\mu_0$ , with many local maxima and minima [Figs. 2(a), 2(c), 2(d), 2(f), 2(g), and 2(i)], and a qualitatively similar behavior is also observed when  $X_0$  is random [Figs. 3(a), 3(c), 3(d), 3(f), 3(g), and 3(i)]. Therefore, increasing the amount of presynaptic spontaneous activity may result in both improving or deteriorating the accuracy of decoding the stimulus, depending on the particular parameter conditions. For example, from Fig. 2(c) we observe that the Fisher information can be initially decreasing or increasing with respect to  $\mu_0$  depending on whether the observation time window is small ( $t^* = 0.007$  and  $0.02$  s) or large ( $t^* = 0.04$  s).

If we ignore local extremes, the overall tendency is that the Fisher information decreases with respect to  $\mu_0$  due to the variability of the underlying process, which is linearly proportional to  $\mu_0$ . When  $k = 0$ , however, the Fisher information has a mild overall increasing tendency (Fig. 4). This is because more spikes can be observed in the same time window due to the higher spontaneous activity, while the variability of the process remains constant (Fig. 4).

It is interesting to look at the mutual effect of both the spontaneous activity and the time window on the Fisher information [see Figs. 2(a) and 3(a), where  $X_0 = 0$  and  $X_0$  is random, respectively]. The white dashed lines displayed in the figures mark all points where the time window is  $t^* = nB/\mu(s)$ ,  $n \in \mathbb{N}$ , which approximately correspond to the local maxima of the Fisher information with respect to  $t^*$  for fixed  $\mu_0$  derived when  $\sigma_0^2 = \sigma^2(s) = 0$ , as discussed in more detail in Sec. III B. If the parameters of the system are chosen to be near a local maximum of  $J(s)$  with respect to  $t^*$  and then the observation time window is taken shorter, the loss in the decoding accuracy can be to some extent compensated by increasing the level of spontaneous activity  $\mu_0$ . Similarly, if the level of spontaneous activity is increased, by shortening the observation window we may get almost the original decoding accuracy.

##### 3. Fisher information with respect to the diffusion coefficient $\sigma^2(s)$

If  $X_0$  is random, increasing the diffusion parameter  $\sigma^2(s)$  through increasing  $k$  or  $m$  always yields a lower Fisher information [Figs. 3(d)–3(i)]. On the other hand, when  $X_0 = 0$ , in some cases the Fisher information is initially increasing with respect to  $k$  and  $m$  [Figs. 2(d)–2(i)], meaning that a positive level of noise enhances the signal. There are also cases, though, where the Fisher information is strictly decreasing with respect to  $k$  or  $m$ . Whether  $J(s)$  is increasing or decreasing in a given situation depends on the level of spontaneous activity. If the spontaneous drift  $\mu_0$  is approximately around

$$\mu_0 \approx \frac{nB}{t^*} - \frac{A}{1 + \exp[-b(s - s_0)]}, \quad n \in \mathbb{N}, \quad (9)$$

then increasing  $\sigma^2(s)$  has the effect of deteriorating the decoding accuracy. On the other hand, if the spontaneous drift




FIG. 2. Fisher information  $J(0)$  about the stimulus intensity  $s = 0$  based on the number of spikes  $N(t^*)$  in a time window of length  $t^*$ . The observation time window starts with a spike and the initial value of the membrane potential at the beginning of the time window is thus  $X_0 = 0$ . In all cases, the diffusion coefficient is  $\sigma^2(s) = k\mu(s) + m$ . (a)–(c) Fisher information as a function of the spontaneous firing rate  $\mu_0$  and the duration of the time window  $t^*$  when  $k = 0.01$  and  $m = 0.5$ . The Fisher information is nonmonotonic with respect to both  $\mu_0$  and  $t^*$ . (d)–(f) Fisher information as a function of the spontaneous firing rate  $\mu_0$  and the absolute coefficient  $m$  of the diffusion parameter  $\sigma^2(s)$  when  $t^* = 0.025$  s and  $k = 0.01$ . (g)–(i) Fisher information as a function of the spontaneous firing rate  $\mu_0$  and the coefficient of proportionality  $k$  of the diffusion parameter  $\sigma^2(s)$  when  $t^* = 0.025$  s and  $m = 0.5$ . For some values of  $\mu_0$  [dashed and dot-dashed lines in (e) and (f)], the Fisher information is initially increasing with respect to  $k$  and  $m$  and thus with respect to the fluctuation level of the membrane potential. The white dashed lines in (a), (d), and (g) mark the points satisfying  $t^* = nB/\mu(s)$ , which approximately correspond to the local maxima of  $J(s)$  with respect to  $t^*$ , where  $B = 1$  is the membrane potential threshold and  $n = 1, 2, \dots$ .

$\mu_0$  is around

$$\mu_0 \approx \frac{2n(n+1)B}{(2n+1)t^*} - \frac{A}{1 + \exp[-b(s-s_0)]}, \quad n \in \mathbb{N}, \quad (10)$$

and  $\sigma^2(s)$  is small, then increasing  $\sigma^2(s)$  through  $k$  or  $m$  may improve the decoding accuracy. Finally, increasing  $k$  or  $m$  causes the local extremes of the Fisher information with respect to  $t^*$  and  $\mu_0$  to become smaller and for a sufficiently large  $\sigma^2(s)$  they are almost negligible.

### B. Comparison with the limit case $\sigma^2 = 0$

The counterintuitive nonmonotonic behavior of the Fisher information can be explained by looking at the limit case

$\sigma_0^2 = \sigma^2(s) = 0$ , when the process  $X(t)$  is deterministic. The only randomness in the system is then the initial value of the membrane potential  $X_0$  when the observation time window does not start with a spike.

Denote by  $T_n$  the time from  $t_0$  to the  $n$ th following spike. If  $X_0 = x_0$  is known,  $T_n = t_n$  is deterministic and equal to

$$t_n = (nB - x_0)/\mu(s). \quad (11)$$

If  $X_0$  is random, the CDF of  $T_n$  is given by

$$\begin{aligned} F_{T_n}(t) &= \mathbb{P}\left(\frac{nB - X_0}{\mu(s)} \leq t\right) \\ &= \mathbb{P}[X_0 \geq nB - \mu(s)t] = 1 - F_{X_0}[nB - \mu(s)t], \end{aligned} \quad (12)$$






FIG. 3. Fisher information  $J(0)$  about the stimulus intensity  $s = 0$  when the time window starts at  $t_0$  and the initial value of the membrane potential  $X_0$  is thus random. In all cases, the coefficient diffusion is  $\sigma^2(s) = k\mu(s) + m$ . (a)–(c) Fisher information as a function of the spontaneous firing rate  $\mu_0$  and the time window  $t^*$  when  $k = 0.01$  and  $m = 0.5$ . The Fisher information is nonmonotonic with respect to both  $\mu_0$  and  $t^*$ . (d)–(f) Fisher information as a function of  $\mu_0$  and  $m$  when  $t^* = 0.025$ s and  $k = 0.01$ . (g)–(i) Fisher information as a function of  $\mu_0$  and  $k$  when  $t^* = 0.025$ s and  $m = 0.5$ . For any choice of  $\mu_0$ , the Fisher information is decreasing in  $k$  and  $m$ . The white dashed lines in (a), (d), and (g) mark the points satisfying  $t^* = nB/\mu(s)$ , which approximately correspond to the local maxima of the Fisher information with respect to  $t^*$ , where  $B = 1$  is the membrane potential threshold and  $n \in \mathbb{N}$ .

where  $F_{X_0}$  is the CDF of  $X_0$ . Consequently, we can write

$$p_{N(t^*)}(n) = \mathbb{P}(T_n \leq t^* < T_{n+1}) = \begin{cases} F_{X_0}[B - \mu(s)t^*], & n = 0 \\ F_{X_0}[(n+1)B - \mu(s)t^*] - F_{X_0}[nB - \mu(s)t^*], & n \geq 1. \end{cases} \quad (13)$$

If  $\sigma_0^2 = 0$ ,  $X_0$  is uniformly distributed over the interval  $[0, B]$ , so we obtain

$$p_{N(t^*)}(n) = \begin{cases} 0, & t^* \in (0, \frac{(n-1)B}{\mu(s)}) \\ 1 - n + \frac{\mu(s)t^*}{B}, & t^* \in [\frac{(n-1)B}{\mu(s)}, \frac{nB}{\mu(s)}) \\ n + 1 - \frac{\mu(s)t^*}{B}, & t^* \in [\frac{nB}{\mu(s)}, \frac{(n+1)B}{\mu(s)}) \\ 0, & t^* \in [\frac{(n+1)B}{\mu(s)}, \infty) \end{cases} \quad (14)$$

for  $n \in \{1, 2, \dots\}$ , and for  $n = 0$  we have

$$p_{N(t^*)}(0) = \begin{cases} 1 - \frac{\mu(s)t^*}{B}, & t^* \in (0, \frac{B}{\mu(s)}) \\ 0, & t^* \in [\frac{B}{\mu(s)}, \infty). \end{cases} \quad (15)$$

The Fisher information about  $s$  based on observing  $N(t^*)$  can be calculated as

$$J(s) = \frac{[\mu'(s)t^*]^2}{[(n+1)B - \mu(s)t^*][nB - \mu(s)t^*]} \quad \text{if } t^* \in \left(\frac{nB}{\mu(s)}, \frac{(n+1)B}{\mu(s)}\right) \quad (16)$$

and is illustrated in Fig. 5(a). It is a piecewise function that is defined on every interval  $[nB/\mu(s), (n+1)B/\mu(s)]$ ,




FIG. 4. Fisher information  $J(0)$  about the stimulus intensity  $s = 0$  as a function of the spontaneous firing rate  $\mu_0$  when the diffusion coefficient is independent of stimulation, namely,  $\sigma^2(s) = \sigma_0^2 = 2$ . (a) Fisher information when the observation time window starts with a spike. (b) Fisher information when the observation window starts at  $t_0$ . Three lengths of the observation window are used:  $t^* = 0.007$  s (red solid line),  $t^* = 0.02$  s (blue dashed line), and  $t^* = 0.04$  s (green dotted line). If the local extremes are ignored, the Fisher information has a mild overall increasing tendency in all the cases, suggesting that the decoding accuracy improves with increasing spontaneous firing rate  $\mu_0$ .

$n = 0, 1, 2, \dots$ , and for  $t^* = nB/\mu(s)$ ,  $n \in \{1, 2, \dots\}$  goes to infinity. The minimum Fisher information on the interval  $t^* \in [nB/\mu(s), (n+1)B/\mu(s)]$  is achieved at

$$t_{\min}^* = \frac{2n(n+1)B}{(2n+1)\mu(s)}, \quad n \in \mathbb{N}. \quad (17)$$

The discontinuities are caused by the shape of the probabilities  $p_{N(t^*)}(n) = \mathbb{P}[N(t^*) = n]$ , illustrated in Fig. 5(c). The probabilities  $p_{N(t^*)}(n)$  are piecewise linear and the derivatives with respect to  $s$  are not defined at  $t^* = nB/\mu(s)$ ,  $n \in \mathbb{N}$ . At these points also  $p_{N(t^*)}(n-1)$  and  $p_{N(t^*)}(n+1)$  tend to zero and the corresponding terms in the formula for the Fisher information tend to infinity. At  $t^* = nB/\mu(s)$ , the probability of observing  $n$  spikes is equal to one.

Now we compare the Fisher information  $J(s)$  for the limit case  $\sigma_0^2 = \sigma^2(s) = 0$  [Fig. 5(a)] with the Fisher information  $J(s)$  for the previously studied cases  $\sigma_0^2 > 0$  and  $\sigma^2(s) > 0$  [Fig. 5(b)]. It can be seen that the local maxima of the Fisher information for  $\sigma_0^2 > 0$  and  $\sigma^2(s) > 0$  are located near  $t^* = nB/\mu(s)$ ,  $n \in \mathbb{N}$ . This can also be confirmed in Figs. 2(a), 2(d), 2(g), 3(a), 3(d), and 3(g), where the white dashed lines, corresponding to  $t^* = nB/\mu(s)$ , are very close




FIG. 5. (a) Fisher information  $J(0)$  about the stimulus intensity  $s = 0$  as a function of the length of the time window  $t^*$  for the limit case of the zero diffusion coefficient  $\sigma_0^2 = \sigma^2(s) = 0$ . The Fisher information goes to infinity for all  $t^*$  satisfying  $t^* = nB/\mu(s)$ ,  $n \in \{1, 2, \dots\}$ . (b) Fisher information  $J(0)$  when  $\sigma_0^2 > 0$  and  $\sigma^2(s) > 0$ . The local maxima are located approximately at  $t^* = nB/\mu(s)$ . (c) Probabilities of observing  $n$  spikes  $p_{N(t^*)}(n) = \mathbb{P}[N(t^*) = n]$  as functions of  $t^*$  for the limit case  $\sigma_0^2 = \sigma^2(s) = 0$ . At  $t^* = nB/\mu(s)$ , the probability of observing  $n$  spikes is equal to 1 and all the other outcomes have zero probability. (d) Probabilities  $p_{N(t^*)}(n)$  as functions of  $t^*$  when  $\sigma_0^2 > 0$  and  $\sigma^2(s) > 0$ . At  $t^* = nB/\mu(s)$ , the probability of observing  $n$  spikes is nearly one and decreases when the diffusion coefficient  $\sigma^2(s)$  is increased. All figures were obtained for the spontaneous firing rate  $\mu_0 = 50$  Hz.

to the local maxima of the Fisher information with respect to  $t^*$ . In Fig. 5(b) we can also notice that if the value of  $\sigma^2(s)$  is increased by increasing  $k$  or  $m$ , the local maxima have the tendency to vanish.

We can also see the correspondence between the shape of the Fisher information  $J(s)$  and the probabilities  $p_{N(t^*)}(n)$ . As  $\sigma^2(s)$  increases, the probabilities  $p_{N(t^*)}(n)$  become smoother with not so sharp transitions between the increasing and the decreasing part. Also, the maximum of the probability  $p_{N(t^*)}(n)$  is less than 1 [Fig. 5(d)], which results in smaller fluctuations of the Fisher information.

We can conclude that the nonmonotonic behavior of the Fisher information is caused by the discrete character of  $N(t^*)$ . If the randomness of  $X(t)$  in the form of  $\sigma^2(s)$  is not too high, there are only a few possible outcomes of  $N(t^*)$ . Hence the Fisher information is the sum of only a small number of nonzero terms in the form  $[\partial p_{N(t^*)}(n)/\partial s]^2/p_{N(t^*)}(n)$ . Therefore, a big change in any of them is strongly reflected in the Fisher information. Note that an individual term has a big contribution if the respective probability is close to zero, which typically happens around  $t^* = nB/\mu(s)$ . The local maxima and minima of the Fisher information with respect to  $t^*$  appear periodically with the period approximately equal to  $B/\mu(s)$ , which is the mean interspike interval. Hence, this phenomenon is not restricted only to short time windows.

**IV. DISCUSSION**

This analysis reveals that even in neuronal models as simple as the perfect integrate-and-fire model the effect of spontaneous activity, noise, and the length of the observation window might be highly nontrivial. In spite of the relative simplicity, the question of neuronal coding is often studied at this or at a similar level of complexity [54–58], because it allows analytical calculations. Therefore, it is possible to identify the reasons for the nonmonotonic behavior of the Fisher information in our work. Since the perfect integrate-and-fire model is based on Brownian motion, it can be also applied on a wide variety of physical problems beyond the field of neuroscience, implying that the described phenomena might appear also there.

Although analogous calculations for more detailed biophysical models, such as models including leakage or setups considering the activity of a group of neurons, which are the directions in which the analysis should be extended in the future, are technically prohibitive, this fact cannot be used to imply that the described effects do not occur therein. On the contrary, the most important result presented here, i.e., that the Fisher information is nonmonotonic with respect to the observation window  $t^*$ , is caused by the discrete nature of the counting distribution. Thus it is not specific for the chosen model and should be largely independent of the amount of biophysical detail taken into account. Our expectation concerning the role of  $\mu_0$ ,  $k$ , and  $m$  in leaky integrate-and-fire models is that they could have an even stronger effect, since

they may help to move the process from the subthreshold to the suprathreshold regime (in the case of  $\mu_0$ ) or increase the chance that  $X(t)$  crosses the threshold in the subthreshold regime (in the case of  $k$  and  $m$ ), which is the classical phenomenon of stochastic resonance [30].

We assumed one particular form of  $\mu(s)$  given by the logistic function. However, the observed phenomena are more general. When analyzing the Fisher information with respect to  $t^*$ ,  $k$ , and  $m$ , we fix  $\mu_0$  and  $s$ , so the value of  $\mu(s)$  is constant and the functional form of  $\mu(s)$  plays no role. When we study the behavior of the Fisher information with respect to  $\mu_0$ , the assumed form of  $\mu(s)$  might have some influence. Nevertheless, our results with respect to  $\mu_0$  can also be obtained for other types of  $\mu(s)$ , for example, for the linear relationship  $\mu(s) = \mu_0 + s$ ,  $s \geq 0$ .

**V. CONCLUSION**

We studied the accuracy of the rate code with respect to several key parameters of the neuronal system and showed that the relationships are nontrivial. Altogether, our results may be summarized in the following four points. (i) Using a longer time window may not necessarily improve the decoding accuracy. (ii) Although the presynaptic spontaneous activity might seem a disturbing element, since it bears no information about the stimulus, its presence at a certain level might improve the decoding accuracy. (iii) The loss in the decoding accuracy caused by the change of the time window can be partially compensated if it is accompanied by an appropriate change of the level of spontaneous activity and vice versa. For example, a shorter time window may sometimes give almost the same accuracy if the level of spontaneous activity is increased by an optimal amount. (iv) If the time window begins with a spike, the decoding accuracy might be improved by increasing the fluctuation of the membrane potential.

**ACKNOWLEDGMENTS**

This work was supported by the Czech Science Foundation Project No. 15-08066S and by the Joint Research Project between Austria and the Czech Republic through Grant No. 7AMB17AT048.

**APPENDIX: DISTRIBUTION OF  $N(t^*)$**

Here we show how the distribution of  $N(t^*)$  can be derived. The first step to obtain the distribution of  $N(t^*)$  is to find the distribution of  $T_n$ , the time from  $t_0$  to the  $n$ th following spike.

*Distribution of  $T_n$ .* Since the Wiener process is a time and space homogeneous process, the distribution of  $T_n$ , the  $n$ th passage time of  $X(t)$  to a constant threshold  $B$ , is the same as the first-passage time of the same process  $X(t)$  to the threshold  $nB$ . Thus,  $T_n|X_0$ , the time to the  $n$ th spike given a fixed starting position  $X_0 = x_0$ , follows an inverse Gaussian distribution  $IG[(nB - x_0)/\mu(s), (nB - x_0)^2/\sigma^2(s)]$  with the CDF

$$F_{T_n|X_0}(t|x_0) = \Phi\left(-\frac{nB - x_0 - \mu(s)t}{\sigma(s)\sqrt{t}}\right) + \exp\left(\frac{2(nB - x_0)\mu(s)}{\sigma^2(s)}\right)\Phi\left(-\frac{nB - x_0 + \mu(s)t}{\sigma(s)\sqrt{t}}\right). \tag{A1}$$

Multiplying the PDF of  $T_n|X_0$  by  $f_{X_0}(x)$ , given by (6) with the parameters  $\mu_0$  and  $\sigma_0^2$ , and integrating over all possible  $x_0$  values, we obtain the unconditional PDF of  $T_n$ ,

$$f_{T_n}(t) = \int_{-\infty}^B f_{T_n|X_0}(t|x)f_{X_0}(x)dx = \frac{\mu(s)\sigma_0^2 - 2\mu_0\sigma^2(s)}{B\sigma_0^2} \exp\left(\frac{2\mu_0[\mu_0\sigma^2(s)t - \mu(s)\sigma_0^2t + (n-1)B\sigma_0^2]}{\sigma_0^4}\right) \\ \times \left[ \exp\left(\frac{2\mu_0B}{\sigma_0^2}\right) \Phi\left(-\frac{nB\sigma_0^2 + 2\mu_0\sigma^2(s)t - \mu(s)\sigma_0^2t}{\sigma_0^2\sigma(s)\sqrt{t}}\right) - \Phi\left(-\frac{(n-1)B\sigma_0^2 + 2\mu_0\sigma^2(s)t - \mu(s)\sigma_0^2t}{\sigma_0^2\sigma(s)\sqrt{t}}\right) \right] \\ + \frac{\mu(s)}{B} \left[ \Phi\left(\frac{nB - \mu(s)t}{\sigma(s)\sqrt{t}}\right) - \Phi\left(\frac{(n-1)B - \mu(s)t}{\sigma(s)\sqrt{t}}\right) \right]. \quad (\text{A2})$$

When  $n = 1$ , the distribution of  $T_1$  is identical to the distribution of the first-spike latency reported in [59].

*Distribution of  $N(t^*)$ .* Consider first the case when the membrane potential at time  $t_0$  is fixed and known, i.e.,  $X_0 = x_0$ . The conditional probability of observing  $n$  spikes in the time window of length  $t^*$  given that  $X_0 = x_0$  can be written as

$$p_{N(t^*)|X_0}(n|x_0) = \mathbb{P}[N(t^*) = n|X_0 = x_0] = \mathbb{P}(T_n \leq t^*|X_0 = x_0) - \mathbb{P}(T_{n+1} \leq t^*|X_0 = x_0). \quad (\text{A3})$$

Plugging (A1) into (A3), we obtain

$$p_{N(t^*)|X_0}(n|x_0) = \Phi\left(-\frac{nB - x_0 - \mu(s)t^*}{\sigma(s)\sqrt{t^*}}\right) - \Phi\left(-\frac{(n+1)B - x_0 - \mu(s)t^*}{\sigma(s)\sqrt{t^*}}\right) + \exp\left(\frac{2\mu(s)[nB - x_0]}{\sigma^2(s)}\right) \\ \times \left[ \Phi\left(-\frac{nB - x_0 + \mu(s)t^*}{\sigma(s)\sqrt{t^*}}\right) - \exp\left(\frac{2\mu(s)B}{\sigma^2(s)}\right) \Phi\left(-\frac{(n+1)B - x_0 + \mu(s)t^*}{\sigma(s)\sqrt{t^*}}\right) \right] \quad (\text{A4})$$

for  $n = 1, 2, \dots$ . If  $n = 0$ , we have

$$p_{N(t^*)|X_0}(0|x_0) = 1 - \Phi\left(-\frac{B - x_0 - \mu(s)t^*}{\sigma(s)\sqrt{t^*}}\right) - \exp\left(\frac{2(B - x_0)\mu(s)}{\sigma^2(s)}\right) \Phi\left(-\frac{B - x_0 + \mu(s)t^*}{\sigma(s)\sqrt{t^*}}\right). \quad (\text{A5})$$

If the value of  $X_0$  is not known, the straightforward way to get the unconditional density of  $N(t^*)$  is to calculate the joint density of  $N(t^*)$  and  $X_0$  and integrate over all possible values of  $X_0$ , that is,

$$p_{N(t^*)}(n) = \int_{-\infty}^B \mathbb{P}[N(t^*) = n|X_0 = x]f_{X_0}(x)dx, \quad (\text{A6})$$

using (A4) and (6). Unfortunately, the closed form of (A6) cannot be obtained and numerical integration must be done.

To avoid possible numerical issues, we suggest computing  $p_{N(t^*)}(n)$  as

$$p_{N(t^*)}(n) = \begin{cases} \int_0^{t^*} f_{T_n}(\tau)[1 - F_{T_1}(t - \tau)]d\tau, & n \geq 1 \\ 1 - F_{T_1}(t^*), & n = 0, \end{cases} \quad (\text{A7})$$

which has the numerical advantage of performing the numerical integration only over a finite interval.

- 
- [1] E. D. Adrian, *The Basis of Sensation* (Norton, New York, 1928).  
 [2] W. Gerstner and W. M. Kistler, *Spiking Neuron Models* (Cambridge University Press, Cambridge, 2002).  
 [3] F. Theunissen and J. P. Miller, *J. Comput. Neurosci.* **2**, 149 (1995).  
 [4] R. Kobayashi and S. Shinomoto, *Phys. Rev. E* **75**, 011925 (2007).  
 [5] Y. Mochizuki and S. Shinomoto, *Phys. Rev. E* **89**, 022705 (2014).  
 [6] W. Bialek, F. Rieke, R. R. De Ruyter Van Steveninck, and D. Warland, *Science* **252**, 1854 (1991).  
 [7] M. N. Shadlen and W. T. Newsome, *Curr. Opin. Neurobiol.* **4**, 569 (1994).  
 [8] M. W. Oram, M. C. Wiener, R. Lestienne, and B. J. Richmond, *J. Neurophysiol.* **81**, 3021 (1999).  
 [9] A. L. Jacobs, G. Fridman, R. M. Douglas, N. M. Alam, P. E. Latham, G. T. Prusky, and S. Nirenberg, *Proc. Natl. Acad. Sci. USA* **106**, 5936 (2009).  
 [10] M. Ainsworth, S. Lee, M. O. Cunningham, R. D. Traub, N. J. Kopell, and M. A. Whittington, *Neuron* **75**, 572 (2012).  
 [11] S. Thorpe, D. Fize, and C. Marlot, *Nature (London)* **381**, 520 (1996).  
 [12] J. Feng, *J. Phys. A* **34**, 7475 (2001).  
 [13] S. Wu, S.-i. Amari, and H. Nakahara, *Neural Comput.* **14**, 999 (2002).  
 [14] S. D. Wilke and C. W. Eurich, *Neural Comput.* **14**, 155 (2002).  
 [15] N. Gordon, T. M. Shackleton, A. R. Palmer, and I. Nelken, *J. Neurosci. Methods* **169**, 391 (2008).  
 [16] F. Montani, E. Phoka, M. Portesi, and S. R. Schultz, *Physica A* **392**, 3066 (2013).  
 [17] M. Deger, T. Schwalger, R. Naud, and W. Gerstner, *Phys. Rev. E* **90**, 062704 (2014).  
 [18] S. Koyama and L. Kostal, *Math Biosci. Eng.* **11**, 63 (2014).  
 [19] P. Dayan and L. F. Abbott, *Theoretical Neuroscience* (MIT Press, Cambridge, 2001).  
 [20] N. Brunel and J.-P. Nadal, *Neural. Comput.* **10**, 1731 (1998).



- [21] I. Dean, N. S. Harper, and D. McAlpine, *Nat. Neurosci.* **8**, 1684 (2005).
- [22] L. Kostal, P. Lansky, and S. Pilarski, *J. Neural Eng.* **12**, 036012 (2015).
- [23] S. Pilarski and O. Pokora, *BioSystems* **136**, 11 (2015).
- [24] R. E. Spinney, J. T. Lizier, and M. Prokopenko, *Phys. Rev. E* **94**, 022135 (2016).
- [25] R. L. Winslow and M. B. Sachs, *Hearing Res.* **35**, 165 (1988).
- [26] M. Bethge, D. Rotermund, and K. Pawelzik, *Neural Comput.* **14**, 2317 (2002).
- [27] D. H. Johnson and W. Ray, *J. Comput. Neurosci.* **16**, 129 (2004).
- [28] P. Lansky and P. E. Greenwood, *BioSystems* **89**, 10 (2007).
- [29] M. Stemmler, *Network* **7**, 687 (1996).
- [30] L. Gammaitoni, P. Hänggi, P. Jung, and F. Marchesoni, *Rev. Mod. Phys.* **70**, 223 (1998).
- [31] P. E. Greenwood, L. M. Ward, and W. Wefelmeyer, *Phys. Rev. E* **60**, 4687 (1999).
- [32] F. Chapeau-Blondeau, S. Blanchard, and D. Rousseau, *Phys. Rev. E* **74**, 031102 (2006).
- [33] L. A. Gatys, A. S. Ecker, T. Tschumatchenko, and M. Bethge, *Phys. Rev. E* **91**, 062707 (2015).
- [34] I. Bashkirtseva, A. B. Neiman, and L. Ryashko, *Phys. Rev. E* **91**, 052920 (2015).
- [35] J. Zylberberg and E. Shea-Brown, *Phys. Rev. E* **92**, 062707 (2015).
- [36] M. Levakova, M. Tamborrino, L. Kostal, and P. Lansky, *Neural Comput.* **28**, 2162 (2016).
- [37] S. Amari and H. Nakahara, *Neural Comput.* **17**, 839 (2005).
- [38] P. E. Greenwood and P. Lansky, *Biol. Cybern.* **92**, 199 (2005).
- [39] E. J. G. Pitman, *Some Basic Theory for Statistical Inference* (Chapman and Hall, London, 1979).
- [40] C. R. Rao, *Linear Statistical Inference and Its Applications*, 2nd ed. (Wiley, New York, 2002).
- [41] G. Gerstein and B. Mandelbrot, *Biophys. J.* **4**, 41 (1964).
- [42] D. R. Cox and H. D. Miller, *The Theory of Stochastic Processes* (Chapman and Hall, London, 1965).
- [43] M. T. Giraudo, P. E. Greenwood, and L. Sacerdote, *Neural Comput.* **23**, 1743 (2011).
- [44] D. R. Cox and P. A. W. Lewis, *The Statistical Analysis of Series of Events* (Methuen, London, 1966).
- [45] S. A. Frank, *Biol. Direct* **8**, 31 (2013).
- [46] M. Chastrette, T. Thomas-Danguin, and E. Rallet, *Chem. Senses* **23**, 181 (1998).
- [47] L. Nizami, *Hearing Res.* **167**, 13 (2002).
- [48] J.-P. Rospars, P. Lansky, A. Duchamp, and P. Duchamp-Viret, *Eur. J. Neurosci.* **18**, 1135 (2003).
- [49] S. Durant, C. W. G. Clifford, N. A. Crowder, N. S. C. Price, and M. R. Ibbotson, *J. Opt. Soc. Am. A* **24**, 1529 (2007).
- [50] A. Grémiaux, T. Nowotny, D. Martinez, P. Lucas, and J.-P. Rospars, *Brain Res.* **1434**, 123 (2012).
- [51] H. C. Tuckwell, *Introduction to Theoretical Neurobiology, Vol.2: Nonlinear and Stochastic Theories* (Cambridge University Press, Cambridge, 1988), p. 138.
- [52] K. Miura, Y. Tsubo, M. Okada, and T. Fukai, *J. Neurosci.* **27**, 13802 (2007).
- [53] B. Sengupta, S. B. Laughlin, and J. E. Niven, *PLoS Comput. Biol.* **9**, e1003263 (2013).
- [54] B. Lindner, *Phys. Rev. E* **69**, 022901 (2004).
- [55] D. Andrieux and T. Monnai, *Phys. Rev. E* **80**, 021933 (2009).
- [56] R. D. Vilela and B. Lindner, *Phys. Rev. E* **80**, 031909 (2009).
- [57] F. Müller-Hansen, F. Droste, and B. Lindner, *Phys. Rev. E* **91**, 022718 (2015).
- [58] R. Mankin and A. Rekker, *Phys. Rev. E* **94**, 062103 (2016).
- [59] M. Tamborrino, S. Ditlevsen, and P. Lansky, *Lifetime Data Anal.* **21**, 331 (2015).