

Effective drift velocity from turbulent transport by vorticity

Hussein Aluie ^{1,2,*} Shikhar Rai ¹ Hao Yin ¹ Aarne Lees,^{1,2} Dongxiao Zhao ¹
Stephen M. Griffies ^{3,4} Alistair Adcroft ^{3,4} and Jessica K. Shang ^{1,2}

¹*Department of Mechanical Engineering, University of Rochester, Rochester, New York 14625, USA*

²*Laboratory for Laser Energetics, University of Rochester, Rochester, New York 14625, USA*

³*NOAA/Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey 08544, USA*

⁴*Princeton University Program in Atmospheric and Oceanic Sciences, Princeton, New Jersey 08544, USA*



(Received 1 June 2022; accepted 14 September 2022; published 3 October 2022)

We highlight the differing roles of vorticity and strain in the transport of coarse-grained scalars at length scales larger than ℓ by smaller scale (subscale) turbulence. We use the first term in a multiscale gradient expansion due to Eyink [J. Fluid Mech. **549**, 159 (2006)], which exhibits excellent correlation with the exact subscale physics when the partitioning length ℓ is any scale smaller than that of the spectral peak. We show that unlike subscale strain, which acts as an anisotropic diffusion/antidiffusion tensor, subscale vorticity's contribution is solely a conservative advection of coarse-grained quantities by an eddy-induced nondivergent velocity, $\mathbf{v}_*$, that is proportional to the curl of vorticity. Therefore, material (Lagrangian) advection of coarse-grained quantities is accomplished not by the coarse-grained flow velocity, $\bar{\mathbf{u}}_\ell$, but by the effective velocity, $\bar{\mathbf{u}}_\ell + \mathbf{v}_*$, the physics of which may improve commonly used LES models.

DOI: [10.1103/PhysRevFluids.7.104601](https://doi.org/10.1103/PhysRevFluids.7.104601)

I. INTRODUCTION

Basic considerations from fluid dynamics (e.g., [1]) indicate that the distance between particles in a laminar flow is determined by the strain. Vorticity merely imparts a rotation on their separation vector $\mathbf{r}$ without affecting its magnitude. This behavior can be seen by considering the velocity, $\mathbf{u}$, difference between particles P and Q at positions $\mathbf{x}$ and $\mathbf{x} + \mathbf{r}$, respectively,

$$\mathbf{u}_Q - \mathbf{u}_P = \delta \mathbf{u} = \mathbf{u}(\mathbf{x} + \mathbf{r}) - \mathbf{u}(\mathbf{x}) = \mathbf{r} \cdot \nabla \mathbf{u} \Big|_{\mathbf{x}} + \cdots, \quad (1)$$

where a Taylor-series expansion is justified for short distances $|\mathbf{r}|$ over which the flow is sufficiently smooth. In the Lagrangian frame of P at $\mathbf{x}$, the separation from Q evolves as

$$\frac{D\mathbf{r}}{Dt} = \delta \mathbf{u} = \mathbf{r} \cdot \mathbf{S} + \underbrace{\mathbf{r} \cdot \boldsymbol{\Omega}}_{\frac{1}{2} \boldsymbol{\omega} \times \mathbf{r}}, \quad (2)$$

where the velocity gradient tensor, $\nabla \mathbf{u} = \mathbf{S} + \boldsymbol{\Omega}$, has been decomposed into the symmetric strain rate tensor $\mathbf{S} = [\nabla \mathbf{u} + (\nabla \mathbf{u})^T]/2$ and the antisymmetric vorticity tensor $\boldsymbol{\Omega} = [\nabla \mathbf{u} - (\nabla \mathbf{u})^T]/2 = -\frac{1}{2} \epsilon_{ijk} \omega_k$. Here $\boldsymbol{\omega} = \nabla \times \mathbf{u}$ is vorticity and ϵ_{ijk} is the Levi-Civita symbol. Taking an inner product of Eq. (2) with $\mathbf{r}$,

$$\frac{1}{2} \frac{D|\mathbf{r}|^2}{dt} = \mathbf{r} \cdot \mathbf{S} \cdot \mathbf{r}, \quad (3)$$

*hussein@rochester.edu




FIG. 1. Schematic of how eddy-induced advection by a nondivergent velocity, $\mathbf{v}_* \sim \nabla \times \bar{\boldsymbol{\omega}}_\ell$, (blue arrow) from Eq. (43) can arise due to subscale vorticity (red). In this example, two symmetric but counterrotating eddies (red) are of a size smaller than that of the coarse-graining box (dashed). At length scales larger than that of the box, ℓ , which may represent a grid cell in a simulation or the resolution limit of an observation, these eddies exert an eddy-induced advection with velocity $\mathbf{v}_*$ proportional to the curl of vorticity, depicted by the blue arrow at location $\mathbf{x}$.

shows that the distance is determined by the strain. Vorticity in Eq. (2) acts only to rotate $\mathbf{r}$ without changing its magnitude.

These considerations hinge on the critical assumption that the flow is sufficiently smooth over separations $\mathbf{r}$, which is patently invalid in a turbulent flow for $\mathbf{r}$ at inertial scales [2]. However, a version of this story survives due to the property of scale locality, which justifies an expansion in scale as we shall discuss below. The main result of this paper is Eq. (43), which is an expression for the eddy-induced advection velocity $\mathbf{v}_*$ at length scales larger than ℓ , which may be the size of grid cells in a simulation. The nondivergent velocity, $\mathbf{v}_*$, arises from vortical motions associated with subscale nonlinear interactions. Figure 1 captures the essential insight behind Eq. (43).

In Sec. II we discuss subscale transport, Eyink’s expansion in length scale, and the leading order term. In Sec. III we present empirical support from (1) 3D direct numerical simulation (DNS) of a compressible turbulent flow and (2) two-layer stacked shallow water simulation of geophysical fluid turbulence. Section IV contains the main result, which expresses transport by subscale vorticity as a conservative advection by an effective drift velocity $\mathbf{v}_*$ proportional to the curl of vorticity. Throughout this paper we attempt to make the presentation accessible to readers from various backgrounds.

II. MULTISCALE DYNAMICS

To analyze the dynamics of different scales in a flow, we use the coarse-graining approach, which has proven to be a natural and versatile framework to understand and model scale interactions (e.g., [3,4]). The approach is standard in partial differential equations and distribution theory (e.g., [5,6]). It became common in large eddy simulation (LES) modeling of turbulence thanks to the foundational works of Leonard [7] and Germano [8].

For any field $\mathbf{a}(\mathbf{x})$, a coarse-grained or (low-pass) filtered version of this field, which contains spatial variations at scales $> \ell$, is defined in n dimensions as

$$\bar{\mathbf{a}}_\ell(\mathbf{x}) = \int d^n \mathbf{r} G_\ell(\mathbf{x} - \mathbf{r}) \mathbf{a}(\mathbf{r}). \quad (4)$$

Here G_ℓ is a convolution kernel derived from a “mother” kernel $G(\mathbf{s})$ (borrowing the term from wavelet analysis [9]). $G(\mathbf{s})$ is normalized, $\int d^n \mathbf{s} G(\mathbf{s}) = 1$. It is an even function such that, $\int \mathbf{s} G(\mathbf{s}) d^n \mathbf{s} = 0$, which ensures that local averaging is symmetric. $G(\mathbf{s})$ also has its main support (or variance) over a region of order unity in diameter, $\int d^n \mathbf{s} |\mathbf{s}|^2 G(\mathbf{s}) = O(1)$. The dilated version of the kernel,

$$G_\ell(\mathbf{r}) = \ell^{-n} G(\mathbf{r}/\ell), \quad (5)$$

inherits all those properties, except that its main support is over a region of diameter ℓ . An example is the Top Hat kernel in nondimensional coordinates $\mathbf{s}$,

$$G(\mathbf{s}) = \begin{cases} 1, & \text{if } |s_i| < 1/2 \text{ for } i = 1, \dots, n, \\ 0, & \text{otherwise,} \end{cases} \quad (6)$$

and its dilated version in dimensional coordinates $\mathbf{r}$,

$$G_\ell(\mathbf{r}) = \begin{cases} \ell^{-n}, & \text{if } |r_i| < \ell/2 \text{ for } i = 1, \dots, n, \\ 0, & \text{otherwise.} \end{cases} \quad (7)$$

The scale decomposition in Eq. (4) is essentially a partitioning of scales for a field into large scales ($\gtrsim \ell$), captured by $\bar{\mathbf{a}}_\ell$, and small scales ($\lesssim \ell$), captured by the residual

$$\mathbf{a}'_\ell = \mathbf{a} - \bar{\mathbf{a}}_\ell. \quad (8)$$

In the remainder of this paper, we shall omit subscript ℓ from variables if there is no risk for confusion.

A. Coarse-grained transport

Consider the transport equation of the concentration (per unit volume) $C(\mathbf{x}, t)$:

$$\partial_t C + \nabla \cdot (C \mathbf{u}) = \nabla \cdot [\rho \kappa \nabla \phi]. \quad (9)$$

Here ρ is mass density, $\mathbf{u}$ is the advecting flow velocity, which is possibly compressible. The scalar field $C = \rho \phi$, where ϕ is concentration per unit mass, can be either passive or active, with diffusivity κ , which can be due to the microphysics or an effective (turbulent) diffusivity used in a simulation. Analysis of the incompressible momentum transport follows similar reasoning as below, with C replaced by $\mathbf{u}$ when $\rho = (\text{const})$, so we shall focus on C for simplicity.

We shall call Eq. (9) the “bare” dynamics [10,11] as it governs flow in an experiment or natural system. In a coarse-resolution simulation, such as in implicit or explicit LES [3,12], or when taking spatially underresolved measurements in an experiment, $\mathbf{u}$ and C are not resolved down to the microphysical scales. The equation that governs the coarse-grained scalar $\bar{C}_\ell$ is obtained by filtering Eq. (9) to get (e.g., [13])

$$\partial_t \bar{C}_\ell + \nabla \cdot (\bar{\mathbf{u}}_\ell \bar{C}_\ell) = -\nabla \cdot \bar{\tau}_\ell + \nabla \cdot \overline{(\rho \kappa \nabla \phi)}_\ell, \quad (10a)$$

$$\text{where } \bar{\tau}_\ell = \bar{\tau}_\ell(\mathbf{u}, C) \equiv \overline{(\mathbf{u} C)}_\ell - \bar{\mathbf{u}}_\ell \bar{C}_\ell \quad (10b)$$

is the subscale flux of $\bar{C}_\ell$ [8]. Equation (10) is exact, without any approximation, and describes the scalar transport at length scales larger than ℓ . The last term, $\nabla \cdot \overline{(\rho \kappa \nabla \phi)}_\ell$, was shown mathematically in [14] to have an upper bound proportional to $\alpha_{\text{rms}} \ell^{-2} \phi_{\text{rms}}$ at every location in the flow and every time, where $\alpha = \rho \kappa$. Therefore, it is guaranteed to be negligible for sufficiently large ℓ or sufficiently small α , which was demonstrated numerically in [15]. Therefore, $\nabla \cdot \overline{(\rho \kappa \nabla \phi)}_\ell$ can be dropped from Eq. (10) [but not from the bare Eq. (9)].

An evident role of the subscale flux $\bar{\tau}_\ell(C, \mathbf{u})$ in Eq. (10) is accounting for the influence of unresolved scales C'_ℓ and $\mathbf{u}'_\ell$ on the transport of $\bar{C}_\ell$. Another important role of $\bar{\tau}_\ell(C, \mathbf{u})$ in Eq. (10) that may not be as well appreciated is balancing $\bar{\mathbf{u}}_\ell \bar{C}_\ell$, which can interact with scales $< \ell$ that are approximately within the band $[\ell/2, \ell]$. This is analogous to the term $\rho\kappa\nabla\phi$ balancing $\mathbf{u}C$ at microphysical scales $\sim\ell_\kappa$, below which molecular diffusion dominates advection. To illustrate, assume that at a time t the scalar and velocity fields are fully resolved, $\bar{C}_\ell = C$ and $\bar{\mathbf{u}}_\ell = \mathbf{u}$. Moreover, assume that scale ℓ is sufficiently large, $\ell \gg \ell_\kappa$, such that microphysical diffusivity is negligible and does not damp nonlinear interactions. That small scales are absent, $C'_\ell = 0$ and $\mathbf{u}'_\ell = 0$, does not necessarily imply that the subscale flux $\bar{\tau}_\ell(C, \mathbf{u}) = \overline{(\mathbf{u}C)}_\ell - \mathbf{u}C$ is zero unless there is sufficient scale separation between ℓ and the smallest scales of C and $\mathbf{u}$. Therefore, the general transport of a coarse-grained scalar $\bar{C}_\ell$ is not described by the typical (or bare) transport equation (9) even when C and $\mathbf{u}$ are resolved ($C'_\ell = 0$ and $\mathbf{u}'_\ell = 0$), because coarse graining also alters the representation of their nonlinear interactions, which may be at scales smaller than ℓ .

From definition (10b), it is possible to decompose $\bar{\tau}_\ell(C, \mathbf{u})$ into contributions that are solely due to resolved fields $\bar{C}_\ell$ and $\bar{\mathbf{u}}_\ell$ (e.g., the Leonard term [7,16]) and other contributions that involve unresolved fields C'_ℓ and $\mathbf{u}'_\ell$. In the discussion below, we use the term “subscale” to refer to any contribution from $\bar{\tau}_\ell(C, \mathbf{u})$ because it arises from coarse graining the dynamics at scale ℓ . We shall see that the resolved fields make a dominant contribution to the subscale flux $\bar{\tau}_\ell(C, \mathbf{u})$ due to the property of ultraviolet scale locality.

B. Multiscale expansion

The subscale term $\bar{\tau}_\ell(\mathbf{u}, C)$ does not lend itself to a straightforward intuitive interpretation. To that end, we utilize the multiscale gradient expansion of Eyink [17]. This expansion is accomplished by decomposing a field

$$\mathbf{a}(\mathbf{x}) = \sum_{p=0}^{\infty} \mathbf{a}^{[p]}(\mathbf{x}) \quad (11)$$

into a sum of band-passed fields [17]

$$\mathbf{a}^{[p]} = \bar{\mathbf{a}}_{\ell_p} - \bar{\mathbf{a}}_{\ell_{p-1}}, \quad p = 1, 2, \dots, \quad (12a)$$

$$\mathbf{a}^{[0]} = \bar{\mathbf{a}}_{\ell_0} \quad (12b)$$

for a sequence of scales $\ell_p = \lambda^{-p}\ell_0$ with the nondimensional number $\lambda > 1$, e.g., $\lambda = 2$. ℓ_0 can be a characteristic large scale such as the domain size. A similar decomposition can be done for the subscale flux,

$$\bar{\tau}_\ell(\mathbf{u}, C) = \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \tau_\ell^{[p,q]}(\mathbf{u}, C), \quad (13a)$$

$$\text{with } \tau_\ell^{[p,q]}(\mathbf{u}, C) = \overline{(\mathbf{u}^{[p]}C^{[q]})}_\ell - \overline{(\mathbf{u}^{[p]})}_\ell \overline{(C^{[q]})}_\ell. \quad (13b)$$

The term $\tau_\ell^{[p,q]}(\mathbf{u}, C)$ represents the subscale contribution from scalar and velocity scales within bands $[q]$ and $[p]$, respectively. Eyink [4,17] proved that the series (13) converges absolutely and at a geometric rate under the condition that the scaling exponents of increments of $\delta\mathbf{u}(r) \sim r^{\sigma^u}$ and $\delta C(r) \sim r^{\sigma^C}$ satisfy $\sigma^u > 0$ and $\sigma^C > 0$.¹

¹More precisely, the condition is that $\sigma_p^u > 0$ and $\sigma_p^C > 0$ if increments scale as $\langle |\delta\mathbf{u}(r)|^p \rangle^{1/p} \sim u_{\text{rms}} A_p (r/\ell_0)^{\sigma_p^u}$ and $\langle |\delta C(r)|^p \rangle^{1/p} \sim C_{\text{rms}} B_p (r/\ell_0)^{\sigma_p^C}$ for some dimensionless constants A_p and B_p , where $\langle \dots \rangle$ is a space average. The condition has to hold for at least $p = 2$, with higher values of p representing more

Exponents σ'' and σ^C reflect the smoothness of their respective fields (see Fig. 1 in [10] and the associated discussion), with larger values of σ implying smoother (or, heuristically, less turbulent) fields. For example, the Kolmogorov-Obukhov theory [18–20] predicts $\sigma'' = \sigma^C = 1/3$ for velocity and passive scalar increments over scales within the inertial-convective range of a high Reynolds number turbulent flow that is statistically homogeneous. In contrast, for low Reynolds number flows, even if they are chaotic, we can expect $\sigma'' \geq 1$. Ignoring intermittency corrections, the scaling exponent σ of increments is related to the power-law scaling of a spectrum, $E(k)$, of either $\mathbf{u}$ or C , via $E(k) \sim k^{-2\sigma-1}$. Therefore, the condition $\sigma > 0$ corresponds to a wave number scaling of spectra that decays faster than k^{-1} in wave number, which is a fairly weak condition and is expected to hold in most flows. The conditions are the same as those required for $\bar{\tau}_\ell(\mathbf{u}, C)$ to be ultraviolet scale-local [4,21,22], which means that length scales $\delta \ll \ell$ make a negligible contribution to $\bar{\tau}_\ell(\mathbf{u}, C)$,

$$|\bar{\tau}_\ell(C'_\delta, \mathbf{u}'_\delta)| \ll |\bar{\tau}_\ell(C, \mathbf{u})|. \quad (14)$$

Under these conditions of convergence, the leading order term in Eq. (13) is the dominant contribution to the series, which justifies the approximation

$$\bar{\tau}_\ell(\mathbf{u}, C) \approx \tau_\ell^{[0,0]}(\mathbf{u}, C) = \bar{\tau}_\ell(\bar{\mathbf{u}}_\ell, \bar{C}_\ell). \quad (15)$$

This approximation corresponds to the well-known similarity (or Bardina) model [3,23]. To readers who are perhaps less familiar with LES or coarse graining, the term

$$\bar{\tau}_\ell(\bar{\mathbf{u}}_\ell, \bar{C}_\ell) = \overline{(\bar{\mathbf{u}}_\ell \bar{C}_\ell)_\ell} - \overline{(\bar{\mathbf{u}}_\ell)_\ell} \overline{(\bar{C}_\ell)_\ell} \quad (16)$$

may seem insignificant, especially considering that if one were to replace operation $\overline{(\dots)_\ell}$ with a Reynolds (or ensemble) average, expression (16) would be identically zero. However, operation $\overline{(\dots)_\ell}$ is a decomposition in length scale, which is inherently different from Reynolds averaging. To aid in the conceptual understanding of expression (16), assume that $\overline{(\dots)_\ell}$ is a sharp-spectral projection in wave number space, $\overline{(\dots)_\ell} = (\dots)^{<K}$. Applying $(\dots)^{<K}$ removes all wave numbers larger than $K = \ell^{-1}$ (i.e., small length scales) while keeping wave numbers smaller than K intact. Then

$$\bar{\tau}_\ell(\bar{\mathbf{u}}_\ell, \bar{C}_\ell) = (\mathbf{u}^{<K} C^{<K})^{<K} - (\mathbf{u}^{<K})^{<K} (C^{<K})^{<K} \quad (17a)$$

$$= (\mathbf{u}^{<K} C^{<K})^{<K} - \mathbf{u}^{<K} C^{<K}, \quad (17b)$$

using the sharp projection property, $((\dots)^{<K})^{<K} = (\dots)^{<K}$, to arrive at the last expression. The term $\mathbf{u}^{<K} C^{<K}$ in Eq. (17b), being a quadratic product, contains wave numbers $< 2K$. Of these, the contribution $(\mathbf{u}^{<K} C^{<K})^{<K}$ from wave numbers $< K$ is subtracted in Eq. (17b), such that $\bar{\tau}_\ell(\bar{\mathbf{u}}_\ell, \bar{C}_\ell)$ represents wave numbers within the band $[K, 2K]$ or, equivalently, between length scales ℓ and $\ell/2$. As discussed in previous work on scale locality [4,24], this band of scales is expected to make the dominant contribution to the subscale flux $\bar{\tau}_\ell(\mathbf{u}, C)$ if the spectrum decays faster than k^{-1} in wave number over the range of scales $< \ell$. This is essentially the condition under which Eq. (15) was derived. Physically, a spectral decay faster than k^{-1} implies that dyadic wave number band $[k, 2k]$ has more energy than band $[2k, 4k]$ (e.g., [22,25]), which justifies retaining the smallest wave numbers via the leading order term in expansion (13) [22]. A spectral decay faster than k^{-1} is a fairly weak condition and is expected in most flows (turbulent or laminar), but may fail, for example, if ℓ approaches scales of the spectral peak or larger. We emphasize that filtering with a sharp-spectral kernel in Eq. (17) is merely for conceptual understanding. Unlike the Top Hat

stringent conditions. Note that a power-law scaling of increments is not required; the condition is still satisfied if increments decay faster than any power law with $r \rightarrow 0$, for example exponentially.

or Gaussian, a sharp-spectral kernel is not positive semidefinite in physical space, which violates physical realizability conditions [26], rendering subscale energy or coarse-grained mass density negative in physical space [4,14].

C. Relation to increments and gradients

Using the usual definition of an increment,

$$\delta f(\mathbf{x}; \mathbf{r}) = f(\mathbf{x} + \mathbf{r}) - f(\mathbf{x}), \quad (18)$$

the subscale flux (10b) can be rewritten exactly in terms of δC and $\delta \mathbf{u}$ [4,24,27]:

$$\bar{\tau}(C, \mathbf{u}) = \langle \delta C \delta \mathbf{u} \rangle_\ell - \langle \delta C \rangle_\ell \langle \delta \mathbf{u} \rangle_\ell. \quad (19)$$

Equation (19) is exact and follows directly from Eq. (10b) without requiring any assumption about scale locality or the scaling exponents, where

$$\langle \delta f \rangle_\ell(\mathbf{x}) \equiv \int d^n \mathbf{r} G_\ell(-\mathbf{r}) \delta f(\mathbf{x}; \mathbf{r}) \quad (20)$$

is a local average around $\mathbf{x}$ over all separations $\mathbf{r}$ weighted by the kernel G_ℓ . A spatially localized (or compact) kernel effectively limits the average in Eq. (20) to separations $|\mathbf{r}| \leq \ell/2$. Note that $G_\ell(-\mathbf{r}) = G_\ell(\mathbf{r})$ for even kernels, such as a Gaussian or a Top Hat. Relation (19) is a key step and reveals the connection between the subscale physics and increments (spatial variations) over distances smaller than ℓ . This connection is underscored by noting that expression (20) equals (minus) the high-pass field containing scales $< \ell$,

$$\int d^n \mathbf{r} G_\ell(-\mathbf{r}) \delta f(\mathbf{x}; \mathbf{r}) = -[f(\mathbf{x}) - \bar{f}_\ell(\mathbf{x})] = -f'_\ell(\mathbf{x}). \quad (21)$$

We may also connect these considerations with the argument presented in the introduction, where increments reflect gradients [17], including strain and vorticity of the subscale flow.

Approximating the subscale flux by the leading order term in Eq. (15) allows us to use identity (19) for the filtered quantities,

$$\bar{\tau}(\bar{C}_\ell, \bar{\mathbf{u}}_\ell) = \langle \delta \bar{C}_\ell \delta \bar{\mathbf{u}}_\ell \rangle_\ell - \langle \delta \bar{C}_\ell \rangle_\ell \langle \delta \bar{\mathbf{u}}_\ell \rangle_\ell. \quad (22)$$

Since a filtered field $\bar{f}_\ell(\mathbf{x})$ is smooth, we can Taylor expand its increments around $\mathbf{x}$,

$$\delta \bar{f}_\ell(\mathbf{x}; \mathbf{r}) = \bar{f}_\ell(\mathbf{x} + \mathbf{r}) - \bar{f}_\ell(\mathbf{x}) \approx \mathbf{r} \cdot \nabla \bar{f}_\ell(\mathbf{x}) + \dots, \quad (23)$$

where we neglect higher order terms. Substituting the first term in the Taylor expansion of each of $\delta \bar{C}$ and $\delta \bar{\mathbf{u}}$ into Eq. (22) gives in n dimensions

$$\bar{\tau}_\ell(\bar{C}, \bar{u}_i) = [\partial_k \bar{C}_\ell][\partial_m \overline{(u_i)}_\ell][\langle r_k r_m \rangle_\ell - \langle r_k \rangle_\ell \langle r_m \rangle_\ell] \quad (24a)$$

$$= [\partial_k \bar{C}_\ell][\partial_m \overline{(u_i)}_\ell] \left[\frac{1}{n} \delta_{km} \ell^2 \int d^n \mathbf{s} G(\mathbf{s}) |\mathbf{s}|^2 \right] \quad (24b)$$

$$= \frac{1}{n} M_2 \ell^2 \partial_k \overline{(u_i)}_\ell \partial_k \bar{C}_\ell. \quad (24c)$$

In deriving the second line, we used the symmetry of the kernel such that $\langle r_k \rangle_\ell = 0$. In the final expression, the mother kernel's second moment, $M_2 = \int d^n \mathbf{s} G(\mathbf{s}) |\mathbf{s}|^2$, depends solely on the shape of kernel G and, in particular, is independent of scale ℓ . For the Top Hat in Eq. (6), $M_2 = n/12$ in n dimensions.

D. Summary and interpretation

From the derivation above, we see that the final expression in Eq. (24c) represents the leading order contribution to $\bar{\tau}_\ell(C, \mathbf{u})$. Therefore, to the extent expression (24c) faithfully approximates $\bar{\tau}_\ell(C, \mathbf{u})$, it also represents the subscale physics. How can expression (24c), which involves only coarse-grained (or resolved) quantities, represent subscale interactions? It is analogous to inferring the value of a function $f(x_0 + h)$ a distance h away from x_0 based solely on its local properties at x_0 via a Taylor series expansion, the accuracy of which depends on smoothness properties of f . Here the expansion is done in length scale [17] rather than in space to arrive at Eq. (24c). The analog to smoothness is ultraviolet scale locality, which is guaranteed under weak spectral scaling conditions [4,21,22,24].

Expression (24c) corresponds to the nonlinear (or Clark) model [28], which is a closure of the subscale flux using a product of gradients in LES modeling and has been shown to be an excellent (*a priori*) approximation of the subscales in several previous studies (e.g., [13,29,30]). Some studies have also offered viable ideas for utilizing it in *a posteriori* LES [23,31,32]. Below we provide additional *a priori* support from simulations of 3D compressible turbulence and two-layer stacked shallow water simulations of geophysical fluid turbulence.

In the discussion above, we followed Eyink's multiscale gradient expansion [17] to derive expression (24c), which is more general than the standard derivations of the nonlinear model found in the LES literature (e.g., [2,23,28,29,32–34]). Since Eyink's multiscale gradient expansion is convergent, it provides a rigorous foundation to justify neglecting higher order terms. Another advantage of using Eyink's expansion is in bringing to the fore the underlying physics, including the assumptions under which the series (13) converges and, therefore, the conditions under which the approximation may fail. Moreover, being a systematic expansion, it allows for future improvements by retaining higher order terms or by using a closed form of those terms following the recent work of Johnson [35,36].

Approximation (24c) provides us with physical insight into the mechanisms by which the subscales transport $\bar{C}_\ell$. Below we show that subscale vorticity's contribution is an eddy-induced drift velocity. Prior to doing so, we discuss how well Eq. (24c) approximates $\bar{\tau}_\ell(\mathbf{u}, C)$ from simulations.

III. EMPIRICAL SUPPORT

Approximating $\bar{\tau}_\ell(\mathbf{u}, C)$ by expression (24c) has significant support in the LES literature from *a priori* tests of the nonlinear model. It has been shown to exhibit excellent agreement when C is taken as the momentum (e.g., [29,34]), as the temperature field in the atmospheric boundary layer [13,30], and as a passive scalar [37,38].

In this section we present additional supporting evidence for active scalars, when C represents (1) density in a 3D compressible turbulent flow and (2) layer thickness in a two-layer stacked shallow water simulation of geophysical fluid turbulence. The correlation is generally better than 0.9 at inertial scales smaller than the scale at which the spectrum peaks.

A. Compressible turbulence

We carry out a direct numerical simulation (DNS) of forced compressible turbulence in a periodic box of nondimensional size 2π with grid resolution 1024^3 . From the simulation data, we test how well the expression in Eq. (24c) approximates $\bar{\tau}_\ell(\mathbf{u}, C)$ with C replaced by the density field, ρ , which is an active scalar. The DNS follows the configuration of [39] and solves the fully compressible Navier-Stokes equations using our DiNuSUR code [15,39]:

$$\partial_t \rho + \partial_j(\rho u_j) = 0, \quad (25)$$

$$\partial_t(\rho u_i) + \partial_j(\rho u_i u_j) = -\partial_i P + \partial_j \sigma_{ij} + \rho F_i, \quad (26)$$

$$\partial_t(\rho E) + \partial_j(\rho E u_j) = -\partial_j(P u_j) + \partial_j\left[2\mu u_i(S_{ij} - \frac{1}{3}S_{kk}\delta_{ij})\right] - \partial_j q_j + \rho u_i F_i - \mathcal{RL}. \quad (27)$$




FIG. 2. Left to right: visualization of x , y , and z components, respectively, of momentum, $\rho \mathbf{u}$, in physical space from the 1024^3 DNS. These images are shown at an instant of time after the flow has reached steady state. Shocks can be seen as discontinuities. The spectrum of this flow is shown in Fig. 15 in the Appendix.

Here $E = |\mathbf{u}|^2/2 + e$ is total energy per unit mass, where e is specific internal energy, P is thermodynamic pressure, μ is dynamic viscosity, $\mathbf{q} = -\kappa \nabla T$ is the heat flux with a thermal conductivity κ and temperature T . Both dynamic viscosity and thermal conductivity are spatially variable, where $\mu(\mathbf{x}) = \mu_0(T(\mathbf{x})/T_0)^{0.76}$. Thermal conductivity is set to satisfy a Prandtl number $\text{Pr} = c_p \mu / \kappa = 0.7$, where $c_p = R \gamma / (\gamma - 1)$ is the specific heat with specific gas constant R and $\gamma = 5/3$ for a monoatomic gas. We use the ideal gas equation of state, $P = \rho R T$. We stir the flow using an external acceleration field F_i , and $\mathcal{RL} = \rho u_i F_i$ represents radiation losses from internal energy (see [39,40]). $S_{ij} = (\partial_j u_i + \partial_i u_j)/2$ is the symmetric rate of strain tensor and σ_{ij} is the deviatoric (traceless) viscous stress,

$$\sigma_{ij} = 2 \mu (S_{ij} - \frac{1}{3} S_{kk} \delta_{ij}). \quad (28)$$

We solve the above equations using the pseudo-spectral method with 2/3rd dealiasing. We advance in time using the fourth-order Runge-Kutta scheme with a variable time step.

The acceleration, $\mathbf{F}$, is similar to that in [41]. In Fourier space, the acceleration is defined as

$$\widehat{F}_i(\mathbf{k}) = \widehat{f}_j(\mathbf{k}) P_{ij}^\zeta(\mathbf{k}), \quad (29)$$

where $\widehat{\mathbf{f}}$ is constructed from independent Ornstein-Uhlenbeck stochastic processes [42]. The projection operator

$$P_{ij}^\zeta(\mathbf{k}) = \zeta \delta_{ij} + (1 - 2\zeta) \frac{k_i k_j}{|\mathbf{k}|^2} \quad (30)$$

allows for controlling the ratio of solenoidal ($\nabla \cdot \mathbf{F} = 0$) to dilatational ($\nabla \times \mathbf{F} = 0$) components of the forcing using the parameter ζ . When $\zeta = 0$, the forcing is purely dilatational and when $\zeta = 1$, the forcing is purely solenoidal. For the simulation we use here, shown in Fig. 2, we set $\zeta = 0.01$ to simulate turbulence with high compressibility, such that dilatation, $\nabla \cdot \mathbf{u} \neq 0$, is significant.

Our simulated flow has significant compressibility, which can be seen from Fig. 2 and inferred from the following three time-averaged quantities. (1) First, we use the ratio $(\nabla \cdot \mathbf{u})_{\text{rms}}/(\nabla \times \mathbf{u})_{\text{rms}} = 5.61$ in our flow, which measures compressibility at small scales. (2) We also Helmholtz decompose the velocity field into the dilatational, $\nabla \times \mathbf{u}^d = 0$, and solenoidal, $\nabla \cdot \mathbf{u}^s = 0$, components, $\mathbf{u} = \mathbf{u}^d + \mathbf{u}^s$. The ratio of dilatational kinetic energy, $K^d = \langle \rho u_i^d u_i^d / 2 \rangle$, to solenoidal kinetic energy, $K^s = \langle \rho u_i^s u_i^s / 2 \rangle$, yields a measure of compressibility at large scales, which in our flow is $K^d/K^s = 4.80$. (3) We also calculate the turbulence Mach number, $M_t = \langle |\mathbf{u}|^2 \rangle^{1/2} / c = 0.23$, where $c = 2.18$ is the mean nondimensional sound speed. Spectra of density and velocity are plotted in Fig. 15 in the Appendix.

We use the Top Hat kernel in Eq. (7) for the coarse graining. Figure 3 shows a 2D slice from the 3D flow at the instant of time shown in Fig. 2, comparing the exact $\bar{\tau}_\ell(u_x, \rho)$ at $\ell = 0.19635$




FIG. 3. A 2D slice at $x = 0$ from a snapshot of the 3D compressible turbulence DNS, comparing the (left) exact $\bar{\tau}_\ell(u_x, \rho)$ at $\ell = 0.19635$ with (right) its approximation from Eq. (24c), $\tau_m = \frac{1}{3}M_2 \ell^2 \partial_k \bar{\rho} \partial_k \bar{u}_x$.

(domain size is 2π) with its approximation (24c). The other two components are shown in the Appendix Fig. 16. All indicate an excellent agreement. Figure 4 (left column) shows a more quantitative evaluation of all three components of the subscale transport approximation, where we can see that the approximation is remarkably accurate almost everywhere but underestimates $\bar{\tau}_\ell(u_x, \rho)$ at locations of strong shocks. This underestimation indicates that higher order terms in the expansion (13) are needed for better agreement at those locations since these rare but strong discontinuities have a significant contribution from scales $< \ell$ and are an extreme test case for the approximation. Indeed, at a location x_0 of a shock, the increment scales as $\delta u(x_0; r) = u(x_0 + r) - u(x_0) \sim r^\sigma$, with $\sigma \approx 0$ for distances r larger than the viscous shock width (see Fig. 1 in [10] and the associated discussion). This value is at the margin of validity of the $\sigma > 0$ condition² for Eyink’s expansion (13) to converge.

Figure 4 (right column) shows the joint probability density function (PDF) between $\bar{\tau}_\ell(\mathbf{u}, \rho)$ and its approximation, with correlations exceeding 0.9. In Figs. 11–13 in the Appendix, we show similar comparisons at other length scales. The correlation coefficient between two data sets, A and B , is defined as

$$r(A, B) = \frac{\langle AB \rangle - \langle A \rangle \langle B \rangle}{[(\langle A^2 \rangle - \langle A \rangle^2)(\langle B^2 \rangle - \langle B \rangle^2)]^{1/2}}, \quad (31)$$

where $\langle \dots \rangle$ is a space-time average over the entire domain and time after which the flow has become statistically stationary. The joint PDF in Fig. 4 uses the so-called “z-scores” of τ_m and $\bar{\tau}_\ell$, denoted by τ_m^* and τ^* (all other figures show the actual $\bar{\tau}_\ell$ and τ_m values). The z-score of A is defined as

$$A^* = \frac{A - \langle A \rangle}{(\langle (A - \langle A \rangle)^2 \rangle)^{1/2}}, \quad (32)$$

which helps focus on the spatio-temporal variation patterns after subtracting the mean and normalizing by the standard deviation. For completeness, we plot the joint PDFs of the actual τ_m and $\bar{\tau}_\ell$ in Fig. 14 in the Appendix. Using z-scores in the joint PDF only affects the slope of the linear fit

²Formally, the condition $\sigma > 0$ for Eyink’s expansion to converge is not based on the scaling of an increment $\delta u(x; r)$ at a single point x . Rather, it is based on the scaling of structure functions, which are spatial averages of increments.




FIG. 4. Comparison of $\bar{\tau}_\ell(\rho, \mathbf{u})$ with its approximation $\tau_m = \frac{1}{3} M_2 \ell^2 \partial_k \bar{\rho} \partial_k \bar{\mathbf{u}}$ at $\ell = 0.19635$. (a) x component of τ and τ_m along a diagonal transect through the domain from a single snapshot. (b) Joint PDF between τ^* and τ_m^* , where the asterisk superscript indicates z-scores, which emphasizes trends [see Eq. (32) and associated discussion in the text]. Color bar indicates the number of grid points on a logarithmic scale, proportional to the probability. Black line is the best linear fit, and the correlation squared, r^2 , quantifies the fraction of variability in τ that is captured by τ_m . (c), (d) Same as (a) and (b) but for the y component. (e), (f) Same as (a) and (b) but for the z component.

between τ_m and $\bar{\tau}_\ell$. The correlation coefficient is the same whether using the actual values or their z-scores,

$$r(A^*, B^*) = r(A, B). \quad (33)$$

Therefore, z-scores show how well the approximation τ_m is able to capture the exact subscale physics $\bar{\tau}_\ell$ up to a proportionality constant.

Figure 5 plots the correlation coefficient as a function of ℓ , which has values exceeding 0.90 for scales ℓ smaller than that of the peak and decreasing to ≈ 0.85 for scales $\ell = \pi$ (half the domain




FIG. 5. Correlation coefficient r between $\bar{\tau}_\ell(\rho, \mathbf{u})$ and its approximation $\tau_m = \frac{1}{3} M_2 \ell^2 \partial_k \bar{\rho} \partial_k \bar{\mathbf{u}}$ as a function of scale ℓ . (a) x component; (b) y component; (c) z component.

size). These strong correlations justify our usage of expression (24c) as a proxy for $\bar{\tau}_\ell(\mathbf{u}, \rho)$ to gain insight into its underlying mechanics.

B. Geophysical fluid turbulence

We also ran simulations of the two-layer stacked shallow water equations (34) and (35) using the ocean general circulation model MOM6 [43,44]. We simulated an ocean current [45] that is baroclinically unstable [46,47], resulting in mesoscale eddy generation and geostrophic turbulence, which we visualize in Fig. 7.

Details of the simulation setup can be found in [45]. The equations solved in MOM6 are

$$\frac{\partial \mathbf{u}_m}{\partial t} + (f + \omega) \hat{\mathbf{z}} \times \mathbf{u}_m = -\nabla(M_m + |\mathbf{u}_m|^2/2) - \nabla \cdot \mathbf{T} - \delta_{m2} C_D |\mathbf{u}_2| \mathbf{u}_2, \quad (34)$$

$$\frac{\partial h_m}{\partial t} + \nabla \cdot (h_m \mathbf{u}_m) = (3 - 2m)[\gamma(\bar{\eta}_{3/2}^x - \eta_{3/2,\text{ref}}) - \nabla \cdot (K_h \nabla \eta_{3/2})]. \quad (35)$$

The subscript $m = 1, 2$ labels the top and bottom layers, respectively, and repeated indices are not summed in the thickness equation (35). $\eta_{1/2}$ is the vertical position of the top layer interface




FIG. 6. Initial condition for two-layer stacked shallow water simulation following [45]. The top layer has zonal (east-west) velocity with a Gaussian profile peaking at ≈ 0.35 m/s (positive to the east) while the bottom layer is initially quiescent. Bottom topography is flat. The surface has initial height $\eta_{1/2} = 0$ m. The interface height $\eta_{3/2}$ between the two layers is initialized to the reference $\eta_{3/2,\text{ref}}$ shown to satisfy geostrophic balance. $\eta_{3/2}$ is allowed to evolve freely while being relaxed back to $\eta_{3/2,\text{ref}}$ at the rate γ .




FIG. 7. Visualization of the relative vorticity normalized by the Coriolis parameter f at the domain's midlatitude from our simulation of an eastward flowing ocean current that is baroclinically unstable, resulting in eddy generation and turbulence. The simulation solves the two-layer shallow water equations, with the top and bottom panels showing the respective top and bottom layers. Time increases from left to right: panels (a1) and (a2) are from day 60, (b1) and (b2) from day 93, and (c1) and (c3) from day 180. Our analysis uses model data after day 169, when the flow has become statistically steady.

(surface height), $\eta_{3/2}$ is the height of the interface between the two layers, $\mathbf{u}_m$ is the horizontal velocity, ω is vertical component of relative vorticity, $f = f_0 + \beta y$ is the Coriolis parameter for Earth's rotation (i.e., the planetary vorticity), where f_0 is the value of f at southernmost boundary ($y = 0$) and β is the linear rate of change of f along meridional direction. $\mathbf{T}$ is the horizontal stress tensor parameterized by Smagorinsky biharmonic viscosity [48], C_D is the dimensionless bottom drag coefficient and $M_1 = g \eta_{1/2}$ and $M_2 = M_1 + (g \frac{\Delta \rho}{\rho_0}) \eta_{3/2}$ are the pressure of the layers 1 and 2 normalized by ρ_0 , the density of the top layer. $\Delta \rho$ is the density difference between the two layers, h_m is the layer thickness, K_h is the interface height diffusivity coefficient, and γ is a rate that is proportional to the mass flux between the layers [45]. K_h allows for the dissipation of available potential energy [49] and γ forces the interface height between the two layers to reference zonal mean state $\eta_{3/2, \text{ref}}$. Here $(\cdot)^x$ represents a zonal average (i.e., in the east-west direction). The flow is able to attain a statistically steady state in the absence of direct wind forcing in Eq. (34) by nudging the layer interface $\eta_{3/2}$ to the reference profile $\eta_{3/2, \text{ref}}$ in Eq. (35), which drives the flow by providing a source of potential energy [45], mimicking the sloping of isopycnals by Ekman pumping in the real ocean [47].

The flat bottom domain is 1200 km $\times$ 1600 km in horizontal extent and is discretized on a Cartesian grid with a grid-cell size of 5 km. The total depth is 2 km and the bottom surface is flat. It is zonally periodic and subject to free-slip walls at the northern and southern boundaries. The initial conditions have a zonal jet 200 km wide with a Gaussian profile peaking at ≈ 0.35 m/s for top layer and bottom layer is quiescent, similar to initial conditions of [45]. The interface between the two layers is such that it satisfies geostrophic balance and is as shown in Fig. 6. The Coriolis parameter at




FIG. 8. Comparing the (left) exact $\bar{\tau}_\ell(\mathbf{u}, h)$ at $\ell = 100$ km with (right) its approximation from Eq. (24c), $\tau_m = \frac{1}{2} M_2 \ell^2 \partial_k \bar{h} \partial_k \bar{\mathbf{u}}$, from a single snapshot. Top panels are for the zonal (eastward) component. Bottom panels are for the meridional (northward) component. Gray bands at northern and southern boundaries are excluded from our analysis to avoid details of the coarse graining next to boundaries.

the southernmost boundary is $f_0 = 6.49 \times 10^{-5} \text{ s}^{-1}$ and $\beta = \partial_y f = 2 \times 10^{-11} \text{ m}^{-1} \text{ s}^{-1}$. The value of $\gamma = 1.1574 \times 10^{-6} \text{ s}^{-1}$ and $K_h = 8000 \text{ m}^2 \text{ s}^{-1}$. The simulation is run for 180 days, sufficiently long for the geostrophic turbulence to develop as a result of baroclinic instabilities. For our results, we analyze 100 snapshots taken every 2.5 hours starting from day 169. Spectra of $\mathbf{u}_1$ and h_1 are shown in Appendix Fig. 21. For our purpose, we only analyze the fields from the top layer since the flow in the bottom layer is much weaker. Hereafter we drop the layer subscript since all results pertain to the top layer.

We compare $\bar{\tau}_\ell(h, \mathbf{u})$ with its approximation $\frac{1}{2} \ell^2 M_2 \partial_k \bar{h} \partial_k \bar{\mathbf{u}}$ in Eq. (24c). We use a graded Top Hat kernel following our previous work analyzing oceanic flow [50, 51],

$$G_\ell(r) = 0.5 - 0.5 \tanh[(r - \ell/2)/10], \quad (36)$$

where ℓ denotes the kernel width. This kernel is essentially the same as the Top Hat in Eq. (7), but with smoothed edges to avoid discretization noise from the nonuniform grids [50] commonly used in general circulation models. Since the model domain is flat, r is a simple Euclidean distance as in Eq. (7), without complications due to Earth's surface curvature [52]. Figure 8 compares the $\bar{\tau}_\ell(\mathbf{u}, h)$ at $\ell = 100$ km with its approximation (24c). It indicates an excellent agreement. Figure 9 (left column) shows a more quantitative evaluation of both components of the subscale transport approximation, where we can see that the approximation is remarkably accurate almost everywhere. Figure 9 (right column) shows the joint PDFs between $\bar{\tau}_\ell(\mathbf{u}, \rho)$ and its approximation




FIG. 9. Comparison of $\bar{\tau}_\ell(\mathbf{h}, \mathbf{u})$ with its approximation $\tau_m = \frac{1}{2} M_2 \ell^2 \partial_k \bar{h} \partial_k \bar{\mathbf{u}}$ at $\ell = 100$ km. (a) Zonal (eastward) component of τ and τ_m along the transect $y = 802.5$ km from a snapshot on day 171 of the simulation. (b) Joint PDF for all space-time points between τ^* and τ_m^* , where the asterisk superscript indicates Z scores, which emphasizes trends [see Eq. (32) and associated discussion in the text]. Color bar indicates the number of grid points on a logarithmic scale, proportional to the probability. Red line is the best linear fit, and the correlation squared, r^2 , quantifies the fraction of variability in τ that is captured by τ_m .

for both components (eastward and northward directions). They show correlations exceeding 0.95. In Figs. 18–20 in the Appendix, we show similar comparisons at other length scales of 50 km, 200 km, and 400 km. The joint PDF in Fig. 9 uses the z-scores of $\bar{\tau}_\ell$ and τ_m , denoted by τ^* and τ_m^* and (all other figures show the actual $\bar{\tau}_\ell$ and τ_m values). As mentioned in Eq. (33), the correlation coefficient between τ^* and τ_m^* is the same as that between $\bar{\tau}_\ell$ and τ_m . We use z-scores to focus on how well the approximation τ_m is able to capture the exact subscale physics $\bar{\tau}_\ell$ up to a proportionality constant. For completeness, we plot the joint PDFs of the actual τ_m and $\bar{\tau}_\ell$ in Fig. 25 in the Appendix.

Noting that the spectral peak (Fig. 21) is at ≈ 300 km, Fig. 10 plots the correlation coefficients between the exact $\bar{\tau}_\ell(\mathbf{u}, h)$ and its approximation (24c) as a function of ℓ , which has values exceeding 0.90 for scales ℓ smaller than the that of the spectral peak and decreasing to ≈ 0.75 for scales $\ell = 400$ km. These high correlation values justify our usage of expression (24c) as a proxy for $\bar{\tau}_\ell(\mathbf{u}, h)$ to gain insight into its underlying mechanics at least for scales ℓ smaller than that of the spectral peak.

IV. TRANSPORT BY SUBSCALE VORTICITY AND STRAIN

Insight from Eq. (24c) into how the subscale physics contributes to large-scale transport can be gleaned by using a fundamental result from [53–55], which showed in different physical systems the importance of scalar transport by an antisymmetric skew diffusion tensor, or equivalently an eddy-induced advective flux. We elaborate on this idea in this section, which contains the main result of the paper in Eq. (43).




FIG. 10. Correlation coefficient r between $\bar{\tau}_\ell(h, \mathbf{u})$ and its approximation $\tau_m = \frac{1}{2} M_2 \ell^2 \partial_k \bar{h} \partial_k \bar{\mathbf{u}}$ as a function of scale ℓ . (a) Zonal (eastward) direction; (b) meridional (northward) direction.

Replacing $\bar{\tau}_\ell(\mathbf{u}, C)$ in Eq. (10) by expression (24c), we have

$$\partial_i \bar{C}_\ell + \nabla \cdot (\bar{\mathbf{u}}_\ell \bar{C}_\ell) = \nabla \cdot (-A \ell^2 \partial_j \bar{\mathbf{u}}_\ell \partial_j \bar{C}_\ell), \quad (37)$$

with the constant $A = M_2/n$. Therefore, transport of $\bar{C}_\ell$ by the subscale flux,

$$R(C) = \nabla \cdot \underbrace{[(-A \ell^2 \partial_j \bar{\mathbf{u}}_\ell) \partial_j \bar{C}_\ell]}_{K_{ij}}, \quad (38)$$

resembles that of generalized transport tensor

$$K_{ij} = -A \ell^2 \partial_j \bar{u}_i. \quad (39)$$

Note that K_{ij} is a function of the length scale ℓ through both the ℓ^2 factor as well as the coarse-grained velocity $\bar{\mathbf{u}}_\ell$. For notational brevity we drop the ℓ when writing individual components of $\bar{\mathbf{u}}_\ell$.

Physical properties of the subscale transport tensor K_{ij} can be illuminated by separating it into a symmetric part K_{ij}^{sym} representing subscale strain, and antisymmetric part (also called skew-symmetric) K_{ij}^{skew} representing subscale vorticity. The tensor is rewritten as

$$K_{ij} = -A \ell^2 \partial_j \bar{u}_i = -A \ell^2 \left[\underbrace{\frac{1}{2}(\partial_j \bar{u}_i + \partial_i \bar{u}_j)}_{\bar{s}_{ij}} + \underbrace{\frac{1}{2}(\partial_j \bar{u}_i - \partial_i \bar{u}_j)}_{\bar{\Omega}_{ij}} \right] = K_{ij}^{\text{sym}} + K_{ij}^{\text{skew}} \quad (40)$$

with vorticity and strain underlying kinematically distinct flow processes (e.g., [55]). Note that the symmetric part satisfies $K_{ij}^{\text{sym}} = K_{ji}^{\text{sym}}$ while the antisymmetric part satisfies $K_{ij}^{\text{skew}} = -K_{ji}^{\text{skew}}$.

Transport due to the subscale strain,

$$R^{\text{sym}}(C) = \partial_i [K_{ij}^{\text{sym}} \partial_j \bar{C}_\ell], \quad (41)$$

can be thought of in the local eigenframe reference of K_{ij}^{sym} as anisotropic diffusion/antidiffusion of $\bar{C}_\ell$ in directions that are not necessarily parallel to $\nabla \bar{C}_\ell$. Such transport³ was analyzed in detail in [13]. Positive eigenvalues are associated with a stretching flow direction (diffusion) while negative eigenvalues are associated with a contracting flow direction (antidiffusion). If the underlying flow is nonsolenoidal, $\nabla \cdot \mathbf{u} \neq 0$, such as in our simulations presented here, then K_{ij}^{sym} can also account for isotropic diffusion or coalescence due to the subscale physics.

³The tensor K^{sym} would have to be positive semidefinite to qualify as a diffusion tensor in the strict sense.

To understand the role of subscale vorticity, represented by the antisymmetric (or skew-symmetric) tensor K_{ij}^{skew} , we follow the analysis of [53–55] and rewrite

$$R^{\text{skew}}(C) = \partial_i [K_{ij}^{\text{skew}} \partial_j \bar{C}_\ell] = [\partial_i K_{ij}^{\text{skew}}] \partial_j \bar{C}_\ell = \partial_j [\underbrace{\partial_i K_{ij}^{\text{skew}}}_{-(\mathbf{v}_*)_j} \bar{C}_\ell], \quad (42)$$

where the second equality follows from the chain rule and noting that the product of K_{ij}^{skew} (antisymmetric tensor) with $\partial_i \partial_j \bar{C}_\ell$ (symmetric tensor) vanishes. Similarly, the third equality follows from $\partial_i \partial_j K_{ij}^{\text{skew}} = 0$. The interpretation of $\mathbf{v}_*$ is an eddy-induced velocity arising from the stirring induced by subscale vortical motions. It is straightforward to show that

$$\mathbf{v}_* = -\nabla \cdot \mathbf{K}^{\text{skew}} = \frac{1}{2} A \ell^2 \nabla \times (\nabla \times \bar{\mathbf{u}}_\ell), \quad (43)$$

which is the main result of this paper. An example of how it can arise in swirling flow is sketched in Fig. 1. Note that this eddy-induced velocity is solenoidal,

$$\nabla \cdot \mathbf{v}_* = 0, \quad (44)$$

due to the antisymmetry of K_{ij}^{skew} .

A. Summary and interpretation

The above result is a manifestation of the equivalence between transport by (1) flux due to an antisymmetric diffusion tensor and (2) an advective flux [53–55]. It suggests that coarse-grained simulations such as LES may need to solve

$$\partial_t \bar{C}_\ell + \nabla \cdot [(\bar{\mathbf{u}}_\ell + \mathbf{v}_*) \bar{C}_\ell] = -\nabla \cdot [\mathbf{J}(C)] \quad (45)$$

to represent the unresolved (subgrid) vorticity physics self-consistently.

$\mathbf{J}(C)$ in Eq. (45) can represent traditional subgrid models such as turbulent (Smagorinsky) diffusion [56], $\mathbf{J}(C) = -\alpha_{\text{turb}} \nabla \bar{C}_\ell$. The evaluation of $\mathbf{v}_* \sim \nabla \times (\nabla \times \bar{\mathbf{u}}_\ell)$ presents no additional cost since the resolved velocity $\bar{\mathbf{u}}_\ell$ is known. Figures 17–26 in the Appendix show the joint PDFs of $\mathbf{v}_*$ and $\bar{\mathbf{u}}_\ell$ at different length scales ℓ , which demonstrate that $\mathbf{v}_*$ and the coarse velocity $\bar{\mathbf{u}}_\ell$ are often of a comparable magnitude.

Our interpretation of $\mathbf{v}_*$ as an eddy-induced velocity representing subscale vortical motions can be justified by appealing to a relation by Johnson⁴ [35,36],

$$\bar{\tau}_\ell(u_i, C) = 2 \int_0^\ell d\delta \delta \overline{(\partial_k \bar{u}_i)_\delta \partial_k \bar{C}_\delta} / \sqrt{\ell^2 - \delta^2}. \quad (46)$$

Equation (46) highlights that subscale flux $\bar{\tau}_\ell(\mathbf{u}, C)$ is due to the cumulative contribution of subscale velocity and scalar gradients at all scales $< \ell$. Since subscale velocity gradients can be decomposed into strain and vorticity, and since expression (24c) is the dominant contribution, we are justified in interpreting $\mathbf{v}_*$ as an eddy-induced velocity representing subscale vortical motions.

V. CONCLUSION

We have shown that unlike subscale strain, which acts as an anisotropic diffusion/antidiffusion tensor, subscale vorticity’s contribution at leading order is solely a conservative advection of coarse-grained scalars by an eddy-induced velocity $\mathbf{v}_*$ proportional to the curl of vorticity. Our analysis relied on the leading order term in Eyink’s multiscale gradient expansion, which coincides with the nonlinear model and is known to be an excellent *a priori* representation of the subscale physics. Here we provided additional evidence for this excellent agreement from a three-dimensional compressible


⁴Johnson’s relation (46) is exact only when filtering with a Gaussian kernel.


turbulence simulation and from a two-layer shallow water model of geophysical turbulence. Since the convergence of Eyink’s expansion and, therefore, the dominance of the leading order term relies on ultraviolet scale locality, our results and conclusions may not hold at length scales larger than those of the spectral peak.

While the focus of this paper was on the transport of scalars, a similar analysis may also apply to the transport of momentum. Indeed, this is straightforward for incompressible flows where approximation (24c) has been shown to work very well for momentum transport [28,29,34]. Johnson’s recent work on the energy cascade [36] seems to indicate that approximation (24c) may be deficient in its magnitude (by a factor $\approx \frac{1}{2}$) compared to the true momentum flux. Yet, since the two exhibit excellent spatiotemporal correlation as was shown in [29,34], a simple rescaling of expression (24c) by a space-time independent coefficient (possibly ≈ 2) would suffice to match the true momentum flux. Once approximation (24c) (possibly $\times 2$) is accepted as representative of the subscale flux, the same argument we presented here follows through for momentum after replacing C with $\mathbf{u}$ in Eq. (24c). In flows with significant density variations such as compressible turbulence, the subscale flux of momentum has a somewhat different functional form than in Eq. (10b) due to the density-weighted filtering required to satisfy the inviscid criterion [14,57]. Yet we conjecture that a result similar to what we derived here is still possible for subscale momentum flux in variable-density flows, which was shown to be scale-local in [24].

ACKNOWLEDGMENTS

We thank Brandon Reichl at GFDL and the reviewers for valuable feedback and suggestions. This research was funded by U.S. DOE Grant No. DE-SC0020229. Partial support from U.S. NSF Grants No. PHY-2020249, No. OCE-2123496, U.S. NASA Grant No. 80NSSC18K0772, and U.S. NNSA Grant No. DE-NA0003856 is acknowledged. H.A. was also supported by U.S. DOE Grants No. DE-SC0014318, DE-SC0019329, NSF Grant No. PHY-2206380, and U.S. NNSA Grant No. DE-NA0003914. J.K.S. was also supported by U.S. DOE Grant No. DE-SC0019329, U.S. NNSA Grant No. DE-NA0003914, and U.S. NSF Grant No. CBET-2143702. Computing time was provided by NERSC under Contract No. DE-AC02-05CH11231 and NASA’s HEC Program through NCCS at Goddard Space Flight Center. MOM6 is publicly available and can be accessed at [60]. The coarse-graining analysis was performed using codes similar to FlowSieve, which is publicly available at [61]. Other data used here may be obtained from the corresponding author upon reasonable request.


 FIG. 11. Same as in Fig. 4 for $\ell = 0.3927$.

FIG. 12. Same as Fig. 4 for $\ell = 0.7854$.




FIG. 13. Same as Fig. 4 for $\ell = 1.5708$.

APPENDIX

1. Additional supporting material from the compressible turbulence simulation
2. Additional supporting material from the shallow water simulation




FIG. 14. Similar to the joint PDFs of the z-scores in Fig. 4, here we show the joint PDFs of the actual $\bar{\tau}_\ell(\rho, \mathbf{u})$ and its approximation $\tau_m = \frac{1}{3}M_2\ell^2\partial_k\bar{\rho}\partial_k\bar{\mathbf{u}}$. Top row (a)–(c) shows the joint PDFs between τ and τ_m at $\ell = 0.19635$. Second row (d)–(f) is for $\ell = 0.3927$. Third row (g)–(i) is for $\ell = 0.7854$. Bottom row (j)–(l) is for $\ell = 1.5708$. Left, middle, and right columns show the x , y , and z components, respectively.




FIG. 15. Spectra of (left) velocity, $E^u(k)$, and (right) density, $E^\rho(k)$. Both decay with wave number k sufficiently faster than the k^{-1} required for the multiscale gradient expansion to converge. Here $E^u(k) = \sum_{|\mathbf{k}|-k|<0.5} \frac{1}{2} |\hat{\mathbf{u}}(\mathbf{k})|^2$ and $E^\rho(k) = \sum_{|\mathbf{k}|-k|<0.5} |\hat{\rho}(\mathbf{k})|^2$, where $\hat{\mathbf{u}}(\mathbf{k})$ and $\hat{\rho}(\mathbf{k})$ are Fourier coefficients.




FIG. 16. Same as in Fig. 3, comparing the (left column) exact $\bar{\tau}_\ell(\rho, \mathbf{u})$ with (right column) its approximation for the (top row) y components and (bottom row) z components.








FIG. 17. Joint PDFs of $\mathbf{v}_*$ and $\bar{\mathbf{u}}$. Color bar indicates the number of grid points on a logarithmic scale, proportional to the probability. Slope of best linear fit and correlation squared, r^2 , are shown in the top left of each panel. The best linear fits are omitted in the the plots due to small slopes. The rms values are shown in the bottom right of each panel. Top row (a)–(c) shows the joint PDFs between $\mathbf{v}_*$ and $\bar{\mathbf{u}}$ at $\ell = 0.19635$. Second row (d)–(f) is for $\ell = 0.3927$. Third row (g)–(i) is for $\ell = 0.7854$. Bottom row (j)–(l) is for $\ell = 1.5708$. Left, middle, and right columns show the x , y , and z components, respectively. This figure highlights that the magnitudes of $\mathbf{v}_*$ and the coarse velocity $\bar{\mathbf{u}}_\ell$ are often comparable.


 FIG. 18. Same as in Fig. 9 for $\ell = 50$ km.

 FIG. 19. Same as in Fig. 9 for $\ell = 200$ km.

FIG. 20. Same as in Fig. 9 for $\ell = 400$ km.FIG. 21. Spectra of (left) u and (right) h computed from the two-layer stacked shallow water simulation. These are filtering spectra, calculated by filtering at successive length scales as described in [58,59].




FIG. 22. Same as in Fig. 8 for $\ell = 50$ km.




FIG. 23. Same as in Fig. 8 for $\ell = 200$ km.



FIG. 24. Same as in Fig. 8 for $\ell = 400$ km.

FIG. 25. Similar to the joint PDFs of the z-scores in Fig. 9, here we show the joint PDFs of the actual $\bar{\tau}_\ell(h, \mathbf{u})$ and its approximation $\tau_m = \frac{1}{2} M_2 \ell^2 \partial_k \bar{h} \partial_k \bar{\mathbf{u}}$, without normalization. Top and bottom panels show zonal (eastward) and meridional (northward) components, respectively. (a1) and (a2) are for $\ell = 50$ km, (b1) and (b2) for 100 km, (c1) and (c2) for 200 km, (d1) and (d2) for 400 km.




FIG. 26. Joint probability density of $\mathbf{v}_*$ and $\bar{\mathbf{u}}$. The colorbars are in log scale. The top panels show zonal (eastward) component and the bottom panels show meridional (northward) component. (a1) and (a2) are for $\ell = 50$ km, (b1) and (b2) are for $\ell = 100$ km, (c1) and (c2) are for $\ell = 200$ km, (d1) and (d2) are for $\ell = 400$ km. The red lines are linear regression fits of which the slope and goodness of fit (r^2) are in the text in each panel. The rms values are shown in the bottom right of each panel. This figure highlights that $\mathbf{v}_*$ is often comparable in magnitude relative to the coarse velocity $\bar{\mathbf{u}}_\ell$.

- [1] P. K. Kundu, I. M. Cohen, and D. R. Dowling, *Fluid Mechanics* (Academic Press, Boston, 2015).
- [2] S. B. Pope, *Turbulent Flows* (Cambridge University Press, New York, 2000).
- [3] C. Meneveau and J. Katz, Scale-invariance and turbulence models for large-eddy simulation, *Annu. Rev. Fluid Mech.* **32**, 1 (2000).
- [4] G. L. Eyink, Locality of turbulent cascades, *Physica D* **207**, 91 (2005).
- [5] R. S. Strichartz, *A Guide to Distribution Theory and Fourier Transforms* (World Scientific, New Jersey, 2003).
- [6] L. C. Evans, *Partial Differential Equations* (American Mathematical Society, Providence, RI, 2010).
- [7] A. Leonard, Energy cascade in large-eddy simulations of turbulent fluid flows, *Adv. Geophys.* **18**, A237 (1974).
- [8] M. Germano, Turbulence: The filtering approach, *J. Fluid Mech.* **238**, 325 (1992).
- [9] I. Daubechies, *Ten Lectures on Wavelets* (SIAM, Philadelphia, 1992).
- [10] H. Aluie, Coarse-grained incompressible magnetohydrodynamics: Analyzing the turbulent cascades, *New J. Phys.* **19**, 025008 (2017).
- [11] G. L. Eyink, Review of the Onsager “ideal turbulence” theory, [arXiv:1803.02223](https://arxiv.org/abs/1803.02223).
- [12] F. F. Grinstein, L. G. Margolin, and W. J. Rider, *Implicit Large Eddy Simulation* (Cambridge University Press, Cambridge, 2007), Vol. 10.
- [13] C. W. Higgins, M. B. Parlange, and C. Meneveau, The heat flux and the temperature gradient in the lower atmosphere, *Geophys. Res. Lett.* **31** (2004).
- [14] H. Aluie, Scale decomposition in compressible turbulence, *Physica D* **247**, 54 (2013).
- [15] D. Zhao and H. Aluie, Inviscid criterion for decomposing scales, *Phys. Rev. Fluids* **3**, 054603 (2018).
- [16] M. Germano, A proposal for a redefinition of the turbulent stresses in the filtered Navier–Stokes equations, *Phys. Fluids* **29**, 2323 (1986).
- [17] G. L. Eyink, Multi-scale gradient expansion of the turbulent stress tensor, *J. Fluid Mech.* **549**, 159 (2006).
- [18] A. N. Kolmogorov, The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers, *Dokl. Akad. Nauk SSSR* **30**, 299 (1941).

- [19] A. M. Obukhov, The structure of the temperature field in a turbulent flow, *Izv. Akad. Nauk. SSSR, Ser. Geogr. Geophys.* **13**, 58 (1949).
- [20] S. Corrsin, On the spectrum of isotropic temperature fluctuations in an isotropic turbulence, *J. Appl. Phys.* **22**, 469 (1951).
- [21] G. L. Eyink and H. Aluie, Localness of energy cascade in hydrodynamic turbulence. I. Smooth coarse graining *Phys. Fluids* **21**, 115107 (2009).
- [22] H. Aluie and G. L. Eyink, Localness of energy cascade in hydrodynamic turbulence. II. Sharp spectral filter, *Phys. Fluids* **21**, 115108 (2009).
- [23] J. Bardina, J. H. Ferziger, and W. C. Reynolds, Improved subgrid-scale models for large-eddy simulation, *13th Fluid and Plasma Dynamics Conference* (American Institute of Aeronautics and Astronautics, Reston, VA, 1980).
- [24] H. Aluie, Compressible Turbulence: The Cascade and Its Locality, *Phys. Rev. Lett.* **106**, 174502 (2011).
- [25] M. Frazier, M. W. Frazier, B. Jawerth, and G. Weiss, *Littlewood-Paley Theory and the Study of Function Spaces*, Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics Vol. 79 (American Mathematical Society, Providence, RI, 1991).
- [26] B. Vreman, B. Geurts, and H. Kuerten, Realizability conditions for the turbulent stress tensor in large-eddy simulation, *J. Fluid Mech.* **278**, 351 (1994).
- [27] P. Constantin, E. Weinan, and E. S. Titi, Onsager’s conjecture on the energy conservation for solutions of Euler’s equation, *Commun. Math. Phys.* **165**, 207 (1994).
- [28] R. A. Clark, J. H. Ferziger, and W. C. Reynolds, Evaluation of subgrid-scale models using an accurately simulated turbulent flow, *J. Fluid Mech.* **91**, 1 (1979).
- [29] S. Liu, C. Meneveau, and J. Katz, On the properties of similarity subgrid-scale models as deduced from measurements in a turbulent jet, *J. Fluid Mech.* **275**, 83 (1994).
- [30] F. Porté-Agel, M. B. Parlange, C. Meneveau, and W. E. Eichinger, A priori field study of the subgrid-scale heat fluxes and dissipation in the atmospheric surface layer, *J. Atmos. Sci.* **58**, 2673 (2001).
- [31] B. Kosović, Subgrid-scale modelling for the large-eddy simulation of high-Reynolds-number boundary layers, *J. Fluid Mech.* **336**, 151 (1997).
- [32] F. Bouchet, Parameterization of two-dimensional turbulence using an anisotropic maximum entropy production principle, [arXiv:cond-mat/0305205](https://arxiv.org/abs/cond-mat/0305205).
- [33] A. Leonard, Large-eddy simulation of chaotic convection and beyond, in *35th Aerospace Sciences Meeting and Exhibit* (AIAA, Reno, Nevada, 1997), p. 204.
- [34] V. Borue and S. A. Orszag, Local energy flux and subgrid-scale statistics in three-dimensional turbulence, *J. Fluid Mech.* **366**, 1 (1998).
- [35] P. L. Johnson, Energy Transfer from Large to Small Scales in Turbulence by Multiscale Nonlinear Strain and Vorticity Interactions, *Phys. Rev. Lett.* **124**, 104501 (2020).
- [36] P. L. Johnson, On the role of vorticity stretching and strain self-amplification in the turbulence energy cascade, *J. Fluid Mech.* **922**, A3 (2021).
- [37] H. S. Kang and C. Meneveau, Passive scalar anisotropy in a heated turbulent wake: New observations and implications for large-eddy simulations, *J. Fluid Mech.* **442**, 161 (2001).
- [38] S. G. Chumakov, A priori study of subgrid-scale flux of a passive scalar in isotropic homogeneous turbulence, *Phys. Rev. E* **78**, 036313 (2008).
- [39] A. Lees and H. Aluie, Baropycnal work: A mechanism for energy transfer across scales, *Fluids* **4**, 92 (2019).
- [40] S. Jagannathan and D. A. Donzis, Reynolds and Mach number scaling in solenoidally-forced compressible turbulence using high-resolution direct numerical simulations, *J. Fluid Mech.* **789**, 669 (2016).
- [41] C. Federrath, R. S. Klessen, and W. Schmidt, The density probability distribution in compressible isothermal turbulence: Solenoidal versus compressive forcing, *Astrophys. J.* **688**, L79 (2008).
- [42] V. Eswaran and S. B. Pope, An examination of forcing in direct numerical simulations of turbulence, *Comput. Fluids* **16**, 257 (1988).
- [43] A. Adcroft, W. Anderson, V. Balaji, C. Blanton, M. Bushuk, C. O. Dufour, J. P. Dunne, S. M. Griffies, R. Hallberg, M. J. Harrison *et al.*, The GFDL global ocean and sea ice model OM4.0: Model description and simulation features, *J. Adv. Model. Earth Syst.* **11**, 3167 (2019).

- [44] S. M. Griffies, A. Adcroft, and R. W. Hallberg, A primer on the vertical Lagrangian-remap method in ocean models based on finite volume generalized vertical coordinates, *J Adv. Model Earth Syst.* **12**, e2019MS001954 (2020).
- [45] R. Hallberg, Using a resolution function to regulate parameterizations of oceanic mesoscale eddy effects, *Ocean Modelling* **72**, 92 (2013).
- [46] N. A. Phillips, Energy transformations and meridional circulations associated with simple baroclinic waves in a two-level, quasi-geostrophic model, *Tellus* **6**, 274 (1954).
- [47] G. K. Vallis, *Atmospheric and oceanic fluid dynamics: Fundamentals and large-scale circulation*, 2nd ed. (Cambridge University Press, Cambridge, 2017), <https://doi.org/10.1017/9781107588417>.
- [48] S. M. Griffies and R. W. Hallberg, Biharmonic friction with a Smagorinsky-like viscosity for use in large-scale eddy-permitting ocean models, *Mon. Weather Rev.* **128**, 2935 (2000).
- [49] P. R. Gent and J. C. McWilliams, Isopycnal mixing in ocean circulation models, *J. Phys. Oceanogr.* **20**, 150 (1990).
- [50] H. Aluie, M. Hecht, and G. K. Vallis, Mapping the energy cascade in the North Atlantic Ocean: The coarse-graining approach, *J. Phys. Oceanogr.* **48**, 225 (2018).
- [51] S. Rai, M. Hecht, M. Maltrud, and H. Aluie, Scale of oceanic eddy killing by wind from global satellite observations, *Sci. Adv.* **7**, eabf4920 (2021).
- [52] H. Aluie, Convolutions on the sphere: Commutation with differential operators, *Int. J. Geomath.* **10**, 9 (2019).
- [53] H. K. Moffatt, Transport effects associated with turbulence with particular attention to the influence of helicity, *Rep. Prog. Phys.* **46**, 621 (1983).
- [54] J. F. Middleton and J. W. Loder, Skew fluxes in polarized wave fields, *J. Phys. Oceanogr.* **19**, 68 (1989).
- [55] S. M. Griffies, The Gent–McWilliams skew flux, *J. Phys. Oceanogr.* **28**, 831 (1998).
- [56] J. Smagorinsky, General circulation experiments with the primitive equations: I. The basic experiment, *Mon. Weather Rev.* **91**, 99 (1963).
- [57] D. Livescu, Turbulence with large thermal and compositional density variations, *Annu. Rev. Fluid Mech.* **52**, 309 (2020).
- [58] M. Sadek and H. Aluie, Extracting the spectrum of a flow by spatial filtering, *Phys. Rev. Fluids* **3**, 124610 (2018).
- [59] M. Buzzicotti, B. A. Storer, S. M. Griffies, and H. Aluie, A coarse-grained decomposition of surface geostrophic kinetic energy in the global ocean, *Earth and Space Science Open Archive* (2021), p. 1–58, <https://doi.org/10.1002/essoar.10507290.1>.
- [60] <https://github.com/mom-ocean/MOM6>.
- [61] <https://github.com/husseinaluie/FlowSieve>.