

Order α_s^4 QCD Corrections to Z and τ Decays

P. A. Baikov

Institute of Nuclear Physics, Moscow State University, Moscow 119899, Russia

K. G. Chetyrkin* and J. H. Kühn

Institut für Theoretische Teilchenphysik, Universität Karlsruhe, D-76128 Karlsruhe, Germany

(Received 11 January 2008; published 3 July 2008)

Using recently developed methods for the evaluation of five-loop amplitudes in perturbative QCD, corrections of order α_s^4 for the nonsinglet part of the cross section for electron-positron annihilation into hadrons and for the decay rates of the Z boson and the τ lepton into hadrons are evaluated. The new terms lead to a significant stabilization of the perturbative series, to a reduction of the theory uncertainty in the strong coupling constant α_s , as extracted from these measurements, and to a small shift of the central value, moving the two central values closer together. The agreement between two values of α_s measured at vastly different energies constitutes a striking test of asymptotic freedom. Combining the results from Z and τ decays we find $\alpha_s(M_Z) = 0.1198 \pm 0.0015$ as one of the most precise and presently only result for the strong coupling constant in order α_s^4 .

DOI: [10.1103/PhysRevLett.101.012002](https://doi.org/10.1103/PhysRevLett.101.012002)

PACS numbers: 12.38.Bx, 13.35.Dx, 13.85.Lg

The strong coupling constant α_s is one of the three fundamental gauge coupling constants of the standard model (SM) of particle physics. Its precise determination is one of the most important aims of particle physics. Experiments at different energies allow us to test the predictions for its energy dependence based on renormalization group equations; the comparison of the results obtained from different processes leads to critical tests of the theory and potentially to the discovery of physics beyond the standard model. Last but not least, the convergence of the three gauge coupling constants related by $SU(3) \times SU(2) \times U(1)$ to a common value, after evolving them to high energies, allows us to draw conclusions about the possibility of embedding the SM in the framework of a grand unified theory.

One of the most precise and theoretically safe determinations of α_s is based on measurements of the cross section for electron-positron annihilation into hadrons. These have been performed in the low-energy region between 2 and 10 GeV and, in particular, at and around the Z resonance at 91.2 GeV. Conceptually closely related is the measurement of the semileptonic decay rate of the τ lepton, leading to a determination of α_s at a scale below 2 GeV.

From the theoretical side, in the framework of perturbative QCD, these rates and cross sections are evaluated as inclusive rates into massless quarks and gluons [1,2]. (Power suppressed mass effects are well under control for e^+e^- annihilation, both at low energies and around the Z resonance, and for τ decays [3–8], and the same applies to mixed QCD and electroweak corrections [9,10].)

The ratio $R(s) \equiv \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$ is expressed through the absorptive part of the correlator of the electromagnetic current j_μ :

$$R(s) = 12\pi \text{Im}\Pi(-s - i\epsilon), \quad (1)$$

$$3Q^2\Pi(Q^2) = i \int d^4x e^{iq \cdot x} \langle 0 | T j_\mu(x) j^\mu(0) | 0 \rangle, \quad (2)$$

with $Q^2 = -q^2$. It is also convenient to introduce the Adler function as

$$D(Q^2) = -12\pi^2 Q^2 \frac{d}{dQ^2} \Pi(Q^2) = \int_0^\infty \frac{Q^2 R(s) ds}{(s+Q^2)^2},$$

$$R(s) = D(s) - \pi^2 \beta_0^2 \left\{ \frac{d_1}{3} a_s^3 + \left(d_2 + \frac{5}{6\beta_0} d_1 \beta_1 \right) a_s^4 \right\} + \dots$$

We define the perturbative expansions

$$D(Q^2) = \sum_{i=0}^{\infty} d_i a_s^i(Q^2), \quad R(s) = \sum_{i=0}^{\infty} r_i a_s^i(s), \quad (3)$$

where $a_s \equiv \alpha_s/\pi$ and the normalization scale is set to $\mu^2 = Q^2$ or to $\mu^2 = s$ for the Euclidian and Minkowskian functions, respectively. The results for generic values of μ can be easily recovered with standard RG techniques.

Note that the first three terms of the perturbative series for D and R coincide. Starting from r_3 , terms proportional π^2 arise which can be predicted from those of lower order. It has been speculated that these “ π^2 -terms,” also called “kinematical terms,” might constitute a major part of the full higher order corrections (see, e.g., [11,12] and references therein); however, the validity of this hypothesis can only be established by the full calculation. Indeed, for the scalar correlator this assumption has been shown to fail [13].

For the vector correlator the terms of order a_s^2 and a_s^3 have been evaluated nearly 30 and about 15 years ago [14–16], respectively. The a_s^4 corrections are conveniently classified according to their power of n_f , with n_f denoting the number of light quarks. The $a_s^4 n_f^3$ term is part of the

“renormalon chain,” the evaluation of the next term, of order $a_s^4 n_f^2$, was a test case for the techniques used extensively in this paper and, furthermore, led to useful insights into the structure of the perturbative series already [17].

The complete five-loop calculation requires the evaluation of about 20 thousand diagrams (we have used QGRAF [18] for their automatic generation). Using “infrared rearrangement” [19], the R^* operation [20] and the prescriptions formulated in [21] to algorithmically resolve the necessary combinatorics, it is possible to express the absorptive part of the five-loop diagrams in terms of four-loop massless propagator integrals.

These integrals can be reduced to a sum of 28 master integrals with rational functions of the space-time dimension D as coefficients. The latter ones were fully reconstructed after evaluating sufficiently many terms of the $1/D$ expansion [22] of their representation proposed in [23]. This direct and largely automatic procedure required enormous computing resources and was performed using a parallel version [24] of FORM [25].

In this Letter we present the results for the so-called “nonsinglet” diagrams. These are sufficient for a complete description of τ decays. For e^+e^- annihilation through a virtual photon they correspond to the dominant terms proportional $\sum_i Q_i^2$. The singlet contributions proportional $(\sum_i Q_i)^2$ arise for the first time in $\mathcal{O}(\alpha_s^3)$. They are known to be small, and will be evaluated at a later point. Similar comments apply to the singlet contributions in Z decays.

The analytic result for the five-loop term in the Adler function is given by (we suppress the trivial factor $3\sum_f Q_f^2$ throughout)

$$\begin{aligned}
 d_4 = n_f^3 & \left[-\frac{6131}{5832} + \frac{203}{324}\zeta_3 + \frac{5}{18}\zeta_5 \right] \\
 & + n_f^2 \left[\frac{1045381}{15552} - \frac{40655}{864}\zeta_3 + \frac{5}{6}\zeta_3^2 - \frac{260}{27}\zeta_5 \right] \\
 & + n_f \left[-\frac{13044007}{10368} + \frac{12205}{12}\zeta_3 - 55\zeta_3^2 \right. \\
 & \left. + \frac{29675}{432}\zeta_5 + \frac{665}{72}\zeta_7 \right] + \frac{144939499}{20736} - \frac{5693495}{864}\zeta_3 \\
 & + \frac{5445}{8}\zeta_3^2 + \frac{65945}{288}\zeta_5 - \frac{7315}{48}\zeta_7. \quad (4)
 \end{aligned}$$

The knowledge of d_4 leads straightforwardly to R at order α_s^4 , for brevity given below in numerical form:

$$\begin{aligned}
 R = 1 + a_s & + (1.9857 - 0.1152n_f)a_s^2 \\
 & + (-6.63694 - 1.20013n_f - 0.00518n_f^2)a_s^3 \\
 & + (-156.61 + 18.77n_f - 0.7974n_f^2 + 0.0215n_f^3)a_s^4. \quad (5)
 \end{aligned}$$

It is also instructive to explicitly display the genuine five-loop contributions to d_4 [underlined in (6) and (7)] and the “kinematical” terms originating from the analytic continu-

ation:

$$\begin{aligned}
 r_3 = \underline{18.2} - 24.9 + (\underline{-4.22} + 3.02)n_f \\
 + (\underline{-0.086} + 0.091)n_f^2, \quad (6)
 \end{aligned}$$

$$\begin{aligned}
 r_4 = \underline{135.8} - 292.4 + (\underline{-34.4} + 53.2)n_f + (\underline{1.88} - 2.67)n_f^2 \\
 + (\underline{-0.010} + 0.032)n_f^3. \quad (7)
 \end{aligned}$$

Since it will presumably take a long time until the next term of the perturbative series will be evaluated, it is of interest to investigate the predictive power of various optimization schemes empirically. Using the principles of “fastest apparent convergence” (FAC) [26] or of “minimal sensitivity” (PMS) [27], which happen to coincide in this order, the central values of the predictions [11,28]

$$d_4^{\text{pred}}(n_f = 3, 4, 5) = 27 \pm 16, 8 \pm 18, -8 \pm 44$$

differ significantly from the exact result,

$$d_4^{\text{exact}}(n_f = 3, 4, 5) = 49.08, 27.39, 9.21. \quad (8)$$

However, within the error estimates [28], predicted and exact values are in agreement. The picture changes, once these estimates are used to predict the coefficient r_4 . Although sizable cancellations between “dynamical” and “kinematical” terms are observed for the individual n_f coefficients in (7) the predictions for the final results are significantly closer (in relative sense) to the results of the exact calculation:

$$\begin{aligned}
 r_4^{\text{pred}}(n_f = 3, 4, 5) & = -129 \pm 16, -112 \pm 30, -97 \pm 44, \\
 r_4^{\text{exact}}(n_f = 3, 4, 5) & = -106.88, -92.898, -79.98.
 \end{aligned}$$

(This is in striking contrast to the case of the scalar correlator, where the predictions for the dynamical terms work well, but, as a consequence of the strong cancellations between dynamical and kinematical terms fail in the Minkowskian region [13].)

Using FAC and the exact result for d_4 , the coefficients d_5 and r_5 can be predicted (following [11,28]) for $n_f = 3, 4, 5$, namely

$$d_5^{\text{pred}}(n_f = 3, 4, 5) = 275, 152, 89, \quad (9)$$

$$r_5^{\text{pred}}(n_f = 3, 4, 5) = -505, -134, 168. \quad (10)$$

These terms may become of relevance for the International Linear Collider (ILC) running in the GIGA-Z mode with an anticipated precision of $\delta\alpha_s = 0.0005\text{--}0.0007$ [29], and already today for the analysis of τ decays.

From the combined analysis of data for $\sigma(e^+e^- \rightarrow \text{hadrons})$ in the region between 3 and 10 GeV a value

$$\alpha_s(9 \text{ GeV}) = 0.182 \pm 0.033 \quad (11)$$

has been obtained recently [30]. The shift in α_s from the inclusion of the α_s^4 term amounts to $\delta\alpha_s(9 \text{ GeV}) = 0.003$

and is thus irrelevant compared to the large experimental error.

The situation is different for Z decays. The analysis of the electroweak working group [31] is based on Eq. (4) with $n_f = 5$, including term up to $\mathcal{O}(\alpha_s^3)$ and leads to

$$\alpha_s(M_Z)^{\text{NNLO}} = 0.1185 \pm 0.0026. \quad (12)$$

Since additional corrections (mixed QCD-electroweak or mass terms) are only weakly α_s dependent we may consider $R(s = M_Z^2)$ as a pseudo-observable:

$$R(s = M_Z^2) = 1.03904 \pm 0.00087. \quad (13)$$

Including the α_s^4 term leads to a shift $\delta\alpha_s(M_Z) = 0.0005$

$$\alpha_s(M_Z)^{\text{NNNLO}} = 0.1190 \pm 0.0026^{\text{exp}}, \quad (14)$$

The theory error may either be conservatively based on the shift produced by the last term (0.0005) or on the scale variation with $\mu/M_Z = \frac{1}{3}-3$, leading to ± 0.0002 and can be neglected in both cases.

Higher orders are of particular relevance in the low-energy region, for example, in τ decays. The correction from perturbative QCD to the ratio

$$R_{\tau,V+A} = \frac{\Gamma(\tau \rightarrow \text{hadrons}_{S=0} + \nu_\tau)}{\Gamma(\tau \rightarrow l + \bar{\nu}_l + \nu_\tau)} \\ = 3|V_{ud}|^2 S_{\text{EW}}(1 + \delta_0 + \delta'_{\text{EW}} + \delta_2 + \delta_{\text{NP}}) \quad (15)$$

is given by

$$1 + \delta_0 = 2 \int_0^{M_\tau^2} \frac{ds}{M_\tau^2} \left(1 - \frac{s}{M_\tau^2}\right)^2 \left(1 + \frac{2s}{M_\tau^2}\right) R(s). \quad (16)$$

In the subsequent analysis we will use $S_{\text{EW}} = 1.0198 \pm 0.0006$ and $\delta'_{\text{EW}} = 0.001$ for the electroweak corrections, $\delta_2 = (-4.4 \pm 2.0) \times 10^{-4}$ for light quark mass effects, $\delta_{\text{NP}} = (-4.8 \pm 1.7) \times 10^{-3}$ for the nonperturbative effects [2,32] and $V_{ud} = 0.97418 \pm 0.00027$ [33].

The perturbative quantity δ_0 can be evaluated in fixed order perturbation theory or with ‘‘contour improvement’’ as proposed in [34,35]

$$\delta_0^{\text{FO}} = a_s + 5.202a_s^2 + 26.366a_s^3 + 127.079a_s^4, \quad (17)$$

$$\delta_0^{\text{CI}} = 1.364a_s + 2.54a_s^2 + 9.71a_s^3 + 64.29a_s^4. \quad (18)$$

[To obtain the α_s -dependent coefficients in Eq. (18) we follow [34–37] and use $\alpha_s(M_\tau) = 0.334$ as reference value.] For the subsequent analysis we will use as starting point $\delta_0^{\text{exp}} = 0.1998 \pm 0.0043_{\text{exp}}$ as obtained from [36] and $R_{\tau,V+A} = 3.471 \pm 0.011$ which in turn is based on the ‘‘universality-improved’’ electronic branching ratio $B_e = (17.818 \pm 0.032)\%$ and the world average of the ratio of strange hadronic and electronic widths 0.1686 ± 0.0047 . The new values of $\alpha_s(M_\tau)$ in dependence on the choice of d_4 (with the previous estimate and the new exact result) are summarized in Table I.

TABLE I. Results for $\alpha_s(M_\tau)$ for different values of d_4 . The first line displays the α_s^3 results [with the α_s^4 terms set to zero in Eqs. (17) and (18)]. The second line uses the previously predicted value for d_4 , the last one uses the exact result (4). The first error is the experimental one; the second (theoretical) uncertainty in the value of α_s corresponds to changing the renormalization scale μ as follows $\mu^2/M_\tau^2 = 0.4-2$.

	$\alpha_s^{\text{FO}}(M_\tau)$	$\alpha_s^{\text{CI}}(M_\tau)$
...	$0.337 \pm 0.004 \pm 0.03$	$0.354 \pm 0.006 \pm 0.02$
$d_4 = 25$	$0.325 \pm 0.004 \pm 0.02$	$0.347 \pm 0.006 \pm 0.009$
$d_4 = 49.08$	$0.322 \pm 0.004 \pm 0.02$	$0.342 \pm 0.005 \pm 0.01$

As stated above the theory error for α_s from Z decays is small compared to the experimental uncertainties. The situation is more problematic for τ decays and to some extent the theory error remains to be matter of choice. As anticipated in [28] it decreases significantly, once α_s^4 terms are included. However, the difference between the two methods stabilizes (this was checked in [28] by adding an estimate for the α_s^5 term) and must be considered as irreducible uncertainty. Given the input specified above we obtain as our final result

$$\alpha_s(M_\tau) = 0.332 \pm 0.005_{\text{exp}} \pm 0.015_{\text{th}}. \quad (19)$$

For the central value we take the mean value of FO and CI. For the theory error we take half of the difference between two methods (0.01) plus (module of) the estimated correction from α_s^5 term (-0.005), the latter being based on $d_5 = 275$ [see Eq. (9)].

Applying four-loop running and matching [38–41] to (19) we arrive at

$$\alpha_s(M_Z) = 0.1202 \pm 0.0006_{\text{exp}} \pm 0.0018_{\text{th}} \pm 0.0003_{\text{evol}} \\ = 0.1202 \pm 0.0019. \quad (20)$$

Here the evolution error receives contributions from the uncertainties in the c -quark mass [0.00003 , $m_c(m_c) = 1.286(13)$ GeV [42]] and the b -quark mass [0.00001 , $m_b(m_b) = 4.164(25)$ GeV [42]], the matching scale (0.0001 , μ varied between $0.7m_q(m_q)$ and $3.0m_q(m_q)$), the four-loop truncation in the matching expansion (0.0001) and the four-loop truncation in the RGE equation (0.0003). (For the last two errors the size of the shift due the highest known perturbative term was treated as systematic uncertainty.) The errors are added in quadrature.

Summary: The exact result for the α_s^4 term in the Adler function allows to extract the strong coupling constant from Z and τ decays with high precision. Including the exact α_s^4 leads to small shifts of the central values and to a significant reduction of the theory uncertainty. Note that the shifts in $\alpha_s(M_Z)$ from Z and τ decays are opposite in sign and move the values in the *proper* direction, decreasing, thus, the current slight mismatch between two independent determinations of α_s .

The final results

$$\alpha_s(M_Z)|_Z = 0.1190 \pm 0.0026,$$

$$\alpha_s(M_Z)|_\tau = 0.1202 \pm 0.0019$$

from these two observables, although based on measurements of vastly different energy scales, are in remarkable agreement. This constitutes a striking test of asymptotic freedom in QCD. The two values can be combined to

$$\alpha_s(M_Z) = 0.1198 \pm 0.0015.$$

This is one of the most precise and presently only result in order α_s^4 .

We thank G. Quast for useful discussions and M. Steinhauser for reading the manuscript and good advice. This work was supported by the Deutsche Forschungsgemeinschaft in the Sonderforschungsbereich/Transregio No. SFB/TR-9 “Computational Particle Physics,” by INTAS (Grant No. 03-51-4007) and by RFBR (Grants No. 05-02-17645, No. 08-02-01451). The computer calculations were partially performed on the HP No. XC4000 super computer of the federal state Baden-Württemberg at the High Performance Computing Center Stuttgart (HLRS) under the grant “ParFORM.”

*Permanent address: Institute for Nuclear Research, Russian Academy of Sciences, Moscow 117312, Russia.

- [1] K. G. Chetyrkin, J. H. Kühn, and A. Kwiatkowski, *Phys. Rep.* **277**, 189 (1996), and references therein.
- [2] M. Davier, A. Hocker, and Z. Zhang, *Rev. Mod. Phys.* **78**, 1043 (2006), and references therein.
- [3] K. G. Chetyrkin, J. H. Kühn, and M. Steinhauser, *Phys. Lett. B* **371**, 93 (1996).
- [4] K. G. Chetyrkin, J. H. Kühn, and M. Steinhauser, *Nucl. Phys.* **B482**, 213 (1996).
- [5] K. G. Chetyrkin and J. H. Kühn, *Phys. Lett. B* **248**, 359 (1990).
- [6] K. G. Chetyrkin and J. H. Kühn, *Nucl. Phys.* **432**, 337 (1994).
- [7] K. G. Chetyrkin, R. V. Harlander, and J. H. Kühn, *Nucl. Phys.* **B586**, 56 (2000); **B634**, 413(E)(2002).
- [8] P. A. Baikov, K. G. Chetyrkin, and J. H. Kühn, *Nucl. Phys. B, Proc. Suppl.* **135**, 243 (2004).
- [9] A. Czarnecki and J. H. Kühn, *Phys. Rev. Lett.* **77**, 3955 (1996).
- [10] R. Harlander, T. Seidensticker, and M. Steinhauser, *Phys. Lett. B* **426**, 125 (1998).
- [11] A. L. Kataev and V. V. Starshenko, *Mod. Phys. Lett. A* **10**, 235 (1995).
- [12] D. V. Shirkov, *Theor. Math. Phys.* **127**, 409 (2001).
- [13] P. A. Baikov, K. G. Chetyrkin, and J. H. Kühn, *Phys. Rev. Lett.* **96**, 012003 (2006).
- [14] K. G. Chetyrkin, A. L. Kataev, and F. V. Tkachov, *Phys. Lett. B* **85**, 277 (1979).
- [15] S. G. Gorishnii, A. L. Kataev, and S. A. Larin, *Phys. Lett. B* **259**, 144 (1991).
- [16] L. R. Surguladze and M. A. Samuel, *Phys. Rev. Lett.* **66**, 560 (1991); **66**, 2416(E)(1991).
- [17] P. A. Baikov, K. G. Chetyrkin, and J. H. Kühn, *Phys. Rev. Lett.* **88**, 012001 (2001).
- [18] P. Nogueira, *J. Comput. Phys.* **105**, 279 (1993).
- [19] A. A. Vladimirov, *Theor. Math. Phys.* **43**, 417 (1980).
- [20] K. G. Chetyrkin and V. A. Smirnov, *Phys. Lett. B* **144**, 419 (1984).
- [21] K. G. Chetyrkin, *Phys. Lett. B* **391**, 402 (1997).
- [22] P. A. Baikov, *Phys. Lett. B* **634**, 325 (2006).
- [23] P. A. Baikov, *Phys. Lett. B* **385**, 404 (1996).
- [24] M. Tentyukov *et al.*, arXiv:cs.sc/0407066.
- [25] J. A. M. Vermaseren, arXiv:math-ph/0010025.
- [26] G. Grunberg, *Phys. Rev. D* **29**, 2315 (1984).
- [27] P. M. Stevenson, *Phys. Rev. D* **23**, 2916 (1981).
- [28] P. A. Baikov, K. G. Chetyrkin, and J. H. Kühn, *Phys. Rev. D* **67**, 074026 (2003).
- [29] M. M. Winter, ECFA-DESY Linear Collider Note Report No. LC-PHSM-2001-016, 2001.
- [30] J. H. Kühn, M. Steinhauser, and T. Teubner, *Phys. Rev. D* **76**, 074003 (2007).
- [31] J. Alcaraz *et al.* (LEP Collaboration, ALEPH Collaboration, DELPHI Collaboration, L3 Collaboration, OPAL Collaboration, and LEP Electroweak Working Group), arXiv:0712.0929.
- [32] S. Schael *et al.* (ALEPH), *Phys. Rep.* **421**, 191 (2005).
- [33] W. -M. Yao *et al.*, *J. Phys. G* **33**, 1 (2006), and 2007 partial update for the 2008 edition.
- [34] A. A. Pivovarov, *Z. Phys. C* **53**, 461 (1992).
- [35] F. Le Diberder and A. Pich, *Phys. Lett. B* **286**, 147 (1992).
- [36] M. Davier, A. Hocker, and Z. Zhang, *Nucl. Phys. B, Proc. Suppl.* **169**, 22 (2007).
- [37] S. Groote, J. G. Korner, and A. A. Pivovarov, *Mod. Phys. Lett. A* **13**, 637 (1998).
- [38] T. van Ritbergen, J. A. M. Vermaseren, and S. A. Larin, *Phys. Lett. B* **400**, 379 (1997).
- [39] M. Czakon, *Nucl. Phys.* **B710**, 485 (2005).
- [40] K. G. Chetyrkin, J. H. Kühn, and C. Sturm, *Nucl. Phys.* **B744**, 121 (2006).
- [41] Y. Schroder and M. Steinhauser, *J. High Energy Phys.* **01** (2006) 051.
- [42] J. H. Kühn, M. Steinhauser, and C. Sturm, *Nucl. Phys.* **B778**, 192 (2007).