

Observation of the Doubly Strange b Baryon Ω_b^-

- V. M. Abazov,³⁶ B. Abbott,⁷⁵ M. Abolins,⁶⁵ B. S. Acharya,²⁹ M. Adams,⁵¹ T. Adams,⁴⁹ E. Aguilera,⁶ M. Ahsan,⁵⁹ G. D. Alexeev,³⁶ G. Alkhazov,⁴⁰ A. Alton,^{64,*} G. Alverson,⁶³ G. A. Alves,² M. Anastasoaie,³⁵ L. S. Ancu,³⁵ T. Andeen,⁵³ B. Andrieu,¹⁷ M. S. Anzelc,⁵³ M. Aoki,⁵⁰ Y. Arnoud,¹⁴ M. Arov,⁶⁰ M. Arthaud,¹⁸ A. Askew,⁴⁹ B. Åsman,⁴¹ A. C. S. Assis Jesus,³ O. Atramentov,⁴⁹ C. Avila,⁸ F. Badaud,¹³ L. Bagby,⁵⁰ B. Baldin,⁵⁰ D. V. Bandurin,⁵⁹ P. Banerjee,²⁹ S. Banerjee,²⁹ E. Barberis,⁶³ A.-F. Barfuss,¹⁵ P. Bargassa,⁸⁰ P. Baringer,⁵⁸ J. Barreto,² J. F. Bartlett,⁵⁰ U. Bassler,¹⁸ D. Bauer,⁴³ S. Beale,⁶ A. Bean,⁵⁸ M. Begalli,³ M. Begel,⁷³ C. Belanger-Champagne,⁴¹ L. Bellantoni,⁵⁰ A. Bellavance,⁵⁰ J. A. Benitez,⁶⁵ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²³ I. Bertram,⁴² M. Besançon,¹⁸ R. Beuselinck,⁴³ V. A. Bezzubov,³⁹ P. C. Bhat,⁵⁰ V. Bhatnagar,²⁷ C. Biscarat,²⁰ G. Blazey,⁵² F. Blekman,⁴³ S. Blessing,⁴⁹ K. Bloom,⁶⁷ A. Boehlein,⁵⁰ D. Boline,⁶² T. A. Bolton,⁵⁹ E. E. Boos,³⁸ G. Borissov,⁴² T. Bose,⁷⁷ A. Brandt,⁷⁸ R. Brock,⁶⁵ G. Brooijmans,⁷⁰ A. Bross,⁵⁰ D. Brown,⁸¹ X. B. Bu,⁷ N. J. Buchanan,⁴⁹ D. Buchholz,⁵³ M. Buehler,⁸¹ V. Buescher,²² V. Bunichev,³⁸ S. Burdin,^{42,†} T. H. Burnett,⁸² C. P. Buszello,⁴³ J. M. Butler,⁶² P. Calfayan,²⁵ S. Calvet,¹⁶ J. Cammin,⁷¹ E. Carrera,⁴⁹ W. Carvalho,³ B. C. K. Casey,⁵⁰ H. Castilla-Valdez,³³ S. Chakrabarti,¹⁸ D. Chakraborty,⁵² K. M. Chan,⁵⁵ A. Chandra,⁴⁸ E. Cheu,⁴⁵ F. Chevallier,¹⁴ D. K. Cho,⁶² S. Choi,³² B. Choudhary,²⁸ L. Christofek,⁷⁷ T. Christoudias,⁴³ S. Cihangir,⁵⁰ D. Claes,⁶⁷ J. Clutter,⁵⁸ M. Cooke,⁵⁰ W. E. Cooper,⁵⁰ M. Corcoran,⁸⁰ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ S. Crépé-Renaudin,¹⁴ V. Cuplov,⁵⁹ D. Cutts,⁷⁷ M. Ćwiołek,³⁰ H. da Motta,² A. Das,⁴⁵ G. Davies,⁴³ K. De,⁷⁸ S. J. de Jong,³⁵ E. De La Cruz-Burelo,³³ C. De Oliveira Martins,³ K. DeVaughan,⁶⁷ J. D. Degenhardt,⁶⁴ F. Déliot,¹⁸ M. Demarteau,⁵⁰ R. Demina,⁷¹ D. Denisov,⁵⁰ S. P. Denisov,³⁹ S. Desai,⁵⁰ H. T. Diehl,⁵⁰ M. Diesburg,⁵⁰ A. Dominguez,⁶⁷ H. Dong,⁷² T. Dorland,⁸² A. Dubey,²⁸ L. V. Dudko,³⁸ L. Duflot,¹⁶ S. R. Dugad,²⁹ D. Duggan,⁴⁹ A. Duperrin,¹⁵ J. Dyer,⁶⁵ A. Dyshkant,⁵² M. Eads,⁶⁷ D. Edmunds,⁶⁵ J. Ellison,⁴⁸ V. D. Elvira,⁵⁰ Y. Enari,⁷⁷ S. Eno,⁶¹ P. Ermolov,^{38,††} H. Evans,⁵⁴ A. Evdokimov,⁷³ V. N. Evdokimov,³⁹ A. V. Ferapontov,⁵⁹ T. Ferbel,⁷¹ F. Fiedler,²⁴ F. Filthaut,³⁵ W. Fisher,⁵⁰ H. E. Fisk,⁵⁰ M. Fortner,⁵² H. Fox,⁴² S. Fu,⁵⁰ S. Fuess,⁵⁰ T. Gadfort,⁷⁰ C. F. Galea,³⁵ C. Garcia,⁷¹ A. Garcia-Bellido,⁷¹ V. Gavrilov,³⁷ P. Gay,¹³ W. Geist,¹⁹ W. Geng,^{15,65} C. E. Gerber,⁵¹ Y. Gershtein,⁴⁹ D. Gillberg,⁶ G. Ginther,⁷¹ N. Gollub,⁴¹ B. Gómez,⁸ A. Goussiou,⁸² P. D. Grannis,⁷² H. Greenlee,⁵⁰ Z. D. Greenwood,⁶⁰ E. M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,²⁵ S. Grünendahl,⁵⁰ M. W. Grünewald,³⁰ F. Guo,⁷² J. Guo,⁷² G. Gutierrez,⁵⁰ P. Gutierrez,⁷⁵ A. Haas,⁷⁰ N. J. Hadley,⁶¹ P. Haefner,²⁵ S. Hagopian,⁴⁹ J. Haley,⁶⁸ I. Hall,⁶⁵ R. E. Hall,⁴⁷ L. Han,⁷ K. Harder,⁴⁴ A. Harel,⁷¹ J. M. Hauptman,⁵⁷ J. Hays,⁴³ T. Hebbeker,²¹ D. Hedin,⁵² J. G. Hegeman,³⁴ A. P. Heinson,⁴⁸ U. Heintz,⁶² C. Hensel,^{22,‡} K. Herner,⁷² G. Hesketh,⁶³ M. D. Hildreth,⁵⁵ R. Hirosky,⁸¹ J. D. Hobbs,⁷² B. Hoeneisen,¹² H. Hoeth,²⁶ M. Hohlfeld,²² S. Hossain,⁷⁵ P. Houben,³⁴ Y. Hu,⁷² Z. Hubacek,¹⁰ V. Hynek,⁹ I. Iashvili,⁶⁹ R. Illingworth,⁵⁰ A. S. Ito,⁵⁰ S. Jabeen,⁶² M. Jaffré,¹⁶ S. Jain,⁷⁵ K. Jakobs,²³ C. Jarvis,⁶¹ R. Jesik,⁴³ K. Johns,⁴⁵ C. Johnson,⁷⁰ M. Johnson,⁵⁰ D. Johnston,⁶⁷ A. Jonckheere,⁵⁰ P. Jonsson,⁴³ A. Juste,⁵⁰ E. Kajfasz,¹⁵ J. M. Kalk,⁶⁰ D. Karmanov,³⁸ P. A. Kasper,⁵⁰ I. Katsanos,⁷⁰ D. Kau,⁴⁹ V. Kaushik,⁷⁸ R. Kehoe,⁷⁹ S. Kermiche,¹⁵ N. Khalatyan,⁵⁰ A. Khanov,⁷⁶ A. Kharchilava,⁶⁹ Y. M. Kharzeev,³⁶ D. Khatidze,⁷⁰ T. J. Kim,³¹ M. H. Kirby,⁵³ M. Kirsch,²¹ B. Klima,⁵⁰ J. M. Kohli,²⁷ E. V. Komissarov,^{36,††} J.-P. Konrath,²³ A. V. Kozelov,³⁹ J. Kraus,⁶⁵ T. Kuhl,²⁴ A. Kumar,⁶⁹ A. Kupco,¹¹ T. Kurča,²⁰ V. A. Kuzmin,³⁸ J. Kvita,⁹ F. Lacroix,¹³ D. Lam,⁵⁵ S. Lammers,⁷⁰ G. Landsberg,⁷⁷ P. Lebrun,²⁰ W. M. Lee,⁵⁰ A. Leflat,³⁸ J. Lellouch,¹⁷ J. Li,^{78,††} L. Li,⁴⁸ Q. Z. Li,⁵⁰ S. M. Lietti,⁵ J. K. Lim,³¹ J. G. R. Lima,⁵² D. Lincoln,⁵⁰ J. Linnemann,⁶⁵ V. V. Lipaev,³⁹ R. Lipton,⁵⁰ Y. Liu,⁷ Z. Liu,⁶ A. Lobodenko,⁴⁰ M. Lokajicek,¹¹ P. Love,⁴² H. J. Lubatti,⁸² R. Luna,³ A. L. Lyon,⁵⁰ A. K. A. Maciel,² D. Mackin,⁸⁰ R. J. Madaras,⁴⁶ P. Mättig,²⁶ C. Magass,²¹ A. Magerkurth,⁶⁴ P. K. Mal,⁸² H. B. Malbouisson,³ S. Malik,⁶⁷ V. L. Malyshev,³⁶ Y. Maravin,⁵⁹ B. Martin,¹⁴ R. McCarthy,⁷² A. Melnitchouk,⁶⁶ L. Mendoza,⁸ P. G. Mercadante,⁵ Y. P. Merekov,³⁶ M. Merkin,³⁸ K. W. Merritt,⁵⁰ A. Meyer,²¹ J. Meyer,^{22,‡} J. Mitrevski,⁷⁰ R. K. Mommsen,⁴⁴ N. K. Mondal,²⁹ R. W. Moore,⁶ T. Moulik,⁵⁸ G. S. Muanza,²⁰ M. Mulhern,⁷⁰ O. Mundal,²² L. Mundim,³ E. Nagy,¹⁵ M. Naimuddin,⁵⁰ M. Narain,⁷⁷ N. A. Naumann,³⁵ H. A. Neal,⁶⁴ J. P. Negret,⁸ P. Neustroev,⁴⁰ H. Nilsen,²³ H. Nogima,³ S. F. Novaes,⁵ T. Nunnemann,²⁵ V. O'Dell,⁵⁰ D. C. O'Neil,⁶ G. Obrant,⁴⁰ C. Ochando,¹⁶ D. Onoprienko,⁵⁹ J. Orduna,³³ N. Oshima,⁵⁰ N. Osman,⁴³ J. Osta,⁵⁵ R. Otec,¹⁰ G. J. Otero y Garzón,⁵⁰ M. Owen,⁴⁴ P. Padley,⁸⁰ M. Pangilinan,⁷⁷ N. Parashar,⁵⁶ S.-J. Park,^{22,‡} S. K. Park,³¹ J. Parsons,⁷⁰ R. Partridge,⁷⁷ N. Parua,⁵⁴ A. Patwa,⁷³ G. Pawloski,⁸⁰ B. Penning,²³ M. Perfilov,³⁸ K. Peters,⁴⁴ Y. Peters,²⁶ P. Pétroff,¹⁶ M. Petteni,⁴³ R. Piegaia,¹ J. Piper,⁶⁵ M.-A. Pleier,²² P. L. M. Podesta-Lerma,^{33,§} V. M. Podstavkov,⁵⁰ Y. Pogorelov,⁵⁵ M.-E. Pol,² P. Polozov,³⁷ B. G. Pope,⁶⁵ A. V. Popov,³⁹ C. Potter,⁶ W. L. Prado da Silva,³ H. B. Prosper,⁴⁹ S. Protopopescu,⁷³ J. Qian,⁶⁴ A. Quadt,^{22,‡} B. Quinn,⁶⁶ A. Rakitine,⁴² M. S. Rangel,² K. Ranjan,²⁸ P. N. Ratoff,⁴² P. Renkel,⁷⁹ P. Rich,⁴⁴ J. Rieger,⁵⁴ M. Rijssenbeek,⁷² I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷⁶ S. Robinson,⁴³ R. F. Rodrigues,³

M. Rominsky,⁷⁵ C. Royon,¹⁸ A. Rozhdestvenski,³⁶ P. Rubinov,⁵⁰ R. Ruchti,⁵⁵ G. Safronov,³⁷ G. Sajot,¹⁴
A. Sánchez-Hernández,³³ M. P. Sanders,¹⁷ B. Sanghi,⁵⁰ G. Savage,⁵⁰ L. Sawyer,⁶⁰ T. Scanlon,⁴³ D. Schaile,²⁵
R. D. Schamberger,⁷² Y. Scheglov,⁴⁰ H. Schellman,⁵³ T. Schliephake,²⁶ S. Schlobohm,⁸² C. Schwanenberger,⁴⁴
A. Schwartzman,⁶⁸ R. Schwienhorst,⁶⁵ J. Sekaric,⁴⁹ H. Severini,⁷⁵ E. Shabalina,⁵¹ M. Shamim,⁵⁹ V. Shary,¹⁸
A. A. Shchukin,³⁹ R. K. Shrivpuri,²⁸ V. Siccardi,¹⁹ V. Simak,¹⁰ V. Sirotenko,⁵⁰ P. Skubic,⁷⁵ P. Slattery,⁷¹ D. Smirnov,⁵⁵
G. R. Snow,⁶⁷ J. Snow,⁷⁴ S. Snyder,⁷³ S. Söldner-Rembold,⁴⁴ L. Sonnenschein,¹⁷ A. Sopczak,⁴² M. Sosebee,⁷⁸
K. Soustruznik,⁹ B. Spurlock,⁷⁸ J. Stark,¹⁴ J. Steele,⁶⁰ V. Stolin,³⁷ D. A. Stoyanova,³⁹ J. Strandberg,⁶⁴ S. Strandberg,⁴¹
M. A. Strang,⁶⁹ E. Strauss,⁷² M. Strauss,⁷⁵ R. Ströhmer,²⁵ D. Strom,⁵³ L. Stutte,⁵⁰ S. Sumowidagdo,⁴⁹ P. Svoisky,⁵⁵
A. Sznajder,³ P. Tamburello,⁴⁵ A. Tanasijczuk,¹ W. Taylor,⁶ B. Tiller,²⁵ F. Tissandier,¹³ M. Titov,¹⁸ V. V. Tokmenin,³⁶
I. Torchiani,²³ D. Tsybychev,⁷² B. Tuchming,¹⁸ C. Tully,⁶⁸ P. M. Tuts,⁷⁰ R. Unalan,⁶⁵ L. Uvarov,⁴⁰ S. Uvarov,⁴⁰
S. Uzunyan,⁵² B. Vachon,⁶ P. J. van den Berg,³⁴ R. Van Kooten,⁵⁴ W. M. van Leeuwen,³⁴ N. Varelas,⁵¹ E. W. Varnes,⁴⁵
I. A. Vasilyev,³⁹ P. Verdier,²⁰ L. S. Vertogradov,³⁶ Y. Vertogradova,³⁶ M. Verzocchi,⁵⁰ D. Vilanova,¹⁸
F. Villeneuve-Seguier,⁴³ P. Vint,⁴³ P. Vokac,¹⁰ M. Voutilainen,⁶⁷^{II} R. Wagner,⁶⁸ H. D. Wahl,⁴⁹ M. H. L. S. Wang,⁵⁰
J. Warchol,⁵⁵ G. Watts,⁸² M. Wayne,⁵⁵ G. Weber,²⁴ M. Weber,⁵⁰[¶] L. Welty-Rieger,⁵⁴ A. Wenger,^{23,**} N. Wermes,²²
M. Wetstein,⁶¹ A. White,⁷⁸ D. Wicke,²⁶ M. Williams,⁴² G. W. Wilson,⁵⁸ S. J. Wimpenny,⁴⁸ M. Wobisch,⁶⁰ D. R. Wood,⁶³
T. R. Wyatt,⁴⁴ Y. Xie,⁷⁷ S. Yacoob,⁵³ R. Yamada,⁵⁰ W.-C. Yang,⁴⁴ T. Yasuda,⁵⁰ Y. A. Yatsunenko,³⁶ H. Yin,⁷ K. Yip,⁷³
H. D. Yoo,⁷⁷ S. W. Youn,⁵³ J. Yu,⁷⁸ C. Zeitnitz,²⁶ S. Zelitch,⁸¹ T. Zhao,⁸² B. Zhou,⁶⁴ J. Zhu,⁷² M. Zielinski,⁷¹
D. Ziemska,⁵⁴ A. Ziemska,⁵⁴^{††} L. Zivkovic,⁷⁰ V. Zutshi,⁵² and E. G. Zverev³⁸

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Universidade Federal do ABC, Santo André, Brazil⁵Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil⁶University of Alberta, Edmonton, Alberta, Canada,*Simon Fraser University, Burnaby, British Columbia, Canada,**York University, Toronto, Ontario, Canada,**and McGill University, Montreal, Quebec, Canada*⁷University of Science and Technology of China, Hefei, People's Republic of China⁸Universidad de los Andes, Bogotá, Colombia⁹Center for Particle Physics, Charles University, Prague, Czech Republic¹⁰Czech Technical University, Prague, Czech Republic¹¹Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic¹²Universidad San Francisco de Quito, Quito, Ecuador¹³LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France¹⁵CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France¹⁶LAL, Université Paris-Sud, IN2P3/CNRS, Orsay, France¹⁷LPNHE, IN2P3/CNRS, Universités Paris VI and VII, Paris, France¹⁸CEA, Irfu, SPP, Saclay, France¹⁹IPHC, Université Louis Pasteur, CNRS/IN2P3, Strasbourg, France²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France²¹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany²²Physikalisches Institut, Universität Bonn, Bonn, Germany²³Physikalisches Institut, Universität Freiburg, Freiburg, Germany²⁴Institut für Physik, Universität Mainz, Mainz, Germany²⁵Ludwig-Maximilians-Universität München, München, Germany²⁶Fachbereich Physik, University of Wuppertal, Wuppertal, Germany²⁷Panjab University, Chandigarh, India²⁸Delhi University, Delhi, India²⁹Tata Institute of Fundamental Research, Mumbai, India³⁰University College Dublin, Dublin, Ireland³¹Korea Detector Laboratory, Korea University, Seoul, Korea³²SungKyunKwan University, Suwon, Korea³³CINVESTAV, Mexico City, Mexico

³⁴*FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands*³⁵*Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands*³⁶*Joint Institute for Nuclear Research, Dubna, Russia*³⁷*Institute for Theoretical and Experimental Physics, Moscow, Russia*³⁸*Moscow State University, Moscow, Russia*³⁹*Institute for High Energy Physics, Protvino, Russia*⁴⁰*Petersburg Nuclear Physics Institute, St. Petersburg, Russia*⁴¹*Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden*⁴²*Lancaster University, Lancaster, United Kingdom*⁴³*Imperial College, London, United Kingdom*⁴⁴*University of Manchester, Manchester, United Kingdom*⁴⁵*University of Arizona, Tucson, Arizona 85721, USA*⁴⁶*Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA*⁴⁷*California State University, Fresno, California 93740, USA*⁴⁸*University of California, Riverside, California 92521, USA*⁴⁹*Florida State University, Tallahassee, Florida 32306, USA*⁵⁰*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*⁵¹*University of Illinois at Chicago, Chicago, Illinois 60607, USA*⁵²*Northern Illinois University, DeKalb, Illinois 60115, USA*⁵³*Northwestern University, Evanston, Illinois 60208, USA*⁵⁴*Indiana University, Bloomington, Indiana 47405, USA*⁵⁵*University of Notre Dame, Notre Dame, Indiana 46556, USA*⁵⁶*Purdue University Calumet, Hammond, Indiana 46323, USA*⁵⁷*Iowa State University, Ames, Iowa 50011, USA*⁵⁸*University of Kansas, Lawrence, Kansas 66045, USA*⁵⁹*Kansas State University, Manhattan, Kansas 66506, USA*⁶⁰*Louisiana Tech University, Ruston, Louisiana 71272, USA*⁶¹*University of Maryland, College Park, Maryland 20742, USA*⁶²*Boston University, Boston, Massachusetts 02215, USA*⁶³*Northeastern University, Boston, Massachusetts 02115, USA*⁶⁴*University of Michigan, Ann Arbor, Michigan 48109, USA*⁶⁵*Michigan State University, East Lansing, Michigan 48824, USA*⁶⁶*University of Mississippi, University, Mississippi 38677, USA*⁶⁷*University of Nebraska, Lincoln, Nebraska 68588, USA*⁶⁸*Princeton University, Princeton, New Jersey 08544, USA*⁶⁹*State University of New York, Buffalo, New York 14260, USA*⁷⁰*Columbia University, New York, New York 10027, USA*⁷¹*University of Rochester, Rochester, New York 14627, USA*⁷²*State University of New York, Stony Brook, New York 11794, USA*⁷³*Brookhaven National Laboratory, Upton, New York 11973, USA*⁷⁴*Langston University, Langston, Oklahoma 73050, USA*⁷⁵*University of Oklahoma, Norman, Oklahoma 73019, USA*⁷⁶*Oklahoma State University, Stillwater, Oklahoma 74078, USA*⁷⁷*Brown University, Providence, Rhode Island 02912, USA*⁷⁸*University of Texas, Arlington, Texas 76019, USA*⁷⁹*Southern Methodist University, Dallas, Texas 75275, USA*⁸⁰*Rice University, Houston, Texas 77005, USA*⁸¹*University of Virginia, Charlottesville, Virginia 22901, USA*⁸²*University of Washington, Seattle, Washington 98195, USA*

(Received 29 August 2008; published 5 December 2008)

We report the observation of the doubly strange b baryon Ω_b^- in the decay channel $\Omega_b^- \rightarrow J/\psi \Omega^-$, with $J/\psi \rightarrow \mu^+ \mu^-$ and $\Omega^- \rightarrow \Lambda K^- \rightarrow (p\pi^-)K^-$, in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. Using approximately 1.3 fb^{-1} of data collected with the D0 detector at the Fermilab Tevatron Collider, we observe $17.8 \pm 4.9(\text{stat}) \pm 0.8(\text{syst})$ Ω_b^- signal events at a mass of $6.165 \pm 0.010(\text{stat}) \pm 0.013(\text{syst})$ GeV. The significance of the observed signal is 5.4σ , corresponding to a probability of 6.7×10^{-8} of it arising from a background fluctuation.

The Ω^- baryon, composed of three strange quarks, has played an important historical role in our understanding of the basic structure of matter. Its discovery in 1964 [1] at a mass predicted from SU(3) symmetry breaking was a great success for the theory [2]. The Ω_b^- (bss) (charge conjugate states are assumed throughout this Letter) is a predicted heavy cousin of the Ω^- with a b quark replacing one of the three strange quarks. While the Ω^- has $J^P = 3/2^+$, the ground state Ω_b^- is expected to have $J^P = 1/2^+$, a mass between 5.94–6.12 GeV and a lifetime such that $0.55 < \tau(\Omega_b^-)/\tau(B^0) < 1.10$ [3].

In this Letter, we report the first observation of the Ω_b^- baryon, fully reconstructed from its decay $\Omega_b^- \rightarrow J/\psi \Omega^-$, with $J/\psi \rightarrow \mu^+ \mu^-$, $\Omega^- \rightarrow \Lambda K^-$, and $\Lambda \rightarrow p \pi^-$. The analysis is based on a data sample of 1.3 fb^{-1} collected in $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ with the D0 detector [4] at the Fermilab Tevatron Collider. The detector components most relevant to this analysis are the central tracking system and the muon spectrometer. The central tracking system consists of a silicon microstrip tracker (SMT) and a central fiber tracker (CFT) inside a 2 Tesla superconducting solenoid. The SMT is optimized for tracking and vertexing over the pseudorapidity region $|\eta| < 3$ while the CFT has coverage for $|\eta| < 2$. A liquid argon and uranium calorimeter provides coverage up to $|\eta| < 4.2$. The muon spectrometer covers $|\eta| < 2$.

The $\Omega_b^- \rightarrow J/\psi \Omega^- \rightarrow J/\psi \Lambda K^- \rightarrow J/\psi p \pi^- K^-$ decay topology is similar to that of the $\Xi_b^- \rightarrow J/\psi \Xi^- \rightarrow J/\psi \Lambda \pi^- \rightarrow J/\psi p \pi^- \pi^-$ decay with Ω^- in place of Ξ^- and K^- in place of π^- . Consequently, the reconstruction of the J/ψ and Λ and their selection discussed below follow closely the analysis that led to the first direct observation of the Ξ_b^- baryon [5]. However, in this analysis we use a multivariate selection for the Ω^- owing to the smaller signal to background ratio compared to that for the Ξ^- in the Ξ_b^- analysis. We use the PYTHIA Monte Carlo (MC) program [6] to generate Ω_b^- and the EVTGEN program [7] to simulate Ω_b^- decays. The Ω_b^- mass and lifetime are set to be 6.052 GeV and 1.54 ps, respectively. The generated events are subjected to a GEANT [8] based D0 detector simulation, and to the same reconstruction and selection programs as the data. We optimize the Ω^- selection using MC Ω_b^- events for the signal and a sample of $J/\psi(\Lambda K^+)$ data (referred to below as wrong-sign events) for the background, while keeping the $J/\psi \Omega^-$ data blinded. Once all selection criteria have been determined, we apply them to the $J/\psi \Omega^-$ data.

We begin the event selection by reconstructing $J/\psi \rightarrow \mu^+ \mu^-$ candidates from two oppositely charged muons with transverse momentum (p_T) greater than 1.5 GeV that are compatible with being from a common vertex. Muons are identified by tracks reconstructed in the central tracking system that are matched with either track segments in the muon spectrometer or calorimeter energy deposits consistent with a minimum ionizing particle.

Events must have a well-reconstructed $p\bar{p}$ interaction point that we take to be the Ω_b^- production vertex and a $J/\psi \rightarrow \mu^+ \mu^-$ candidate in the mass window $2.75 < M_{\mu\mu} < 3.40 \text{ GeV}$. Events with J/ψ candidates are reprocessed with a version of the track reconstruction algorithm that increases the efficiency for tracks with low p_T and high impact parameters.

We form $\Lambda \rightarrow p \pi^-$ candidates from two oppositely charged particles, each with $p_T > 0.2 \text{ GeV}$, that are consistent with having originated from a common vertex. The two tracks are required to have a total of no more than two hits in the tracking system before the reconstructed $p \pi^-$ vertex. The impact parameter significance (the impact parameter with respect to the $p\bar{p}$ interaction point divided by its uncertainty) must exceed four for at least one of the tracks and three for the other. The track with the higher p_T is assumed to be the proton. MC studies show that this assignment leads to the correct combination nearly 100% of the time. Furthermore, we require the Λ transverse decay length to be greater than 4 times its uncertainty and the proper decay length to exceed 10 times its uncertainty, where the transverse decay length is the distance between the production and decay vertices in the transverse plane while the proper decay length is the transverse decay length corrected by the Lorentz boost calculated from $p_T(\Lambda)$. Λ candidates must have reconstructed masses between 1.108 and 1.126 GeV.

We combine Λ candidates with negatively charged particles (assumed to be kaons) to form $\Omega^- \rightarrow \Lambda K^-$ decay candidates. The Λ and the kaon are required to have a common vertex. The Ω^- candidates must have transverse decay length significances greater than four and the uncertainties of the proper decay lengths less than 0.5 cm. These two requirements reduce backgrounds from combinatorics and mismeasured tracks. The Ξ^- baryon has a mass of 1.322 GeV [9] and decays into $\Lambda \pi^-$. If the kaon mass is assigned to the pion, this decay could be a major background for $\Omega^- \rightarrow \Lambda K^-$. To eliminate this background, we remove candidates with $\Lambda \pi^-$ mass less than 1.34 GeV. Figure 1(a) shows the mass distribution of the reconstructed $\Omega^- \rightarrow \Lambda K^-$ candidates after these selections. The distribution of wrong-sign ΛK^+ events is also shown. An excess of events above the background around the expected Ω^- mass of 1.672 GeV [9] is visible in the distribution of the right-sign ΛK^- events.

To further enhance the Ω^- signal over the combinatorial background, kinematic variables associated with daughter particle momenta, vertices, and track qualities are combined using boosted decision trees (BDT) [10,11]. The ΛK^- mass distribution after the BDT selection is shown in Fig. 1(b). The BDT selection retains 87% of the Ω^- signal while rejecting 89% of the background. The enhanced Ω^- mass peak is evident in the distribution. A ΛK^- pair is considered to be a Ω^- candidate if its mass is in the range of 1.662–1.682 GeV.




FIG. 1 (color online). The invariant mass distribution of the ΛK pair before (a) and after (b) the BDT selection. Solid circles are from the right-sign ΛK^- events while the histogram is from the wrong-sign ΛK^+ events without any additional normalization.

To select $\Omega_b^- \rightarrow J/\psi \Omega^-$ candidates, we develop selection criteria using the MC Ω_b^- events as the signal and the data wrong-sign events as the background. The background events are formed by combining J/ψ candidates with ΛK^+ pairs with mass between 1.662 and 1.682 GeV. We form $\Omega_b^- \rightarrow J/\psi \Omega^-$ decay candidates from J/ψ and Ω^- pairs that are consistent with being from a common vertex. We require the uncertainty of the Ω_b^- proper decay length to be less than 0.03 cm and impose a minimum p_T cut of 6 GeV on the Ω_b^- candidates. Finally, J/ψ and Ω^- daughters from the Ω_b^- decays are expected to be boosted in the direction of the Ω_b^- ; therefore, we require the opening angle in the transverse plane between the J/ψ and the Ω^- to be less than $\pi/2$.

We then apply the above selections to the right-sign events in the data to search for the Ω_b^- baryon in the mass window between 5.6 and 7.0 GeV. This range is chosen since 5.624 GeV is the mass of the lightest b baryon, the Λ_b , and the upper limit of 7.0 GeV is nearly 1 GeV higher than the predicted Ω_b^- mass [11]. We calculate the Ω_b^- candidate mass using the formula $M(\Omega_b^-) = M(J/\psi \Omega^-) - M(\mu^+ \mu^-) - M(\Lambda K^-) + \hat{M}(J/\psi) + \hat{M}(\Omega^-)$. Here $M(J/\psi \Omega^-)$, $M(\mu^+ \mu^-)$, and $M(\Lambda K^-)$ are the reconstructed masses while $\hat{M}(J/\psi)$ and $\hat{M}(\Omega^-)$ are taken from Ref. [9]. This calculation improves the mass resolution of the MC Ω_b^- events from 0.080 GeV to 0.034 GeV. In the mass search window, we observe 79 candidates in the data with the mass distribution shown in Fig. 2(a). An excess of events near 6.2 GeV is apparent. No such structure, however, is seen in the corresponding mass distribution [Fig. 2(b)] of the 30 wrong-sign events.

Assuming the excess is due to the Ω_b^- production, we fit Ω_b^- candidate masses with the hypothesis of a Gaussian signal plus a flat background using an unbinned likelihood method. We fix the Gaussian width to 0.034 GeV, the width of the MC Ω_b^- signal. The fit gives an Ω_b^- mass of $6.165 \pm$




FIG. 2 (color online). (a) The $M(\Omega_b^-)$ distribution of the Ω_b^- candidates after all selection criteria. The dotted curve is an unbinned likelihood fit to the model of a constant background plus a Gaussian signal. The mass distributions for the wrong-sign background (b), the Ω^- sideband events (c), and the Λ sideband events (d).

0.010(stat) GeV and a yield of 17.8 ± 4.9 (stat) signal events. To assess the significance of the excess, we first determine the likelihood \mathcal{L}_{s+b} of the signal plus background fit above and then repeat the fit with only the background contribution to find a new likelihood \mathcal{L}_b . The logarithmic likelihood ratio $\sqrt{2 \ln(\mathcal{L}_{s+b}/\mathcal{L}_b)}$ yields a statistical significance of 5.4σ , equivalent to a probability of 6.7×10^{-8} that the background could fluctuate with a significance equal to or greater than what is observed. Fitted yields for positively and negatively charged candidates are 6.2 ± 3.1 (stat) Ω_b^+ and 12.0 ± 3.9 (stat) Ω_b^- , respectively.

Various checks have been performed to ensure that the observed resonance is genuine. (1) We apply the event selection to data events in the sidebands of the reconstructed Ω^- and Λ resonances separately. The $J/\psi(p\pi^-)K^-$ mass distributions of these sideband events are shown in Figs. 2(c) and 2(d). No significant excess is present in either distribution. (2) We investigate the possibility of a false signal due to residual b hadron backgrounds by applying the final Ω_b^- selection to MC $B^- \rightarrow J/\psi K^{*-} \rightarrow J/\psi K_S^0 \pi^-$, $\Xi_b^- \rightarrow J/\psi \Xi^-$, and $\Lambda_b \rightarrow J/\psi \Lambda$ samples with equivalent integrated luminosities significantly greater than that of the analyzed data. No indication of any resonance is observed in the reconstructed $J/\psi \Omega^-$ mass distribution. (3) We check the mass distributions of the Ω_b^- decay products. For Ω_b^- candidates within a $\pm 3\sigma$ mass window around the observed peak, we relax the mass requirements on the Ω^- and Λ candidates and perform a fit to each mass distribution. The numbers of the Ω^- and Λ candidates from the fits are consistent with the observed number of Ω_b^- signal events. (4) We replace the BDT selection with individual cuts on the most important variables according to the BDT optimization and confirm the existence of a peak with a comparable event yield but a

higher background at a mass consistent with that observed using the BDT. (5) We test the robustness of the peak by varying selections such as the Ξ^- veto, Λ and Ω^- mass windows, Λ transverse decay requirements, BDT selection, and the requirement on $p_T(\Omega_b^-)$. All the above studies confirm the existence of the resonance.

Potential sources of systematic uncertainties on the measured Ω_b^- mass include event selection, signal and background models, and momentum scale. Varying the selection criteria and applying a set of cuts on individual kinematic variables lead to a maximum change of 0.012 GeV in the fitted mass. Using a linear function as the background model results in negligible change in the mass. Varying the Gaussian width in the signal model between 0.028 and 0.040 GeV changes the fitted mass by at most 0.003 GeV. When a tighter selection is applied to enhance signal over background, we can float the width of the signal model in the fit. This leads to a mass shift of 0.002 GeV and a fitted signal width of 0.033 ± 0.010 GeV, consistent with the MC expectation. To study the effect of the track momentum scale uncertainty on the measured Ω_b^- mass, we reconstruct the higher statistics $\Lambda_b \rightarrow J/\psi \Lambda$ decays and measure the Λ_b mass for different minimum p_T requirements on the Λ_b daughter particles. We compare these measurements to the world average value of the Λ_b mass [9] and take the maximum deviation of 0.004 GeV as a systematic uncertainty. Adding in quadrature, we get a total systematic uncertainty of 0.013 GeV to obtain a measured Ω_b^- mass: $6.165 \pm 0.010(\text{stat}) \pm 0.013(\text{syst})$ GeV. Similarly, we estimate the systematic uncertainty on the Ω_b^- yield by varying the signal and background models in the fit. The observed maximum change of 0.8 is assigned as the systematic uncertainty on the yield: $17.8 \pm 4.9(\text{stat}) \pm 0.8(\text{syst})$. In all these studies, the signal significance remains above 5σ .

Figure 3 shows the distribution of the proper decay length of the Ω_b^- candidates observed in the $\pm 3\sigma$ mass

window around the fitted Ω_b^- mass. The distribution of the MC Ω_b^- signal plus the data background events is also shown. The background distribution is modeled using events in the Ω_b^- sidebands of 5.8–6.0 and 6.4–6.6 GeV. Despite the low statistics, the data distribution contains significantly more positive than negative decay lengths as expected and consistent with a weakly decaying b hadron.

We calculate the Ω_b^- production rate relative to that of the Ξ_b^- [5]. The selection efficiency ratio $\epsilon(\Omega_b^- \rightarrow J/\psi \Omega^-)/\epsilon(\Xi_b^- \rightarrow J/\psi \Xi^-)$ is found to be $1.5 \pm 0.2(\text{stat})$ assuming inclusive Ω^- and Ξ^- decays. The higher efficiency for the Ω_b^- is due primarily to a harder p_T spectrum of the kaon from the Ω^- decay than that of the pion from the Ξ^- decay and a shorter lifetime of the Ω^- compared to the Ξ^- . By using the reported Ξ_b^- events [5] and the observed Ω_b^- yield here, we estimate

$$\mathcal{R} = \frac{f(b \rightarrow \Omega_b^-)}{f(b \rightarrow \Xi_b^-)} \frac{\text{Br}(\Omega_b^- \rightarrow J/\psi \Omega^-)}{\text{Br}(\Xi_b^- \rightarrow J/\psi \Xi^-)}$$

to be $\mathcal{R} = 0.80 \pm 0.32(\text{stat})^{+0.14}_{-0.22}(\text{syst})$. Here $f(b \rightarrow \Omega_b^-)$ and $f(b \rightarrow \Xi_b^-)$ are the fractions of b quarks that hadronize to form Ω_b^- and Ξ_b^- , respectively. The systematic uncertainty includes contributions from the signal yields as well as the efficiency ratio. Using $\Gamma(\Omega_b^- \rightarrow J/\psi \Omega^-)/\Gamma(\Xi_b^- \rightarrow J/\psi \Xi^-) = 9.8$ [12], the central values of $\tau(\Xi_b^-) = 1.42^{+0.28}_{-0.24}$ ps [9], the \mathcal{R} value above, and $\tau(\Omega_b^-)$ in the range of 0.83–1.67 ps [3], we obtain $f(b \rightarrow \Omega_b^-)/f(b \rightarrow \Xi_b^-) \approx 0.07\text{--}0.14$.

In summary, by analyzing 1.3 fb^{-1} of data collected by the D0 experiment in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV at the Fermilab Tevatron Collider, we have made the first observation of the doubly strange b baryon Ω_b^- in the fully reconstructed decay mode $\Omega_b^- \rightarrow J/\psi \Omega^-$ with $J/\psi \rightarrow \mu^+ \mu^-$, $\Omega^- \rightarrow \Lambda K^-$ and $\Lambda \rightarrow p \pi^-$. We measure the Ω_b^- mass to be $6.165 \pm 0.010(\text{stat}) \pm 0.013(\text{syst})$ GeV. The significance of the observed signal is greater than 5σ .

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (U.S.); CEA and CNRS/IN2P3 (France); FASI, Rosatom, and RFBR (Russia); CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC (United Kingdom); MSMT and GACR (Czech Republic); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); and the Alexander von Humboldt Foundation (Germany).




FIG. 3 (color online). The distribution of the proper decay length of the Ω_b^- candidates in the $\pm 3\sigma$ mass window around the observed peak along with the expected distribution from the MC Ω_b^- signal with a lifetime 1.54 ps plus the data background events.

*Visitor from Augustana College, Sioux Falls, SD, USA.

[†]Visitor from The University of Liverpool, Liverpool, U.K.

[‡]Visitor from II. Physikalisches Institut, Georg-August-University, Göttingen, Germany.

^{\$}Visitor from ECFM, Universidad Autonoma de Sinaloa, Culiacán, Mexico.

[¶]Visitor from Helsinki Institute of Physics, Helsinki, Finland.

[¶]Visitor from Universität Bern, Bern, Switzerland.

^{**}Visitor from Universität Zürich, Zürich, Switzerland.

^{††}Deceased.

- [1] V.E. Barnes *et al.*, Phys. Rev. Lett. **12**, 204 (1964).
- [2] M. Gell-Mann and Y. Ne'eman, *The Eightfold Way* (W.A. Benjamin, New York, 1964).
- [3] X. Liu *et al.*, Phys. Rev. D **77**, 014031 (2008); M. Karliner *et al.*, arXiv:0804.1575; E. E. Jenkins, Phys. Rev. D **77**, 034012 (2008); R. Roncaglia, D. B. Lichtenberg, and E. Predazzi, Phys. Rev. D **52**, 1722 (1995); N. Mathur, R. Lewis, and R. M. Woloshyn, Phys. Rev. D **66**, 014502 (2002); D. Ebert, R. N. Faustov, and V. O. Galkin, Phys.

Rev. D **72**, 034026 (2005); T. Ito, M. Matsuda, and Y. Matsui, Prog. Theor. Phys. **99**, 271 (1998).

[4] V. M. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **565**, 463 (2006).

[5] V. M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **99**, 052001 (2007).

[6] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).

[7] D. J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001).

[8] R. Brun and F. Carminati, CERN Program Library Writeup Report No. W5013, 1993 (unpublished).

[9] C. Amsler *et al.*, Phys. Lett. B **667**, 1 (2008).

[10] L. Breiman *et al.*, *Classification and Regression Trees* (Wadsworth, Stamford, 1984).

[11] A. Höcker *et al.*, arXiv:physics/0703039.

[12] H.-Y. Cheng, Phys. Rev. D **56**, 2799 (1997).