

Generalized Kapchinskij-Vladimirskij Distribution and Envelope Equation for High-Intensity Beams in a Coupled Transverse Focusing Lattice

Hong Qin,¹ Moses Chung,² and Ronald C. Davidson¹

¹*Plasma Physics Laboratory, Princeton University, Princeton, New Jersey 08543, USA*

²*Accelerator Physics Center, Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*

(Received 26 August 2009; published 25 November 2009)

In an uncoupled lattice, the Kapchinskij-Vladimirskij (KV) distribution function first analyzed in 1959 is the only known exact solution of the nonlinear Vlasov-Maxwell equations for high-intensity beams including self-fields in a self-consistent manner. The KV solution is generalized here to high-intensity beams in a coupled transverse lattice using the recently developed generalized Courant-Snyder invariant for coupled transverse dynamics. This solution projects to a rotating, pulsating elliptical beam in transverse configuration space, determined by the generalized matrix envelope equation.

DOI: 10.1103/PhysRevLett.103.224802

PACS numbers: 29.27.Bd

Modern high-intensity beams have many important applications ranging from high energy density physics and ion-beam-driven fusion to high-flux neutron sources and light sources. It is becoming increasingly important to understand the self-field effects of high-intensity beams including self-electric and self-magnetic fields in a fully self-consistent manner, from the nonlinear Vlasov-Maxwell equations [1]. In an uncoupled lattice, the Kapchinskij-Vladimirskij (KV) distribution function analyzed in 1959 [2] is the only known exact self-consistent solution of the nonlinear Vlasov-Maxwell equations for high-intensity beams. In practical accelerators and beam transport systems, the transverse coupling between the horizontal and vertical directions, induced by error fields and misalignments, is always a significant effect [3–8]. Strong coupling of the transverse dynamics is introduced intentionally in certain types of cooling channels [9] and in the final focusing system for high energy density physics experiments [10], as well as in the conceptual design of the Möbius accelerator [11]. In this Letter, we generalize the KV solution to describe high-intensity beam dynamics in a coupled transverse focusing lattice using the recently developed generalized Courant-Snyder invariant [12,13] for coupled transverse dynamics.

In a coupled transverse focusing lattice, the Vlasov-Maxwell equations that govern the evolution of the distribution function f of a high-intensity beam and the corresponding space-charge potential ψ are [1]

$$\frac{\partial f}{\partial s} + \mathbf{v} \cdot \frac{\partial f}{\partial \mathbf{x}} - (\nabla \psi + \kappa_q \mathbf{x}) \cdot \frac{\partial f}{\partial \mathbf{v}} = 0, \quad (1)$$

$$\nabla^2 \psi = \frac{-2\pi K_b}{N_b} \int f dv_x dv_y. \quad (2)$$

Here, particle motion in the beam frame is assumed to be nonrelativistic, ψ is the space-charge potential normalized by $\gamma_b^3 m \beta_b^2 c^2 / q_b$, $\beta_b c$ is the directed beam velocity in the longitudinal direction, $\gamma_b = (1 - \beta_b^2)^{-1/2}$ is the relativistic

mass factor, s is the time variable normalized by $1/\beta_b c$, $K_b = 2N_b q_b^2 / \gamma_b^3 m \beta_b^2 c^2$ is the beam self-field perveance, $N_b = \int f dx dy dv_x dv_y$ is the line density, $\mathbf{x} = (x, y)^T$ represents the normalized transverse displacement of a beam particle, $\mathbf{v} = d\mathbf{x}/ds = (v_x, v_y)^T = (\dot{x}, \dot{y})^T$ is the normalized transverse velocity in the beam frame, and $\kappa_q \mathbf{x}$ is the coupled linear focusing force. In Eq. (1)

$$\kappa_q = \begin{pmatrix} \kappa_{qx} & \kappa_{qxy} \\ \kappa_{qyx} & \kappa_{qy} \end{pmatrix} \quad (3)$$

is the matrix of coupling coefficients, κ_{qx} and κ_{qy} are the focusing coefficients for the quadrupole lattice, and $\kappa_{qxy} = \kappa_{qyx}$ are the coupling coefficients produced by the skew-quadrupole component of the lattice. In general, the coupled linear focusing force can also depend on transverse velocity, as in the case of a solenoidal lattice, which can be transformed into the form of Eqs. (1)–(3) if we choose the local Lamor frame [1,13]. For simplicity of presentation, we consider here only the coupling due to skew quadrupoles given by Eq. (3). The $-\nabla \psi$ term in Eq. (1) describes the self-field force, and is nonlinearly coupled to f through Eq. (2). Equations (1) and (2) form a set of nonlinear integro-differential equations, whose analytical solutions are difficult to find in general.

For the case of an uncoupled lattice, i.e., $\kappa_{qxy} = \kappa_{qyx} = 0$, Eqs. (1) and (2) admit a remarkable solution known as the Kapchinskij-Vladimirskij (KV) distribution [2], which has played an important role in high-intensity beam physics [14–17]. The KV distribution function is constructed as a function of the Courant-Snyder (CS) invariants of the transverse dynamics [18]. Since the CS invariants are valid for linear, uncoupled transverse forces, the KV distribution must self-consistently generate a linear, uncoupled space-charge force. The KV distribution indeed satisfies this requirement. It is given by [1,2]

$$f_{KV} = \frac{N_b}{\pi^2 \epsilon_x \epsilon_y} \delta\left(\frac{I_x}{\epsilon_x} + \frac{I_y}{\epsilon_y} - 1\right), \quad (4)$$

$$I_x = \frac{x^2}{w_x^2} + (w_x \dot{x} - x \dot{w}_x)^2, \quad I_y = \frac{y^2}{w_y^2} + (w_y \dot{y} - y \dot{w}_y)^2. \quad (5)$$

Here, I_x and I_y are the CS invariants for the x - and y -motions, respectively, ε_x and ε_y are the constant transverse emittances, and w_x and w_y are the envelope functions satisfying the envelope equations,

$$\ddot{w}_x + \kappa_x w_x = w_x^{-3}, \quad \ddot{w}_y + \kappa_y w_y = w_y^{-3}, \quad (6)$$

$$\kappa_x = \kappa_{qx} - \frac{2K_b}{a(a+b)}, \quad \kappa_y = \kappa_{qy} - \frac{2K_b}{b(a+b)}, \quad (7)$$

$$a \equiv \sqrt{\varepsilon_x} w_x, \quad b \equiv \sqrt{\varepsilon_y} w_y. \quad (8)$$

The density profile in the transverse configuration space projected by the distribution function f_{KV} in Eq. (4) is given by

$$n(x, y, s) = \int dx dy f_{KV} = \begin{cases} N_b / \pi ab = \text{const}, & 0 \leq x^2/a^2 + y^2/b^2 < 1, \\ 0, & 1 < x^2/a^2 + y^2/b^2. \end{cases} \quad (9)$$

which corresponds to a constant-density beam with elliptical cross section and pulsating transverse dimensions a and b [see Fig. 1(a)]. The associated space-charge potential inside the beam, determined from Eq. (2), is given by

$$\psi = \frac{-K_b}{a+b} \left(\frac{x^2}{a} + \frac{y^2}{b} \right), \quad 0 \leq x^2/a^2 + y^2/b^2 < 1. \quad (10)$$

The KV distribution (4) reduces the original nonlinear Vlasov-Maxwell equations (1) and (2) to the two envelope equations in Eq. (6) for w_x and w_y , or equivalently, for $a = \sqrt{\varepsilon_x} w_x$ and $b = \sqrt{\varepsilon_y} w_y$ [Eq. (8)]. As the only known solution of the nonlinear Vlasov-Maxwell equations (1) and (2), the KV distribution and the associated envelope equations provide very important elementary theoretical




FIG. 1 (color online). Beam cross sections for the KV distribution. (a) Uncoupled lattice: the cross section is determined by $0 \leq x^2/a^2 + y^2/b^2 < 1$; and (b) coupled lattice: the cross section is determined by $0 \leq \mathbf{x}^T w^{-1} w^{-1T} \mathbf{x} < \varepsilon$.

tools for our understanding of high-intensity beam dynamics [14–17]. The KV distribution in Eq. (4) is constructed from the exact dynamical invariants I_x and I_y in Eq. (5), and constitutes an exact solution of the Vlasov equation (1), which also generates the uncoupled linear space-charge force assumed *a priori*.

We now show how to generalize this KV solution to the case of coupled transverse dynamics when $\kappa_{qxy} = \kappa_{qyx} \neq 0$, using the recently developed generalized CS invariant for coupled transverse lattice [12,13]. In the coupled case, the generalized KV distribution that solves the nonlinear Vlasov-Maxwell system (1) and (2) projects to a rotating, pulsating beam with elliptical cross section in transverse configuration space with constant density inside the beam. Both the dimensions a and b , and the tilt angle θ are functions of $s = \beta_b ct$ [see Fig. 1(b)], in contrast with the pulsating upright elliptical beam cross section for the uncoupled case [see Fig. 1(a)]. The rotating, pulsating beam with elliptical cross section in transverse configuration space, and constant density inside the beam, generates a coupled linear space-charge force of the form

$$-\nabla\psi = -\kappa_s \mathbf{x}, \quad \kappa_s = \begin{pmatrix} \kappa_{sx} & \kappa_{sxy} \\ \kappa_{syx} & \kappa_{sy} \end{pmatrix}, \quad (11)$$

where $\kappa_{sxy} = \kappa_{syx}$, which allows us to apply the generalized CS invariant for the coupled transverse dynamics. The exact form of κ_s will be determined self-consistently [see Eq. (24)]. Our strategy is to use the generalized CS invariant to construct a generalized KV solution of the Vlasov equation (1), which also projects to a rotating, pulsating elliptical beam with constant density inside the beam. In this manner, a self-consistent solution of the nonlinear Vlasov-Maxwell equations (1) and (2) is found for high-intensity beams in a coupled transverse focusing lattice.

For a charged particle subject to the coupled linear focusing force and the coupled linear space-charge force

$$-\nabla\psi - \kappa_q \mathbf{x} = -\kappa \mathbf{x}, \quad \kappa = \kappa_q + \kappa_s, \quad (12)$$

the generalized CS invariant is given by [12,13]

$$I_{CS} = \mathbf{x}^T w^{-1} w^{-1T} \mathbf{x} + (\dot{\mathbf{x}}^T w^T - \mathbf{x}^T \dot{w}^T)(w \dot{\mathbf{x}} - \dot{w} \mathbf{x}), \quad (13)$$

where

$$w = \begin{pmatrix} w_1 & w_2 \\ w_3 & w_4 \end{pmatrix}$$

is the 2×2 envelope matrix determined from the matrix envelope equation

$$\dot{w} + w \kappa = (w^{-1})^T w^{-1} (w^{-1})^T. \quad (14)$$

Since I_{CS} is an invariant of the particle dynamics, any function of I_{CS} is a solution of the Vlasov equation (1). However, in order to solve the nonlinear Vlasov-Maxwell equations (1) and (2), the distribution function must generate the coupled linear space-charge force of the form in Eq. (11) as well. For this purpose, we select the distribution

function to be the following generalized KV distribution

$$f_{KV} = \frac{N_b |w|}{A \varepsilon \pi} \delta\left(\frac{I_{CS}}{\varepsilon} - 1\right). \quad (15)$$

Here, N_b and ε are constants, where N_b is the line-density, and ε is the transverse emittance. Moreover, $|w|$ is the determinant of the envelope matrix w , and A is the area of the beam cross section determined by $|w|$ and ε . Both $|w|$ and A are functions of $s = \beta_b c t$. The beam density profile in transverse configuration space is

$$\begin{aligned} n(x, y, s) &= \int d\dot{x}d\dot{y} f_{KV} = \int d\left(\frac{r^2}{\varepsilon}\right) \frac{N_b}{A} \delta\left(\frac{I_{CS}}{\varepsilon} - 1\right) \\ &= \begin{cases} N_b/A, & 0 \leq \mathbf{x}^T w^{-1} w^{-1T} \mathbf{x} < \varepsilon, \\ 0, & \varepsilon < \mathbf{x}^T w^{-1} w^{-1T} \mathbf{x}. \end{cases} \end{aligned} \quad (16)$$

In the above calculation, the velocity integration with respect to $d\dot{x}d\dot{y}$ is carried out in the new velocity coordinates (p, q) through the transformation

$$d\dot{x}d\dot{y} = \frac{1}{|w|} dpdq = \frac{2\pi}{|w|} r dr, \quad (17)$$

$$p \equiv w_1 \dot{x} + w_2 \dot{y} - \dot{w}_1 x - \dot{w}_2 y, \quad (18)$$

$$q \equiv w_3 \dot{x} + w_4 \dot{y} - \dot{w}_3 x - \dot{w}_4 y, \quad (19)$$

$$r^2 \equiv p^2 + q^2. \quad (20)$$

The density profile $n(x, y, s)$ obtained in Eq. (16) is indeed of the desired form. That is, $n(x, y, s)$ is constant inside the ellipse defined by

$$\mathbf{x}^T \beta^* \mathbf{x} = \varepsilon, \quad \beta^* \equiv w^{-1} w^{-1T}, \quad (21)$$

and $n(x, y, s) = 0$ outside the ellipse. The ellipse defined by Eq. (21) is pulsating and rotating. Its transverse dimensions $a(s)$ and $b(s)$, and tilt angle $\theta(s)$ depend on $s = \beta_b c t$ and are determined from the matrix β^* . Because β^* is obviously real, symmetric, and positive definite, the two eigenvectors v_1 and v_2 of β^* are orthogonal with two positive eigenvalues λ_1 and λ_2 . It is an elementary result [19] that the transverse dimensions of the ellipse are given by $a = \sqrt{\varepsilon/\lambda_1}$ and $b = \sqrt{\varepsilon/\lambda_2}$, and the tilt angle θ is that of v_1 . The principal axis theorem [19] states that the diagonalizing matrix Q of β^* can be constructed as $Q = (v_1, v_2)$ with $Q^{-1} = Q^T$ and

$$Q^{-1} \beta^* Q = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}.$$

We now introduce the rotating frame

$$\begin{pmatrix} X \\ Y \end{pmatrix} = Q^{-1} \begin{pmatrix} x \\ y \end{pmatrix}.$$

The ellipse in (X, Y) coordinates is given

$$\frac{X^2}{a^2} + \frac{Y^2}{b^2} = 1, \quad (22)$$

and the self-field force is

$$-\begin{pmatrix} \partial\psi/\partial X \\ \partial\psi/\partial Y \end{pmatrix} = \frac{2K_b}{a+b} \begin{pmatrix} 1/a & 0 \\ 0 & 1/b \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix}. \quad (23)$$

Transforming back to (x, y) coordinate, the self-field force can be expressed as

$$\begin{aligned} -\begin{pmatrix} \partial\psi/\partial x \\ \partial\psi/\partial y \end{pmatrix} &= -\kappa_s \begin{pmatrix} x \\ y \end{pmatrix}, \\ \kappa_s &= \frac{-2K_b}{a+b} Q \begin{pmatrix} 1/a & 0 \\ 0 & 1/b \end{pmatrix} Q^{-1}. \end{aligned} \quad (24)$$

The coupled linear space-charge coefficient κ_s is a function of the envelope matrix w and the constant emittance ε . When Eq. (24) is substituted back into Eq. (12), the envelope equation (14) becomes a closed nonlinear matrix equation for the envelope matrix w . Therefore, we have succeeded in finding a class of self-consistent solutions of the nonlinear Vlasov-Maxwell equations for high-intensity beams in a coupled transverse focusing lattice. The solution reduces to a nonlinear matrix ordinary differential equation for the envelope matrix w , which determines the geometry of the pulsating and rotating beam ellipse. The matrix envelope equation (14) can be numerically solved in a straightforward manner. We note that the self-consistent solution constructed for the coupled lattice has one emittance ε in the transverse directions [11], whereas in an uncoupled lattice, the standard KV distribution contains two emittances, i.e., ε_x and ε_y . This should not come as a surprise because a coupled lattice is more complex than an uncoupled lattice, and it is natural for the self-consistent solution to have less freedom in a coupled lattice than in an uncoupled lattice. In accelerators and storage rings with coupling, a single emittance in the transverse directions implies an equilibrium between the x direction and the y direction, which can be reached in certain situations, but not always. Therefore, the self-consistent distribution constructed only applies to those cases where such an equilibrium is reached, such as in a strongly coupled system or in the lattice of transport lines.

As a specific example, we consider a periodic quadrupole FODO lattice with the middle magnet being misaligned by a small angle ξ . The misaligned magnet induces a skew-quadrupole component of the form [4] $\kappa_{qxy} = \kappa_{qyx} = \kappa_q \sin 2\xi$. The strength of the quadrupole component of the misaligned magnet is reduced to $\kappa_{qx} = -\kappa_{qy} = \kappa_q \cos 2\xi$. The normalized quadrupole focusing field is $\kappa_q \equiv q_b B'_q / \gamma_b m \beta_b c^2 = 15$ with a filling factor $\eta = 0.15$. The misalignment is $\xi = 11.4^\circ$, and the normalized self-field perveance is $K_b/\varepsilon = 0.1$. The matrix envelope equation (14) has been solved numerically to find a matched solution. The numerical result, plotted in




FIG. 2 (color online). Beam cross sections as a function of $s/S = \beta_b ct/S$ over the interval $0 \leq s/S \leq 1$. The dynamics of the beam pulsation and rotation is evident from the figure.

Fig. 2, shows the beam cross section as a function of $s/S = \beta_b ct/S$, where S is the lattice period. The dynamics of the beam pulsation and rotation is clearly demonstrated in the plots. The rotation dynamics result in a wobbling motion of the tilt angle between $\theta = 14.28^\circ$ at $s/S = 0$ and $\theta = 81.35^\circ$ at $s/S = 0.5$. As expected, in the rotating frame the transverse dimensions a and b of the beam ellipse oscillate with time. Note that the dynamics of beam rotation and pulsation is matched with the lattice period.

In conclusion, the KV distribution function, the exact self-consistent solution of the nonlinear Vlasov-Maxwell equations for high-intensity charged particle beams in an uncoupled focusing lattice including self-electric and self-magnetic fields, has been generalized to describe high-intensity beam dynamics in a coupled transverse focusing lattice using the recently developed generalized Courant-Snyder invariant [12,13] for coupled transverse dynamics. The fully self-consistent solution reduces the nonlinear Vlasov-Maxwell equations to a nonlinear matrix ordinary differential equation for the envelope matrix w , which determines the geometry of the pulsating and rotating

beam ellipse. This result provides us with a new theoretical tool to investigate the dynamics of high-intensity beams in a coupled transverse lattice.

This research was supported by the U. S. Department of Energy.

-
- [1] R. C. Davidson and H. Qin, *Physics of Intense Charged Particle Beams in High Energy Accelerators* (World Scientific, Singapore, 2001).
 - [2] I. Kapchinskij and V. Vladimirskij, in *Proceedings of the International Conference on High Energy Accelerators and Instrumentation* (CERN Scientific Information Service, Geneva, 1959).
 - [3] D. A. Edwards and M. J. Syphers, *An Introduction to the Physics of High-Energy Accelerators* (Wiley, New York, 1993).
 - [4] J. J. Barnard, in *Proceedings of the 1995 Particle Accelerators Conference* (IEEE, Piscataway, NJ, 1996), p. 3241.
 - [5] Y. Cai, *Phys. Rev. E* **68**, 036501 (2003).
 - [6] R. A. Kishek, J. J. Barnard, and D. P. Grote, in *Proceedings of the 1999 Particle Accelerator Conference* (IEEE, Piscataway, NJ, 1999), p. 1761.
 - [7] J. J. Barnard and R. Losic, in *Proceedings of the 20th International Linac Conference* (2001), p. MOE12.
 - [8] F. J. Sacherer, Ph.D. thesis, Univ. of California, Berkeley, 1968.
 - [9] M. Chung, H. Qin, and R. C. Davidson, *Phys. Rev. ST Accel. Beams* (to be published).
 - [10] B. G. Logan *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **577**, 1 (2007).
 - [11] R. Talman, *Phys. Rev. Lett.* **74**, 1590 (1995).
 - [12] H. Qin and R. C. Davidson, *Phys. Plasmas* **16**, 050705 (2009).
 - [13] H. Qin and R. C. Davidson, *Phys. Rev. ST Accel. Beams* **12**, 064001 (2009).
 - [14] T. S. Wang and L. Smith, *Part. Accel.* **12**, 247 (1982).
 - [15] I. Hofmann, L. J. Laslett, L. Smith, and I. Haber, *Part. Accel.* **13**, 145 (1983).
 - [16] J. Struckmeier and I. Hofmann, *Part. Accel.* **39**, 219 (1992).
 - [17] C. Chen, R. Parker, and R. C. Davidson, *Phys. Rev. Lett.* **79**, 225 (1997).
 - [18] E. Courant and H. Snyder, *Ann. Phys. (N.Y.)* **3**, 1 (1958).
 - [19] G. Strang, *Linear Algebra and Its Applications* (Harcourt Brace Jovanovich, San Diego, 1988), 3rd ed.