

Nonclassical Correlations from Randomly Chosen Local Measurements

Yeong-Cherng Liang,^{1,*} Nicholas Harrigan,² Stephen D. Bartlett,¹ and Terry Rudolph³

¹*School of Physics, The University of Sydney, Sydney, New South Wales 2006, Australia*

²*Department of Physics, Imperial College London, London SW7 2BW, United Kingdom*

³*Institute for Mathematical Sciences, Imperial College London, London SW7 2BW, United Kingdom*

(Received 15 September 2009; published 2 February 2010)

We show that correlations inconsistent with any locally causal description can be a generic feature of measurements on entangled quantum states. Specifically, spatially separated parties who perform local measurements on a maximally entangled state using randomly chosen measurement bases can, with significant probability, generate nonclassical correlations that violate a Bell inequality. For n parties using a Greenberger-Horne-Zeilinger state, this probability of violation rapidly tends to unity as the number of parties increases. We also show that, even with both a randomly chosen two-qubit pure state and randomly chosen measurement bases, a violation can be found about 10% of the time. Among other applications, our work provides a feasible alternative for the demonstration of Bell inequality violation without a shared reference frame.

DOI: 10.1103/PhysRevLett.104.050401

PACS numbers: 03.65.Ud, 03.65.Ta, 03.67.Mn

One of the most remarkable features of quantum mechanics is that distant measurements can exhibit correlations that are inconsistent with any locally causal description (LCD) [1]. These quantum correlations are signatures of entanglement [2], and serve as a resource [3] for a range of classically impossible information processing tasks such as quantum key distribution [4], teleportation [5], and reduced communication complexity [6].

Not all quantum states are useful for quantum information processing tasks. Even for entangled states, nonclassical correlations are not an immediate consequence of the entanglement present in such systems (see Ref. [7] and references therein). They also depend crucially on the choice of measurements to which they are subjected, and specifically how these measurements are correlated between parties. For example, in the standard scenarios to violate a Bell-Cluser-Horne-Shimony-Holt (Bell-CHSH) inequality [1,8] or more generally an n -party inequality using a Greenberger-Horne-Zeilinger (GHZ) state, a maximum violation is achieved by all parties using specific measurements in the common x - y plane [9]. In an experiment, this requires some care [10]; for two parties using polarization-encoded single-photon pairs transmitted through long birefringent optical fibers, the random rotation of the polarization of each photon is first compensated through some method of alignment in order to set this frame [11]. With larger numbers of parties, greater violations can be achieved but more parties require an increasing complexity of alignments.

Several approaches to circumvent this problem have been studied. Encoded entangled states that are invariant with respect to some collective unitary operations [12,13] can be used, but generally require much more complicated state preparation as well as joint measurements on multiple spins. Alternatively, an alignment of frames can be per-

formed through the coherent exchange of quantum systems [11,14] or by supplementing each system with a small quantum reference frame of bounded size [15]. Such solutions are resource intensive given that this alignment consumes many of these quantum resources that could otherwise be used to generate useful correlations.

Here, we prove that such resource-intensive approaches are unnecessary, and that nonclassical correlations occur with high probability even without any alignment of local measurements. Specifically, we investigate the detection of nonlocal correlations wherein each party uses randomly chosen measurement bases. Ruling out LCD in this scenario cannot be achieved deterministically for a general entangled quantum state [13]. However, we demonstrate that the probability of observing correlations inconsistent with LCD using randomly chosen measurement bases can be remarkably high. Although one might naively expect that increasing the number of parties participating in the test (and the corresponding complexity of alignments) would only complicate matters, we show that the probability of demonstrating a violation rapidly approaches unity as the number of parties increases. Finally, we demonstrate the possibility of two parties violating a Bell inequality using both randomly chosen measurements and a randomly chosen pure state.

We consider a typical scenario of an n -party Bell experiment using spin-1/2 particles, wherein a verifier prepares $N \gg 1$ copies of the n -partite GHZ state

$$|\text{GHZ}\rangle_n = \frac{1}{\sqrt{2}}(|0\rangle_1|0\rangle_2 \cdots |0\rangle_n + |1\rangle_1|1\rangle_2 \cdots |1\rangle_n), \quad (1)$$

and distributes them to n parties. Each particle is subjected to a local measurement with two possible outcomes, ± 1 . The measurement for each particle is chosen randomly, with equal probability, from a set of two measurement

bases, defined for the k th party by a pair of spatial directions $\Omega_{s_k}^{[k]}(\theta, \phi)$, with $s_k = 1, 2$ labeling the bases. For spins measured via a Stern-Gerlach device, these directions correspond to the orientation of the magnetic field in the measurement apparatus. However, an equivalent picture in terms of spatial directions can be used for any quantum systems described by a two-dimensional Hilbert space (e.g., the polarization encoding of single photons) via the Bloch sphere. After the completion of all measurements, the n parties return to the verifier a list containing only the label s_k of which measurement they performed for each particle and the value ± 1 of the outcome they obtained for that measurement.

A relevant Bell inequality for this scenario is the n -partite Mermin-Ardehali-Belinskiĭ-Klyshko (MABK) inequality [16–18],

$$\mathcal{S}_{\text{MABK}}^{(n)} = \left| \sum_{s_1, \dots, s_n=1}^2 \beta(s_1, \dots, s_n) E(s_1, \dots, s_n) \right| \leq 2^{n-(1/2)}, \quad (2)$$

where

$$\beta(s_1, \dots, s_n) = \sum_{k_1, \dots, k_n = \pm 1} \cos \left[\frac{\pi}{4} \left(n + 1 - \sum_{l=1}^n k_l \right) \right] \prod_{j=1}^n k_j^{s_j - 1}. \quad (3)$$

Here, the n -partite correlation function $E(s_1, \dots, s_n)$ is defined to be the expectation value of the product of measurement outcomes (± 1) given that the k th party measured their spin in the basis s_k . When $n = 2$, the MABK inequality is equivalent to the familiar Bell-CHSH inequality $\mathcal{S}_{\text{CHSH}} = |E(1, 1) + E(1, 2) + E(2, 1) - E(2, 2)| \leq 2$ [1,8].

In a standard Bell experiment, the maximal violations for a given Bell inequality are achieved by choosing the local orientations $\Omega_1^{[k]}$ and $\Omega_2^{[k]}$ to be perpendicular [9], i.e., $\Omega_1^{[k]} \cdot \Omega_2^{[k]} = 0$, and by fixing the relative orientations of these measurements with the other experimentalists (and the verifier who prepared the systems). The latter necessitates the use of a shared reference frame; in what follows, we will see what can be achieved without a shared reference frame.

We will first consider the completely unconstrained case, wherein each party k chooses both members of their pair of directions $\Omega_{s_k}^{[k]}$ for $s_k = 1, 2$ independently and uniformly from the set of all possible directions. We refer to this case as random isotropic measurements (RIM). In such a scenario, we determine the probability $p_{\text{MABK}}^{|\text{GHZ}_n\rangle}$ that the measurement statistics received by the verifier will give rise to a violation of the MABK inequality of Eq. (2), given by

$$p_{\text{MABK}}^{|\text{GHZ}_n\rangle} = \frac{1}{(4\pi)^{2n}} \int \prod_{\substack{k=1, \dots, n \\ s_k=1,2}} d\Omega_{s_k}^{[k]} f_n^{\text{MABK}}(\{\Omega_{s_k}^{[k]}\}_{k=1}^n), \quad (4)$$

where $d\Omega_{s_k}^{[k]}/4\pi$ is the Haar measure associated with mea-

surement direction $\Omega_{s_k}^{[k]}$ and $f_n^{\text{MABK}}(\{\Omega_{s_k}^{[k]}\}_{k=1}^n)$ is a function that returns 1 if orientations $\{\Omega_{s_k}^{[k]}\}_{k=1}^n$ give rise to a correlation (i.e., measurement statistics), that violates the MABK inequality and 0 otherwise. These integrals simply pick out those measurement directions that will give a violation, normalizing them against the total volume of possible measurement directions.

For the detection of nonclassical correlation, we can allow the verifier to make use of the freedom in relabeling all measurement settings and/or outcomes and/or parties [17,19]. To appreciate the importance of this fact, consider as an example the following choice of local measurement for the $n = 2$ case: $\{\sigma_z, \sigma_x\}$ for Alice and $\{\frac{1}{\sqrt{2}}(\sigma_x + \sigma_z), \frac{1}{\sqrt{2}}(\sigma_x - \sigma_z)\}$ for Bob. Suppose that Alice has labeled her measurements such that $\sigma_z \leftrightarrow s_1 = 1$ and $\sigma_x \leftrightarrow s_1 = 2$, while Bob has labeled his measurements such that $\frac{1}{\sqrt{2}}(\sigma_x + \sigma_z) \leftrightarrow s_2 = 1$ and $\frac{1}{\sqrt{2}}(\sigma_x - \sigma_z) \leftrightarrow s_2 = 2$, then the verifier will find that as it is, $\mathcal{S}_{\text{CHSH}}$ vanishes, and hence does not lead to a Bell-CHSH inequality violation. However, if the verifier is also given the freedom to relabel the measurement settings, they could use, instead, the following labeling for Alice's measurements: $\sigma_x \leftrightarrow s_1 = 1$ and $\sigma_z \leftrightarrow s_1 = 2$. As opposed to the original labeling, this new labeling unveils measurement statistics that violate the Bell-CHSH inequality maximally. We say that two Bell inequalities are equivalent if they can be obtained from each other by relabeling the measurement settings and/or outcomes and/or parties [17,19]. In calculating the probability of observing a violation, we clearly want to include all such equivalent inequalities.

We numerically calculate $p_{\text{MABK}}^{|\text{GHZ}_n\rangle}$, using the relabeling strategy to test a given set of experimental data against all 2^{n+1} equivalent MABK inequalities [20]; see Fig. 1. The special case of $n = 2$ can be solved analytically (see




FIG. 1 (color online). Probability of finding a nonclassical n -partite GHZ correlation. The two sets of data at the bottom and top are, respectively, for correlation generated from random isotropic measurements (RIM) and random orthogonal measurements (ROM). The markers are for the scenarios when the verifier has access to (1) the 2^{n+1} MABK inequalities (Δ for RIM, \times for ROM), and (2) the 2^{2^n} WWZB inequalities (∇ for RIM, $+$ for ROM).

Ref. [21] for details) and is found to be $2(\pi - 3) \approx 28.3\%$ (see also Ref. [22] for contrasting results). For $n > 2$, the probability is less than this amount but increases monotonically for $n > 3$ up to the limit of our analysis, $n = 15$. Note, however, it appears that this probability is asymptotically approaching a value that is less than unity.

The case of $n = 2$ appears somewhat anomalous. The reason is that, for $n = 2$, the equivalence class of Bell-CHSH inequalities obtained by relabeling $\mathcal{S}_{\text{MABK}}^{(2)}$ (or $\mathcal{S}_{\text{CHSH}}$) is both necessary and sufficient for determining if a given set of experimental statistics generated by performing two binary-outcome measurements per site admits a LCD [23]. In other words, in the space of measurement statistics generated by performing such measurements, all nonclassical correlations can be detected by this equivalence class of inequalities.

For $n > 2$, there are (i) Bell inequalities with n -partite correlation functions that are not of the MABK form (2) and (ii) Bell inequalities involving less-than- n -partite correlation functions. To determine if a given correlation for $n > 2$ is incompatible with any LCD, we need to test the measurement statistics against the complete set of Bell inequalities that are relevant to the particular experiment. For a small number of parties, this can be achieved efficiently using linear programming (see [21,24]) or by checking the correlation directly against the complete set of relevant Bell inequalities. Our results are summarized in Table I.

These results clearly indicate that the probability of a violation rapidly approaches unity with increasing n . The input to the linear program, however, scales exponentially with n [21], thus making it intractable to determine the probability of violation reliably for larger values of n . To obtain some insight into the behavior, we can restrict our attention to the extensive set of 2^{2^n} n -partite correlation inequalities discovered independently by Werner and Wolf [17] and Żukowski and Brukner [18] (WWZB). While violation of these inequalities is still sufficient to rule out LCD, the converse is generally not true. Thus, using this set of inequalities yields a lower bound on the probability that a given correlation is nonclassical.

To test if a given correlation satisfies all of the WWZB inequalities is equivalent to testing if it satisfies the follow-

ing nonlinear inequality [17,18]

$$\sum_{k_1, \dots, k_n = \pm 1} \left| \sum_{s_1, \dots, s_n = 1}^2 \prod_{j=1}^n k_j^{s_j-1} E(s_1, \dots, s_n) \right| \leq 2^n. \quad (5)$$

We have numerically computed the probability of finding a randomly generated correlation from RIM to violate inequality (5) for $n \leq 15$. As can be seen in Fig. 1, this probability generally increases with n and is above 50% for $n = 15$. However, it is inconclusive whether this probability asymptotes to unity or not for large n .

Finally, we consider another mechanism by which parties who do not share a reference frame can increase the probability of observing a violation. So far we have considered each of the two local measurement bases for each party to be chosen independently and isotropically. However, because the optimal violations are typically found when the pair of local measurement directions are orthogonal [9], we can consider a random selection of measurement bases under the constraint that each local pair are orthogonal. We have repeated the above calculations with this constraint (i.e., that $\Omega_1^{[k]} \cdot \Omega_2^{[k]} = 0$); we refer to this distribution as random orthogonal measurements (ROM). The results reveal a significant increase in the probability of a violation, as shown in Fig. 1 and Table I. For larger values of n , our results for the WWZB inequalities also strongly suggest that the probability asymptotes to unity for large n .

Our results so far have all made use of a fixed maximally entangled pure state (1) (although clearly the local bases in which this state is defined are irrelevant for our result). We can also consider how the probability of observing a violation of LCD varies with the degree of entanglement of the distributed quantum state. For $n = 2$, a pure quantum state can always be written as $|\Psi\rangle = \cos\theta|0\rangle_1|0\rangle_2 + \sin\theta|1\rangle_1|1\rangle_2$ for some local bases. Numerically, we can compute the probability of violation as a function of entanglement (for details, see [21]). With this result, we can determine the probability of observing a violation of LCD using both randomly chosen measurement bases and a randomly chosen state. We sample two-qubit pure states randomly and uniformly from $\mathbb{C}^2 \otimes \mathbb{C}^2$ according to the SU(4) Haar measure. Given that our results are independent of local choice of bases, we can reproduce this distribution by sampling Schmidt coefficients as for single-qubit mixed states from the uniform Bloch ball [25]. The relevant measure is the Hilbert-Schmidt measure on a single qubit, i.e., $P_{\text{HS}}(r) = 24r^2$, where $r = \frac{\cos 2\theta}{2}$ is the length of the Bloch vector (Ref. [25], p. 354). The probability of violation for a randomly chosen pure two-qubit state can then be determined by numerical integration to be, respectively, about 5.3% for RIM and 10.1% for ROM.

Conclusion and other directions.—We introduced the idea of performing a Bell experiment using randomly chosen measurement bases, with the aim of demonstrating nonclassical correlations in the absence of a shared reference frame. We applied this idea to the n -partite GHZ state

TABLE I. Probability of finding a nonclassical correlation from the n -partite GHZ state for the scenario where each party is allowed to perform binary projective measurements in two randomly chosen measurement bases. The number of data points (run) used to compute the probabilities for RIM (ROM) is included in the third (fifth) row of the table.

n	2	3	4	5	6
RIM	28.3185%	74.6899%	94.2380%	99.5926%	99.97%
$\frac{\text{run}}{10^5}$		5000	500	25	0.8
ROM	41.2982%	96.2073%	99.9757%	99.9999%	100.00%
$\frac{\text{run}}{10^5}$	5000	5000	500	25	0.8

and showed that the probability of finding a nonclassical correlation rapidly tends to unity as the number of parties increases. Naively, one might have expected that this chance would diminish with increasing n , due to the increasing chance of misalignment of the experimenters' measurement bases (or, alternatively, by considering the fragility of the GHZ state to dephasing). Our results clearly show otherwise. We have thus shown that without a shared reference frame, a Bell inequality violation can still be demonstrated reliably without resorting to complicated state preparation or consumption of expensive quantum resources. This, of course, significantly reduces the technical requirements for experimentally violating a Bell inequality, performing quantum key distribution based on such violations, or establishing large-scale quantum networks.

Our work also represents the first systematic study of the set of correlation derivable from a quantum resource such as the n -partite GHZ state. Specifically, our results indicate that the correlations obtained by performing projective measurements on this quantum resource are almost ubiquitously nonclassical (this can be rigorously quantified by considering a volume measure in the space of GHZ-attainable-correlations induced by RIM and ROM). Given that measurements and the resulting correlations play an important role in many quantum information processing tasks, the tools and ideas that we have introduced here, suitably generalized, may shed some light on the origin of the power of other quantum resources such as the highly entangled cluster state. In particular, they may add new insights to the computational power of correlations recently discussed in Ref. [26].

The results presented here motivate several additional research directions. First, one may attempt to determine a new intuition for our result of increasing probability of violation with the number of parties n , by relating this probability to the average magnitude of the MABK violation, which also increases with n [21]. Second, for many quantum information processing tasks, violation of the inequalities considered in the present work may not be sufficient, as other types of multipartite entanglement (perhaps different from that of the GHZ state) may be required that are not quantified by these inequalities. To this end, it will be interesting to investigate the efficiency of random measurements against inequalities that detect other, more general types of multipartite entanglement [27]. Finally, for real-world applications of these results, a critical issue is to determine how robust they are against decoherence.

Y.C.L. acknowledges fruitful discussion with Nicolas Brunner, Tamás Vértesi, Stefano Pironio, Mafalda Almeida, Valerio Scarani, Eric Cavalcanti, and Daniel Cavalcanti. S.D.B. acknowledges support from the ARC. T.R. and N.H. acknowledge the support of the EPSRC.

- [1] J.S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, Cambridge, 2004).
- [2] R. Horodecki *et al.*, *Rev. Mod. Phys.* **81**, 865 (2009).
- [3] J. Barrett *et al.*, *Phys. Rev. A* **71**, 022101 (2005); N. Brunner, N. Gisin, and V. Scarani, *New J. Phys.* **7**, 88 (2005).
- [4] A.K. Ekert, *Phys. Rev. Lett.* **67**, 661 (1991); J. Barrett, L. Hardy, and A. Kent, *ibid.* **95**, 010503 (2005); A. Acín, N. Gisin, and Ll. Masanes, *ibid.* **97**, 120405 (2006); Ll. Masanes, *ibid.* **102**, 140501 (2009).
- [5] C.H. Bennett *et al.*, *Phys. Rev. Lett.* **70**, 1895 (1993); R. Horodecki, M. Horodecki, and P. Horodecki, *Phys. Lett. A* **222**, 21 (1996).
- [6] Č. Brukner *et al.*, *Phys. Rev. Lett.* **92**, 127901 (2004).
- [7] R.F. Werner and M.M. Wolf, *Quantum Inf. Comput.* **1**, 1 (2001); Y.-C. Liang and A.C. Doherty, *Phys. Rev. A* **75**, 042103 (2007); Ll. Masanes, Y.-C. Liang, and A.C. Doherty, *Phys. Rev. Lett.* **100**, 090403 (2008).
- [8] J.F. Clauser *et al.*, *Phys. Rev. Lett.* **23**, 880 (1969).
- [9] V. Scarani and N. Gisin, *J. Phys. A* **34**, 6043 (2001).
- [10] A. Aspect, arXiv:quant-ph/0402001.
- [11] S.D. Bartlett, T. Rudolph, and R. W. Spekkens, *Rev. Mod. Phys.* **79**, 555 (2007).
- [12] S.D. Bartlett, T. Rudolph, and R. W. Spekkens, *Phys. Rev. Lett.* **91**, 027901 (2003).
- [13] A. Cabello, *Phys. Rev. A* **68**, 042104 (2003); A. Cabello, *Phys. Rev. Lett.* **91**, 230403 (2003).
- [14] T. Rudolph and L. Grover, *Phys. Rev. Lett.* **91**, 217905 (2003).
- [15] F. Costa *et al.*, *New J. Phys.* **11**, 123007 (2009).
- [16] N.D. Mermin, *Phys. Rev. Lett.* **65**, 1838 (1990); S.M. Roy and V. Singh, *ibid.* **67**, 2761 (1991); M. Ardehali, *Phys. Rev. A* **46**, 5375 (1992); A.V. Belinskiĭ and D.N. Klyshko, *Phys. Usp.* **36**, 653 (1993); N. Gisin and H. Bechmann-Pasquinucci, *Phys. Lett. A* **246**, 1 (1998).
- [17] R.F. Werner and M.M. Wolf, *Phys. Rev. A* **64**, 032112 (2001).
- [18] M. Żukowski and Č. Brukner, *Phys. Rev. Lett.* **88**, 210401 (2002).
- [19] Ll. Masanes, *Quantum Inf. Comput.* **3**, 345 (2003); D. Collins and N. Gisin, *J. Phys. A* **37**, 1775 (2004).
- [20] All distinct inequalities in the MABK class can be generated by applying the 2^n different permutations ($s_k \rightarrow s_k$ or $s_k \rightarrow 3 - s_k$ for all $k = 1, 2, \dots, n$) to the coefficients $\beta(s_1, \dots, s_n)$ of Eq. (3) defining the inequality.
- [21] See supplementary material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.104.050401> for our analytic derivation of the result for $n = 2$, our linear programming approach to the problem of identifying nonclassical correlations, and the plot for probability of violation vs bipartite pure state entanglement.
- [22] P. Lougovski and S. J. van Enk, *Phys. Rev. A* **80**, 034302 (2009).
- [23] A. Fine, *Phys. Rev. Lett.* **48**, 291 (1982).
- [24] D. Gosal *et al.*, *Phys. Rev. A* **70**, 042106 (2004).
- [25] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States: An Introduction to Quantum Entanglement* (Cambridge University Press, Cambridge, 2006).
- [26] J. Anders and D.E. Browne, *Phys. Rev. Lett.* **102**, 050502 (2009).
- [27] D. Collins *et al.*, *Phys. Rev. Lett.* **88**, 170405 (2002); M. Seevinck and G. Svetlichny, *ibid.* **89**, 060401 (2002).

*ycliang@physics.usyd.edu.au