

Time-Dependent Mean Field Theory for Quench Dynamics in Correlated Electron Systems

Marco Schiró¹ and Michele Fabrizio^{1,2}

¹International School for Advanced Studies (SISSA), and CRS Democritos, CNR-INFM, via Bonomea 265, 34136 Trieste Italy

²The Abdus Salam International Centre for Theoretical Physics (ICTP), Post Office Box 586, I-34014 Trieste, Italy

(Received 2 May 2010; published 10 August 2010)

A simple and very flexible variational approach to the out-of-equilibrium quantum dynamics in strongly correlated electron systems is introduced through a time-dependent Gutzwiller wave function. As an application, we study the simple case of a sudden change of the interaction in the fermionic Hubbard model and find at the mean-field level an extremely rich behavior. In particular, a dynamical transition between small and large quantum quench regimes is found to occur at half-filling, in accordance with the analysis of Eckstein *et al.*, *Phys. Rev. Lett.* **103**, 056403 (2009), obtained by dynamical mean-field theory, that turns into a crossover at any finite doping.

DOI: 10.1103/PhysRevLett.105.076401

PACS numbers: 71.10.Fd, 05.30.Fk, 05.70.Ln

Introduction.—Triggered by the enormous advances in the physics of ultracold atomic gases [1], time-dependent nonequilibrium phenomena in strongly interacting quantum systems have recently become of greatest interest. The possibility of artificially engineering many-particle quantum states with tunable interactions and almost perfect isolation from the environment gives the chance of probing directly in the time domain the quantum dynamics following an external perturbation [2]. While early experiments focused mainly on bosonic systems [3] or fermionic condensates [4], the recent experimental realization of a fermionic Mott insulator [5] has opened the way to investigate out-of-equilibrium phenomena in electron systems too [6]. From a theoretical perspective, these experiments raise several intriguing questions touching quantum dynamics at its roots. Indeed, when driven out of equilibrium, interacting quantum systems can display peculiar dynamical behaviors or even be trapped into metastable configurations [7] that differ completely from their equilibrium counterpart. The simplest way one could imagine to induce a nontrivial dynamics is through a so-called *quantum quench*. Here the system is first prepared in the ground state of some given Hamiltonian $\mathcal{H}_i$, and then suddenly let evolved under the action of a new Hamiltonian $\mathcal{H}_f$. Recently, quantum quenches have been the subject of vast literature focusing on integrable systems [8,9], one-dimensional models [10–12], or systems close to a quantum criticality [13]. The interest in these classes of non-equilibrium problems relies both on the dynamics itself [14], as well as on the long-time properties where the issue of thermalization, or its lack of, is still highly debated [15,16]. For what concerns strongly correlated electrons in more than one dimension, the subject is still largely unexplored and progress has been made only recently. The single band fermionic Hubbard model is likely the simplest lattice model of correlated electrons embodying the competition between metallic and insulating behavior driven by a local Hubbard repulsion U . Its Hamiltonian reads

$$\mathcal{H}(t) = -\sum_{\sigma} \sum_{\langle i,j \rangle} t_{ij} c_{i\sigma}^{\dagger} c_{j\sigma} + U(t) \sum_i n_{i\uparrow} n_{i\downarrow}. \quad (1)$$

In two pioneering works [17,18], the response of a Fermi sea to a sudden switch-on of the Hubbard U has been studied in infinite dimensions using, respectively, the flow-equation method and the dynamical mean-field theory (DMFT). Results suggest the existence of two different regimes in the real-time dynamics depending on the final interaction strength U_f . At weak coupling [17], the systems are trapped at long times into a quasistationary regime where correlations are more effective than in equilibrium. This *prethermalization* phenomenon has been confirmed by DMFT results [18], which further indicate a true dynamical transition above a critical U_{fc} towards another regime with pronounced oscillations in the dynamics of physical quantities. These intriguing results have been so far restricted to a quench starting from a noninteracting system ($U_i = 0$) and, more importantly, limited to rather short accessible time scales and weak quenches, thus leaving open many important issues.

A simple and flexible approach, although less rigorous than, e.g., DMFT, is thus extremely desirable and this is actually the aim of the present work. Specifically, here we propose a variational approach to the out-of-equilibrium dynamics of a correlated electron system based on a proper extension of the Gutzwiller wave function. We apply this technique to study the interaction quench in the Hubbard model, where we find a rich behavior featuring a transition in the real-time dynamics at a critical quench line $U_{fc}(U_i)$, in accordance with Ref. [18] at $U_i = 0$. Remarkably, a finite doping completely washes out this transition, leaving behind only a crossover from weak to strong coupling.

Variational approach to quantum dynamics.—Many basic concepts in the theory of strongly correlated systems, like, e.g., the Brinkman-Rice scenario for the Mott transition [19], have been originated from calculations based on a very simple and physically transparent variational approach introduced in the 1960's by Gutzwiller. This

approach has been so far applied only at equilibrium or at most in the linear response regime [20], but it turns out to be so flexible to allow for full out-of-equilibrium calculations.

For simplicity, we assume initially a many-body wave function $|\Psi_0\rangle$, which, for times $t > 0$, is let evolve with a Hamiltonian $\mathcal{H}$ that includes sizable on-site interactions. In the spirit of the Gutzwiller approach, we make the following variational ansatz for the time-dependent wave function $|\Psi_{\text{exact}}(t)\rangle = e^{-i\mathcal{H}t}|\Psi_0\rangle$

$$|\Psi_{\text{exact}}(t)\rangle \simeq |\Psi(t)\rangle = \prod_i e^{-iS_i(t)} \mathcal{P}_i(t) |\Phi(t)\rangle, \quad (2)$$

where $|\Phi(t)\rangle$ is a time-dependent Slater determinant. $\mathcal{P}_i(t)$ is a Hermitian operator that acts on the Hilbert space of site i and controls the weights of the local electronic configurations. $S_i(t)$ is also Hermitian and we assume it depends on some variables $\phi_{i\alpha}(t)$ such that

$$\frac{\partial}{\partial \phi_{i\alpha}} e^{-iS_i} = -i\mathcal{O}_{i\alpha} e^{-iS_i},$$

where $\mathcal{O}_{i\alpha}$ is any local Hermitian operator. Since (2) is just a variational ansatz, it does not solve the full Schrödinger equation. Our proposal is to determine the variational parameters by requiring: (i) that the Heisenberg equations of motion of the local operators $\mathcal{O}_{i\alpha}$ are satisfied when averaging over (2); (ii) that the average energy $E = \langle \Psi(t) | \mathcal{H} | \Psi(t) \rangle$ is, as it should be, conserved during the evolution. Since, by definition,

$$\frac{\partial}{\partial \phi_{i\alpha}} e^{iS_i} \mathcal{H} e^{-iS_i} = i e^{iS_i} [\mathcal{O}_{i\alpha}, \mathcal{H}] e^{-iS_i},$$

it follows that

$$\begin{aligned} \frac{\partial \mathcal{O}_{i\alpha}}{\partial t} &= -i \langle \Psi_{\text{exact}}(t) | [\mathcal{O}_{i\alpha}, \mathcal{H}] | \Psi_{\text{exact}}(t) \rangle \\ &\equiv -i \langle \Psi(t) | [\mathcal{O}_{i\alpha}, \mathcal{H}] | \Psi(t) \rangle = -\frac{\partial E}{\partial \phi_{i\alpha}}, \end{aligned} \quad (3)$$

where the equivalence is our variational assumption. Within the Gutzwiller approximation [21], which is exact in the limit of infinite coordination lattices, $E = \langle \Phi(t) | \mathcal{H}_*(t) | \Phi(t) \rangle$, where $\mathcal{H}_*(t)$ is a noninteracting Hamiltonian that depends on all time-dependent variational parameters defining $\mathcal{P}_i$ and S_i . In general, these parameters can be expressed in terms of $\phi_{i\alpha}$ and $\mathcal{O}_{i\alpha}$. If we impose that $|\Phi(t)\rangle$ is the solution of the Schrödinger equation

$$-i \frac{\partial}{\partial t} |\Phi(t)\rangle = \mathcal{H}_*(t) |\Phi(t)\rangle, \quad (4)$$

and furthermore that

$$\frac{\partial \phi_{i\alpha}}{\partial t} = \frac{\partial E}{\partial \mathcal{O}_{i\alpha}}, \quad (5)$$

conservation of energy follows automatically. Therefore, $\phi_{i\alpha}$ and $\mathcal{O}_{i\alpha}$ act like conjugate variables and the energy functional E as their effective Hamiltonian.

As a simple application of the above variational scheme, we assume $\mathcal{H}$ to be the Hubbard model (1) at half-filling with $U(t \leq 0) = U_i \geq 0$ and $U(t > 0) = U_f > U_i$, and furthermore we limit our analysis to homogeneous paramagnetic wave functions. In the limit of infinite coordination lattices, one can compute exactly average values on the variational wave function provided the following conditions are imposed [21,22]

$$\langle \Phi(t) | \mathcal{P}_i^2(t) | \Phi(t) \rangle = 1, \quad \langle \Phi(t) | \mathcal{P}_i^2(t) c_{i\sigma}^\dagger c_{i\sigma} | \Phi(t) \rangle = \frac{1}{2}.$$

We assume $\mathcal{P}_i(t) = \sum_{n=0}^2 \lambda_{i,n}(t) \mathcal{P}_{i,n}$, where $\mathcal{P}_{i,n}$ is the projector at site i onto configurations with $n = 0, \dots, 2$ electrons and $S_i(t) = \sum_{n=0}^2 \phi_{i,n}(t) \mathcal{P}_{i,n}$, which implies that $\phi_{i,n}(t)$ plays the role of the conjugate variable of the occupation probability $P_{i,n} = \langle \Psi(t) | \mathcal{P}_{i,n} | \Psi(t) \rangle$. From the constraints above it follows that $P_{i,0} = P_{i,2}$ and $P_{i,1} = 1 - 2P_{i,2}$. We define $P_{i,2} = (1 - \cos\theta_i)/4$ and set $\phi_{i,0} = \phi_{i,2} = \phi_i$ while $\phi_{i,1} = 0$. Using θ_i and ϕ_i as variational parameters, one finds the average energy [21]

$$E = \frac{U_f}{4} \sum_i (1 - \cos\theta_i(t)) - \sum_{ij} w_{ij}(t) \sin\theta_i(t) \cos\phi_i(t) \sin\theta_j(t) \cos\phi_j(t), \quad (6)$$

where $w_{ij}(t) = t_{ij} \sum_\sigma \langle \Phi(t) | c_{i\sigma}^\dagger c_{j\sigma} + \text{H.c.} | \Phi(t) \rangle$. One recognizes in (6) the mean-field energy of an Ising model in a transverse field

$$\mathcal{H}_I = \frac{U_f}{4} \sum_i (1 - \sigma_i^z) - \sum_{ij} w_{ij}(t) \sigma_i^x \sigma_j^x, \quad (7)$$

where $\langle \sigma_i^z \rangle = \cos\theta_i$ and $\langle \sigma_i^x \rangle = \sin\theta_i \cos\phi_i$. This connection can be established rigorously at the variational level [23], and agrees with the Z_2 -slave-spin theory recently introduced [24,25]. Therefore, it is not surprising that the equations of motion that we obtain through Eqs. (3) and (5) are just those of the Ising model $\langle \partial_t \sigma_i^a \rangle = -i \langle [\sigma_i^a, \mathcal{H}_I] \rangle$ within the mean field. In other words, under the above assumption of homogeneous and paramagnetic wave functions, a quantum quench in the half-filled Hubbard model is equivalent, within the Gutzwiller variational scheme, to a quench in a Ising model in the presence of a transverse field. In particular, if $|\Phi(t)\rangle$ is taken to be the half-filled Fermi sea, then $w_{ij}(t) = w$ and (7) is the conventional ferromagnetic Ising model with constant and uniform exchange w and transverse field $U_f/4$. Quantum quenches of the transverse field have been recently investigated in one dimension [9] and on a fully connected lattice [26]. In the following we assume that the system is prepared in the metallic variational wave function that optimizes (1) with $U = U_i < U_c$, where U_c is the variational estimate of the Mott transition.

This corresponds to initial values $\phi_i(0) = 0$ and $\cos\theta_i(0) = U_i/U_c \equiv u_i$ for the coupled equations: $2\dot{\phi} = U_c \cos\theta \cos^2\phi - U_f$ and $2\dot{\theta} = U_c \sin\theta \sin\phi \cos\phi$. We note that, apart from the trivial case in which

$U_f = U_i$, these equations admit a nontrivial stationary solution $\theta = 0$ and $\cos^2 \phi = U_f/U_c = u_f$, which is compatible with the initial conditions only when $u_f = u_{fc} = (1 + u_i)/2$. It turns out that u_{fc} identifies a dynamical critical point that separates two different regimes similarly to a simple pendulum. When $u_f < u_{fc}$, $2\phi(t)$ oscillates around the origin, while, for $u_f > u_{fc}$, it performs a cyclic motion around the whole circle.

In order to characterize the different regimes, we focus on three physical quantities, the double occupancy $D(t) = (1 - \cos \theta(t))/4$, the quasiparticle residue $Z(t) = \sin^2 \theta(t) \cos^2 \phi(t)$ and their period of oscillation, $\mathcal{T}$.

While detailed calculations will be presented elsewhere [23], in the rest of the Letter we just sketch the results of the mean-field dynamics. Let us start from the weak coupling side $u_i < u_f < u_{fc}$, see top panel in Fig. 1, where both $D(t)$ and $Z(t)$ display small oscillations. Their amplitude and period *increase* with the strength of the quench $\delta u = u_f - u_i$, the latter reading $\mathcal{T} = \frac{4\sqrt{2}\mathcal{K}(k)}{\sqrt{Z(0)}}$, where $\mathcal{K}(k)$ is the complete elliptic integral of the first kind with argument $k^2 = 4u_f \delta u / Z(0)$. For $\delta u \rightarrow 0$ we find a linear increase $\mathcal{T} \simeq \mathcal{T}_0(1 + u_i \delta u / Z(0))$ with $\mathcal{T}_0 = 4\pi/\sqrt{Z(0)}$.

Conversely, when quenching above the critical value, $u_f > u_{fc}$, a novel strong-coupling dynamical behavior emerges. Here oscillations become faster, their period $\mathcal{T} = 4\mathcal{K}(1/k)/\sqrt{u_f \delta u}$ now decreases as a function of δu . In particular, for $u_f \gg u_i$, we get $\mathcal{T} \simeq \frac{2\pi}{u_f}$ smoothly matching the atomic limit result. The oscillation amplitude of $D(t)$ decreases with u_f , which results into a frozen dynamics in the infinite quench limit [7], while quasipar-

ticle weight $Z(t)$ still shows large oscillations even for $u_f \rightarrow \infty$, mainly reflecting the unbounded dynamics of the phase $\phi(t)$.

Remarkably, the weak and the strong-coupling regimes are separated by a critical quench line u_{fc} at which mean-field dynamics exhibits *exponential relaxation*. Indeed, upon approaching this line from both sides, the period $\mathcal{T}$ diverges logarithmically, $\mathcal{T} \simeq \frac{4}{\sqrt{Z(0)}} \log(\frac{1}{|u_f - u_{fc}|})$. Right at criticality, $u_f = u_{fc}$, the mean-field dynamics can be integrated exactly. The result gives $D(t) = D(0)(1 - \tanh^2(t/\tau_*))$; the double occupation relaxes exponentially to $\bar{D} = 0$ pushing also $\bar{Z} \rightarrow 0$, with a characteristic time scale $\tau_* = 2/\sqrt{Z(0)}$ that increases upon approaching the Mott Insulator.

We now turn to discuss long-time average properties of the Gutzwiller mean-field dynamics that we define as $\bar{O} = \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t dt' O(t')$. The analytical expressions [23] of $\bar{D}$ and $\bar{Z}$ are shown in Fig. 2. At weak coupling, we find $\bar{D} = D(0)[1 + \frac{u_f}{u_i} \frac{(E(k) - \mathcal{K}(k))}{\mathcal{K}(k)}]$, where $E(k)$ is the complete elliptic integral of the second kind and k^2 is the same as before. In addition, due to energy conservation, the knowledge of $\bar{D}$ completely fixes the average quasiparticle weight $\bar{Z} = Z(0) + 8u_f(\bar{D} - D(0))$.

It is interesting to consider first the small quench limit $\delta u \rightarrow 0$. We find that, given $D(0) = (1 - u_i)/4$ the initial equilibrium value, $\bar{D} \simeq D(0) - \delta u/4 = (1 - u_f)/4$, namely, tends to the equilibrium value corresponding to the final interaction. Hence the prethermalization result [17] for the quasiparticle weight $\bar{Z}$ immediately follows. Indeed, quenching from a noninteracting Fermi sea, $u_i = 0$ hence $Z(0) = 1$, we find that the nonequilibrium $\bar{Z}$ is reduced twice more than its equilibrium value at U_f .




FIG. 1 (color online). Left Panels: Gutzwiller mean-field dynamics at half-filling for quasiparticle weight $Z(t)$ (black line) and double occupancy $D(t)$ (dashed red line) for quantum quenches from $u_i = 0.25$ to $u_f = 0.35$ (top panel) and $u_f = 1.25$ (bottom panel). Right Panel: Period of oscillations at half-filling and for a finite doping $\delta \neq 0$. Note the logarithmic singularity at u_{fc} for $\delta = 0$ (see main text) which is cut-off by finite doping.




FIG. 2 (color online). Average double occupancy $\bar{D}$ (top) and quasiparticle weight $\bar{Z}$ (bottom) as a function of u_f at fixed $u_i = 0.0, 0.5$. We show results at half-filling (full lines) that display a sharp transition at u_{fc} , as well as at finite doping (dashed lines) where only a crossover remains. We also plot the zero temperature equilibrium results for $\delta = 0$ (red points).

For large quenches, the average double occupancy $\bar{D}$ increases as a function of the final interaction u_f , $\bar{D} = D(0) - \frac{1}{2}(u_f - u_i)(1 - E(1/k)/\mathcal{K}(1/k))$, eventually approaches its initial value $D(0)$ for $u_f \rightarrow \infty$. A similar behavior is found for $\bar{Z}$, which is, however, further reduced by a factor 1/2 with respect to the initial value $Z(0)$ due to the freely oscillating behavior of the phase. We find therefore that, for large quantum quenches, the dynamics retains memory of the initial condition and thermalization is prevented by a dynamical blocking.

Finally, for quenches close to u_{fc} , both $\bar{D}$ and $\bar{Z}$ are very small, vanishing as $1/\log|u_f - u_{fc}|$ on approaching the critical point. Namely, u_{fc} not only signals a transition in the dynamics but also identifies the critical interaction at which the quenched system shows genuine Mott insulating behavior.

Away from half-filling, the dynamical equations become more cumbersome [23]. However, key features can be easily derived even without resorting to a numerical integration. In particular, we find that any finite doping turns the half-filled dynamical critical point into a crossover. For instance, the logarithmic singularity of the oscillatory period $\mathcal{T}$ is cutoff by any finite doping, as shown in Fig. 1. As a consequence, the singular behavior of the average values across the half-filling transition is smoothed into a crossover at finite doping; see Fig. 2.

Discussion.—It is worth discussing the above results in light of those recently obtained by DMFT [18]. Remarkably, our variational ansatz (2) seems to catch many nontrivial effects observed in DMFT. In particular the existence of two different regimes separated by a real dynamical transition at u_{fc} , already suggested in [18], clearly emerges from our mean field theory. We note, however, that the suppression of quantum fluctuations, which is at the ground of our results, give rise to an oversimplified periodical dynamics that lacks relaxation. In this respect DMFT, which can treat exactly all local quantum fluctuations, works much better and turns these oscillations into a true relaxation. We guess that a similar result could be obtained from our variational treatment, for example, by allowing fluctuations in the Fermi sea and treating the quantum Ising model beyond the simplest mean field level [24]. Finally, we notice that the Ising analogy provides a simple interpretation of the dynamical transition, at least from a local DMFT-like point of view. Indeed, if we assume that the role of the neglected quantum fluctuations is to provide dissipation for the two-level system described by the local Ising variable, the two dynamical regimes found by DMFT resemble those in the phase diagram of the spin-boson model [27].

Conclusion.—We introduced a variational approach to the out-of-equilibrium dynamics in strongly correlated electron systems. Using a time-dependent Gutzwiller ansatz we address the problem of a sudden change of the interaction in the fermionic Hubbard model finding a dynamical transition at half-filling. Our results provide a

simple and intuitive mean-field theory for the quench dynamics in interacting Fermi systems.

We acknowledge useful discussions with M. Capone, E. Tosatti, A. Millis, and C. Castellani. This work has been supported by Italian Ministry of University and Research, through a PRIN-COFIN Grant.

-
- [1] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
 - [2] M. Greiner, O. Mandel, T. Esslinger, T. Hansch, and I. Bloch, *Nature (London)* **415**, 39 (2002).
 - [3] T. Kinoshita, T. Wenger, and D. S. Weiss, *Nature (London)* **440**, 900 (2006).
 - [4] M. W. Zwierlein, C. H. Schunck, C. A. Stan, S. M. F. Raupach, and W. Ketterle, *Phys. Rev. Lett.* **94**, 180401 (2005).
 - [5] R. Jordens, N. Strohmaier, K. Gunter, H. Moritz, and T. Esslinger, *Nature (London)* **455**, 204 (2008).
 - [6] N. Strohmaier *et al.*, *Phys. Rev. Lett.* **104**, 080401 (2010).
 - [7] A. Rosch, D. Rasch, B. Binz, and M. Vojta, *Phys. Rev. Lett.* **101**, 265301 (2008).
 - [8] M. Eckstein and M. Kollar, *Phys. Rev. Lett.* **100**, 120404 (2008).
 - [9] D. Rossini, A. Silva, G. Mussardo, and G. E. Santoro, *Phys. Rev. Lett.* **102**, 127204 (2009).
 - [10] M. A. Cazalilla, *Phys. Rev. Lett.* **97**, 156403 (2006).
 - [11] C. Kollath, A. M. Läuchli, and E. Altman, *Phys. Rev. Lett.* **98**, 180601 (2007).
 - [12] S. R. Manmana, S. Wessel, R. M. Noack, and A. Muramatsu, *Phys. Rev. Lett.* **98**, 210405 (2007).
 - [13] C. De Grandi, V. Gritsev, and A. Polkovnikov, *Phys. Rev. B* **81**, 012303 (2010).
 - [14] P. Barmettler, M. Punk, V. Gritsev, E. Demler, and E. Altman, *Phys. Rev. Lett.* **102**, 130603 (2009).
 - [15] M. Rigol, V. Dunjko, and M. Olshanii, *Nature (London)* **452**, 854 (2008).
 - [16] G. Biroli, C. Kollath, and A. M. Läuchli, *arXiv:0907.3731v1*.
 - [17] M. Moeckel and S. Kehrein, *Phys. Rev. Lett.* **100**, 175702 (2008).
 - [18] M. Eckstein, M. Kollar, and P. Werner, *Phys. Rev. Lett.* **103**, 056403 (2009).
 - [19] W. F. Brinkman and T. M. Rice, *Phys. Rev. B* **2**, 4302 (1970); M. C. Gutzwiller, *Phys. Rev.* **137**, A1726 (1965).
 - [20] G. Seibold and J. Lorenzana, *Phys. Rev. Lett.* **86**, 2605 (2001).
 - [21] M. Fabrizio, *Phys. Rev. B* **76**, 165110 (2007).
 - [22] J. Bünenmann, W. Weber, and F. Gebhard, *Phys. Rev. B* **57**, 6896 (1998).
 - [23] M. Schirò and M. Fabrizio (to be published).
 - [24] S. D. Huber and A. Rüegg, *Phys. Rev. Lett.* **102**, 065301 (2009).
 - [25] A. Rüegg, S. D. Huber, and M. Sigrist, *Phys. Rev. B* **81**, 155118 (2010).
 - [26] A. Das, K. Sengupta, D. Sen, and B. K. Chakrabarti, *Phys. Rev. B* **74**, 144423 (2006).
 - [27] F. B. Anders, R. Bulla, and M. Vojta, *Phys. Rev. Lett.* **98**, 210402 (2007).