

## Sign of Kurtosis near the QCD Critical Point

M. A. Stephanov

*Department of Physics, University of Illinois, Chicago, Illinois 60607, USA*  
(Received 27 April 2011; published 26 July 2011)

We point out that the quartic cumulant (and kurtosis) of the order parameter fluctuations is universally *negative* when the critical point is approached on the crossover side of the phase separation line. As a consequence, the kurtosis of a fluctuating observable, such as, e.g., proton multiplicity, may become smaller than the value given by independent Poisson statistics. We discuss implications for the beam energy scan program at RHIC at BNL.

DOI: 10.1103/PhysRevLett.107.052301

PACS numbers: 25.75.-q, 25.75.Nq, 25.75.Gz, 05.70.Jk

*Introduction.*—Mapping the QCD phase diagram as a function of temperature  $T$  and baryochemical potential  $\mu_B$  is one of the fundamental goals of heavy-ion collision experiments. QCD critical point is a distinct singular feature of the phase diagram. It is a ubiquitous property of QCD models based on the chiral symmetry breaking dynamics (see, e.g., Ref. [1] for a review and further references). Locating the point using first-principles lattice calculations is a formidable challenge (see, e.g., Ref. [2] for a recent review and references). If the critical point is situated in the region accessible to heavy-ion collision experiments, it can be discovered experimentally. The search for the critical point is planned at the Relativistic Heavy Ion Collider (RHIC) at BNL, the Super Proton Synchrotron (SPS) at CERN, the future Facility for Antiproton and Ion Research (FAIR) at GSI, and Nuclotron-based Ion Collider Facility (NICA) in Dubna (see, e.g., Ref. [3]).

The characteristic feature of a critical point is the divergence of the correlation length  $\xi$  and of the magnitude of the fluctuations. The simplest measures of fluctuations in heavy-ion collisions are the variances of the event-by-event observables such as multiplicities or mean transverse momenta of particles. The singular, critical contribution to these variances diverges as (approximately)  $\xi^2$ , and would manifest in a nonmonotonic dependence of such measures as the critical point is passed by during the beam energy scan [4,5]. In realistic heavy-ion collision the divergence of  $\xi$  is cut off by the effects of critical slowing down [5,6], and the estimates of the maximum correlation length are in the range of at most 2–3 fm, compared to the natural 0.5–1 fm away from the critical point. However, higher, non-Gaussian, moments of the fluctuations depend much more sensitively on  $\xi$ , according to Ref. [7]. For example, the 4th moment grows as  $\xi^7$  near the critical point, making it an attractive experimental tool. In this Letter we follow up on the results of Ref. [7] to point out that the sign of the 4th moment could be negative as the critical point is approached from the crossover side of the QCD phase transition.

The sign of various moments have been discussed in the literature in related contexts: see, e.g., discussion of the

sign of the 3rd moment in Ref. [8] or the 6th and 8th moments in Ref. [9] and also numerical lattice calculations in Ref. [10] where the possible sign change of kurtosis is noted.

In this Letter we shall address specifically the sign of the 4th moment (or kurtosis) and do it in a more universal and quantitative way than has been done previously, by using the known parametric form of the universal equation of state near the critical point. We emphasize universality of the behavior of the kurtosis and draw experimental consequences from these results.

*Kurtosis and universal effective potential.*—Let us begin, as in Ref. [1], by describing fluctuations of the order parameter field  $\sigma(\mathbf{x})$  near a critical point using the probability distribution

$$P[\sigma] \sim \exp\{-\Omega[\sigma]/T\}, \quad (1)$$

where  $\Omega$  is the effective action (free energy) functional for the field  $\sigma$ , which can be expanded in powers of  $\sigma$  as well as in the gradients (we chose  $\sigma = 0$  at the minimum):

$$\Omega = \int d^3x \left[ \frac{(\nabla\sigma)^2}{2} + \frac{m_\sigma^2}{2}\sigma^2 + \frac{\lambda_3}{3}\sigma^3 + \frac{\lambda_4}{4}\sigma^4 + \dots \right]. \quad (2)$$

Calculating 2-point correlator  $\langle\sigma(\mathbf{x})\sigma(0)\rangle$  we find that the correlation length  $\xi = m_\sigma^{-1}$ . For the moments of the zero momentum mode  $\sigma_V \equiv \int d^3x \sigma(\mathbf{x})$  in a system of volume  $V$ , we find at tree level

$$\begin{aligned} \kappa_2 = \langle\sigma_V^2\rangle &= VT\xi^2; & \kappa_3 = \langle\sigma_V^3\rangle &= 2\lambda_3 VT^2\xi^6; \\ \kappa_4 = \langle\sigma_V^4\rangle_c &= 6VT^3[2(\lambda_3\xi)^2 - \lambda_4]\xi^8, \end{aligned} \quad (3)$$

where  $\langle\sigma_V^4\rangle_c \equiv \langle\sigma_V^4\rangle - 3\langle\sigma_V^2\rangle^2$  denotes the connected 4th central moment (the 4th cumulant). The critical point is characterized by  $\xi \rightarrow \infty$ . The central observation in Ref. [7] was that the higher moments (cumulants)  $\kappa_3$  and  $\kappa_4$  diverge with  $\xi$  much faster than the quadratic moment  $\kappa_2$ . Here we shall point out that the *sign* of the 4th moment  $\kappa_4$  is negative in a certain sector near the critical point. More precisely, the 4th cumulant is negative when the critical point is approached from the crossover side. Let us demonstrate this in several complementary ways.

A simple way to see why the kurtosis is negative is by following the evolution of the probability distribution of  $\sigma_V$  as we approach the critical point along the crossover line. In Ising scaling coordinates: from above the critical temperature, i.e.,  $t \equiv (T - T_c)/T_c > 0$ , along the line of zero magnetic field  $H = 0$ . Away from the critical point, more precisely for  $\xi^3 \ll V$ , the central limit theorem dictates that the probability distribution of  $\sigma_V$  is Gaussian, with a vanishingly small kurtosis. As we approach the critical point the distribution develops non-Gaussian shape. This intermediate shape is a deformation of the Gaussian towards a two-peak distribution, corresponding to the phase coexistence on the opposite, first-order transition side ( $t < 0$ ) of the critical point. Such a shape is clearly less “peaked” than the Gaussian, and thus corresponds to negative kurtosis.

More quantitatively, the kurtosis vanishes as  $1/V$  at (almost) any point away from the critical point, i.e.,

$$K \equiv \kappa_4/\kappa_2^2 = O(\xi^3/V). \quad (4)$$

The exception is the coexistence line ( $H = 0$ ,  $t < 0$  ray). The distribution there has two peaks of equal height and its kurtosis is  $K = -2 + O(\xi^3/V)$ .

It is important to note that this is only true strictly *on* the coexistence line  $H = 0$ , for the moments measured around the symmetric point of the probability distribution of  $\sigma_V$ , which is actually a dip, not a peak, for  $t < 0$ . At any point close to the coexistence line, i.e., at  $H \neq 0$ ,  $t < 0$ , the kurtosis around the dominant peak is positive.

In the scaling regime (close to but not at the critical point) where  $\xi$  is much greater than the microscopic scale  $a$ , but still much less than the linear size of the system,  $a \ll \xi \ll V^{1/3}$ , the coefficient of  $\xi^3/V$  in Eq. (4) can be expressed in terms of the couplings  $\lambda_i$  using Eq. (3):

$$K = 6(2\lambda_3^2\xi^3 - \lambda_4\xi)\frac{\xi^3}{V}. \quad (5)$$

These couplings, and in fact the shape of the effective potential, are also universal. In particular,  $\lambda_4$  scales with  $\xi$  as  $\lambda_4 = \tilde{\lambda}_4(T\xi)^{-1}$ , where the universal value of  $\tilde{\lambda}_4$  is known approximately to be 4.0 on the crossover line (see, e.g., Ref. [11] for a review). (As in Ref. [7], for simplicity and consistency with our overall level of precision, we neglect the anomalous scaling dimension  $\eta$ , which is only of order few percent.) Since on the crossover line  $\lambda_3 = 0$  and  $\lambda_4 > 0$ , it is clear from Eq. (3) that  $K < 0$ .

Away from the crossover line ( $H = 0$ ,  $t > 0$  ray) the distribution is skewed:  $\lambda_3 \neq 0$ . This makes the kurtosis positive, according to Eq. (5), except for a certain sector around the crossover line.

*Universal equation of state around the critical point.*—To extend this analysis away from the crossover line, i.e., to take into account  $\lambda_3 \neq 0$  in Eq. (5), we need to know the equation of state, in particular,  $\kappa_4$  as a function of both Ising variables: reduced temperature  $t$  and magnetic field

$H$ . In the scaling regime near  $t = H = 0$  this equation of state is also universal. For the Ising model it is known to order  $\varepsilon^3$  in the epsilon expansion as well as numerically.

Before we discuss this universal form, let us keep in mind that the mapping of QCD phase diagram in the  $T, \mu_B$  plane into  $t, H$  plane is not universal. However, this mapping is analytic, i.e., both functions  $t(T, \mu_B)$  and  $H(T, \mu_B)$  are analytic at the critical point, which is mapped into the origin,  $t(T^{\text{cp}}, \mu_B^{\text{cp}}) = H(T^{\text{cp}}, \mu_B^{\text{cp}}) = 0$ .

The standard parametrization, Ref. [12], of the equation of state in the scaling domain near the critical point is in terms of two new scaling variables  $R$  and  $\theta$  (it has been applied in the context of QCD before, Ref. [13]). Denoting the “magnetization” by  $M = \langle \sigma_V \rangle / V$ , we define  $R$  and  $\theta$  as

$$M = R^\beta \theta, \quad t = R(1 - \theta^2). \quad (6)$$

Then the equation of state can be expressed in terms of the single function  $h(\theta)$  as

$$H = R^{\beta\delta} h(\theta). \quad (7)$$

Unlike the explicit form of the singular equation of state  $M = M(t, H)$ , the function  $h(\theta)$  is analytic. It has two zeros. One, at  $\theta = 0$ , corresponds to the crossover line ( $t > 0$ ,  $H = 0$  ray), another, at some  $\theta = \theta_1 > 1$ , corresponds to the coexistence (first-order transition) line ( $t < 0$ ,  $H = 0$  ray). The function  $h(\theta)$  must also be odd since  $M(-H) = -M(H)$ . The simplest function obeying all these requirements is a cubic polynomial

$$h(\theta) = \theta(2 - 3\theta^2), \quad (8)$$

where the value  $\theta_1 = \sqrt{3/2}$  is a good approximation to the universal value for the Ising model [and correct up to  $O(\varepsilon^2)$ ]. The choice (8) is known as the linear parametric model, Ref. [14]. It describes the equation of state with precision quite sufficient for our purposes. The linear parametric model is also known to be exact up to  $O(\varepsilon^3)$ .

Using this parametric equation of state, we can calculate the moments by taking derivatives at fixed  $t$ , up to an overall normalization, unimportant in the present context [it can be fixed by Eq. (3)]. In particular,

$$\kappa_4(t, H) = \left( \frac{\partial^3 M}{\partial H^3} \right)_t. \quad (9)$$

For our purposes, it would be sufficient to use the approximate rational values of critical exponents  $\beta = 1/3$  and  $\delta = 5$ , which are within a few percent of their exact values in three dimensions. The result of Eq. (9) can then be simplified to

$$\kappa_4(t, H) = -12 \frac{81 - 783\theta^2 + 105\theta^4 - 5\theta^6 + 2\theta^8}{R^{14/3}(3 - \theta^2)^3(3 + 2\theta^2)^5}. \quad (10)$$

We represent  $\kappa_4(t, H)$  graphically as a density plot in Fig. 1. We see that the 4th cumulant (and kurtosis) is




FIG. 1 (color online). (a) The density plot of the function  $\kappa_4(t, H)$  given by Eq. (10) obtained using Eq. (9) for the linear parametric model Eqs. (6)–(8) and  $\beta = 1/3$ ,  $\delta = 5$ . The  $\kappa_4 < 0$  region is red, the  $\kappa_4 > 0$  is blue. (b) The dependence of  $\kappa_4$  on  $t$  along the vertical dashed green line on the density plot in (a). This line is the simplest example of a possible mapping of the freeze-out curve (see Fig. 2). The units of  $t$ ,  $H$ , and  $\kappa_4$  are arbitrary.

negative in the sector bounded by two curved rays  $H/t^{\beta\delta} = \pm \text{const}$  (corresponding to  $\theta \approx \pm 0.32$ ).

Also in Fig. 1 we show the dependence of  $\kappa_4$  along a line which could be thought of as representing a possible mapping of the freeze-out trajectory (Fig. 2) onto the  $tH$  plane. Although the absolute value of the peak in  $\kappa_4$  depends on the proximity of the freeze-out curve to the critical point, the ratio of the maximum to minimum along such an  $H = \text{const}$  curve is a universal number, approximately equal to  $-28$  from Eq. (10).

The negative minimum is small relative to the positive peak, but given the large size of the latter, Refs. [7,15], the negative contribution to kurtosis may be significant. In addition, the mapping of the freeze-out curve certainly need not be  $H = \text{const}$ , and the relative size of the positive and negative peaks depends sensitively on that.

The trend described above appears to show in the recent lattice data, Ref. [10], obtained using Padé resummation of the truncated Taylor expansion in  $\mu_B$ . As the chemical potential is increased along the freeze-out curve, the 4th




FIG. 2 (color online). A sketch of the phase diagram of QCD with the freeze-out curve and a possible mapping of the Ising coordinates  $t$  and  $H$ .

moment of the baryon number fluctuations begins to decrease, possibly turning negative, as the critical point is approached (see Fig. 2 in Ref. [10]).

Another observation is that  $-\kappa_4$  grows as we approach the crossover line, corresponding to  $H = 0$ ,  $t > 0$  on the diagram in Fig. 1(a). On the QCD phase diagram the freeze-out point will move in this direction if one reduces the size of the colliding nuclei or selects more peripheral collisions (the freeze-out occurs earlier, i.e., at higher  $T$ , in a smaller system).

*Experimental observables.*—In this section we wish to connect the results for the fluctuations of the order parameter field  $\sigma$  to the fluctuations of the observable quantities. As an example we consider the fluctuations of the multiplicity of given charged particles, such as pions or protons.

For completeness we shall briefly rederive the results of Ref. [7] using a simple model of fluctuations. The model captures the most singular term in the contribution of the critical point to the fluctuation observables. Consider a given species of particle interacting with fluctuating critical mode field  $\sigma$ . The infinitesimal change of the field  $\delta\sigma$  leads to a change of the effective mass of the particle by the amount  $\delta m = g\delta\sigma$ . This could be considered a definition of the coupling  $g$ . For example, the coupling of protons in the sigma model is  $g\sigma\bar{p}p$ . The fluctuations  $\delta f_p$  of the momentum space distribution function  $f_p$  consist of the pure statistical fluctuations  $\delta f_p^0$  around the equilibrium distribution  $n_p$  for a particle of a given mass, which itself fluctuates. This gives

$$\delta f_p = \delta f_p^0 + \frac{\partial n_p}{\partial m} g \delta\sigma. \quad (11)$$

Using this equation we can calculate the most singular contribution from the critical fluctuations to the moments or correlators of  $\delta f_p$ . The fluctuation of the multiplicity  $N = Vd \int_p f_p$  is given by

$$\delta N = \delta N^0 + Vg\delta\sigma d \int_p \frac{\partial n_p}{\partial m}, \quad (12)$$

where  $d$  is the degeneracy factor (e.g., number of spin or charge states of the particle). Neglecting, for clarity and simplicity, the effects of quantum statistics, i.e., assuming  $n_p \ll 1$ , we can use Poisson statistics for  $\delta N^0$ . Using additivity of the cumulants (their defining property), and assuming  $\delta N^0$  and  $\delta\sigma$  are uncorrelated, the contribution of the critical fluctuations can be expressed in terms of the corresponding moments of the critical field  $\sigma$  fluctuations. For example, the contribution to the 4th moment can be expressed as (cf. Refs. [7,15])

$$\langle(\delta N)^4\rangle_c = \langle N \rangle + \langle \sigma_V^4 \rangle_c \left( \frac{gd}{T} \int_p \frac{n_p}{\gamma_p} \right)^4 + \dots, \quad (13)$$

where  $\gamma_p = (dE_p/dm)^{-1}$  is the relativistic gamma factor of a particle with momentum  $\mathbf{p}$  and mass  $m$ . The first term on the right-hand side of Eq. (13) is the Poisson contribution. We neglected  $n_p \ll 1$  in the quantum statistics factor  $(1 \pm n_p)$  for simplicity, and we denoted by “...” other contributions, less singular at the critical point. The model is admittedly crude, but it illustrates the mechanism and correctly captures the most singular contribution near the critical point.

In the region near the critical point where  $\kappa_4 = \langle \sigma_V^4 \rangle_c$  is negative, the 4th cumulant of the fluctuations will be smaller than its Poisson value,  $\langle N \rangle$ . By how much will depend sensitively on the correlation length (as  $\xi^7$ ), i.e., on how close the freeze-out occurs to the critical point, as well as on other factors (for protons, most significantly, on the value of  $\mu_B$ ). We shall not attempt to estimate this effect quantitatively in this Letter. The analysis of Ref. [15] suggests, however, that this effect for protons can be significant compared to the Poisson value already for  $\xi \sim 2$  fm.

Usual caveats apply: other (nontrivial) contributions to moments which do not behave singularly at the critical point can turn out to be relatively large. These include initial geometry fluctuations, jets, and other nonequilibrium effects. In addition, charge conservation effects may impose constraints on certain observables, such as total charge fluctuations. It is beyond the scope of this Letter to estimate these effects. A comprehensive review can be found in Ref. [16]. The size of these background contributions could, in principle, be determined experimentally by performing measurements away from the critical point.

We conclude by asking an obvious question: has the effect of the negative kurtosis been observed? Data from STAR indicate that at  $\sqrt{s} = 19.6$  GeV the ratio  $\kappa_4/\kappa_2$  might be substantially smaller than its Poisson value 1, see Fig. 6 in Ref. [17], while it is very close to 1 at higher  $\sqrt{s}$  (smaller  $\mu_B$ ). Unfortunately, the statistics gathered in the short run at  $\sqrt{s} = 19.6$  GeV are clearly not sufficient to make a reliable conclusion. It would be interesting to see if this effect persists with more statistics at this energy. If confirmed, this result could indicate that the critical point is close, at somewhat larger values of  $\mu_B$  (smaller  $\sqrt{s}$ ). At smaller values of  $\sqrt{s}$  the effect should change sign, increasing kurtosis above its Poisson value.

The author thanks O. Evdokimov for discussions. This work is supported by the DOE Grant No. DE-FG0201ER41195.

- 
- [1] M. A. Stephanov, *Prog. Theor. Phys. Suppl.* **153**, 139 (2004).
  - [2] P. de Forcrand, *Proc. Sci., LAT2009* (2009) 010; S. Gupta, *Proc. Sci., LATTICE2010* (2010) 007.
  - [3] B. Mohanty, *Nucl. Phys.* **A830**, 899C (2009).
  - [4] M. A. Stephanov, K. Rajagopal, and E. V. Shuryak, *Phys. Rev. Lett.* **81**, 4816 (1998).
  - [5] M. A. Stephanov, K. Rajagopal, and E. V. Shuryak, *Phys. Rev. D* **60**, 114028 (1999).
  - [6] B. Berdnikov and K. Rajagopal, *Phys. Rev. D* **61**, 105017 (2000).
  - [7] M. A. Stephanov, *Phys. Rev. Lett.* **102**, 032301 (2009).
  - [8] M. Asakawa, S. Ejiri, and M. Kitazawa, *Phys. Rev. Lett.* **103**, 262301 (2009).
  - [9] B. Friman, F. Karsch, K. Redlich, and V. Skokov, [arXiv:1103.3511](https://arxiv.org/abs/1103.3511).
  - [10] R. V. Gavai and S. Gupta, *Phys. Lett. B* **696**, 459 (2011).
  - [11] M. M. Tsy-pin, *Phys. Rev. Lett.* **73**, 2015 (1994).
  - [12] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena* (Oxford University Press, Oxford, 1996), 3rd ed.
  - [13] C. Nonaka and M. Asakawa, *Phys. Rev. C* **71**, 044904 (2005).
  - [14] P. Schofield, *Phys. Rev. Lett.* **22**, 606 (1969).
  - [15] C. Athanasiou, K. Rajagopal, and M. Stephanov, *Phys. Rev. D* **82**, 074008 (2010).
  - [16] V. Koch, [arXiv:0810.2520](https://arxiv.org/abs/0810.2520).
  - [17] L. Kumar (STAR Collaboration), [arXiv:1101.4310](https://arxiv.org/abs/1101.4310).