

Direct Observation of Two Protons in the Decay of ^{54}Zn

P. Ascher,¹ L. Audirac,¹ N. Adimi,^{1,2} B. Blank,¹ C. Borcea,³ B. A. Brown,⁴ I. Companis,³ F. Delalee,¹ C. E. Demonchy,¹ F. de Oliveira Santos,⁵ J. Giovinazzo,¹ S. Grévy,¹ L. V. Grigorenko,⁶ T. Kurtukian-Nieto,¹ S. Leblanc,¹ J.-L. Pedroza,¹ L. Perrot,⁵ J. Pibernat,¹ L. Serani,¹ P. C. Srivastava,^{5,7} and J.-C. Thomas⁵

¹Centre d'Études Nucléaires de Bordeaux Gradignan—Université Bordeaux I—UMR 5797 CNRS/IN2P3, Chemin du Solarium, BP 120, 33175 Gradignan, France

²Faculté de Physique, USTHB, B.P.32, El Alia, 16111 Bab Ezzouar, Alger, Algeria

³National Institute for Physics and Nuclear Engineering, P.O. Box MG6, Bucharest-Margurele, Romania

⁴Department of Physics and Astronomy, and National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, Michigan 48824-1321, USA

⁵Grand Accélérateur National d'Ions Lourds, CEA/DSM-CNRS/IN2P3, 14076 Caen Cedex 05, France

⁶Flerov Laboratory of Nuclear Reactions, JINR, Dubna 141980, Russian Federation

⁷Nuclear Physics Group, Department of Physics, University of Allahabad, Allahabad 211002, India

(Received 13 April 2011; published 1 September 2011)

The two protons emitted in the decay of ^{54}Zn have been individually observed for the first time in a time projection chamber. The total decay energy and the half-life measured in this work agree with the results obtained in a previous experiment. Angular and energy correlations between the two protons are determined and compared to theoretical distributions of a three-body model. Within the shell model framework, the relative decay probabilities show a strong contribution of the p^2 configuration for the two-proton emission. After ^{45}Fe , the present result on ^{54}Zn constitutes only the second case of a direct observation of the ground state two-proton decay of a long-lived isotope.

DOI: 10.1103/PhysRevLett.107.102502

PACS numbers: 23.50.+z, 27.40.+z, 29.40.Cs, 29.40.Gx

The study of exotic decays is an efficient tool to explore the structure of nuclei at the proton dripline. In particular, the emission of two protons from the ground state of a radioactive nucleus has been searched for since 1960, when two-proton ($2p$) radioactivity was predicted by Goldansky [1] for nuclei beyond the proton dripline, for which one-proton emission is energetically prohibited. This new nuclear decay mode was observed first in the decay of ^{45}Fe in two independent experiments [2,3] and later also in ^{54}Zn [4] and possibly in ^{48}Ni [5]. In these experiments, the ions of interest were deeply implanted in silicon detectors in which the decay was observed. Therefore, only the total decay energy, the half-life, and the absence of β particles from the competing decay by β -delayed-particle emission could be clearly established. In addition, the observation of the daughter decay helped to unambiguously assign the observed decay to $2p$ radioactivity. These experimental results were found in reasonable agreement with predictions from different theoretical models [6], such as the R -matrix theory [7], the shell model embedded in the continuum [8], or the three-body model [9,10].

However, in none of these experiments were the two protons identified separately, while the main physics question in the context of $2p$ radioactivity is how the two protons emitted are correlated in energy and in angle. An answer to that would enable us to investigate the decay dynamics of $2p$ radioactivity and thus reveal details of nuclear structure at the limits of stability. In particular,

these studies should reveal the single-particle ordering and other details of the wave function.

In an experiment performed in 2005 at GANIL (Caen, France), emission of two protons in the decay of ^{45}Fe was observed directly for the first time with a time projection chamber (TPC) [11]. The purpose of this detection setup is to reconstruct the proton tracks in three dimensions. In another experiment performed at Michigan State University (MSU)[12,13], the $2p$ emission in the decay of ^{45}Fe was observed with an optical time projection chamber in which images of ionizing particle trajectories are recorded optically. In this latter work, high statistics data allowed the authors to obtain a first meaningful comparison with a model including the three-body dynamics of the process [9,10].

In this Letter, we report on an experiment where emission of two protons in the decay of ^{54}Zn was observed with the TPC. Angular and energy correlations have been determined. These results allowed a first comparison with theoretical predictions of the three-body model and the nuclear shell model.

The ^{54}Zn nuclei were produced by quasifragmentation of the projectile at GANIL. A primary $^{58}\text{Ni}^{26+}$ beam with an energy of 74.5 MeV/nucleon and an average intensity of 3.5 μA was fragmented in a ^{nat}Ni target (200 μm). The ^{54}Zn fragments were selected by a magnetic-rigidity, energy-loss, and velocity analysis by means of the LISE3 separator including an achromatic energy degrader (500 μm of beryllium).

Two silicon detectors located at the end of the spectrometer allowed us to identify individually the fragments by means of an energy-loss and time-of-flight analysis. The identification parameters were first determined for frequently produced nuclei. Then, the parameters were extrapolated to unambiguously identify the nuclei produced with low probability. Details of this procedure can be found in [14].

The main setting of the spectrometer was optimized for the production and transmission of ^{54}Zn . During a two week experiment, 18 ^{54}Zn nuclei have been produced, as expected according to extrapolations of the cross sections of $^{55,56}\text{Zn}$ [15] and ion beam optical calculations. These ions were finally implanted in the TPC [16].

This detector is based on the principle of a time projection chamber. Ions of interest are implanted in a gas volume (argon 90%—methane 10%) at 750 mbar, where the radioactive decay takes place. The electrons, produced by the slowing down of either the incoming ions or the decay products, drift in the electric field of the TPC towards a set of four gas electron multipliers (GEM) where they are multiplied and finally detected in a two-dimensional strip detector. The analysis of energy signals allows us to reconstruct the tracks of the particles in two dimensions; the drift-time analysis gives the third one. Details can be found in [16].

Among the 18 ^{54}Zn implantation events, only 13 could be correlated in time and space with decays. Five decay events were lost due to the data acquisition dead time and the short half-life of ^{54}Zn . For two decay events, no information about the energy was obtained because the protons emitted did not stop in the active volume of the chamber. For the first event, the range of the protons emitted was very long (more than 8 cm). Such a long range is only compatible with a β -delayed proton emission. For the second event, due to the large momentum acceptance of the LISE3 spectrometer and the large range distribution of ^{54}Zn , the ion was implanted at the entrance of the TPC. The protons were emitted backwards and left the active volume. The other 11 decay signals could be analyzed in detail except one, for which a spurious response of one set of strips did not allow us to extract the relevant information. From this information, the branching ratio for $2p$ emission is determined as $\text{BR} = 92_{-13}^{+6}\%$.

The time difference between an implantation event and its subsequent decay event allowed a determination of the half-life of ^{54}Zn . This spectrum is shown in the left-hand part of Fig. 1. The half-life is determined as $1.59_{-0.35}^{+0.60}$ ms. Measurements of charge signals from the GEMs give access to the total decay energy. The right-hand part of Fig. 1 shows the signal extracted from the third GEM, the decay energy being thus determined to 1.28 ± 0.21 MeV. As can be seen from Table I, all these decay observables are in agreement with those determined in [4]. If we combine the present experimental BR and half-life with




FIG. 1 (color online). Left: Decay-time distribution obtained in the decay of ^{54}Zn , giving rise to a half-life of $1.59_{-0.35}^{+0.60}$ ms. Right: Energy spectrum of the ^{54}Zn decay events determined with the charge signals of one GEM. The total decay energy $E_{2p} = 1.28 \pm 0.21$ MeV is estimated from 11 events in the peak (see text for more details).

the previous values, we obtain $\text{BR} = 90_{-10}^{+5}\%$ and $T_{1/2} = 1.78_{-0.76}^{+0.66}$ ms, leading to a $2p$ partial half-life of $T_{1/2}^{2p} = 1.98_{-0.41}^{+0.73}$ ms.

Observables related to the measurements of individual protons were also determined. As an example, an implantation event spectrum is presented in Fig. 2. The ion enters with an angle of 45° in the chamber and stops at a given (X_0, Y_0) position. The implantation signals are fitted with a Gaussian folded with a straight line. This function is a good approximation of the Bragg peak corresponding to the energy loss of the charged particles inside the gas chamber. It allowed us to determine the implantation position of the ion (top part), which coincides with the emission position of the two protons (middle part). Their tracks in X and Y are fitted with the same function as for the implantation signals: the sum of two foldings of a straight line and a Gaussian, with the main constraint that the energy of a proton is the same along the X and Y direction. Figure 3(a) shows an example of a two-proton emission in two dimensions.

The energy fraction distribution of the individual protons as determined from the energy signals is plotted in Fig. 4 and is found in good agreement with the predictions of the three-body model. As expected in a simultaneous emission, the two protons share the decay energy equally in order to favor the barrier penetration. This theoretical approach [9,10] is the only model of $2p$ radioactivity which takes into account correlations between the two protons.

In a final step, the third component Z of the tracks was obtained. The bottom of Fig. 2 shows the time spectra corresponding to the same event. Only the spectrum on the Y dimension is analyzed because the protons were

TABLE I. Comparison of the experimental decay observables with the values obtained in a previous experiment.

| Experiments | $T_{1/2}$ (ms) | Q_{2p} (MeV) | BR (%) |
|-------------------------|------------------------|-----------------|------------------|
| Blank <i>et al.</i> [4] | $3.2_{-0.8}^{+1.8}$ | 1.48 ± 0.02 | 87_{-17}^{+10} |
| This work | $1.59_{-0.35}^{+0.60}$ | 1.28 ± 0.21 | 92_{-13}^{+6} |




FIG. 2 (color online). Example of energy-loss spectra obtained with the XY strip detector and associated with the implantation of a $2p$ emitter in the TPC. Top: Signals obtained for the implantation of ^{54}Zn : the ion enters in the chamber and stops at a given (X_0, Y_0) depth. The arrows indicate the direction of the incoming beam. The solid line is the fit of the implantation profile whereas the vertical line indicates the determined implantation position. Middle: The decay of ^{54}Zn takes place at the stopping point of the implantation event described above. The two protons are clearly identified. Their tracks are determined by fitting the decay signal with a sum of two foldings of a straight line and a Gaussian. The vertical full line corresponds to the starting point of the trajectories determined from the implantation profile of ^{54}Zn whereas the dashed lines are the two stopping points of the protons trajectories. Bottom: Corresponding time signals. The spectrum on the Y dimension is fitted by a straight line for each proton track, giving the directions of each individual proton.

emitted along the X strips. The spectrum is fitted by a straight line for each proton, giving the third component of each proton track. This drift-time analysis will be described in detail in a subsequent paper. Briefly, due to the short range of the protons, the determination of Δz with the




FIG. 3 (color online). (a) Two-dimensional view of a ^{54}Zn decay event as reconstructed from the XY strip detector. The color corresponds to the energy loss detected by the strips. (b) Same decay event reconstructed in the three-dimensional space.

time signals is not very accurate. Therefore, the time signals were only used to determine whether the proton goes upwards or downwards. Then, we use the theoretical range r of the protons [17] to determine Δz with $\Delta z^2 = r^2 - \Delta x^2 - \Delta y^2$, with Δx and Δy given by the energy signal analysis. The theoretical range was calculated by taking the energy sharing of the protons, as determined from the energy spectra analysis, and the sum energy, as determined from the previous measurements [4].

Among the ten events, seven have been fully reconstructed in the three-dimensional space. For the three remaining, the time signals did not allow us to determine if the protons went up or down. Therefore, we have for these events two possible angles between the two protons: the first one if the protons are emitted in the same hemisphere (up or down), the second one if they are emitted in different hemispheres. Figure 3(b) shows an example of a two-proton emission reconstructed in three dimensions.

The complete analysis of these decay events allowed us to provide angular correlations between the protons. The upper part of Fig. 5 shows each experimental angle,




FIG. 4 (color online). Energy sharing between the two protons emitted in the decay of ^{54}Zn . The dashed line is the energy distribution of the three-body model [10] folded with the response of the detector, which fits well the experimental distribution.

represented by a Gaussian reflecting the angular resolution. The middle part shows the angular distribution obtained. Seven events are represented in the histogram; the three other events, not fully reconstructed, are represented by full lines for the first possibility and by dashed lines for the second. In all cases, these three events are located below




FIG. 5 (color online). Top: Experimental angles between the two protons in the three-dimensional space. Each event is convoluted with a Gaussian representing the angular resolution. Middle: Experimental angular distribution between the two protons. Seven events are represented in the histogram. The dashed and full lines represent two possible angles for three not fully reconstructed events. Bottom: The three lines are the theoretical predictions of the three-body model, each line corresponding to different weights of the p^2 configuration. These model distributions are folded with a Gaussian function representing the detector angular resolution.

90° . Within the three-body model, the angular distribution spectra are calculated for different ℓ^2 configurations of the two emitted protons. The corresponding spectra (bottom part of Fig. 5) show a double-hump structure, with one broad peak centered around 50° and a smaller one at about 145° . When the p^2 contribution becomes negligible, the second hump does not survive. Considering that the experimental distribution shows a double-hump structure, the results can be compared with the model predictions by looking at the ratio between the first and the second hump. From an interpolation of the theoretical ratios, we obtain an experimental p^2 contribution of $30^{+33}_{-21}\%$. This number can be compared to a shell model wave function decomposition over the same p^2 and f^2 contributions using the GPF1A Hamiltonian [18] which yields about 80% for the p^2 contribution.

In the following, we will combine the shell model and the three-body model to determine theoretical half-lives. While the three-body model is adapted to treat the dynamics of the $2p$ emission, the shell model is more appropriate to describe the nuclear structure. Therefore, we use the spectroscopic factors of the shell model and the partial half-lives from the three-body model to determine the relative decay probabilities of the two ℓ^2 configurations, and thus to compare the experimental and the theoretical half-life. Two-proton removal amplitudes of a pair of protons have been calculated using the GPF1A Hamiltonian [18] and are found to be 0.3159, 0.3121, 0.6539, and 0.2631 for the $(0f_{7/2})^2$, $(0f_{5/2})^2$, $(1p_{3/2})^2$, and $(1p_{1/2})^2$ configurations, respectively. In LS coupling, the $S = L = 0$ removal amplitudes are 0.443 and 0.686 for $(0f)^2$ and $(1p)^2$ configurations, respectively. Combining the half-lives calculated by the three-body model for pure configurations [45 and 0.91 ms for $(0f)^2$ and $(1p)^2$, respectively] with the shell model removal amplitudes, we obtain the “shell model corrected” partial half-lives $T_{1/2}(0f^2) = 45/0.443^2 = 230$ ms and $T_{1/2}(1p^2) = 0.91/0.686^2 = 1.9$ ms. The total half-life of the $2p$ emission is therefore calculated by adding the two partial decay amplitudes coherently, with $\sqrt{1/T_{1/2}^{2p}} = \sqrt{1/1.9} + \sqrt{1/230}$, giving $T_{1/2}^{2p} = 1.6$ ms. This value is in excellent agreement with the experimental value obtained in this work ($T_{1/2}^{2p} = 1.98^{+0.73}_{-0.41}$ ms). From the above values, we can now derive the relative decay probability through the $(0p)^2$ configuration which is $P(p^2) = (1/1.9)/[(1/1.9) + (1/230)] = 0.99$. This means that almost all the decay strength goes through the $(0p)^2$ configuration of the wave function. An analysis of the experimental data from ^{45}Fe and ^{54}Zn in the frame work of the models used here will be the subject of a future publication [19].

In summary, we observed directly for the first time the two protons emitted in the decay of ^{54}Zn with a TPC. The half-life and Q value were determined and were found in good agreement with previous experiments and theoretical

models. Energy and angular distributions could be obtained and allowed a first rough comparison with theoretical models giving information about nuclear structure. However, to establish a detailed picture of the decay process, higher statistics of implantation-decay events are needed, which can be obtained in a future experiment. In parallel, improvements of theoretical model predictions are essential to elucidate the decay mechanism which governs two-proton radioactivity.

We would like to thank the whole GANIL and, in particular, the LISE staff and the DAQ group for their support during the experiment. The time projection chamber was financed in part by the Conseil régional d'Aquitaine. B.A.B. obtained support from NFS via Grants No. PHY-0758099 and No. PHY-1068217. Fruitful discussion with N. Smirnova are gratefully acknowledged.

-
- [1] V.I. Goldansky, *Nucl. Phys.* **19**, 482 (1960).
 - [2] J. Giovinazzo *et al.*, *Phys. Rev. Lett.* **89**, 102501 (2002).
 - [3] M. Pfützner *et al.*, *Eur. Phys. J. A* **14**, 279 (2002).
 - [4] B. Blank *et al.*, *Phys. Rev. Lett.* **94**, 232501 (2005).

- [5] C. Dossat *et al.*, *Phys. Rev. C* **72**, 054315 (2005).
- [6] B. Blank and M. Płoszajczak, *Rep. Prog. Phys.* **71**, 046301 (2008).
- [7] B. A. Brown and F. C. Barker, *Phys. Rev. C* **67**, 041304 (2003).
- [8] J. Rotureau, J. Okolowicz, and M. Płoszajczak, *Nucl. Phys. A* **767**, 13 (2006).
- [9] L. V. Grigorenko, R. C. Johnson, I. G. Mukha, I. J. Thompson, and M. V. Zhukov, *Phys. Rev. C* **64**, 054002 (2001).
- [10] L. V. Grigorenko and M. V. Zhukov, *Phys. Rev. C* **68**, 054005 (2003).
- [11] J. Giovinazzo *et al.*, *Phys. Rev. Lett.* **99**, 102501 (2007).
- [12] K. Miernik *et al.*, *Phys. Rev. Lett.* **99**, 192501 (2007).
- [13] K. Miernik, *Eur. Phys. J. A* **42**, 431 (2009).
- [14] C. Dossat *et al.*, *Nucl. Phys. A* **792**, 18 (2007).
- [15] J. Giovinazzo *et al.*, *Eur. Phys. J. A* **11**, 247 (2001).
- [16] B. Blank *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **613**, 65 (2010).
- [17] J. F. Ziegler, *Nucl. Instrum. Methods Phys. Res., Sect. B* **268**, 1818 (2010).
- [18] M. Honma, T. Otsuka, B. A. Brown, and T. Mizusaki, *Eur. Phys. J. A* **25**, Suppl. 1, 499 (2005).
- [19] L. Audirac *et al.* (to be published).