

Exact Exponents for the Spin Quantum Hall Transition in the Presence of Multiple Edge Channels

R. Bondesan,^{1,2,3} I. A. Gruzberg,^{3,4} J. L. Jacobsen,^{2,3,5} H. Obuse,^{3,6,7} and H. Saleur^{1,3,8}

¹*Institute de Physique Théorique, CEA Saclay, F-91191 Gif-sur-Yvette, France*

²*LPTENS, École Normale Supérieure, 24 rue Lhomond, 75231 Paris, France*

³*Institut Henri Poincaré, 11 rue Pierre et Marie Curie, 75231 Paris, France*

⁴*James Franck Institute, University of Chicago, 5640 South Ellis Avenue, Chicago, Illinois 60637, USA*

⁵*Université Pierre et Marie Curie, 4 place Jussieu, 75252 Paris, France*

⁶*Department of Physics, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan*

⁷*Institut für Nanotechnologie, Karlsruhe Institute of Technology, 76021 Karlsruhe, Germany*

⁸*Physics Department, USC, Los Angeles, California 90089-0484, USA*

(Received 22 September 2011; published 19 March 2012)

Critical properties of quantum Hall systems are affected by the presence of extra edge channels—those that are present, in particular, at higher plateau transitions. We study this phenomenon for the case of the spin quantum Hall transition. Using supersymmetry, we map the corresponding network model to a classical loop model, whose boundary critical behavior was recently determined exactly. We verify predictions of the exact solution by extensive numerical simulations.

DOI: [10.1103/PhysRevLett.108.126801](https://doi.org/10.1103/PhysRevLett.108.126801)

PACS numbers: 73.20.Fz, 72.15.Rn, 73.43.Nq

More than 50 years after its discovery, Anderson localization [1] remains a vibrant research field. One central research direction is the physics of Anderson transitions (ATs) [2], including metal-insulator and quantum Hall (QH) type transitions (that is, transitions between different phases of topological insulators). Apart from electronic gases in semiconductor structures, experimental realizations include localization of light [3], cold atoms [4], ultrasound [5], and optically driven atomic systems [6]. Theoretically, the field was strongly boosted by the discovery of unconventional symmetry classes and a complete symmetry classification of disordered systems [2,7–9].

Recently, it was realized that ATs in systems with boundaries may exhibit boundary critical behavior different from, and richer than, the bulk behavior [10–12]. The boundary criticality has served to test conformal invariance at two-dimensional ATs [13–15] and as a strong constraint on possible theories of the integer QH transition [16]. These works often employed the so-called network models [17,18] of ATs for numerical studies. Within a network formulation, the richness of boundary critical behaviors relates to the possibility of having multiple edge channels at the boundary [12,19,20]. Physically, multiple edge channels occur in an integer QH system whenever the filling fraction exceeds one. Some of the results below may directly apply to the physics of higher QH plateaus and transitions between them. However, our description neglects electron interactions at the edge which could be relevant in experimental realizations of the QH effect [21,22].

In this Letter, we study boundary critical properties in the presence of multiple edge states at the so-called spin quantum Hall (SQH) transition [23]. The corresponding network model [23,24] enjoys a very special status. In the bulk, or with reflecting boundaries, the model in its minimal formulation (suitable to describe mean conductances)

can be mapped to classical percolation on a square lattice [25,26]. This mapping determines exact critical properties at the SQH transition [25–29]. In this Letter, we demonstrate that extra edge channels can be straightforwardly included in the mapping. The resulting classical model is not percolation anymore but can nonetheless be formulated as a loop model.

Both network and loop models (or percolation) are lattice regularizations of field-theoretic descriptions of ATs in terms of sigma models on symmetric superspaces [2,8,30]. This connection is thoroughly explained in Ref. [31] and was recently extensively explored by some of us [32]. Through this connection, complete spectra of boundary operators were obtained for the conformal sigma models on superspaces $\mathbb{C}\mathbb{P}^{N+M-1|N}$ with a topological theta angle [33]. In the sigma-model approach, the number of extra edges is related to the exact value of the theta angle, which affects boundary (but not bulk) properties [34]. The case $N = M = 1$ is directly relevant to the SQH effect, and we here apply results of Ref. [33] to obtain exact exponents describing the scaling of the mean boundary point contact conductances in the presence of multiple edges.

We also report extensive numerical simulations of mean conductances in network models on open strips with edge channels on both sides. Conformal invariance (which has been numerically demonstrated for this transition [14]) relates the exponential decay of the mean conductance along the strip to dimensions of certain boundary operators. We extract these dimensions and compare them with the predictions of Ref. [33].

The network model for the SQH effect with extra edge channels is shown in Fig. 1. The bulk of width $2L$ contains alternating up- and down-going columns of links. In addition, m (n) extra columns with the *same* chirality are added




FIG. 1 (color online). Left: Network model with a bulk region of width $2L = 4$, and $m = n = 2$ extra channels at the left and right boundaries. Right: Schematic representation of the decomposition of average node transfer matrices. The two top rows show bulk nodes—see (6)—while the two bottom ones are boundary nodes—see (7).

at the left (right) edge. These extra links can be directed up or down at either edge. We label the four possible variants by $(\mathcal{L} = \pm m, \mathcal{R} = \pm n)$; positive labels $(\mathcal{L}, \mathcal{R})$ mean the same direction of the edge links and the closest bulk link. The links of the network carry doublets of complex fluxes (labeled \uparrow and \downarrow) whose scattering on links is described by matrices uniformly distributed over the $SU(2)$ group. The scattering at the two types of bulk nodes (labeled $S = A$ or B) is described by orthogonal matrices diagonal in spin indices:

$$\mathcal{S}_{S\uparrow} = \mathcal{S}_{S\downarrow} = \begin{pmatrix} (1 - t_S^2)^{1/2} & t_S \\ -t_S & (1 - t_S^2)^{1/2} \end{pmatrix}, \quad (1)$$

with t_S the strength of the quantum tunneling. The SQH transition occurs when $t_A = t_B$. The *boundary* nodes where the fluxes on the extra edge links scatter are described by matrices of the form (1) but with two independent transmission amplitudes, t_L and t_R , one for each edge of the system.

We employ the supersymmetry method for network models [25,35,36], with modifications due to the extra edge links. The row-to-row transfer matrices X and Y (formed by multiplying all node transfer matrices, T_A or T_B for bulk nodes, T_L and T_R for boundary nodes, in a given row) act in the tensor product of bosonic and fermionic Fock spaces defined for each column of links. The columns form sites (labeled i) of a one-dimensional quantum system whose evolution in the vertical (imaginary time t) direction is given by the operator $U = \prod_i^{L_T} (XY)$, where L_T is the number of A nodes (or B nodes) along the t direction (see Fig. 1). With periodic boundary conditions in the t direction, physical quantities, including conductance, may be written as correlation functions, $\langle \dots \rangle \equiv \text{STr}[\dots U]$, and the system is invariant under a global $sl(2|1)$ supersymmetry [25].

Averaging over disorder independently on each link (we denote such averages by overbars) projects the Fock space of bosons and fermions onto the fundamental V (dual-fundamental V^*) three-dimensional irreducible

representation of $sl(2|1)$ on up links (down links) [25]. The average node transfer matrices $\bar{T}_{S,i}$ act in the tensor products of *superspins* $V_i \otimes V_{i+1}^*$ in the bulk and $V_i \otimes V_{i+1}$ or $V_i^* \otimes V_{i+1}^*$ in the extra edge regions. Thus, we have four types of supersymmetric *spin chains*:

$$(m; n): V^{\otimes m} \otimes (V \otimes V^*)^{\otimes L} \otimes (V^*)^{\otimes n}, \quad (2)$$

$$(m; -n): V^{\otimes m} \otimes (V \otimes V^*)^{\otimes L} \otimes V^{\otimes n}, \quad (3)$$

$$(-m; n): (V^*)^{\otimes m} \otimes (V \otimes V^*)^{\otimes L} \otimes (V^*)^{\otimes n}, \quad (4)$$

$$(-m; -n): (V^*)^{\otimes m} \otimes (V \otimes V^*)^{\otimes L} \otimes V^{\otimes n}. \quad (5)$$

All tensor products between neighboring sites of these chains decompose into two $sl(2|1)$ irreducible representations. The averages $\bar{T}_{A,B}$ in the bulk and $\bar{T}_{L,R}$ at the boundaries read

$$\bar{T}_{A,i} = (1 - t_A^2)I + t_A^2 E_i, \quad A \leftrightarrow B, \quad (6)$$

$$\bar{T}_{L,i} = (1 - t_L^2)I + t_L^2 P_{i,i+1}, \quad L \leftrightarrow R, \quad (7)$$

where I is the identity, E_i projects onto the singlet in the decomposition of $V_i \otimes V_{i+1}^*$ (and $V_i^* \otimes V_{i+1}$), and $P_{i,i+1}$ is the graded permutation of states on sites i and $i+1$. The chain (2) with $m = n$ corresponds to a sigma model with topological angle $\Theta = (2n+1)\pi$ and to a transition between the n th and the $(n+1)$ st plateau.

The decompositions (6) and (7) have a natural graphical representation, shown schematically (without coefficients) in Fig. 1 (right). At a bulk node, links can be separated into disjoint lines in two ways, while, at a boundary node, the lines can either avoid each other or cross. Multiplying the transfer matrices to calculate the partition function, the result is the sum of all contributions of dense closed loops of weight one filling the links of the network, weighted by factors of either t_S^2 or $(1 - t_S^2)$ for each node. In the bulk, the loops are percolation hulls, and the loop model is equivalent to bond percolation on a square lattice [25]. The presence of extra edges generalizes this nontrivially, since loops can intersect at the boundary. In this situation, all configurations of the loop model fall into disjoint sectors [37] labeled by the number $\ell + 2k$ of through lines extending throughout the system in the t direction, where $\ell = |\mathcal{L} - \mathcal{R}|$ is the minimal possible number of the through lines for given \mathcal{L} and \mathcal{R} .

In the strip geometry, our model has external links at the top and bottom, where we can insert or extract current. In this geometry, the mean conductance $\bar{g}^{\mathcal{L},\mathcal{R}}$ between the top and bottom contacts is the average number of through lines going from the source to the drain of the current, times 2 for the spin. With the source at the bottom, the minimal number of available through lines is $k_{\min}^{\mathcal{L},\mathcal{R}} = \max(0, \mathcal{L} - \mathcal{R})$, whence

$$\bar{g}^{\mathcal{L},\mathcal{R}} = 2k_{\min}^{\mathcal{L},\mathcal{R}} + 2 \sum_{k=1}^{\infty} kP(k, L_T/L, L_T/\xi), \quad (8)$$

where $P(k, L_T/L, L_T/\xi)$ is the probability (symmetric in \mathcal{L} and \mathcal{R}) that exactly $2k$ “paired” through lines run through the system of size $2L_T$ by $2L + m + n$, and ξ is the bulk correlation length. At the transition, $\xi = \infty$, and, for large L_T/L , we expect that $P(k, L_T/L, 0) \sim e^{-\pi h^{\mathcal{L},\mathcal{R}}(k)L_T/L}$. Conformal invariance at the transition allows us to identify the exponents $h^{\mathcal{L},\mathcal{R}}(k)$ with dimensions of certain boundary operators. In the literature on self-avoiding walks, such exponents are called watermelon exponents and the through lines are called legs. For $L_T/L \gg 1$, the sum in Eq. (8) for \bar{g} is dominated by $k = 1$, and we denote the exponent of the first subleading correction as $\gamma^{\mathcal{L},\mathcal{R}}$:

$$\bar{g}^{\mathcal{L},\mathcal{R}} \sim 2k_{\min}^{\mathcal{L},\mathcal{R}} + C_1 e^{-\pi h^{\mathcal{L},\mathcal{R}}(1)L_T/L} + C_2 e^{-\pi \gamma^{\mathcal{L},\mathcal{R}}L_T/L}. \quad (9)$$

Critical properties of the above geometric loop model, including the exponents $h^{\mathcal{L},\mathcal{R}}(k)$, can be extracted from the anisotropic limit of the spin chains (2)–(5), obtained by taking all $t_S \ll 1$. In this limit, the evolution operator in one unit of time becomes $\overline{XY} \approx \exp(-t_A t_B H)$. For definiteness, we focus on the case (2); the critical ($t_A = t_B$) Hamiltonian H is then

$$H = -u \sum_{i=0}^{m-1} P_{i,i+1} - \sum_{i=m}^{2L+m-2} E_i - v \sum_{i=2L+m-1}^{2L+m+n-2} P_{i,i+1}. \quad (10)$$

The interaction between the superspins in the bulk is anti-ferromagnetic and uniform at the critical point. Interactions between boundary spins are ferromagnetic (compare with [36]), and their magnitudes $u = (t_L/t_A)^2$ and $v = (t_R/t_A)^2$ are kept as arbitrary *positive* numbers. From the diagonalization of H , described in [33] along with full details of the derivation of the results presented below, we obtain the critical exponents $h^{m,n}(k)$ as scaling limits of the lowest eigenvalues in a given sector. A more detailed knowledge of $\bar{g}^{\mathcal{L},\mathcal{R}}$ is needed for predicting $\gamma^{\mathcal{L},\mathcal{R}}$ in Eq. (9) analytically [38], but, in most cases, we can identify its numerical value as $h^{\mathcal{L},\mathcal{R}}(2)$.

The exact expressions for $h^{m,n}(k)$ derived in [33] appear in Table I. There, $h_{r,s} = [(3r - 2s)^2 - 1]/24$, $m = \min(m, n)$, and the parameter r_k is given in terms of m, n , and k for $m \geq n$ (if $m < n$, exchange m and n) by

$$r_k = \frac{6}{\pi} \arccos\left(\frac{\sqrt{3}}{2} \sqrt{\frac{(m+1+k)(n+1-k)}{(m+1)(n+1)}}\right). \quad (11)$$

A few remarks are in order. The exponents are independent of the boundary couplings u and v , implying a boundary renormalization group flow to the stable fixed point with infinite boundary couplings, similar to Ref. [39]. Moreover, since u and v are positive (ferromagnetic), randomness in u and v does not change the exponents. If

TABLE I. The watermelon exponents. k is the number of paired through lines on top of ℓ unpaired ones, and $m = \min(m, n)$. $h_{r,s}$ is the Kac table, and r_k is defined as in Eq. (11).

k	#(legs) = $\ell + 2k$	$h^{m,n}(k)$
0	ℓ	$h_{r_0, r_0} = 0$
1	$\ell + 2$	h_{r_1, r_1}
\vdots	\vdots	\vdots
m	$n + m$	h_{r_m, r_m}
$m + 1$	$n + m + 2$	$h_{1,3} = 1/3$
\vdots	\vdots	\vdots
$m + j$	$n + m + 2j$	$h_{1,1+2j} = j(2j - 1)/3$
\vdots	\vdots	\vdots

the number of legs is ℓ or $m + n + 2j$, with $j > 0$, the exponents equal those of critical percolation with, respectively, 0 or $2j$ hulls through the system (in particular, this is so when $m = 0$ or $n = 0$). However, when the number of legs is between $\ell + 2$ and $m + n$, exponents are highly nontrivial and, remarkably, they are irrational.

The exponents $h^{\mathcal{L},\mathcal{R}}(k)$ are symmetric in \mathcal{L} and \mathcal{R} by invariance of the spectrum under left-right reflection. Further exponent relations follow from symmetries of the critical spin chains (2)–(5). In each case, $V \otimes V^*$ (and $V^* \otimes V$) interacts through E_i , and $V \otimes V$ (and $V^* \otimes V^*$) through P_i , and the u and v can be set to one thanks to the universality. Then, the top-bottom reflection switching V and V^* induces a mapping between the chains (sometimes with different lengths of the bulk region): $(\mathcal{L}, \mathcal{R}) \leftrightarrow (-\mathcal{L} - 1, -\mathcal{R} - 1)$. This implies that

$$h^{\mathcal{L},\mathcal{R}}(k) = h^{-\mathcal{L}-1, -\mathcal{R}-1}(k). \quad (12)$$

In particular, this relates pairwise exponents for the chains (2) and (5), as well as for (3) and (4): $h^{-m, -n}(k) = h^{m-1, n-1}(k)$ and $h^{-m, n}(k) = h^{m-1, -n-1}(k)$. When the total number of legs in the model (5) equals $m + n + 2j$, with $j \geq 0$, the corresponding exponent $h^{m-1, n-1}(k) = h_{1,3+2j} = (j+1)(2j+1)/3$, since, in this case, we must write the total number of legs as $(m-1) + (n-1) + 2j + 2$.

Finally, consider the chain (3). Regarding the leftmost site on the right boundary as part of the bulk and noting that we have identical chiralities at the two boundaries (so that $\ell = m + n$), the $(m + n + 2j)$ -leg exponents are simply $h_{1,2+2j} = j(2j+1)/3$, $j = 0, 1, \dots$, independently of m and n . Results for (4) easily follow from (12): the $(m + n + 2j)$ -leg exponents are $h_{1,2+2j} = j(2j+1)/3$ in both models (3) and (4).

We now present extensive numerical simulations to verify our analytical predictions for the exponents $h^{\mathcal{L},\mathcal{R}}(1)$, the independence on $t_{L,R}$, the symmetry relation (12), and to determine the subleading exponent $\gamma^{\mathcal{L},\mathcal{R}}$. We numerically calculate the conductance of critical SQH

networks with extra channels in the strip geometry with length $2L_T$ and width $2L + m + n$. We parametrize the transmission amplitude in Eq. (1) as $t_A = \sin\theta$ and $t_B = \cos\theta$. In terms of θ , the SQH transition occurs at $\theta = \theta_c \equiv \pi/4$. Similarly, we write $t_{L,R} = \sin\theta_e$, where θ_e can be arbitrarily tuned or chosen randomly in $[0, \pi]$ independently for each boundary node.

The transmission matrix t for the SQH network model in the strip geometry is effectively calculated using the transfer matrix method [18]; the conductance is given by the Landauer formula $g = \text{Tr}tt^\dagger$. Ideally, the exponents $h^{\mathcal{L},\mathcal{R}}(1)$ and $\gamma^{\mathcal{L},\mathcal{R}}$ would be obtained by fitting the data to Eq. (9). However, finite-size effects hamper such an analysis in actual simulations unless $\theta_e = \theta_c$ or θ_e is random. Therefore, we apply finite-size-scaling analysis to systems of various widths [40]. We only simulate networks corresponding to chains (2), (4), and (5), with $\mathcal{L} \leq \mathcal{R}$, so that $k_{\min}^{\mathcal{L},\mathcal{R}} = 0$. Table II summarizes our numerical results and includes analytical predictions for comparison.

First, we focus on systems (2), with $\mathcal{L} = 0$ and $\mathcal{R} \geq 0$. As shown in Table II, numerically obtained $h^{0,\mathcal{R}}(1)$ agrees with analytical results and $\gamma^{0,\mathcal{R}}$ coincides with $h^{0,\mathcal{R}}(2)$ for various \mathcal{R} and θ_e . This suggests that the level-one descendant of $h^{0,\mathcal{R}}(1)$, $h^{0,\mathcal{R}}(1) + 1 = 4/3$, does not contribute to the conductance in these systems. In addition, $h^{0,1}(1) = 1/3$ and $\gamma^{0,1} = 2$ are verified even for random θ_e , confirming our expectation that randomness in the boundary couplings is irrelevant.

Next, we consider the cases $\mathcal{L}, \mathcal{R} > 0$, where $h^{\mathcal{L},\mathcal{R}}(1)$ is irrational. Since the L dependence is weak for $\theta_e = \theta_c$ and random θ_e , we extract several $h^{m,n}(k)$ for these θ_e without finite-size-scaling analysis. We confirm that the numerically obtained exponents $h^{m,n}(1)$ and $\gamma^{m,n}$ agree well with the analytical $h^{m,n}(1)$ and $h^{m,n}(2)$, respectively. Note that there are no analytically predicted exponents with values between $h^{m,n}(1)$ and $h^{m,n}(2)$.

Furthermore, we study the cases $\mathcal{L}, \mathcal{R} < 0$, whose exponents $h^{\mathcal{L},\mathcal{R}}(k)$ are related to those of systems with $\mathcal{L}, \mathcal{R} > 0$ by Eq. (12). Comparing results for $(-m, -n)$ and $(n-1, m-1)$ in Table II (and assuming the $\mathcal{L} \leftrightarrow \mathcal{R}$ symmetry), we see that Eq. (12) for $k = 1, 2$ is verified even if $\theta_e \neq \theta_c$. Finally, when $\mathcal{L} < 0$ and $\mathcal{R} \geq 0$ in Eq. (4), our numerics confirm $h^{\mathcal{L},\mathcal{R}}(1) = 1$ for various cases. Results for the subleading exponent $\gamma^{\mathcal{L},\mathcal{R}}$ are presently inconclusive because of large numerical errors.

Our comprehensive numerical simulations confirm the exact analytical predictions for the leading exponent $h^{\mathcal{L},\mathcal{R}}(1)$ and, in most cases, allow us to identify the first subleading exponent $\gamma^{\mathcal{L},\mathcal{R}}$ as $h^{\mathcal{L},\mathcal{R}}(2)$.

In conclusion, we have considered the SQH transition with extra edge channels. We have mapped the corresponding network model to a classical loop model, whose boundary critical exponents have recently been obtained exactly. Using the mapping, we obtain exact critical exponents at the SQH transition from the exponential decay of the mean

TABLE II. Numerically obtained exponents $h^{\mathcal{L},\mathcal{R}}(1)$ and $\gamma^{\mathcal{L},\mathcal{R}}$ and analytical predictions for $h^{\mathcal{L},\mathcal{R}}(1)$ and $h^{\mathcal{L},\mathcal{R}}(2)$ for various values of $(\mathcal{L}, \mathcal{R})$. ∞ means that the numerical exponents are obtained by finite-size scaling [40]. ‘‘Rand’’ and ‘‘No’’ indicate randomly distributed θ_e and no extra edge channels, respectively. ‘‘...’’ means that numerical estimates of these subleading exponents are unreliable.

\mathcal{L}, \mathcal{R}	Numerical simulations				Analytical	
	$2L$	θ_e	$h^{\mathcal{L},\mathcal{R}}(1)$	$\gamma^{\mathcal{L},\mathcal{R}}$	$h^{\mathcal{L},\mathcal{R}}(1)$	$h^{\mathcal{L},\mathcal{R}}(2)$
0, 0	∞	No	0.3333(12)	2.06(63)	1/3	2
0, 1	∞	θ_c	0.3330(7)	2.00(36)	1/3	2
0, 1	∞	$\pi/10$	0.3341(31)	2.09(47)	1/3	2
0, 1	∞	Rand	0.3332(10)	1.99(55)	1/3	2
0, 10	∞	θ_c	0.3325(24)	1.81(55)	1/3	2
0, 10	∞	$\pi/10$	0.3318(24)	2.03(43)	1/3	2
1, 1	∞	θ_c	0.03775(25)	0.333(5)	0.037 720	1/3
1, 1	∞	$\pi/10$	0.037 79(31)	0.319(13)	0.037 720	1/3
1, 1	∞	Rand	0.037 73(45)	0.343(10)	0.037 720	1/3
2, 2	50	θ_c	0.016 00(2)	0.0737(6)	0.015 906	0.0732
3, 3	50	θ_c	0.008 80(1)	0.0396(2)	0.008 797	0.0377
4, 4	50	θ_c	0.005 62(1)	0.0255(1)	0.005 587	0.0233
1, 2	∞	θ_c	0.0520(25)	0.334(43)	0.052 083	1/3
1, 2	∞	$\pi/10$	0.0517(10)	0.307(16)	0.052 083	1/3
1, 2	50	Rand	0.052 21(3)	0.338(8)	0.052 083	1/3
1, 3	50	θ_c	0.059 86(3)	0.339(9)	0.059 697	1/3
1, 4	50	θ_c	0.064 25(3)	0.34(1)	0.064 421	1/3
2, 3	50	θ_c	0.024 49(10)	0.088(1)	0.024 348	0.0847
2, 3	∞	$\pi/10$	0.024 76(42)	0.086(3)	0.024 348	0.0847
2, 4	50	θ_c	0.029 54(7)	0.095(1)	0.029 589	0.0920
-2, -2	∞	$\pi/3$	0.0377(4)	0.339(10)	0.037 720	1/3
-2, -2	∞	$\pi/10$	0.0372(8)	0.281(4)	0.037 720	1/3
-3, -3	∞	$\pi/3$	0.016 11(9)	0.0749(7)	0.015 906	0.0732
-3, -3	∞	$\pi/10$	0.0160(9)	0.0730(30)	0.015 906	0.0732
-3, -2	∞	$\pi/3$	0.0522(2)	0.333(13)	0.052 083	1/3
-3, -2	∞	$\pi/10$	0.0549(30)	0.336(13)	0.052 083	1/3
-1, 0	∞	θ_c	0.999(9)	...	1	10/3
-2, 0	∞	θ_c	0.999(3)	...	1	10/3
-2, 1	∞	θ_c	0.998(3)	...	1	10/3
-2, 2	∞	θ_c	0.993(1)	...	1	10/3

conductance in the strip geometry. Our extensive numerical simulations confirm the analytical results for the boundary exponents. The demonstrated influence of extra edge channels on boundary critical behavior should be broadly applicable to other QH transitions.

The numerical simulations have been performed using the PADS resource (NSF Grant No. OCI-0821678) and the Teraport Cluster at the Computational Institute at the University of Chicago. The work of J. L. J. and H. S. was supported by a grant from the ANR Projet 2010 Blanc SIMI 4: DIME. H. O. is supported by a Grant-in-Aid from JSPS for Young Scientists. I. A. G. was supported by NSF Grants No. DMR-0448820 and No. DMR-0213745. H. S. thanks C. Candu, N. Read, and V. Schomerus for discussions. H. O. thanks F. Evers for discussions.

- [1] *50 Years of Anderson Localization*, edited by E. Abrahams (World Scientific, Singapore, 2010); E. Abrahams, *Int. J. Mod. Phys. B* **24**, 4150 (2010).
- [2] F. Evers and A. D. Mirlin, *Rev. Mod. Phys.* **80**, 1355 (2008).
- [3] D. S. Wiersma, P. Bartolini, A. Lagendijk, and R. Righini, *Nature (London)* **390**, 671 (1997).
- [4] J. Billy, V. Josse, Z. Zuo, A. Bernard, B. Hambrecht, P. Lugan, D. Clément, L. Sanchez-Palencia, P. Bouyer, and A. Aspect, *Nature (London)* **453**, 891 (2008); G. Roati, C. D'Errico, L. Fattori, C. Fort, M. Zaccanti, G. Modugno, M. Modugno, and M. Inguscio, *ibid.* **453**, 895 (2008).
- [5] S. Faez, A. Strybulevych, J. H. Page, A. Lagendijk, and B. A. van Tiggelen, *Phys. Rev. Lett.* **103**, 155703 (2009).
- [6] G. Lemarié, H. Lignier, D. Delande, P. Szriftgiser, and J. C. Garreau, *Phys. Rev. Lett.* **105**, 090601 (2010).
- [7] A. Altland and M. R. Zirnbauer, *Phys. Rev. B* **55**, 1142 (1997).
- [8] M. R. Zirnbauer, *J. Math. Phys. (N.Y.)* **37**, 4986 (1996).
- [9] P. Heinzner, A. Huckleberry, and M. R. Zirnbauer, *Commun. Math. Phys.* **257**, 725 (2005).
- [10] A. R. Subramaniam, I. A. Gruzberg, A. W. W. Ludwig, F. Evers, A. Mildenberger, and A. D. Mirlin, *Phys. Rev. Lett.* **96**, 126802 (2006).
- [11] A. Mildenberger, A. R. Subramaniam, R. Narayanan, F. Evers, I. A. Gruzberg, and A. D. Mirlin, *Phys. Rev. B* **75**, 094204 (2007).
- [12] H. Obuse, A. Furusaki, S. Ryu, and C. Mudry, *Phys. Rev. B* **78**, 115301 (2008).
- [13] H. Obuse, A. R. Subramaniam, A. Furusaki, I. A. Gruzberg, and A. W. W. Ludwig, *Phys. Rev. Lett.* **98**, 156802 (2007); *Physica (Amsterdam)* **40E**, 1404 (2008).
- [14] H. Obuse, A. R. Subramaniam, A. Furusaki, I. A. Gruzberg, and A. W. W. Ludwig, *Phys. Rev. B* **82**, 035309 (2010).
- [15] S. Ryu, C. Mudry, H. Obuse, and A. Furusaki, *New J. Phys.* **12**, 065005 (2010).
- [16] H. Obuse, A. R. Subramaniam, A. Furusaki, I. A. Gruzberg, and A. W. W. Ludwig, *Phys. Rev. Lett.* **101**, 116802 (2008); F. Evers, A. Mildenberger, and A. D. Mirlin, *ibid.* **101**, 116803 (2008).
- [17] J. T. Chalker and P. D. Coddington, *J. Phys. C* **21**, 2665 (1988).
- [18] B. Kramer, T. Ohtsuki, and S. Kettemann, *Phys. Rep.* **417**, 211 (2005).
- [19] C. Barnes, B. L. Johnson, and G. Kirczenow, *Phys. Rev. Lett.* **70**, 1159 (1993).
- [20] K. Kobayashi, T. Ohtsuki, and K. Slevin, *J. Phys. Soc. Jpn.* **78**, 084708 (2009).
- [21] D. B. Chklovskii, B. I. Shklovskii, and L. I. Glazman, *Phys. Rev. B* **46**, 4026 (1992).
- [22] P. G. Silvestrov and K. B. Efetov, *Phys. Rev. B* **77**, 155436 (2008).
- [23] T. Senthil, J. B. Marston, and M. P. A. Fisher, *Phys. Rev. B* **60**, 4245 (1999).
- [24] V. Kagalovsky, B. Horovitz, Y. Avishai, and J. T. Chalker, *Phys. Rev. Lett.* **82**, 3516 (1999).
- [25] I. A. Gruzberg, A. W. W. Ludwig, and N. Read, *Phys. Rev. Lett.* **82**, 4524 (1999).
- [26] A. R. Subramaniam, I. A. Gruzberg, and A. W. W. Ludwig, *Phys. Rev. B* **78**, 245105 (2008).
- [27] J. Cardy, *Phys. Rev. Lett.* **84**, 3507 (2000).
- [28] E. J. Beamond, J. Cardy, and J. T. Chalker, *Phys. Rev. B* **65**, 214301 (2002).
- [29] A. D. Mirlin, F. Evers, and A. Mildenberger, *J. Phys. A* **36**, 3255 (2003).
- [30] K. B. Efetov, *Supersymmetry in Disorder and Chaos* (Cambridge University Press, Cambridge, England, 1987); A. D. Mirlin, *Phys. Rep.* **326**, 259 (2000); M. R. Zirnbauer, [arXiv:math-ph/0404058](https://arxiv.org/abs/math-ph/0404058).
- [31] N. Read and H. Saleur, *Nucl. Phys.* **B613**, 409 (2001).
- [32] N. Read and H. Saleur, *Nucl. Phys.* **B777**, 263 (2007); C. Candu and H. Saleur, *ibid.* **B808**, 441 (2009); C. Candu, J. L. Jacobsen, N. Read, and H. Saleur, *J. Phys. A* **43**, 142001 (2010); C. Candu, V. Mitev, T. Quella, H. Saleur, and V. Schomerus, *J. High Energy Phys.* **02** (2010) 015.
- [33] R. Bondesan, J. L. Jacobsen, and H. Saleur, *Nucl. Phys.* **B849**, 461 (2011).
- [34] S. Xiong, N. Read and A. D. Stone, *Phys. Rev. B* **56**, 3982 (1997).
- [35] N. Read (unpublished).
- [36] I. A. Gruzberg, N. Read, and S. Sachdev, *Phys. Rev. B* **55**, 10593 (1997).
- [37] P. Martin, *Potts Models and Related Problems in Statistical Mechanics* (World Scientific, Singapore, 1991).
- [38] Conformal invariance implies that the exponential corrections to the conductance are given by conformal dimensions in the theory. $\gamma^{\mathcal{L},\mathcal{R}}$ is then the minimum among $h^{\mathcal{L},\mathcal{R}}(2)$ and dimensions of certain primary and descendant fields in the sector $k = 1$. Unfortunately, the analysis of [33] is not sufficient to identify all the fields contributing to $\bar{g}^{\mathcal{L},\mathcal{R}}$ in the general case.
- [39] I. Affleck and A. W. W. Ludwig, *Phys. Rev. Lett.* **67**, 161 (1991).
- [40] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.108.126801> for details of numerical simulations..