

Anomalous Behavior of Spin Systems with Dipolar Interactions

D. Peter,¹ S. Müller,² S. Wessel,³ and H. P. Büchler¹

¹*Institute for Theoretical Physics III, University of Stuttgart, Stuttgart, Germany*

²*Max Planck Institute for Physics, 80805 Munich, Germany*

³*Institute for Theoretical Solid State Physics, JARA-FIT, and JARA-HPC, RWTH Aachen University, Otto-Blumenthal Strasse 26, D-52056 Aachen, Germany*

(Received 7 March 2012; published 11 July 2012)

We study the properties of spin systems realized by cold polar molecules interacting via dipole-dipole interactions in two dimensions. Using a spin wave theory, that allows for the full treatment of the characteristic long-distance tail of the dipolar interaction, we find several anomalous features in the ground state correlations and the spin wave excitation spectrum, which are absent in their counterparts with short-range interaction. The most striking consequence is the existence of true long-range order at finite temperature for a two-dimensional phase with a broken $U(1)$ symmetry.

DOI: [10.1103/PhysRevLett.109.025303](https://doi.org/10.1103/PhysRevLett.109.025303)

PACS numbers: 67.85.-d, 05.30.Jp, 75.10.Jm, 75.30.Ds

The foundation for understanding the behavior and properties of quantum matter is based on models with short-range interactions. Experimental progress in cooling polar molecules [1] and atomic gases with large magnetic dipole moments [2] has however increased the interest in systems with strong dipole-dipole interactions. While many properties of quantum systems with dipole-dipole interactions derive from our understanding of systems with short-range interactions, the dipole-dipole interaction can give rise to phenomena not present in their short-range counterparts. Prominent examples are the description of dipolar Bose–Einstein condensates, where the contribution of the dipolar interaction cannot be included in the s -wave scattering length [3], and the absence of a first order phase transition with a jump in the density [4]. In this Letter, we demonstrate anomalous behavior in two-dimensional spin systems with dipolar interactions realized by polar molecules in optical lattices.

A remarkable property of cold polar molecules confined into two dimensions is the potential formation of a crystalline phase for strong dipole-dipole interactions [5,6]. In contrast to a Wigner crystal with Coulomb interactions [7], the crystalline phase exhibits the conventional behavior expected for a crystal realized with a short-range repulsion and the characteristic $1/r^3$ behavior of the dipole interaction can be truncated at distances involving several interparticle separations. Several strongly correlated phases have been predicted, which behave in analogy to systems with interactions extending over a finite range, such as a Haldane phase [8], supersolids [9,10], pair supersolids in bilayer systems [11], valence bond solids [12], as well as p -wave superfluidity [13], and self-assembled structures in multilayer setups [14]. On the other hand, it has recently been demonstrated that polar molecules in optical lattices are also suitable for emulating quantum phases of two-dimensional spin models [15–17].

Here, we demonstrate that such spin models with dipole-dipole interactions exhibit several anomalous

features, which are not present in their short-range counterparts. The analysis is based on analytical spin wave theory, which allows for the full treatment of the $1/r^3$ tail of the dipole-dipole interactions. We find that the excitation spectrum exhibits anomalous behavior at low momenta, which gives rise to unconventional dynamic properties of the spin wave excitations. Remarkably, we derive from this anomalous behavior the existence of a long-range ordered ferromagnetic phase at finite temperatures; this finding is consistent with the well-known Mermin–Wagner theorem as the latter does not exclude order for interactions with a $1/r^\alpha$ tail, where $\alpha \leq 4$ [18–20]. Finally, we show that the dipole-dipole interaction gives rise to algebraic correlations even in gapped ground states, in agreement with recent predictions [21,22].

We focus on a set up of polar molecules confined into two dimensions in a square lattice, with each lattice site filled by one polar molecule. A static electric field applied along the z direction splits the rotation levels, and allows us to define a spin $1/2$ system by selecting two states in the rotational manifold. Then, the Hamiltonian reduces to an XXZ model with dipole-dipole interaction between the spins [16]

$$H = \frac{Ja^3}{\hbar^2} \sum_{i \neq j} \frac{\cos\theta S_i^z S_j^z + \sin\theta (S_i^x S_j^x + S_i^y S_j^y)}{|\mathbf{R}_i - \mathbf{R}_j|^3}. \quad (1)$$

Here, the first term accounts for the static dipole-dipole interaction between the different rotational levels with strength $J \cos\theta$, while the last term describes the virtual exchange of a microwave photon between the two polar molecules with strength $J \sin\theta$, and a denotes the lattice spacing. The dependence of the couplings J and θ on the microscopic parameters is discussed in Refs. [16,17,23] and the one-dimensional version of this model has recently been studied in Ref. [24].

Before analyzing this spin model on the square lattice, we present a summary of the phase diagram for its counterpart

with nearest neighbor interactions only. Then, the phase diagram is highly symmetric and exhibits four different phases: (i) an Ising antiferromagnetic phase (I-AF) for $-\pi/4 < \theta < \pi/4$ with an excitation gap, (ii) an XY antiferromagnetic phase (XY-AF) for $\pi/4 < \theta < 3\pi/4$ with a linear excitation spectrum, (iii) an Ising ferromagnetic phase (I-F) for $3\pi/4 < \theta < 5\pi/4$ with an excitation gap, and finally (iv) an XY ferromagnetic phase (XY-F) for $5\pi/4 < \theta < 7\pi/4$ with a linear spectrum.

Next, we analyze the modifications of the phase diagram due to dipole-dipole interactions between the spins within mean-field theory. The main influence is the reduction of the stability for the antiferromagnetic phases, as the next nearest neighbor interaction introduces a weak frustration to the system. The ground state energy per lattice site within mean field reduces to $e_{\text{I-AF}} = J \cos \theta \epsilon_{\mathbf{K}}/4$ and $e_{\text{XY-AF}} = J \sin \theta \epsilon_{\mathbf{K}}/4$ for the antiferromagnetic phases. The summation over the dipole interaction reduces to a dimensionless parameter $\epsilon_{\mathbf{K}} \approx -2.646$, which is related to the dipolar dispersion

$$\epsilon_{\mathbf{q}} = \sum_{j \neq 0} e^{i\mathbf{R}_j \mathbf{q}} \frac{a^3}{|\mathbf{R}_j|^3}, \quad (2)$$

at the corner of the Brillouin zone $\mathbf{K} = (\pi/a, \pi/a)$. In turn, the ferromagnetic phases are enhanced with a mean-field energy $e_{\text{I-F}} = J \cos \theta \epsilon_0/4$ and $e_{\text{XY-F}} = J \sin \theta \epsilon_0/4$ with $\epsilon_0 \approx 9.033$. The modifications to the phase diagram are shown in Fig. 1: first, the Heisenberg points at $\theta = \pi/4, 5\pi/4$ are protected by the SU(2) symmetry and still provide the transition between the Ising and the XY phases. However, the transitions from the ferromagnetic towards the antiferromagnetic phase are shifted to the values $\theta_c = \arctan(\epsilon_{\mathbf{K}}/\epsilon_0) \approx -0.1\pi$ and $\tilde{\theta}_c = \pi + \arctan(\epsilon_0/\epsilon_{\mathbf{K}}) \approx 0.6\pi$.

The dipole dispersion $\epsilon_{\mathbf{q}}$ in Eq. (2) converges very slowly due to the characteristic power law decay of the dipole-dipole interaction. It is this slow decay, which will




FIG. 1 (color online). (a) Mean-field phase diagram for the XXZ model with dipolar interactions, where $\tan \theta$ is the ratio between the XY and the Ising spin couplings. (b) Ground state energy per particle: the dashed lines show the mean-field predictions, while the solid lines include the contributions from the spin waves. At the critical values θ_c and $\tilde{\theta}_c$, the ground state energy exhibits the jump $\Delta e_c \approx 0.14J$ and $\Delta \tilde{e}_c \approx 0.06J$, indicating the potential formation of an intermediate phase.

give rise to several peculiar properties of the system. Therefore, we continue first with a detailed discussion of this dipolar dispersion. The precise determination of $\epsilon_{\mathbf{q}}$ is most conveniently performed using an Ewald summation [7], which transforms the summation over the slowly converging terms with algebraic decay into a summation of exponential factors, i.e.,

$$\begin{aligned} \epsilon_{\mathbf{q}} = & -2\pi a |\mathbf{q}| \operatorname{erfc}(a|\mathbf{q}|/2\sqrt{\pi}) + 4\pi \left(e^{-a^2|\mathbf{q}|^2/4\pi} - \frac{1}{3} \right) \\ & + 2\pi \sum_{i \neq 0} \int_1^\infty \frac{d\lambda}{\lambda^{3/2}} [e^{-\pi\lambda(\mathbf{R}_i/a + a\mathbf{q}/2\pi)^2} \\ & + \lambda^2 e^{-\pi\lambda|\mathbf{R}_i|^2/a^2 + i\mathbf{R}_i \mathbf{q}}], \end{aligned} \quad (3)$$

with $\operatorname{erfc}(x)$ the complementary error function. The important feature of the dipole dispersion is captured by the first term in Eq. (3), which gives rise to a linear and nonanalytic behavior $\epsilon_{\mathbf{q}} \sim \epsilon_0 - 2\pi a |\mathbf{q}|$ for small values $q \ll 1/a$, while all remaining terms are analytic. It is this linear part, which will give rise to several unconventional properties of spin systems in 2D with dipolar interactions, and is a consequence of the slow decay of the dipole-dipole interaction. The summation in the last term converges very quickly and guarantees the periodicity of the dipolar dispersion. The quantitative behavior is shown in Fig. 2(a), and the numerical efficient determination provides $\epsilon_0 \approx 9.033$, and $\epsilon_{\mathbf{K}} = (1/\sqrt{2} - 1)\epsilon_0 \approx -2.646$. Note, that a similar linear dependence appears for continuous systems with dipolar interactions in two dimensions, and gives rise to a so-called anomalous scattering amplitude [13].

Next, we analyze the excitation spectrum above the mean-field ground states within a spin wave analysis. The spin wave analysis is well established [25,26], and its application for a spin system with dipolar interaction is straightforward. Details of the calculation are presented in the Supplemental Material [27] for the XY-AF model. The results are summarized in Table I and shown in Fig. 2.




FIG. 2 (color online). Spin wave excitations with $\Gamma = (0, 0)$, $M = (0, \pi/2)$, and $K = (\pi/a, \pi/a)$ for different θ angles. (a) Spectrum of the I-F phase which also shows the behavior of the dipolar dispersion $\epsilon_{\mathbf{q}}$ for $\theta = -3\pi/4$. (b)–(d) Spectrum for the XY-F, I-AF, and XY-AF phases.

In the following, we present a detailed discussion for each of the four ordered phases.

Ising ferromagnetic phase.—The ferromagnetic mean-field ground state is twofold degenerate with all spins either point up or down, and is the exact ground state for $\theta = \pi$, i.e., $|G\rangle = \prod_i |\downarrow\rangle_i$. Within the spin wave analysis, the ground state is not modified and the excitation spectrum reduces to $E_{\mathbf{q}}^{\text{I-F}}$ (see Table I). The spin waves exhibit an excitation gap Δ : (i) approaching the Heisenberg point at $\theta = -3\pi/4$, the excitation gap vanishes, indicating the instability towards the XY ferromagnet, (ii) in turn, for antiferromagnetic XY couplings, the gap is minimal at \mathbf{K} , vanishes at the mean-field transition point $\tilde{\theta}_c$, and drives an instability towards the formation of antiferromagnetic ordering.

In contrast to any short-range ferromagnetic spin model, the dispersion relation $E_{\mathbf{q}}^{\text{I-F}}$ is not quadratic for small momenta, but rather exhibits a linear behavior, i.e., $E_{\mathbf{q}}^{\text{I-F}} \sim E_0^{\text{I-F}} + \hbar c|\mathbf{q}|$ with velocity $c = -2\pi aJ \sin\theta/\hbar$, which is a consequence of the dipolar interaction in the system. This anomalous behavior strongly influences the dynamics of the spin waves. The dynamical behavior of a single localized spin excitation is shown in Fig. 3(a) for a Gaussian initial state. In order to probe the linear part in the dispersion relation, the width σ of the localization is much larger than the lattice spacing a , and therefore, the dynamics is well described by a continuum description. Instead of the conventional quantum mechanical spreading, one finds a ballistic expansion of a cylindrical wave packet with velocity c . In addition, the dipole-dipole interaction also strongly influences the correlation function. Within conventional perturbation theory, we find algebraic correlations $\langle S_i^x S_j^x \rangle \sim 1/|\mathbf{r}|^3$. This algebraic decay of correlations even in gapped systems is a peculiar property of spin models with long-range interactions [21,22].

XY ferromagnetic phase.—Here, the spins are aligned in the xy plane. Within the spin wave analysis, we obtain the excitation spectrum $E_{\mathbf{q}}^{\text{XY-F}}$ and the modified ground state energy $e_{\text{XY-F}}$. In the low momentum regime, the dispersion relation behaves as $E_{\mathbf{q}}^{\text{XY-F}} \sim \sqrt{|\mathbf{q}|}$, in contrast to the well-known linear Goldstone modes for the broken $U(1)$ symmetry. This anomalous behavior is a peculiar property of the dipolar interaction, and the most crucial consequence




FIG. 3 (color online). Time evolution for localized spin excitations described by the Gaussian wave packet $\psi_0(\mathbf{r}) = e^{-|\mathbf{r}|^2/2\sigma^2}/\sqrt{\pi\sigma^2}$ with $\sigma \gg a$ in the continuum description. (a) For a linear dispersion $c|\mathbf{q}|$ in the I-F phase, the dynamics is described by cylindrical symmetric wave packets (see inset) traveling with velocity c , instead of the conventional quantum mechanical spreading for massive systems. (b) For an anomalous dispersion with $\alpha\sqrt{|\mathbf{q}|}$ in the XY-F phase, the behavior at long times $t \gg \sqrt{\sigma}\alpha$ reduces to a scaling function $\xi(z)$ via $|\psi(x, \tau)|^2 = \xi(x/\tau - 1/2)/\tau^2$ (see inset) using rescaled time $\tau = t\alpha/\sqrt{\sigma}$ and space $x = |\mathbf{r}|/\sigma$ coordinates. It describes a cylindrical symmetric wave front with velocity $\alpha\sqrt{\sigma}$.

is the existence of long-range order for the continuous broken symmetry at finite temperatures even in two dimensions [19]. This property follows immediately from the above spin wave analysis: the order parameter reduces to $m \equiv \Delta m - 1/2 = \langle S_i^x \rangle/\hbar$, where Δm accounts for the suppression of the order parameter by quantum fluctuations. Within spin wave theory, it reduces to $(\Delta m = \langle a_i^\dagger a_i \rangle)$

$$\Delta m = \int \frac{d\mathbf{q}}{v_0} \left[\frac{\cos\theta\epsilon_{\mathbf{q}} + \sin\theta(\epsilon_{\mathbf{q}} - 2\epsilon_0)}{4E_{\mathbf{q}}} \coth\left(\frac{E_{\mathbf{q}}}{2T}\right) - \frac{1}{2} \right].$$

This expression is finite and small: at $T = 0$, the integrand behaves as $\sim 1/\sqrt{|\mathbf{q}|}$ and we find a suppression of the order $\Delta m \approx 0.008$ at $\theta = -\pi/2$. The smallness of this corrections due to quantum fluctuations is a good justification for the validity of the spin wave analysis. On the other hand, even at finite temperatures, the low momentum behavior of the integrand takes the form $\sim T/|\mathbf{q}|$, and provides a finite contribution in contrast to a conventional Goldstone mode, which provides a logarithmic divergence.

The appearance of a long-range order at a finite temperature for a ground state with a broken $U(1)$ symmetry is a peculiar feature of dipole-dipole interactions, which renders the system more mean-field-like. The system

TABLE I. Spin wave excitation spectrum $E_{\mathbf{q}}^\alpha$ and ground state energy e_α .

Ground state α	Spin wave excitation spectrum $E_{\mathbf{q}}^\alpha$	Ground state energy per spin e_α
Ising ferromagnetic phase	$J(\sin\theta\epsilon_{\mathbf{q}} - \cos\theta\epsilon_0)$	$\frac{3J\cos\theta\epsilon_0}{4} + \frac{1}{2} \int \frac{d\mathbf{q}}{v_0} E_{\mathbf{q}}^\alpha(\mathbf{q}) = \frac{J\cos\theta\epsilon_0}{4}$
XY ferromagnetic phase	$J\sqrt{\sin\theta(\epsilon_{\mathbf{q}} - \epsilon_0)(\cos\theta\epsilon_{\mathbf{q}} - \sin\theta\epsilon_0)}$	$\frac{3J\sin\theta\epsilon_0}{4} + \frac{1}{2} \int \frac{d\mathbf{q}}{v_0} E_{\mathbf{q}}^\alpha$
Ising antiferromagnetic phase	$J\sqrt{(\sin\theta\epsilon_{\mathbf{q}+\mathbf{K}} - \cos\theta\epsilon_{\mathbf{K}})(\sin\theta\epsilon_{\mathbf{q}} - \cos\theta\epsilon_{\mathbf{K}})}$	$\frac{3J\cos\theta\epsilon_{\mathbf{K}}}{4} + \frac{1}{2} \int \frac{d\mathbf{q}}{v_0} E_{\mathbf{q}}^\alpha$
XY antiferromagnetic phase	$J\sqrt{\sin\theta(\epsilon_{\mathbf{q}+\mathbf{K}} - \epsilon_{\mathbf{K}})(\cos\theta\epsilon_{\mathbf{q}} - \sin\theta\epsilon_{\mathbf{K}})}$	$\frac{3J\sin\theta\epsilon_{\mathbf{K}}}{4} + \frac{1}{2} \int \frac{d\mathbf{q}}{v_0} E_{\mathbf{q}}^\alpha$

TABLE II. Correlation functions in the XY -F phase predicted by the spin wave analysis and high temperature expansion.

Correlation function	$T = 0$	$0 < T < T_c$	$T_c < T$
$\langle S_i^z S_j^z \rangle$	$\sim \mathbf{r} ^{-5/2}$	$\sim \mathbf{r} ^{-3}$	$\sim \mathbf{r} ^{-3}$
$\langle S_i^y S_j^y + S_i^x S_j^x \rangle - m^2$	$\sim \mathbf{r} ^{-3/2}$	$\sim \mathbf{r} ^{-1}$	$\sim \mathbf{r} ^{-3}$

therefore exhibits a finite temperature transition at a critical temperature T_c into a disordered phase; such a behavior is consistent with the classical XY model with dipolar interactions [19]. The correlation functions determined within spin wave theory and a high temperature expansion are summarized in Table II. Note that the spin wave analysis neglects the influence of vortices. This is well justified here, as the dipolar interactions give rise to a confining of vortices; i.e., the interaction potential between a vortex–antivortex pair increases linearly with the separation between the vortices.

The spin wave dynamics caused by the anomalous dispersion relation $\sim \sqrt{|\mathbf{q}|}$ is shown in Fig. 3(b) for a Gaussian wave packet of width σ . Interestingly, the propagation velocity of the wave packets is proportional to $\sqrt{\sigma}$ and thus faster for broad wave packets, in contrast to the usual dispersion dynamics. This is a consequence of the group velocity $v_{\mathbf{q}} \sim 1/\sqrt{|\mathbf{q}|}$ which is large for the small momentum components involved in the broad wave packets.

Ising antiferromagnetic phase.—Next, we focus on the antiferromagnetic phases and start with the I-AF ground state. Again, the ground state is twofold degenerate on bipartite lattices. We choose the ground state with spin up on sublattice A and spin down on sublattice B , i.e., $|G\rangle = \prod_{i \in A} |\uparrow\rangle_i \prod_{j \in B} |\downarrow\rangle_j$. The spin wave analysis is straightforward (see Supplemental Material [27]), and we obtain the spin wave excitation spectrum $E_{\mathbf{q}}^{\text{I-AF}}$ and ground state energy I-AF (see Table I). The system exhibits an excitation gap as expected for a system with a broken Z_2 symmetry. However, the dipole interactions give rise to an anomalous behavior at small momenta similar to the I-F phase with $E_{\mathbf{q}}^{\text{I-AF}} - E_0^{\text{I-AF}} \sim -\sin\theta|\mathbf{q}|$. Consequently, the dynamics of spin waves at low momenta is analogous to the Ising ferromagnet (see Fig. 3). Within spin wave theory, we obtain that the antiferromagnetic correlations $\langle (-1)^{i-j} S_i^\beta S_j^\beta \rangle$ and the ferromagnetic correlations $\langle S_i^\beta S_j^\beta \rangle$ decay with the power law $\sim 1/|\mathbf{r}|^3$ with $\beta = x, y, z$ determined by the characteristic behavior of the dipole-dipole interaction. The excitation gap vanishes approaching the mean-field critical point θ_c towards the XY -F phase, and also while approaching the Heisenberg point at $\theta = \pi/4$. For the latter, the qualitative behavior of the excitation spectrum changes drastically within a very narrow range of θ [see Fig. 2(c)].

XY antiferromagnetic phase.—Finally, we analyze the properties of the XY -AF phase. In contrast to the XY -F phase, the excitation spectrum $E_{\mathbf{q}}^{\text{XY-AF}}$ exhibits the conventional linear Goldstone mode [see Fig. 2(d)]. This can be

understood, as the antiferromagnetic ordering introduces a cancellation of the dipolar interactions, and provides a behavior in analogy to its short-range counterpart: true long-range order exists only at $T = 0$, while at finite temperature the system exhibits quasi long-range order and eventually undergoes a Kosterlitz–Thouless transition for increasing temperature. Nevertheless, the dipole-dipole interactions give rise to the characteristic algebraic correlations, e.g., $\langle (-1)^{i-j} S_i^z S_j^z \rangle \sim 1/|\mathbf{r}|^3$ for the antiferromagnetic transverse spin correlation at zero temperature.

Finally, we comment on the transitions between the different phases. The spin wave analysis predicts that the excitation spectrum for each phase becomes unstable at the mean-field critical points: For the Heisenberg points at $\theta = \pi/4, 5\pi/4$, such a behavior is expected due to the enhanced symmetry and one indeed finds that at the critical point, the excitation spectrum from the Ising phase coincides with the spectrum from the XY ground state. Consequently, the spin waves provide the same contribution to the ground state energy [see Fig. 1(b)]. In turn, at the instability points θ_c and $\tilde{\theta}_c$, the excitation spectrum from the antiferromagnetic phase is different from the spectrum for the ferromagnetic F phase. Consequently, the ground state energy within the spin wave analysis exhibits a jump [see Fig. 1(a)]. Such a behavior is an indication for the appearance of an intermediate phase. However, this question cannot be conclusively answered within the presented analysis due to the limited validity of spin wave theory close to the transition points. However, the appearance of a first order phase transition can be excluded by arguments similar to Ref. [4].

We thank M. Hermele for helpful discussions. Support by the Deutsche Forschungsgemeinschaft (DFG) within Contract No. SFB/TRR 21 and National Science Foundation under Grant No. NSF PHY05-51164 is acknowledged.

-
- [1] K.-K. Ni *et al.*, *Science* **322**, 231 (2008).
 - [2] A. Griesmaier, J. Werner, S. Hensler, J. Stuhler, and T. Pfau, *Phys. Rev. Lett.* **94**, 160401 (2005).
 - [3] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein, and T. Pfau, *Rep. Prog. Phys.* **72**, 126401 (2009).
 - [4] B. Spivak and S. A. Kivelson, *Phys. Rev. B* **70**, 155114 (2004).
 - [5] H. P. Büchler, E. Demler, M. Lukin, A. Micheli, N. Prokof'ev, G. Pupillo, and P. Zoller, *Phys. Rev. Lett.* **98**, 060404 (2007).
 - [6] G. E. Astrakharchik, J. Boronat, I. L. Kurbakov, and Y. E. Lozovik, *Phys. Rev. Lett.* **98**, 060405 (2007).
 - [7] L. Bonsall and A. A. Maradudin, *Phys. Rev. B* **15**, 1959 (1977).
 - [8] E. G. Dalla Torre, E. Berg, and E. Altman, *Phys. Rev. Lett.* **97**, 260401 (2006).
 - [9] L. Pollet, J. D. Picon, H. P. Büchler, and M. Troyer, *Phys. Rev. Lett.* **104**, 125302 (2010).

- [10] B. Capogrosso-Sansone, C. Trefzger, M. Lewenstein, P. Zoller, and G. Pupillo, *Phys. Rev. Lett.* **104**, 125301 (2010).
- [11] C. Trefzger, C. Menotti, and M. Lewenstein, *Phys. Rev. Lett.* **103**, 035304 (2009).
- [12] L. Bonnes, H. P. Büchler, and S. Wessel, *New J. Phys.* **12**, 053027 (2010).
- [13] N. R. Cooper and G. V. Shlyapnikov, *Phys. Rev. Lett.* **103**, 155302 (2009).
- [14] D.-W. Wang, M. D. Lukin, and E. Demler, *Phys. Rev. Lett.* **97**, 180413 (2006).
- [15] A. Micheli, G. K. Brennen, and P. Zoller, *Nature Phys.* **2**, 341 (2006).
- [16] A. V. Gorshkov, S. R. Manmana, G. Chen, J. Ye, E. Demler, M. D. Lukin, and A. M. Rey, *Phys. Rev. Lett.* **107**, 115301 (2011).
- [17] A. V. Gorshkov, S. R. Manmana, G. Chen, E. Demler, M. D. Lukin, and A. M. Rey, *Phys. Rev. A* **84**, 033619 (2011).
- [18] N. D. Mermin and H. Wagner, *Phys. Rev. Lett.* **17**, 1133 (1966).
- [19] P. Bruno, *Phys. Rev. Lett.* **87**, 137203 (2001).
- [20] J. R. de Sousa, *Eur. Phys. J. B* **43**, 93 (2005).
- [21] X.-L. Deng, D. Porras, and J. I. Cirac, *Phys. Rev. A* **72**, 063407 (2005).
- [22] N. Schuch, J. I. Cirac, and M. M. Wolf, *Commun. Math. Phys.* **267**, 65 (2006).
- [23] S. Müller, Diploma thesis, University of Stuttgart (2010).
- [24] P. Hauke, F. M. Cucchietti, A. Müller-Hermes, M. Bauls, and J. Ignacio, *New J. Phys.* **12**, 113037 (2010).
- [25] R. Kubo, *Phys. Rev.* **87**, 568 (1952).
- [26] A. Auerbach, *Interacting Electrons and Quantum Magnetism* (Springer-Verlag, New York, 1994).
- [27] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.109.025303> for demonstration of the spin wave analysis for the *XY* antiferromagnetic phase of the dipolar spin model