

Proton Spin Structure from Measurable Parton Distributions

Xiangdong Ji,^{1,2,3} Xiaonu Xiong,^{2,4} and Feng Yuan⁴

¹Department of Physics, Shanghai Jiao Tong University,

and Shanghai Key Lab for Particle Physics and Cosmology, Shanghai 200240, People's Republic of China

²Center for High-Energy Physics, Peking University, Beijing 100080, People's Republic of China

³Maryland Center for Fundamental Physics, University of Maryland, College Park, Maryland 20742, USA

⁴Lawrence Berkeley National Laboratory, Nuclear Science Division, Berkeley, California 94720, USA

(Received 25 February 2012; published 10 October 2012)

We present a systematic study of the proton spin structure in terms of measurable parton distributions. For a transversely polarized proton, we derive a polarization sum rule from the leading generalized parton distributions appearing in hard exclusive processes. For a longitudinally polarized proton, we obtain a helicity decomposition from well-known quark and gluon helicity distributions and orbital angular-momentum contributions. The latter are shown to be related to measurable subleading generalized parton distributions and quantum-phase space Wigner distributions.

DOI: 10.1103/PhysRevLett.109.152005

PACS numbers: 13.88.+e, 12.38.Aw, 13.60.Hb, 14.20.Dh

Introduction.—Understanding the internal structure of the proton, its spin structure in particular, has been a driving motive for intense activities in hadron physics in the past two decades. Great progress has been made on both the experimental and theory sides. Studies of deep-inelastic scattering and related hard processes at the electron facilities at SLAC, DESY, CERN, and Jefferson Lab, and of polarized proton-proton collisions at the Relativistic Heavy-Ion Collider, have generated a large body of experimental data, revealing the delicate role of quarks and gluons in the proton spin. These developments have stimulated theoretical advances from a simple parton model description of the nucleon structure to multidimension distributions of partons, including the generalized parton distributions (GPDs), the transverse momentum dependent parton distributions (TMDs), and the quantum phase space Wigner distributions. Together with the advances made in lattice quantum chromodynamics (QCD), these developments have provided us not only deep insights for the partonic structure of the nucleon but also great opportunities to study the strong interaction physics, such as the QCD factorization for hard processes, and the universality of the associated parton distributions. A recent summary on the experimental and theoretical status can be found in Ref. [1].

One of the key developments in understanding the spin structure of the proton is the spin sum rule derived by one of the authors [2], where the total contributions to the spin from the quark and gluons can be measured through their GPDs separately [3]. The partonic interpretation of this spin sum rule is, however, obscure. In particular, for a transversely polarized proton, there appear conflicting partonic interpretations of the spin [4,5]. For longitudinal polarization, one can in principle deduce the quark orbital angular momentum (OAM) by subtracting the quark helicity distribution. However, it has not been possible to

identify a direct probe for the quark OAM in physical processes. Meanwhile, the relation between the gauge-invariant quark OAM and the canonical OAM [6,7] has been a confusing issue in formulating a helicity sum rule with simple physical significance.

In this Letter, we will address the above important questions by systematically seeking a partonic interpretation of the proton spin and the experimental measurability of the relevant distributions. We explain why a simple partonic sum rule exists only for the transverse polarization. We find that the gauge-invariant OAM contribution to the proton helicity is related to twist-2 and -3 GPDs which are measurable in hard exclusive processes. Finally, the canonical OAM distribution in the light-cone gauge is related to a Wigner distribution [8,9], which is accessible through certain hard processes. Our discussions are mainly focused on quarks, but they can be easily extended to gluons.

Our starting point is the matrix element of the QCD AM density $M^{\mu\alpha\beta}$ in the nucleon plane-wave state [6]

$$\begin{aligned} \langle PS | \int d^4\xi M^{\mu\alpha\beta}(\xi) | PS \rangle \\ = J \frac{2S_\rho P_\sigma}{M^2} (2\pi)^4 \delta^4(0) [\epsilon^{\alpha\beta\rho\sigma} P^\mu + \epsilon^{[\alpha\rho\sigma} P^{\beta]} \\ - (\text{trace})] + \dots, \end{aligned} \quad (1)$$

where ξ^μ are the space-time coordinates, P^μ and S^μ ($S \cdot P = 0$, $S^2 = -M^2$) are the four-momentum and polarization of the nucleon, respectively, and $J = 1/2$ and M are the spin and mass, respectively. The $[\alpha \dots \beta]$ indicates antisymmetrization of the two indices. The above equation is fully Lorentz-covariant and can be specialized to any frame of reference. To seek the partonic interpretation, we consider the nucleon in the infinite momentum frame along the z direction and take μ to be the $+$ component [$P^+ = (P^0 + P^3)/\sqrt{2}$]. Because of the antisymmetry between α

and β , the leading component of the angular momentum density comes from $\alpha = +$ and $\beta = \perp = (1, 2)$. This is possible only if the nucleon is transversely polarized (S_\perp) and the matrix element reduces to

$$\langle PS | \int d^4 \xi M^{++\perp} | PS \rangle = J \left[\frac{3(P^+)^2 S^{\perp'}}{M^2} \right] (2\pi)^4 \delta^4(0), \quad (2)$$

where $S^{\perp'} = \epsilon^{-+\perp\rho} S_\rho$ with the convention of $\epsilon^{0123} = 1$. In the above equation, a factor of 2 comes from the first term in the bracket of Eq. (1), whereas the second term contributes to a factor 1 because of the antisymmetric feature of indices α and β .

The longitudinal polarization supports the matrix element of the next-to-leading AM tensor component M^{+12} :

$$\langle PS | \int d^4 \xi \vec{\xi} M^{+12} | PS \rangle = J(2S^+)(2\pi)^4 \delta^4(0), \quad (3)$$

which has one P^+ factor less. Thus the nucleon helicity J is a subleading light-cone quantity, and a partonic interpretation will in general involve parton transverse momentum and correlations.

The above result is contrary to the common intuition about the role of spin-1/2 particle polarization in hard scattering processes: The polarization vector S^μ has the leading light-cone component $S^+ = P^+$ when the nucleon is longitudinally polarized, and the transverse component S^\perp is subleading in the infinite momentum frame.

Transverse-polarization sum rule.—According to Eq. (2), one expects a simple partonic interpretation of the transverse proton polarization from the leading parton distributions. Indeed, the quark AM sum rule derived in terms of the quark distribution $q(x)$ and GPD $E(x, 0, 0)$ is exactly of this type [2]:

$$J_q = \frac{1}{2} \sum_i \int dx x [q_i(x) + E_i(x, 0, 0)], \quad (4)$$

where i sums over different flavors of quarks, and similarly for the gluon AM. We emphasize that this spin sum rule is frame-independent. In Ref. [4], Burkardt has proposed an interesting explanation of the above result in the impact parameter space, in which a transversely polarized nucleon state fixed in the transverse plane generates a spatial asymmetric parton density $q(x, b_\perp)$, which yields to the parton's AM contribution to the transverse spin. Note that the above sum rule is different from that of Leader [5], because the transverse angular \vec{J}_\perp does not commute with the Lorentz boost along the z direction.

To attribute the above sum rule with a simple parton picture, one has to justify that $(x/2)(q(x) + E(x))$ is the transverse AM density in x ; i.e., it is just the contribution to the transverse nucleon spin from partons with longitudinal momentum xP^+ . This can be done easily. Define the quark longitudinal momentum density $\rho^+(x, \xi, S^\perp)$ through

$$\begin{aligned} \rho^+(x, \xi, S^\perp) \\ = x \int \frac{d\lambda}{4\pi} e^{i\lambda x} \langle PS^\perp | \bar{\psi} \left(-\frac{\lambda n}{2}, \xi \right) \gamma^+ \psi \left(\frac{\lambda n}{2}, \xi \right) | PS^\perp \rangle, \end{aligned} \quad (5)$$

where n is the conjugation vector associated with P : $n = (0^+, n^-, 0_\perp)$ with $n \cdot P = 1$. A careful calculation shows that, beside the usual momentum distribution, it has an additional term

$$\begin{aligned} \rho^+(x, \xi, S^\perp) / P^+ = xq(x) + \frac{1}{2} x [q(x) \\ + E(x)] \lim_{\Delta_\perp \rightarrow 0} \frac{S^{\perp'}}{M^2} \partial^{\perp_\epsilon} e^{i\xi_\perp \Delta_\perp}, \end{aligned} \quad (6)$$

where the ξ^\perp dependence comes from the slightly off-forward matrix element, which acts like a “distribution” in a mathematical sense: vanishing normally but nonzero when integrated with some kernels. The parton contribution to the transverse polarization is just the transverse-space moment of $\rho^+(x, \xi, S^\perp)$:

$$\begin{aligned} S_\perp^q(x) = \frac{M^2}{2P^+ S^{\perp'} (2\pi)^2 \delta^2(0)} \int d^2 \xi \xi^\perp \rho^+(x, \xi, S^\perp) \\ = \frac{x}{2} [q(x) + E(x)], \end{aligned} \quad (7)$$

where we have included the contribution from the energy-momentum component $T^{+\perp}$ through Lorentz symmetry.

Helicity sum rule.—Most of the experimental probes on the nucleon spin use the longitudinal polarization, and thus it is natural to explore the nucleon helicity in the parton picture. Considering the z component of the quark AM, we have

$$\begin{aligned} J^3 = \int d^3 \xi \vec{\xi} M^{+12}(\xi) \\ = \int d^3 \xi \left[\bar{\psi} \gamma^+ \left(\frac{\Sigma^3}{2} \right) \psi + \bar{\psi} \gamma^+ [\xi^1 (iD^2) - \xi^2 (iD^1)] \psi \right], \end{aligned} \quad (8)$$

while the quark helicity is well known to have a simple parton density interpretation. However, the quark OAM involves a transverse component of the gluon field and thus is related to three-parton correlations. We notice recent research activities aiming at different decompositions of the nucleon spin [10], which we will not address in this Letter (see also a recent comment on these developments [11]).

Thus a partonic picture of the orbital contribution to the nucleon helicity necessarily involves the parton's transverse momentum. In other words, TMD parton distributions are the right objects for physical measurements and interpretation. In recent years, TMDs and novel effects associated with them have been explored extensively in both theory and experiment [1]. An important theoretical issue related to them is gauge invariance. Whenever a

canonical momentum of color-charged particles appears, the gauge symmetry requires that the gauge potential A^μ must be present simultaneously. This is already true when the parton's longitudinal momentum distribution is considered: In factorization theorems for deep-inelastic scattering, the physical parton represents a gauge-invariant object with a gauge link extended from the location of the parton field to infinity along the conjugating light-cone direction n^μ :

$$\Psi_{\text{LC}}(\xi) = P \left[\exp \left(-ig \int_0^\infty d\lambda n \cdot A(\lambda n + \xi) \right) \right] \psi(\xi), \quad (9)$$

where P indicates path ordering. Therefore, in perturbative diagrams, a parton with momentum $k^+ = xP^+$ represents in fact the sum of all diagrams with longitudinal gluons involved.

When considering the parton's transverse momentum, we also need appropriate gauge links formed of gauge potentials. The choice for the gauge links, however, is scattering-process dependent [12]. As a consequence, there is no unique definition for the TMDs. In practical applications, two choices stand out. The first one uses the same light-cone gauge link as shown in the above. This choice does lead to light-cone singularities, which must be addressed properly in actual calculations [13]. The second choice is a straight-line gauge link along the direction of space-time position ξ^μ :

$$\Psi_{FS}(\xi) = P \left[\exp \left(-ig \int_0^\infty d\lambda \xi \cdot A(\lambda \xi) \right) \right] \psi(\xi). \quad (10)$$

The link reduces to unity in the Fock-Schwinger gauge, $\xi \cdot A(\xi) = 0$. The gauge-invariant parton fields $\Psi(\xi)$ are defined in the infinite momentum frame which is the basis of partonic interpretation.

To investigate parton's OAM contribution to the proton helicity, one also needs their transverse coordinates. The most natural concept is a phase-space Wigner distribution, which was first introduced in Ref. [14]. A Wigner distribution operator for quarks is defined as

$$\hat{W}(\vec{r}, k) = \int \bar{\Psi}(\vec{r} - \xi/2) \gamma^+ \Psi(\vec{r} + \xi/2) e^{ik \cdot \xi} d^4 \xi, \quad (11)$$

where \vec{r} is the quark phase-space position and k the phase-space four-momentum, and Ψ follows the definitions of Eqs. (9) and (10). They represent the two different choices for the gauge links associated with the quark distributions. Including the gauge links in Eqs. (9) and (10) makes the above definition gauge-invariant. However, they do depend on the choice of the gauge link [12], as we will show below. The Wigner distribution can be defined as the expectation value of \hat{W} in the nucleon state:

$$W(k^+ = xP^+, \vec{b}_\perp, \vec{k}_\perp) = \frac{1}{2} \int \frac{d^2 \vec{q}_\perp}{(2\pi)^3} \int \frac{dk^-}{(2\pi)^3} e^{-i\vec{q}_\perp \cdot \vec{b}_\perp} \left\langle \frac{\vec{q}_\perp}{2} \left| \hat{W}(0, k) \right| -\frac{\vec{q}_\perp}{2} \right\rangle, \quad (12)$$

where the nucleon has definite helicity 1/2. The quark's OAM distribution follows from the intuition

$$L(x) = \int (\vec{b}_\perp \times \vec{k}_\perp) W(x, \vec{b}_\perp, \vec{k}_\perp) d^2 \vec{b}_\perp d^2 \vec{k}_\perp, \quad (13)$$

from partons with longitudinal momentum xP^+ .

For our purpose, the most appealing choice is Ψ_{FS} , because it leads to a light-cone AM density both calculable on a lattice and measurable experimentally. To demonstrate this, we need the Taylor expansion

$$\begin{aligned} & \bar{\Psi}_{FS}(-\xi/2) \gamma^+ \Psi_{FS}(\xi/2) \\ &= \sum_{n=0}^{\infty} \bar{\psi}(0) \gamma^+ \vec{D}^{\mu_1} \dots \vec{D}^{\mu_n} \psi(0) \xi_{\mu_1} \dots \xi_{\mu_n}. \end{aligned} \quad (14)$$

It follows that

$$\begin{aligned} & \int x^{n-1} L_{FS}(x) dx \\ &= \frac{1}{\langle PS|PS \rangle} \langle PS| \int d^3 \vec{r} \sum_{i=0}^{n-1} \frac{1}{n} \bar{\psi}(\vec{r}) (in \cdot D)^i (\vec{r}_\perp \times i\vec{D}_\perp) \\ & \quad \times (in \cdot D)^{n-1-i} \psi(\vec{r}) |PS \rangle. \end{aligned} \quad (15)$$

The right-hand side is related to the matrix elements of twist-2 and twist-3 operators, which are extractable from experimental data on twist-3 GPDs [15,16]. Because there are no light-cone nonlocal operators involved, it can also be calculated in lattice QCD [17]. We emphasize that $L_{FS}(x)$ is not the same as the OAM density defined through the generalized AM density in Ref. [18]. The difference is a twist-3 GPD contribution proportional to the gluon field $F^{+\perp}$.

The total OAM sum rule in terms of parton's Wigner distribution is

$$\begin{aligned} & \frac{\langle PS| \int d^3 \vec{r} \bar{\psi}(\vec{r}) \gamma^+ (\vec{r}_\perp \times i\vec{D}_\perp) \psi(\vec{r}) |PS \rangle}{\langle PS|PS \rangle} \\ &= \int (\vec{b}_\perp \times \vec{k}_\perp) W_{FS}(x, \vec{b}_\perp, \vec{k}_\perp) dx d^2 \vec{b}_\perp d^2 \vec{k}_\perp, \end{aligned} \quad (16)$$

which gives a parton picture for the gauge-invariant OAM [2], although the straight-line gauge link destroys the straightforward parton density interpretation.

Other choices of gauge links yield different Wigner distributions and hence different partonic OAM distributions $L(x)$. However, so long as the gauge link between $[-\xi/2, \xi/2]$ is smoothly differentiable, Eq. (16) remains valid. This is one of the important virtues of the gauge-invariant approach. However, for partons with the light-cone gauge link Ψ_{LC} , the above sum rule is invalid, as we shall see below.

Canonical orbital angular momentum.—The quark contribution to the canonical orbital angular momentum was explored in Ref. [7]:

$$l_q(x) = \frac{1}{(2\pi)^2 2P^+ \delta^2(0)} \int \frac{d\lambda}{2\pi} e^{ix\lambda} d^2\xi \langle PS | \bar{\psi} \left(-\frac{\lambda n}{2}, \xi \right) \gamma^+ \times (\xi^1 i\partial^2 - \xi^2 i\partial^1) \psi \left(\frac{\lambda n}{2}, \xi \right) | PS \rangle. \quad (17)$$

This definition represents the canonical OAM in the light-cone gauge $A^+ = 0$ and is not gauge invariant. A gauge-dependent quantity is in principle not measurable experimentally. However, sometimes one can fortunately find its *gauge-invariant extension* (GIE) physically measurable. A GIE of a gauge-variant quantity is a fixed-gauge result gauge-invariantly extrapolated to any other gauge. A GIE of the partial derivative in $A^+ = 0$ gauge is

$$i\partial_{\xi}^{\perp} = iD_{\xi}^{\perp} + \int^{\xi^-} d\eta^- L_{[\xi^-, \eta^-]} g F^{+\perp}(\eta^-, \xi_{\perp}) L_{[\eta^-, \xi^-]}, \quad (18)$$

which is uniquely defined, and $L_{[\xi^-, \eta^-]}$ is the light-cone gauge link connecting ξ^- and η^- . One can plug this into Eq. (17) to obtain a GIE of $l_q(x)$ away from $A^+ = 0$. The covariant derivative term is just what we have discussed before. The second term involves a nonlocal operator along the light cone and has an obscure physical meaning other than in the light-cone gauge. Its matrix element is in principle related to the twist-3 GPDs [15], and an infinite number of moments are involved due to nonlocality. Therefore, we arrive at the interesting conclusion that $l_q(x)$ in the light-cone gauge is actually accessible through twist-2 and -3 GPDs, which is consistent with what Hatta has concluded recently [9].

A clear parton picture emerges through connections between $l_q(x)$ and TMDs and Wigner distributions [8,9]. One can introduce a Wigner distribution with the gauge link in the light-cone direction, $W_{\text{LC}}(x, \vec{b}_{\perp}, \vec{k}_{\perp})$. Integration over the impact parameter space $\int d^2\vec{b}_{\perp} W_{\text{LC}}$ generates quark-spin independent TMDs. It can be shown that the canonical AM distribution in $A^+ = 0$ gauge as defined in Ref. [7] can be obtained from the simple moment of a gauge-invariant Wigner distribution:

$$l_q(x) = \int (\vec{b}_{\perp} \times \vec{k}_{\perp}) W_{\text{LC}}(x, \vec{b}_{\perp}, \vec{k}_{\perp}) d^2\vec{b}_{\perp} d^2\vec{k}_{\perp}. \quad (19)$$

From the discussion of the previous paragraph, this also implies constraints on the moments of Wigner distributions from the GPDs. Finally, the canonical OAM in the light-cone gauge acquires the simple parton sum rule in the quantum phase space [8,9]:

$$l_q = \frac{\langle PS | \int d^3\vec{r} \bar{\psi}(\vec{r}) \gamma^+ (\vec{r}_{\perp} \times i\vec{\partial}_{\perp}) \psi(\vec{r}) | PS \rangle}{\langle PS | PS \rangle} = \int (\vec{b}_{\perp} \times \vec{k}_{\perp}) W_{\text{LC}}(x, \vec{b}_{\perp}, \vec{k}_{\perp}) dx d^2\vec{b}_{\perp} d^2\vec{k}_{\perp}. \quad (20)$$

The measurability of this Wigner distribution will be studied in a future publication [16].

Conclusion.—In summary, we explored systematically parton pictures for the proton spin and achieved a number of important results. For the transverse polarization, we found that it is simple to interpret in terms of parton AM density measurable through twist-2 GPDs. For the nucleon helicity, the gauge-invariant parton picture can be probed through twist-2 and -3 GPDs and is also calculable in lattice QCD. A simpler parton picture in the light-cone gauge can be established through the quantum phase space Wigner distribution and can be measured through either twist-2 and -3 GPDs or directly from Wigner distribution. These results will stimulate further theoretical developments and generate experimental interests to measure, particularly, the parton OAM in hard scattering processes. Phenomenological studies will be presented elsewhere.

We thank M. Burkardt and J. H. Zhang for helpful discussions. This work was partially supported by the U.S. Department of Energy via Grants No. DE-FG02-93ER-40762 and No. DE-AC02-05CH11231 and a Grant (No. 11DZ2260700) from the Office of Science and Technology in Shanghai Municipal Government.

-
- [1] D. Boer *et al.*, arXiv:1108.1713.
 - [2] X. Ji, *Phys. Rev. Lett.* **78**, 610 (1997).
 - [3] X. Ji, *Phys. Rev. D* **55**, 7114 (1997); A. V. Radyushkin, *Phys. Rev. D* **56**, 5524 (1997).
 - [4] M. Burkardt, *Int. J. Mod. Phys. A* **18**, 173 (2003); *Phys. Rev. D* **72**, 094020 (2005).
 - [5] E. Leader, *Phys. Rev. D* **85**, 051501 (2012).
 - [6] R. L. Jaffe and A. Manohar, *Nucl. Phys.* **B337**, 509 (1990).
 - [7] S. Bashinsky and R. L. Jaffe, *Nucl. Phys.* **B536**, 303 (1998); P. Hagler and A. Schafer, *Phys. Lett. B* **430**, 179 (1998).
 - [8] C. Lorce and B. Pasquini, *Phys. Rev. D* **84**, 014015 (2011); C. Lorce, B. Pasquini, X. Xiong, and F. Yuan, *Phys. Rev. D* **85**, 114006 (2012).
 - [9] Y. Hatta, *Phys. Lett. B* **708**, 186 (2012).
 - [10] X.-S. Chen, X.-F. Lu, W.-M. Sun, F. Wang, and T. Goldman, *Phys. Rev. Lett.* **100**, 232002 (2008); M. Wakamatsu, *Phys. Rev. D* **83**, 014012 (2011).
 - [11] X. Ji, Y. Xu, and Y. Zhao, *J. High Energy Phys.* **08** (2012) 082.
 - [12] J. C. Collins, *Phys. Lett. B* **536**, 43 (2002); S. J. Brodsky, D. S. Hwang, and I. Schmidt, *Nucl. Phys.* **B642**, 344 (2002); A. V. Belitsky, X. Ji, and F. Yuan, *Nucl. Phys.* **B656**, 165 (2003).
 - [13] J. C. Collins, *Foundations of Perturbative QCD* (Cambridge University Press, Cambridge, England, 2011); X. Ji, J. P. Ma, and F. Yuan, *Phys. Rev. D* **71**, 034005 (2005).
 - [14] A. V. Belitsky, X. Ji, and F. Yuan, *Phys. Rev. D* **69**, 074014 (2004).
 - [15] A. V. Belitsky and A. V. Radyushkin, *Phys. Rep.* **418**, 1 (2005).
 - [16] X. Ji, X. Xiong, and F. Yuan (to be published).
 - [17] B. U. Musch, P. Hagler, J. W. Negele, and A. Schafer, *Phys. Rev. D* **83**, 094507 (2011).
 - [18] P. Hoodbhoy, X. Ji, and W. Lu, *Phys. Rev. D* **59**, 014013 (1998).