

Bilayers of Rydberg Atoms as a Quantum Simulator for Unconventional Superconductors

J. P. Hague and C. MacCormick

The Open University, Walton Hall, Milton Keynes MK7 6AA, United Kingdom

(Received 12 December 2011; revised manuscript received 3 July 2012; published 28 November 2012)

In condensed matter, it is often difficult to untangle the effects of competing interactions, and this is especially problematic for superconductors. Quantum simulators may help: here we show how exploiting the properties of highly excited Rydberg states of cold fermionic atoms in a bilayer lattice can simulate electron-phonon interactions in the presence of strong correlation—a scenario found in many unconventional superconductors. We discuss the core features of the simulator, and use numerics to compare with condensed matter analogues. Finally, we illustrate how to achieve a practical, tunable implementation of the simulation using “painted spot” potentials.

DOI: [10.1103/PhysRevLett.109.223001](https://doi.org/10.1103/PhysRevLett.109.223001)

PACS numbers: 37.10.Jk, 32.80.Ee, 67.85.-d, 74.20.-z

Cold atom quantum simulators offer an important new approach to the study of correlated electron phenomena without the limitations of computational or analytical techniques. For example, cold atoms have recently been used as quantum simulators to investigate models of condensed matter such as the Hubbard model of strong local Coulomb repulsion, which is important for the understanding of cuprate superconductors [1,2]. This has led to direct observation of important phenomena such as the superfluid to Mott insulator transition [3,4].

Besides the cuprates, there are several superconductors with high transition temperatures, many of which have important roles for electron-phonon interactions, and repulsion driven correlated electron phenomena such as antiferromagnetism. Fulleride superconductors of the family A_3C_{60} have phonon driven transition temperatures [5] of up to 40 K [6], but also exhibit antiferromagnetism at appropriate dopings and structures [7]. Superconductivity in bismuthates with $T_C > 30$ K [8] is probably due to strong couplings of localized electrons to the lattice [9] (as evidenced by a large isotope shift [10,11]). High transition temperatures have also been achieved in the borocarbides [12,13] ($T_C = 23$ K) and chloronitrides [14] ($T_C = 25$ K). The more conventional layered MgB_2 and graphite intercalation compounds are also interesting. Even the cuprate superconductors, where superconductivity is thought by many to be driven by anti-ferromagnetic fluctuations [15], show isotope shifts [16] and other effects such as kinks [17] that may be attributed to a nontrivial interplay between strong correlations and lattice vibrations.

Owing to the importance of electron-phonon interactions in condensed matter, reliable numerical methods have been sought, but even very simplified models [18] are extremely hard to simulate. Simulations either have to deal with the potentially infinite number of phonons associated with even a single electron, or retardation effects if the phonons are integrated out. A key problem in many advanced materials is that electrons are localized to atomic

orbitals and accordingly the Fermi energy is small. This localization means that dimensionless electron-phonon couplings are relatively strong and phonon frequencies can be large at around 10% of the Fermi energy. Even moderate couplings with intermediate frequency phonons can lead to consequences that cannot be predicted with perturbation theory and other analytics. Such couplings can cause additional difficulties for numerics, that are often most efficient in the extreme limits of weak or strong coupling. Recent advances in continuous time quantum Monte Carlo (CTQMC) cluster impurity solvers for dynamical mean field theories offer a state of the art for numerical simulations of Hubbard and Holstein models (for a review, see Ref. [19]). Such simulations are currently limited to ~ 36 site clusters for Hubbard models and 12 site clusters for Holstein models, limiting the range of spatial fluctuations that can be simulated in 2D to a few sites.

Quantum simulation of lattice effects has proved difficult to implement. The existence of quantum simulators with a high degree of control over the form of interactions has the potential to provide significant insight into the subtle interplay between electronic correlation, lattice vibration, and phenomena such as superconductivity and colossal magnetoresistance. We propose an approach for simulating fermionic Hubbard models extended to include strong long-range interactions with lattice vibrations, by discussing Rydberg states of cold atoms in bilayer lattices. Such a simulator is likely to shed light on a wide range of unconventional superconductors with transition temperatures greater than 30 K, and might resolve aspects of ongoing debates on cuprate mechanisms.

A quantum simulator for complex electronic systems such as unconventional superconductors should have the following main characteristics. (1) It must be able to simulate fermions. (2) It should be capable of simulating all filling factors up to and including half filling (1 fermion per lattice site) where the physics of materials with complex phase diagrams is most interesting. (3) All parameters, including the phonon frequency, electron-phonon coupling,

Hubbard U , and hopping t should be highly tunable for all interesting physical regimes.

Several schemes for quantum simulation of interactions between electrons and phonons have been suggested. A proposal to bathe bosonic or fermionic impurities in an optical lattice in a Bose-Einstein condensate (BEC) [20–22] has led to the observation of polaron effects for bosonic impurities [23]. This succeeds in criterion (1) as the scheme can in principle treat very low fermion density. However, it is essential that Fermionic impurities have a negligible effect on the underlying BEC (specifically, fluctuations in the BEC order parameter must be small [20–22]). This limits the schemes to very low fermion density, voiding criterion (2).

Alternate schemes are based on the interaction between phonons and bosonic excitations. Interactions with high-energy phonon states of Rydberg ions are used as a part of the mapping in proposed simulators for spin systems [24,25]. Li and Lesanovsky have discussed structural distortions associated with exciting high energy Rydberg states in cold ion crystals [26]. Rydberg atoms have been proposed as a way of simulating polaron effects in strongly deformable materials [27]. The use of cold polar molecules to obtain Holstein polaron effects has been discussed [28]. These schemes are not extensible to fermions, so do not fulfil criterion (1), although they are highly tunable. The scheme proposed in this Letter goes beyond these as it is capable of simulating interacting Fermions with arbitrary filling factor, while retaining a high level of control over all parameters in the Hamiltonian, thus satisfying all of criteria (1), (2), and (3) necessary to examine the complex phase diagrams of strongly correlated systems in the presence of phonons.

We begin by discussing how electron-phonon interactions can be simulated in a system of cold, highly excited (Rydberg) atoms in a bilayer lattice. Over long ranges, we assume that the Rydberg atoms interact only via dipole-dipole interactions in the strong-coupling Förster regime, $V_{kl} = \mu^2/|\mathbf{R}_k - \mathbf{R}_l|^3$, where μ is the dipole moment on the Rydberg atom and \mathbf{R}_k is the vector to the k th atom in the lattice (see Supplemental Material [29] for a discussion of the origins of this interaction). The dipole moment may be written in terms of the coefficient C_3 as $\mu = \sqrt{2}C_3$, and depends upon the Rydberg states chosen for the experiment. The ground state atoms have no long-range interaction. By coupling the ground $|g\rangle$ and Rydberg $|r\rangle$ states with a laser tuned Δ from the $|g\rangle \rightarrow |r\rangle$ transition and with coupling strength Ω (the Rabi frequency) we mix the states $|g\rangle$ and $|r\rangle$. This technique [30] of dressing the atoms with the laser means that the trapped, ground state atoms acquire the characteristics of the Rydberg state, but in a controllable fashion. In particular, the coefficient C_3 is replaced by an effective interaction coefficient which we write as $C'_3 = (\Omega/2\Delta)C_3$. We note that the use of a different regime of dipole-dipole interactions has been proposed for the simulation of liquid crystalline phases [31].

The Rydberg atoms must now be confined in a bilayer lattice. The “itinerant” layer represents the electrons in a condensed matter problem, and the “phonon” layer generates phonon mediated interactions. The itinerant layer can have any filling, and tunneling between adjacent sites is allowed. It is important that the phonon layer has 1 atom per site and that the tunneling is forbidden—the Mott insulator phase [3]. This is achieved by making the potential barrier in the itinerant layer smaller than in the phonon layer. A further complication arises because the atoms in the phonon layer must be set with low oscillating frequencies, while the itinerant layer must be set up with high phonon frequencies. This can be achieved if the optical lattice in the phonon layer has a special configuration as seen in Fig. 1. This form may be achieved using painted potentials [32].

In the itinerant layer, fermions hop with amplitude t , and experience a local Hubbard U , which originates from scattering from the hard-core potential between the fermions when they share the same lattice site [2]. The Hamiltonian for these interactions is $H_{\text{Hub}} = -t\sum_{\langle i,i'\rangle} c_i^\dagger c_{i'} + U\sum_i n_i n_i$ (n_i is the number operator for fermions on site i , and c_i^\dagger creates a Rydberg atom on site i). Lattice vibrations are introduced by displacing the atoms in the phonon layer. Atomic displacements do not affect the optical lattice, so the vibrations of the atoms are momentum independent Einstein phonons with Hamiltonian, $H_{\text{ph}} = \sum_{\mathbf{k},\nu} \hbar\omega_{\mathbf{k},\nu} (d_{\mathbf{k},\nu}^\dagger d_{\mathbf{k},\nu} + \frac{1}{2})$ and with polarization vectors $\boldsymbol{\xi}_{\mathbf{k},\nu} = \boldsymbol{\xi}_\nu$ in orthogonal directions. Here $\omega_{\mathbf{k},\nu} = \omega_\nu$ is the angular frequency of a phonon in mode ν with momentum \mathbf{k} , and d^\dagger and d create and annihilate phonons.

Small in-plane phonon displacements \mathbf{u}_i cause the interaction between Rydberg states to become $V'_{kl} = -\Omega^2 \mu^2 / 4\Delta^2 |\mathbf{R}_k + \mathbf{u}_k - \mathbf{R}_l - \mathbf{u}_l|^3$, which expands as




FIG. 1 (color online). System of bilayer Rydberg cold atoms for simulation of strong correlations and interactions between fermions and phonons, annotated with Hamiltonian terms. t is the intersite hopping in the itinerant layer, ω_0 is the phonon frequency, g_{ij} is the Rydberg-phonon coupling, and $V_0^{(\text{ph})}$ and $V_0^{(\text{it})}$ are trap depths in the itinerant and phonon layers, respectively.

$$V'(\mathbf{R} + \mathbf{u}) \approx \frac{\Omega^2 \mu^2}{4\Delta^2 |\mathbf{R}|^3} - \frac{3\Omega^2 \mu^2 \mathbf{u} \cdot \hat{\mathbf{R}}}{4\Delta^2 |\mathbf{R}|^4}. \quad (1)$$

The phonons are quantized by substituting $\mathbf{u}_i = \sum_{\mathbf{k}, \nu} \sqrt{\hbar/2NM\omega_{\mathbf{k}, \nu}} \xi_{\mathbf{k}, \nu} (d_{\mathbf{k}, \nu} e^{-i\mathbf{k} \cdot \mathbf{R}_i} + d_{\mathbf{k}, \nu}^\dagger e^{i\mathbf{k} \cdot \mathbf{R}_i})$. Since all sites in the phonon layer are occupied, and the modes momentum independent, a multimode Rydberg-phonon interaction with extended Holstein form is derived,

$$H_{R\text{-ph}} = \frac{3\mu^2 \Omega^2}{4\Delta^2} \left(\frac{\hbar}{2M\omega_0} \right)^{1/2} \sum_{\nu} \sum_{ij} \frac{\hat{\mathbf{R}}_{ij} \cdot \xi_{\nu}}{R_{ij}^4} n_i (d_{j, \nu}^\dagger + d_{j, \nu}). \quad (2)$$

Here, \mathbf{R}_{ij} is a vector between an atom in the itinerant layer at site i , and an atom in the phonon layer at site j , N the number of sites, and M is the mass of the atoms. The presence of multiple phonon modes in the Hamiltonian is interesting since such interactions are difficult to simulate with current numerical techniques.

A further simplification can be made by elongating the potentials in the phonon layer along the direction perpendicular to the planes, so that the Hamiltonian becomes $H_{\text{Holstein}} = \sum_{ij} g_{ij} n_i (d_j^\dagger + d_j)$, with the interaction $g_{ij} = \frac{\Omega^2}{4\Delta^2} \frac{3\mu^2 b}{(b^2 + r_{ij}^2)^{5/2}} \left(\frac{\hbar}{2M\omega_0} \right)^{1/2}$, where r_{ij} is the distance between the projection of sites i and j onto the same layer and b is the interplane distance. To compare with condensed matter analogues, a Holstein model, where the local electron density couples to local optical phonon modes [18], has $g_{ij}^{(\text{Hol})} \sim \delta_{ij}$. In other condensed matter systems, Fröhlich electron-phonon interaction generalized to lattice models (also known as the extended Holstein interaction) has the form $g_{ij}^{(\text{Fr})} \sim \exp(-r_{ij}/R_{sc})(b^2 + r_{ij}^2)^{-3/2}$, where R_{sc} is a screening radius [33].

When the phonon energy is much larger than hopping, the effective instantaneous interaction between Rydberg atoms in the itinerant layer mediated through the phonon layer is $-zt\lambda \sum_m \Phi(\mathbf{r})/\Phi(0)$, where $\Phi(\mathbf{r})/\Phi(0) = \sum_m g_{r,m} g_{0,m} / \sum_m g_{0,m} g_{0,m}$. Figure 2 shows a comparison between the shapes of Rydberg and lattice Fröhlich effective interactions $\Phi^{(\text{Fr})}(x)/\Phi^{(\text{Fr})}(0)$ for various interplane distances and screening radii (a is the intersite distance in the plane). As would be done in the experiment, the near-neighbor interactions are matched by modifying b to get the closest possible correspondence to the Fröhlich interaction that is to be simulated. An excellent correspondence between the shapes of the interactions is seen for interplane distance $b \lesssim a$. The origin of the effective interaction is discussed in the Supplemental Material [29].

To demonstrate the scheme, we use CTQMC calculations [34] to compute the properties of polarons and bipolarons in the bilayer lattice and compare with results from the screened Fröhlich interaction (Fig. 3). The simulations include phonon-mediated interactions, the Hubbard U , and




FIG. 2. Comparison between effective Rydberg-Rydberg interactions in the quantum simulator $\Phi(x)/\Phi(0)$ mediated via phonons in the Rydberg atom system here, and the effective electron-electron interaction $\Phi^{(\text{Fr})}(x)/\Phi^{(\text{Fr})}(0)$ mediated by phonons in a typical condensed matter system with Fröhlich interactions for various interplane distances. A good correspondence between the effective interactions is seen.

also the small direct dipole-dipole interaction between atoms in the itinerant layer, $H_{\text{direct}} = \sum_{i \neq i'} V_{ii'} n_i n_{i'}$ (as discussed in the Supplemental Material [29]). A very close agreement is found between the quantum simulator and Fröhlich system, further demonstrating that the small differences in the tails of the interaction and the residual long-range interactions in the itinerant layer do not affect local pairing.

Finally, we discuss a realistic experimental setup for building the quantum simulator. We recommend that experimentalists stage their investigations by using easier to control bosonic Rydberg atoms before moving on to fermions. A bilayer optical lattice may be set up with painted potentials [32], a powerful technique with a high level of control. Creating a bilayer painted lattice requires two focused horizontal sheets. Gaussian spots of waist $w^{(\text{it})}$ are focused on the itinerant sheet, and pairs of spots with




FIG. 3 (color online). Comparison of polaron and bipolaron energies calculated for the Rydberg quantum simulator (with $b = 1.468a$), and for a screened Fröhlich interaction in the condensed matter analogue. $\lambda = \Phi(0)/2ztM\omega^2$ and z is the in-plane coordination number. To highlight effects of changes in U , values of 0 and $8t$ are used. The small differences in the tails of the interaction do not strongly affect the physics. $k_B T = 0.014t$, $\omega = 0.2t$. Error bars are smaller than the points.




FIG. 4 (color online). Dimensionless parameters for rubidium atoms in the proposed spot potential experiment. All physically relevant regimes are accessible by tuning the spot power in the itinerant and phonon layers ($W = zt$). Further control of λ is possible by changing the detuning Δ .

waist $w^{(\text{ph})}$ separated by a distance $2D$ are painted in the phonon layer. The resulting lattices are filled with Rb atoms and detuned from the 48 S state by $\Delta = 125$ MHz, with $\Omega = 10$ MHz leading to $\mu^2 = 173$ MHz (note that the values found here for the bosonic Rb are representative of all alkali Rydberg atoms). Figure 4 shows the dimensionless parameters calculated for a realistic setup, with $a = 2.62 \mu\text{m}$ and $b = 3.67 \mu\text{m}$. Spot lattices in the itinerant layer have $w^{(\text{it})} = 2.46 \mu\text{m}$, so the potentials overlap leading to a near sinusoidal potential with $V_0^{(\text{it})} = 70.1$ Hz and energy level spacing in the itinerant layer of 153 Hz, $U = 5.34$ Hz, $t = 4.83$ Hz, so a one band model will be simulated. Phonon frequencies are easily changed in the phonon layer by tuning D , and can therefore be changed independently. As an example, with $V_0^{(\text{ph})} = 578$ Hz, $w^{(\text{ph})} = 0.655 \mu\text{m}$, and $D = 0.999w^{(\text{ph})}$, $\omega_0^{(\text{ph})} = 13.8$ Hz and $\lambda = 1.00$ is achieved—challenging but possible with a state of the art diffraction-limited optical arrangement. A simpler 1D version of the experiment that only requires a single horizontal sheet can be performed using two coupled chains (a phonon chain and an itinerant chain) and is recommended as a starting place when implementing the experiment. In 1D, we find equally good correspondence between the quantum simulator and condensed matter analogue.

In the bilayer system, the interesting physics are encoded in the momentum distribution of the gas held in the itinerant layer. What we want to study is the correlation function $C(\mathbf{k}, \sigma; -\mathbf{k}, -\sigma)$. To extract this, the momentum distribution must be observable. In the experiments in Ref. [35], this was accomplished using a time-of-flight method, where the momentum distribution of the roaming atoms is mapped onto the spatial location at the time of imaging.

Since it is the purpose of the proposed system to understand phase diagrams, we briefly note the effect that

different interaction types have in forming and modifying these phases. Repulsive Hubbard U promotes a Mott insulating state and antiferromagnetism close to half filling, and may promote superconductivity through spin fluctuations. It may also control the form of the superconductivity, reducing or eliminating s -wave pairing. Direct in-plane V_{ij} may promote charge ordering and suppress superconductivity if it is repulsive, and could enhance superconductivity when attractive. Interaction with phonons promotes BCS superconductivity at weak coupling, and a BEC of bipolarons for large λ .

In this Letter, we have shown how systems of cold Rydberg atoms in a bilayer can be used as a simulator for electron-phonon interactions in the presence of strong electronic correlation, of the type found in many unconventional superconductors. We have carried out the mapping to an extended Hubbard-Holstein model and used numerics to demonstrate the simulator is capable of reproducing the pairing and polaron physics of standard electron-phonon models. Furthermore, we have described how the simulator can be implemented using contemporary techniques. The proposed system goes well beyond the possibilities of previous quantum simulators for the simulation of interactions with lattice vibrations. In particular, we can reliably simulate half filled fermion systems relevant to cuprate and other unconventional superconductors, tune all parameters directly through the optical lattice, and can easily include multiple phonon modes.

J. P. H. acknowledges EPSRC Grant No. EP/H015655/1 and C. M. Grant No. EP/F031130/1. We thank I. Lesanovsky, M. Bruderer, F. Herrera, A. Kowalczyk, N. Braithwaite, S. Bergamini, S. Alexandrov, and J. Samson for useful discussions, and especially thank Pavel Kornilovitch for long-standing collaboration on QMC and polarons.

-
- [1] J. Hubbard, *Proc. R. Soc. A* **276**, 238 (1963).
 - [2] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
 - [3] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, *Nature (London)* **415**, 39 (2002).
 - [4] G. K. Campbell, J. Mun, M. Boyd, P. Medley, A. E. Leanhardt, L. G. Marcassa, D. E. Pritchard, and W. Ketterle, *Science* **313**, 649 (2006).
 - [5] A. Hebard, M. J. Rosseinsky, R. C. Haddon, D. W. Murphy, S. H. Glarum, T. T. M. Palstra, A. P. Ramirez, and A. R. Kortan, *Nature (London)* **350**, 600 (1991).
 - [6] T. T. M. Palstra, O. Zhou, Y. Iwasa, P. E. Sulewski, R. M. Fleming, and B. R. Zegarski, *Solid State Commun.* **93**, 327 (1995).
 - [7] A. Y. Ganin *et al.*, *Nature (London)* **466**, 221 (2010).
 - [8] R. J. Cava, B. Batlogg, J. J. Krajewski, R. Farrow, L. W. Rupp, A. E. White, K. Short, W. F. Peck, and T. Kometani, *Nature (London)* **332**, 814 (1988).

- [9] I. B. Bischofs, V. N. Kostur, and P. B. Allen, *Phys. Rev. B* **65**, 115112 (2002).
- [10] B. Batlogg, R. Cava, L. Rupp, A. Mjlsce, J. Krajewski, J. Remeika, W. Peck, A. Cooper, and G. Espinosa, *Phys. Rev. Lett.* **61**, 1670 (1988).
- [11] D. G. Hinks, D. R. Richards, B. Dabrowski, D. T. Marx, and A. W. Mitchell, *Nature (London)* **335**, 419 (1988).
- [12] R. J. Cava *et al.*, *Nature (London)* **367**, 146 (1994).
- [13] Z. Hossain, L. C. Gupta, C. Mazumdar, R. Nagarajan, S. K. Dhar, C. Godart, C. Levy-Clement, B. D. Padalia, and R. Vijayaraghavan, *Solid State Commun.* **92**, 341 (1994).
- [14] H. Kawaji, S. Yamanaka, and K. Hotehama, *Nature (London)* **392**, 580 (1998).
- [15] P. W. Anderson, *The Theory of Superconductivity in the Cuprates* (Princeton University, Princeton, NY, 1997).
- [16] G. M. Zhao, H. Keller, and K. Conder, *J. Phys. Condens. Matter* **13**, R569 (2001).
- [17] A. Lanzara *et al.*, *Nature (London)* **412**, 510 (2001).
- [18] T. Holstein, *Ann. Phys. (N.Y.)* **8**, 325 (1959).
- [19] E. Gull, A. Millis, A. Lichtenstein, A. Rubtsov, M. Troyer, and P. Werner, *Rev. Mod. Phys.* **83**, 349 (2011).
- [20] M. Bruderer, A. Klein, S. R. Clark, and D. Jaksch, *Phys. Rev. A* **76**, 011605(R) (2007).
- [21] M. Bruderer, T. Johnson, S. Clark, D. Jaksch, A. Posazhennikova, and W. Belzig, *Phys. Rev. A* **82**, 043617 (2010).
- [22] A. Privitera and W. Hofstetter, *Phys. Rev. A* **82**, 063614 (2010).
- [23] B. Gadway, D. Pertot, R. Reimann, and D. Schneble, *Phys. Rev. Lett.* **105**, 045303 (2010).
- [24] D. Porras and J. I. Cirac, *Phys. Rev. Lett.* **92**, 207901 (2004).
- [25] M. Müller, L. Liang, I. Lesanovsky, and P. Zoller, *New J. Phys.* **10**, 093009 (2008).
- [26] W. Li and I. Lesanovsky, *Phys. Rev. Lett.* **108**, 023003 (2012).
- [27] J. P. Hague and C. McCormick, *New J. Phys.* **14**, 033019 (2012).
- [28] F. Herrera and R. V. Krems, *Phys. Rev. A* **84**, 051401 (2011).
- [29] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.109.223001> for more information on the Rydberg-Rydberg interaction, an introduction to phonon mediated interactions, the CTQMC algorithm, and properties of painted spot potentials.
- [30] N. Henkel, R. Nath, and T. Pohl, *Phys. Rev. Lett.* **104**, 195302 (2010).
- [31] J. Quintanilla, S. T. Carr, and J. J. Betouras, *Phys. Rev. A* **79**, 031601(R) (2009).
- [32] K. Henderson, C. Ryu, C. McCormick, and M. G. Boshier, *New J. Phys.* **11**, 043030 (2009).
- [33] A. S. Alexandrov and P. E. Kornilovitch, *Phys. Rev. Lett.* **82**, 807 (1999).
- [34] J. P. Hague and P. E. Kornilovitch, *Phys. Rev. B* **80**, 054301 (2009).
- [35] W. S. Bakr, A. Peng, M. E. Tai, R. Ma, J. Simon, J. I. Gillen, S. Folling, L. Pollet, and M. Greiner, *Science* **329**, 547 (2010).