

Erratum: Topological Hall Effect in the A Phase of MnSi [Phys. Rev. Lett. 102, 186602 (2009)]

A. Neubauer, C. Pfleiderer, B. Binz, A. Rosch, R. Ritz, P.G. Niklowitz, and P. Böni

(Received 29 April 2013; published 16 May 2013)

DOI: [10.1103/PhysRevLett.110.209902](https://doi.org/10.1103/PhysRevLett.110.209902)

PACS numbers: 72.15.Gd, 72.25.-b, 72.80.Ga, 75.30.-m, 99.10.Cd

As part of an extensive study of the pressure dependence of the topological Hall signal associated with the Skyrmion lattice phase in MnSi [1,2], we have recently noticed a computational error in the value of the emergent magnetic field per Skyrmion in MnSi, $B_{\text{eff}} \approx -2.5$ T, stated in the Letter. The correct value is $B_{\text{eff}} \approx -13.15$ T and thus roughly a factor of 5 larger. The formalism presented in the Letter is unchanged and correct. In combination with new experimental insights reported in Refs. [1,2] the larger value of B_{eff} reconciles data reported in Ref. [3]. For detailed information, we refer to Refs. [1,2].

- [1] R. Ritz, M. Halder, C. Franz, A. Bauer, M. Wagner, R. Bamler, A. Rosch, and C. Pfleiderer, *Phys. Rev. B* **87**, 134424 (2013).
- [2] R. Ritz, M. Halder, C. Franz, A. Bauer, M. Wagner, and C. Pfleiderer, *Nature (London)* **497**, 231 (2013).
- [3] M. Lee, W. Kang, Y. Onose, Y. Tokura, and N. P. Ong, *Phys. Rev. Lett.* **102**, 186601 (2009).