

Emergent Irreversibility and Entanglement Spectrum Statistics

Claudio Chamon,¹ Alioscia Hamma,² and Eduardo R. Mucciolo³

¹*Department of Physics, Boston University, Boston, Massachusetts 02215, USA*

²*Center for Quantum Information, Institute for Interdisciplinary Information Sciences, Tsinghua University, Beijing 100084, People's Republic of China*

³*Department of Physics, University of Central Florida, Orlando, Florida 32816, USA*

(Received 13 November 2013; revised manuscript received 4 April 2014; published 16 June 2014)

We study the problem of irreversibility when the dynamical evolution of a many-body system is described by a stochastic quantum circuit. Such evolution is more general than a Hamiltonian one, and since energy levels are not well defined, the well-established connection between the statistical fluctuations of the energy spectrum and irreversibility cannot be made. We show that the entanglement spectrum provides a more general connection. Irreversibility is marked by a failure of a disentangling algorithm and is preceded by the appearance of Wigner-Dyson statistical fluctuations in the entanglement spectrum. This analysis can be done at the wave-function level and offers an alternative route to study quantum chaos and quantum integrability.

DOI: 10.1103/PhysRevLett.112.240501

PACS numbers: 03.67.Mn, 03.65.Ta, 05.45.Mt, 05.70.-a

In closed quantum systems, evolution is unitary and both irreversibility and nonintegrability are elusive notions. Because of unitarity, evolution is always stable under errors in initial conditions. Thus, in quantum mechanics irreversibility is defined by the vanishing of the probability (known as fidelity) of returning to an initial state under arbitrarily small imperfections in the Hamiltonian during the reversed time evolution [1]. Nonintegrability is associated to a Wigner-Dyson distribution of the energy-level spacings that shows level repulsion [2] and nonintegrable Hamiltonians in this context are irreversible. Integrable Hamiltonians, instead, tend to show clustering of energy levels but can be either reversible or irreversible [3,4]. When the time evolution is not governed by a Hamiltonian, or when the Hamiltonian is time dependent, energy levels are not well defined and these associations cease to be meaningful. How can one relate nonintegrability and irreversibility in these more general cases of quantum evolution?

In this Letter we show that one can answer this question by looking at the wave function alone. This route allows one to study generic quantum evolutions even when energy is not well defined. We show that by studying the level statistics of the entanglement spectrum one can determine whether the evolution is irreversible or not through a protocol that we call entanglement cooling. It turns out that the onset of irreversibility is marked by the presence of Wigner-Dyson statistics in the entanglement spectrum.

The quantum system we consider contains n qubits and evolves unitarily from an initial factorized state of the form $|\Psi_0\rangle = |\psi_1\rangle \otimes |\psi_2\rangle \otimes \dots \otimes |\psi_n\rangle$ where each single-qubit state is defined as $|\psi_j\rangle = \cos(\theta_j/2)|0\rangle + \sin(\theta_j/2)e^{i\phi_j}|1\rangle$, with θ_j and ϕ_j arbitrary. Formally, the evolution is obtained by applying a unitary matrix U to the

state vector, $|\Psi_t\rangle = U|\Psi_0\rangle = \sum_x \Psi_t(x)|x\rangle$, where the states $|x\rangle \equiv |x_1 x_2 \dots x_n\rangle$ form the computational basis, with $x_j = 0, 1$ for $j = 1, \dots, n$. Using the language of quantum computing, we assume that this unitary matrix is represented by gates. We recall that the two-qubit CNOT gate and arbitrary one-qubit rotations are sufficient for universal quantum computing [5]. In what follows, we shall restrict the gates to the permutation group, which is a subgroup of the unitary group. The restriction to the permutation subgroup of unitary transformations allows for a much more efficient computation of the state of the system as it evolves with gates. In particular, we consider the unitary gates in the set $\mathcal{I}_3 = \{SWAP, CNOT, Toffoli\}$ depicted in Fig. 1. We build a stochastic quantum circuit $U = \prod_k^M U_k$




FIG. 1. Reversible gates used in the quantum stochastic evolutions. The output values of the gates are defined when a , b , and c take values 0 and 1. Top row: The two-qubit gates SWAP and CNOT. Bottom row: The different variations of the three-qubit Toffoli gates. We note that different variations of the CNOT and Toffoli gates can be obtained from one fixed variation plus NOT gates.

by drawing randomly with uniform probability pairs or triplets of qubits and a random gate $U_k \in \mathcal{I}_3$ with probability $1/3$. We remark that the Toffoli gate alone is sufficient for universal classical computation [6]. We also consider more restricted (and nonuniversal) circuits obtained by employing only gates in the set $\mathcal{I}_2 = \{\text{SWAP}, \text{CNOT}\}$.

At each step k of the circuit, the n qubits are partitioned into subsystems (A, B) with n_A and n_B qubits, and the entanglement properties of the system are obtained through the singular values $\lambda_k > 0$, $k = 1, \dots, r$, which result from the Schmidt decomposition [7,8] of the state $|\Psi_t\rangle = \sum_{k=1}^r \lambda_k |\psi_{t(k)}^A\rangle \otimes |\psi_{t(k)}^B\rangle$. The reduced density matrices $\rho_A = \text{tr}_B(|\Psi_t\rangle\langle\Psi_t|)$ and $\rho_B = \text{tr}_A(|\Psi_t\rangle\langle\Psi_t|)$ have eigenvalues $\{p_k = \lambda_k^2\}$. These p_k define a probability distribution whose Rényi entropies are defined as [9]

$$S_q(n_A, n_B) = \frac{1}{1-q} \log_2 \sum_{k=1}^r p_k^q, \quad (1)$$

with $\sum_{k=1}^r p_k = 1$. The zeroth Rényi entropy is related to the rank, namely, the number r of nonzero singular values, $S_0 = \log_2 r$. The $q = 1$ Rényi entropy is the Shannon entropy measuring the amount of information in the distribution $\{p_k\}$: $S_1 = -\sum_k p_k \log_2 p_k$.

What happens to entanglement during the evolution with gates? One can show that, under a generic stochastic random circuit, entanglement grows linearly with time, and then saturates to its maximum possible value [10,11]. This occurs typically, meaning that the probability of having a different outcome is zero in the thermodynamic limit. A similar behavior is obtained also for the restricted quantum evolutions considered here, whether one uses two- or three-qubit gates. The saturation value is typically reached after about $M \sim n^2$ transformations. In Fig. 2, we see a numerical simulation of the protocol used, with

both two-qubit and three-qubit gates, which confirms this scenario. We call this part of the protocol “entanglement heating.”

Because entanglement increases with the number of gates, in order to revert the evolution to return back to the initial state, it is natural to attempt an algorithm that completely disentangles the system. The entanglement entropies provide a natural metric to use in a minimization process. If one is able to remove all the entanglement while recording the moves that led to the decreases, one builds one possible reverse algorithm that takes the system from the final state back to the initial (product) state. In practice, we implement such a disentangling or “entropy cooling” algorithm as follows. We attempt a gate, chosen at random, and compute the change in entanglement entropy. Then we decide whether or not to accept this gate into the sequence according to a Metropolis algorithm: if the entanglement goes down, we always take this move; if not, we take it with a certain probability, which we decrease as function of the number of attempts (similarly to simulated annealing, but applied to entanglement entropy and not energy). More precisely, we use as the optimization function the sum of the entanglement entropies over all bipartitions of the system into n_A and n_B consecutive qubits with $n_A + n_B = n$, namely, $\mathcal{S}_q = \sum_{n_A=1}^{n-1} S_q(n_A, n_B)$. The reason for this choice is that a single bipartition is sensitive only to gates that act on qubits in both subsystems A and B . However, if one considers the sums over the entanglement for all bipartitions, one is sensitive to all reductions in entanglement, no matter where the gates act.

The resulting $q = 1$ Rényi entropy as a function of the gate number for a given sequence of the algorithm using *only* the gates in \mathcal{I}_2 is shown in Fig. 2(a). The data show two examples of the entanglement evolution for two particular “heating” and “cooling” runs, as well as the average of 128 different realizations with random initial product states for 16 qubits, with each θ_j randomly picked




FIG. 2 (color online). Evolution of the entanglement entropy S_1 obtained from the bipartition of the qubit string at the middle ($n_A = n_B = 8$) as a function of the number of applied gates. First, M_h reversible gates are randomly applied (heating period); second, a Metropolis algorithm is used to reverse the evolution and restore zero entropy (cooldown period). The shorter solid black line ($M_h = 512$) and the longer red line ($M_h = 4096$) result from averaging S_1 over 128 initial-state realizations. The faded brown lines show the evolution of S_1 for two typical realizations. (a) Only two-qubit gates are used. Upper inset: The heating period. Lower inset: Shifted curves, showing that the cooling, on average, is independent of the duration of the heating period. (b) A mixture of two-qubit and Toffoli gates is used. Inset: Detail of the transition between heating and cooling periods. The dashed line indicates the transition point.

from the interval $[0, \pi]$, and $\phi_j = 0, \pi$ (thus focusing on real wave functions). We show data for the case when the system is entangled with 512 and with 4096 gates. The system is “cooled” by minimizing S_0 (similar results are obtained when minimizing S_2). Notice that the “cooling” time for the average curve does not depend on how long the system was “heated,” provided that the same maximum entanglement entropy is reached. The disentangling algorithm works for *all* individual realizations of the protocol. We were *always* able to reverse to a completely factorized tensor product state with zero entanglement. This is quite remarkable, because the success of the algorithm does not depend at all on the amount of the entanglement produced. So one may wonder whether *every* quantum circuit can be reversed with such a cooling protocol.

To answer this question, consider now the case when entanglement entropy “heating” involves the gates in \mathcal{I}_3 . Then, apply the disentangling algorithm using the same set of gates. For *all* realizations studied, we find that it is *never* possible to completely disentangle the state using the Metropolis protocol described above [12]. In Fig. 2(b) we show two typical realizations of the heating and cooling protocol, with a random initial product state and 512 random gate sequences for the heating phase. We also show the average over 128 realizations.

And yet, by only looking at the amount of entanglement generated upon “heating,” we cannot tell whether the evolution is reversible by the cooling algorithm. As we have shown, by heating with either two-qubit gates or a mixture of two-qubit and Toffoli gates, one rapidly reaches an almost maximally entangled state. Nevertheless, only for the former are we able to reverse the system back into a product-state form. Indeed, it is known that most states in the Hilbert space are maximally entangled, and that generic quantum evolutions will eventually lead to an almost maximally entangled state [10,13–15]. This happens even under quantum quench with an integrable Hamiltonian [16].

What is in the entanglement, which is not the entanglement entropy, that tells us whether a quantum evolution is reversible or not? The answer lies in the statistics of the levels $\{p_k\}$ in the entanglement spectrum. We have computed the entanglement spectrum of the qubit string at the end of the heating period. The spectrum is obtained from the singular values resulting from the Schmidt decomposition of the quantum state upon bipartitioning of the qubit string in the middle (i.e., $n_A = n_B = n/2$). The spectrum is first unfolded to yield a constant density before the statistical analysis is performed (see Supplemental Material [17]). In Fig. 3(a) we show the distribution of the spacings between adjacent singular values for reversible cases (heating period performed with \mathcal{I}_2 gates) and irreversible ones (heating period performed with \mathcal{I}_3 gates). The difference is striking: while the data points for the irreversible case match quite closely the distribution of




FIG. 3 (color online). (a) The distribution of the spacing between consecutive unfolded singular values obtained from a bipartition at the middle of a $n = 16$ qubit string at the end of the heating period ($M_h = 512$). Solid black circles are for two-qubit gate heating and open red squares are for heating with a mixture of two-qubit and Toffoli gates. Solid green line: GOE prediction. Dashed blue line: Poisson distribution. Dotted-dashed maroon line: Semi-Poisson distribution. Inset: The tail of the distributions. (b) The average spectral rigidity $\Delta_3(L)$ obtained from the same spectra (a linear fit is used for the semi-Poisson line). A total of 5000 realizations were used to compute the averages.

spacings of the Gaussian orthogonal ensemble (GOE) of random matrices [18], the data points for the reversible case show a weaker repulsion and follow the so-called semi-Poisson statistics, which has been proposed for the energy spectra of systems at metal-insulator transitions [19]. The difference in behavior is also manifest in the spectral rigidity function $\Delta_3(L)$, which measures, for a given interval L , the least-square deviation of the spectral staircase from the best-fitting straight line [20]. In Fig. 3(b), long-range correlations are much stronger in the irreversible case, with the data points also falling close to the GOE prediction. For the reversible case, the singular values are much less correlated and the spectrum much less rigid. This indicates that the statistical fluctuations of the entanglement spectra of irreversible systems are similar to those observed in the energy spectrum of the so-called quantum chaotic systems [2].

In Fig. 4 we also show the entanglement level statistics for the particular case when one starts with initial factorized states of the form $|\chi^{(k)}\rangle = |\psi_1^{(k)}\rangle \otimes |\psi_2^{(k)}\rangle \otimes \cdots \otimes |\psi_n^{(k)}\rangle$, $k = 1, 2$, where $|\psi_1^{(1)}\rangle = |0\rangle$ and $|\psi_j^{(1)}\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$ for $j = 2, \dots, n$, and $|\psi_j^{(2)}\rangle = (|0\rangle - |1\rangle)/\sqrt{2}$ for $j = 1, \dots, n$. We evolve these $n = 16$ -bit states with




FIG. 4 (color online). The statistical fluctuations of the entanglement spectrum for initial product states $|\chi^{(1)}\rangle$ (orange empty squares) and $|\chi^{(2)}\rangle$ (blue full circles) entangled with a mixture of two-qubit and Toffoli gates ($n = 16$, $M_h = 512$). The solid green line is the GOE prediction. (a) Distribution of unfolded singular value spacings. Inset: Distribution tails. (b) Average spectral rigidity $\Delta_3(L)$. Statistical averages were performed over 5000 realizations.

$M = 512$ gates chosen randomly from the set \mathcal{I}_3 . The data in Fig. 4 clearly conform to the GOE statistics, and we observe that the disentangling algorithm again fails, indicating that reversing the computation is extremely difficult.

In quantum mechanics, irreversibility, chaos, nonintegrability and thermalization are phenomena often associated with one another. Unfortunately, some of these notions are ill defined, such as integrability and lack thereof, and the associations are either weak or plagued by counterexamples. For instance, irreversibility can be associated to both chaotic and nonchaotic Hamiltonians [3,4] and there are nonintegrable systems that do not thermalize [21]. Moreover, some of these concepts are only defined in the context of time-independent Hamiltonian evolutions. For instance, the energy levels of a chaotic Hamiltonian show Wigner-Dyson statistics.

In this Letter we presented an alternative approach to the question of irreversibility and complex behavior in quantum systems that works purely at the wave-function level. We did so by studying the eigenvalues of the reduced density matrix of a subsystem, the so-called entanglement spectrum [22]. We showed that (i) a disentangling Metropolis algorithm provides a firm notion of reversibility, namely, the evolution can be inverted if the state can be disentangled, and (ii) irreversibility arises when the level statistics of the entanglement spectrum of a subsystem is Wigner-Dyson.

On the other hand, in the example we studied where the spectrum did not follow Wigner-Dyson statistics, we were always capable of reverting the evolution, even with zero knowledge about the quantum circuit. It is remarkable that the length of the reverted circuit does not depend on the length of the initial circuit, as long as the maximum entanglement entropy is reached. In the disentangling algorithm, we obtained similar results with the Rényi entropy S_2 . This is remarkable because S_2 is an observable that can be measured [23,24], for example, in optical lattices with ultracold atomic gases [25].

The results of this work motivate several questions and applications. The method presented here is applicable to any kind of quantum evolution, regardless of whether it comes from a quantum circuit, a time-dependent or -independent Hamiltonian system, or an open quantum system. First of all, we can examine the behavior of the entanglement level spacing statistics in integrable Hamiltonian models both in the ground state or during the time evolution after a quantum quench. Using techniques such as the density matrix renormalization group, one can study these models once integrability is broken. We believe that our approach can shed new light on the notion of integrability and lack thereof in quantum systems. The possibility of studying quantum systems away from equilibrium and their universal properties in dynamical phase transitions [26] and many-body localization [27–29] is another feature of the method that only involves wave functions. Similarly, we can study the behavior of the entanglement level spacing statistics at critical points of integrable and nonintegrable systems [30,31]. The adiabaticity of time-dependent quantum processes [32] can be examined under the lens of the entanglement spectrum as well, with potential applications to adiabatic quantum computing. Moreover, one can study how the complexity of the entanglement spectrum is related to quantum algorithms capable of giving an exponential speedup [33]. Under the same lens of the entanglement spectrum, one should study the typicality of quantum chaos in random states [34,35]. Entanglement is very ubiquitous in the Hilbert space [14,34], and while this feature has been crucial to show the typicality of thermalization in closed quantum systems [36], this also means that entanglement entropy is unable to characterize quantum irreversibility, and the difference between integrable and nonintegrable systems. Our results show that the understanding of complex quantum behavior lies in the statistics of the fluctuations of the entanglement level spacing.

We acknowledge financial support from the U.S. National Science Foundation through Grants No. CCF 1116590 and No. CCF 1117241, and by the National Basic Research Program of China Grants No. 2011CBA00300, No. 2011CBA00301, the National Natural Science Foundation of China Grants No. 61033001 and No. 61361136003.

- [1] A. Peres, *Phys. Rev. A* **30**, 1610 (1984).
- [2] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics* (Springer Verlag, New York, 1991).
- [3] T. Gorin, T. Prosen, T. H. Seligman, and M. Žnidarič, *Phys. Rep.* **435**, 33 (2006).
- [4] G. Benenti and G. Casati, *Phys. Rev. E* **79**, 025201(R) (2009).
- [5] D. P. DiVincenzo, *Phys. Rev. A* **51**, 1015 (1995).
- [6] E. Fredkin and T. Toffoli, *Int. J. Theor. Phys.* **21**, 219 (1982).
- [7] A. Ekert and P. L. Knight, *Am. J. Phys.* **63**, 415 (1995).
- [8] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer Academic, Dordrecht, 1995).
- [9] A. Rényi, in *Proceedings of the 4th Berkeley Symposium on Mathematical Statistics and Probability* (University of California Press, Berkeley, 1961), Vol. **1**, p. 547.
- [10] A. Hamma, S. Santra, and P. Zanardi, *Phys. Rev. Lett.* **109**, 040502 (2012).
- [11] C. Chamon and E. R. Mucciolo, *Phys. Rev. Lett.* **109**, 030503 (2012).
- [12] The gates used in \mathcal{I}_3 are just a subgroup of the full unitary group and are not universal. A circuit comprising a universal set of gates for quantum computation—such as CNOT and general one-qubit rotations—produces states that also fail to be disentangled. A systematic study of the full unitary group will be presented elsewhere.
- [13] D. N. Page, *Phys. Rev. Lett.* **71**, 1291 (1993).
- [14] J. Emerson *et al.*, *Science* **302**, 2098 (2003).
- [15] W. G. Brown, Y. S. Weinstein, and L. Viola, *Phys. Rev. A* **77**, 040303(R) (2008).
- [16] P. Calabrese and J. Cardy, *J. Stat. Mech.* (2005) P04010.
- [17] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.112.240501> for a detailed description of the unfolding procedure.
- [18] M. L. Mehta, *Random Matrices* (Academic Press, Amsterdam, 2004), 3rd ed.
- [19] E. B. Bogomolny, U. Gerland, and C. Schmit, *Phys. Rev. E* **59**, R1315 (1999).
- [20] F. J. Dyson and M. L. Mehta, *J. Math. Phys. (N.Y.)* **4**, 701 (1963).
- [21] C. Gogolin, M. P. Müller, and J. Eisert, *Phys. Rev. Lett.* **106**, 040401 (2011).
- [22] Many authors use this terminology for the logarithms of such eigenvalues, particularly in the context of entangling Hamiltonians in condensed matter systems.
- [23] P. Horodecki and A. Ekert, *Phys. Rev. Lett.* **89**, 127902 (2002).
- [24] D. A. Abanin and E. Demler, *Phys. Rev. Lett.* **109**, 020504 (2012).
- [25] M. Greiner, O. Mandel, T. Esslinger, T. W. Hansch, and I. Bloch, *Nature (London)* **415**, 39 (2002).
- [26] A. Polkovnikov, A. K. Sengupta, A. Silva, and M. Vengalattore, *Rev. Mod. Phys.* **83**, 863 (2011).
- [27] D. M. Basko, I. L. Aleiner, and B. L. Altshuler, *Ann. Phys. (Amsterdam)* **321**, 1126 (2006).
- [28] A. Pal and D. A. Huse, *Phys. Rev. B* **82**, 174411 (2010).
- [29] S. Iyer, V. Oganesyan, G. Refael, and D. A. Huse, *Phys. Rev. B* **87**, 134202 (2013).
- [30] A. Lakshminarayan and V. Subrahmanyam, *Phys. Rev. A* **71**, 062334 (2005).
- [31] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, *Rev. Mod. Phys.* **80**, 517 (2008).
- [32] H. T. Quan and W. H. Zurek, *New J. Phys.* **12**, 093025 (2010).
- [33] K. Maity and A. Lakshminarayan, *Phys. Rev. E* **74**, 035203 (R) (2006).
- [34] S. J. Szarek, E. Werner, and K. Życzkowski, *J. Phys. A* **44**, 045303 (2011).
- [35] Vinayak and M. Žnidarič, *J. Phys. A* **45**, 125204 (2012).
- [36] S. Popescu, A. J. Short, and A. Winter, *Nat. Phys.* **2**, 754 (2006).