



Slow Kinetics of Brownian Maxima

E. Ben-Naim¹ and P. L. Krapivsky²

¹Theoretical Division and Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

²Department of Physics, Boston University, Boston, Massachusetts 02215, USA

(Received 3 May 2014; published 17 July 2014)

We study extreme-value statistics of Brownian trajectories in one dimension. We define the maximum as the largest position to date and compare maxima of two particles undergoing independent Brownian motion. We focus on the probability $P(t)$ that the two maxima remain ordered up to time t and find the algebraic decay $P \sim t^{-\beta}$ with exponent $\beta = 1/4$. When the two particles have diffusion constants D_1 and D_2 , the exponent depends on the mobilities, $\beta = (1/\pi) \arctan \sqrt{D_2/D_1}$. We also use numerical simulations to investigate maxima of multiple particles in one dimension and the largest extension of particles in higher dimensions.

DOI: 10.1103/PhysRevLett.113.030604

PACS numbers: 05.40.Jc, 02.50.Cw, 02.50.Ey, 05.40.Fb

Consider a pair of particles undergoing independent Brownian motion in one dimension [1]. These two particles do not meet with a probability that decays as $t^{-1/2}$ in the long-time limit. This classical first-passage behavior holds for Brownian particles with arbitrary diffusion constants. It holds even for particles undergoing symmetric Lévy flights [2,3] and has numerous applications [3,4]. Here, we generalize this ubiquitous first-passage behavior to maxima of Brownian particles. Figure 1 shows that the maximal position of each particle forms a staircase, and it illustrates that unlike the position, the maximum is a non-Markovian random variable [5,6]. We find that two such staircases do not intersect with probability P that is inversely proportional to the one-fourth power of time, $P \sim t^{-1/4}$, in the long-time limit. If the particles move with diffusion constants D_1 and D_2 , the two maxima remain ordered during the time interval $(0, t)$ with the slowly decaying probability

$$P \sim t^{-\beta} \quad \text{where} \quad \beta = \frac{1}{\pi} \arctan \sqrt{\frac{D_2}{D_1}}. \quad (1)$$

In this Letter, we obtain this result analytically and investigate numerically related problems involving multiple maxima and diffusion in higher dimensions.

Anomalous relaxation with nontrivial persistence exponents [7–9], enhanced transport due to disorder [10,11], and anomalous diffusion due to exclusion [12,13] are dynamical phenomena that were recently demonstrated in experiments involving Brownian particles. Understanding the nonequilibrium statistical physics of these diffusion processes is closely intertwined with the characteristic behavior of extreme fluctuations and the statistics of extreme values [14–19].

We first establish Eq. (1) for two Brownian particles having the same diffusion constant D . Let us denote the positions of the particles at time t by $x_1(t)$ and $x_2(t)$, and without loss of generality, we assume $x_1(0) > x_2(0)$. We

define the maximum of the first particle $m_1(t)$ to be its rightmost position up to time t ; similarly, $m_2(t)$ is the maximal position of the second particle. Our goal is to find the probability $P(t)$ that the two maxima remain ordered $m_1(\tau) > m_2(\tau)$ for all $0 \leq \tau \leq t$.

The two maxima remain ordered if and only if $m_1(\tau) > x_2(\tau)$ at all times $0 \leq \tau \leq t$. Hence, to find P , there is no need to keep track of the maximum m_2 , and it suffices to consider only the position x_2 . As a further simplification, we focus on the distance of each particle from the maximum m_1 and introduce the variables

$$u = m_1 - x_1 \quad \text{and} \quad v = m_1 - x_2. \quad (2)$$

By definition, both distances are positive, $u \geq 0$ and $v \geq 0$. The transformation (2) maps the four variables (two positions and two maxima) onto the two relevant variables (two distances). Since the positions x_1 and x_2 undergo simple diffusion, the distances u and v also undergo simple diffusion in the domain $u > 0$ and $v > 0$. Hence, the probability density $\rho(u, v, t)$ obeys the diffusion equation




FIG. 1 (color online). Space-time diagram of the positions (thin lines) and the ordered maxima (thick lines) of two Brownian particles.

$\partial_u \rho = D \nabla^2 \rho$ with $\nabla^2 = \partial_u^2 + \partial_v^2$ along with the boundary conditions $\rho|_{v=0} = 0$ and $(\partial_u - \partial_v)\rho|_{u=0} = 0$. The boundary $v = 0$ is absorbing so that position x_2 does not exceed maximum m_1 . The second boundary condition (see Supplemental Material [20] for derivation) guarantees that there is no current through the line $u = 0$, and it also takes into account the upward drift along the boundary $u = 0$. Indeed, when the maximum increases, $m_1 \rightarrow m_1 + \delta m$, one distance remains the same, $u = 0$, but the second distance increases, $v \rightarrow v + \delta m$ (Fig. 2).

The probability $P(t)$ is the integral of the probability density, $P(t) = \int_0^\infty \int_0^\infty dudv \rho(u, v, t)$. This quantity equals the survival probability of a “composite” particle with coordinates (u, v) that is undergoing Brownian motion in two dimensions. This composite particle starts somewhere along the boundary $u = 0$, and it diffuses in the domain $u > 0$ and $v > 0$. The particle slips along the boundary $u = 0$, but it is annihilated when it reaches the boundary $v = 0$ (Fig. 2).

In general, the probability P depends on the initial coordinates $u(0)$ and $v(0)$. It is convenient to compute the probability P directly rather than through the probability density $\rho(u, v, t)$. With the shorthand notations $X \equiv u(0)$ and $Y \equiv v(0)$, the probability $P \equiv P(X, Y, t)$ obeys the standard diffusion equation [21,22]

$$\frac{\partial P}{\partial t} = D \nabla^2 P \quad (3)$$

when $X > 0$ and $Y > 0$, where $\nabla^2 = \partial_X^2 + \partial_Y^2$ is the Laplace operator. The initial condition is $P = 1$ in the region $X \geq 0$ and $Y > 0$, and the boundary conditions are $P|_{Y=0} = 0$ and $(\partial_X + \partial_Y)P|_{X=0} = 0$. The former reflects that the boundary $Y = 0$ is absorbing, and the second is a consequence of the upward drift (see Supplemental Material [20] for details). Our problem corresponds to the special case $X = 0$ and $Y = x_1(0) - x_2(0)$.




FIG. 2 (color online). Sample trajectory of the composite particle. There is upward drift along the boundary $u = 0$ and the boundary $v = 0$ is absorbing.

In terms of the polar coordinates $R = \sqrt{X^2 + Y^2}$ and $\theta = \arctan(Y/X)$, the probability $P \equiv P(R, \theta, t)$ obeys the diffusion equation (3) with

$$\nabla^2 = \frac{\partial^2}{\partial R^2} + \frac{1}{R} \frac{\partial}{\partial R} + \frac{1}{R^2} \frac{\partial^2}{\partial \theta^2}.$$

The first boundary condition is simply $P|_{\theta=0} = 0$. The second boundary condition becomes

$$\left(R \frac{\partial P}{\partial R} - \frac{\partial P}{\partial \theta} \right) \Big|_{\theta=\pi/2} = 0, \quad (4)$$

where we have utilized $\partial_X = \cos \theta \partial_R - R^{-1} \sin \theta \partial_\theta$ and $\partial_Y = \sin \theta \partial_R + R^{-1} \cos \theta \partial_\theta$.

To solve for the probability P , we first note that both the diffusion equation (3) and the boundary condition (4) are invariant under the scaling transformation $(X, Y) \rightarrow (\alpha X, \alpha Y)$ and $t \rightarrow \alpha^2 t$. As a result, the quantity $R^2/(Dt)$ is the only dimensionless combination of the variables R, D, t . Of course, the probability P is dimensionless, and we thus expect $P(R, \theta, t) \sim (R^2/Dt)^\beta f(\theta)$ in the long-time limit. We now substitute this expression into (3) and observe that the left-hand side vanishes in the long-time limit. Consequently, the function f obeys $f'' + (2\beta)^2 f = 0$. Next, we choose the solution $f(\theta) = \sin(2\beta\theta)$ to satisfy the boundary condition $S|_{\theta=0} = 0$. We, thus, find (see, also, [22])

$$P(R, \theta, t) \sim \left(\frac{R^2}{Dt} \right)^\beta \sin(2\beta\theta) \quad (5)$$

in the limit $t \rightarrow \infty$. Next, we substitute (5) into (4) and observe that the second boundary condition is obeyed when $\tan(\beta\pi) = 1$. Thus, $\beta = 1/4$, and we arrive at the slow kinetics (Fig. 3)

$$P \sim t^{-1/4}. \quad (6)$$




FIG. 3 (color online). The probability P versus time t . Shown are Monte Carlo simulation results obtained from 10^8 independent realizations (circles). Also shown as reference (line) is the theoretical prediction (6).

Importantly, the decay exponent is an eigenvalue of the angular component of the Laplace operator, and it is specified by the boundary conditions. We note that the behavior (6) also characterizes the probability that a particle diffusing on a plane avoids a semi-infinite needle [23] and that similar kinetics are found for diffusion in shear flows [24,25] and random acceleration processes [26,27].

Consider now the general case where the two particles have diffusion constants D_1 and D_2 . The transformation $(x_1, x_2) \rightarrow (\hat{x}_1, \hat{x}_2)$ with $(\hat{x}_1, \hat{x}_2) = (x_1/\sqrt{D_1}, x_2/\sqrt{D_2})$ maps this anisotropic Brownian motion onto isotropic Brownian motion in two dimensions. The maxima are also rescaled, $(m_1, m_2) \rightarrow (\hat{m}_1, \hat{m}_2)$ with $(\hat{m}_1, \hat{m}_2) = (m_1/\sqrt{D_1}, m_2/\sqrt{D_2})$. The two maxima remain ordered, $m_1 > m_2$, as long as

$$\sigma \hat{m}_1 > \hat{m}_2 \quad \text{with} \quad \sigma = \sqrt{D_1/D_2}. \quad (7)$$

Thus, we expect that the exponent depends on the ratio of diffusion constants, $\beta \equiv \beta(D_1/D_2)$. When one particle is immobile, the problem simplifies. If $D_1 = 0$, the maxima remain ordered if the particles do not meet, $x_1(0) > x_2(t)$, and, hence, $P \sim t^{-1/2}$. In the complementary case $D_2 = 0$, an immobile particle cannot overtake a maximum set by a mobile particle and $P = 1$. The limiting values are, therefore, $\beta(0) = 1/2$ and $\beta(\infty) = 0$, and since the exponent should be a monotonic function of the ratio D_1/D_2 , we deduce $0 \leq \beta \leq 1/2$.

The above analysis is straightforward to generalize if instead of (2) we use the distances $u = \hat{m}_1 - \hat{x}_1$ and $v = \sigma \hat{m}_1 - \hat{x}_2$. Again, diffusion takes place in the domain $u > 0$ and $v > 0$, and the boundary conditions are $P|_{Y=0} = 0$ and $(\partial_X + \sigma \partial_Y)P|_{X=0} = 0$. In polar coordinates, the latter boundary condition reads $(\sigma R \partial_R - \partial_\theta)S|_{\theta=\pi/2} = 0$. Using this boundary condition and the probability P given by (5), we deduce $\sigma \tan(\beta\pi) = 1$ and thus, obtain our main result (1).

The exponent is rational for special values of the diffusion constants, for instance, $\beta(1/3) = 1/3$, $\beta(1) = 1/4$, and $\beta(3) = 1/6$. Exponent β varies continuously with the ratio D_1/D_2 (Fig. 4). Unlike the universal first-passage behavior $t^{-1/2}$ characterizing positions of Brownian particles, the behavior of the probability P is not universal. Further, first-passage kinetics of maxima of mobile Brownian particles are generally slower compared with first-passage kinetics of positions since $\beta < 1/2$. As expected, the limiting values are $\beta(0) = 1/2$ and $\beta(\infty) = 0$, and the limiting behaviors are $1/2 - \beta \approx \frac{1}{\pi} \sqrt{D_1/D_2}$ when $D_1 \ll D_2$ and $\beta \approx \frac{1}{\pi} \sqrt{D_2/D_1}$ for $D_2 \ll D_1$.

When the diffusion constants differ, there are two separate exponents $\beta(D_1/D_2)$ and $\beta(D_2/D_1)$. Equation (1) together with the trigonometric identity $\arctan(x) + \arctan(1/x) = \pi/2$ give the relationship [28]

$$\beta\left(\frac{D_1}{D_2}\right) + \beta\left(\frac{D_2}{D_1}\right) = \frac{1}{2}. \quad (8)$$




FIG. 4 (color online). The exponent β versus the ratio D_1/D_2 . The line corresponds to the theoretical curve (1) and the dots to results of Monte Carlo simulations with 10^7 independent realizations.

Let β_{\max} be the larger of these two exponents and β_{\min} be the smaller one. When the particles have different mobilities, there are two different relaxation processes that govern lead changes between two Brownian maxima: the faster relaxation $P \sim t^{-\beta_{\max}}$ occurs when a particle tries to overtake the maxima set by a less mobile particle, and the slower relaxation $P \sim t^{-\beta_{\min}}$ occurs in the complementary case. The two particles keep exchanging the lead, and interestingly, this process exhibits alternating kinetics: slow—fast—slow—fast *ad infinitum*.

One anticipates that the asymptotic behavior (1) applies to a broad class of diffusion processes. As a test, we performed Monte Carlo simulations (see also Refs. [29,30]) of discrete-time random walks in one dimension with two different implementations: (i) a random walk on a lattice where all step lengths have the same size, and (ii) a random walk on a line where the step lengths are chosen from a uniform distribution with compact support. In both cases, the simulation results are in excellent agreement with the theoretical predictions. The simulation results shown in Figs. 3 and 4 correspond to random walks on a line.

For n particles undergoing Brownian motion, there are three natural generalizations of the probability P . First is the probability A_n that all n maxima remain perfectly ordered, that is, n staircases as in Fig. 1 never intersect; for positions of Brownian particles, this problem dates back to [31]. Second is the probability B_n that the rightmost staircase is never overtaken; the corresponding problem for positions was studied in [32]. Third is the probability C_n that the leftmost staircase never overtakes another maxima [33]. We expect all three quantities to decay as power laws,

$$A_n \sim t^{-\alpha_n}, \quad B_n \sim t^{-\beta_n}, \quad C_n \sim t^{-\gamma_n}, \quad (9)$$

with exponents α_n , β_n , and γ_n that depend on the number of particles n . Table I lists results of Monte Carlo

TABLE I. The exponents α_n , β_n , and γ_n defined in Eq. (9) versus the number of particles n (all particles have the same diffusion constant D). Also shown as a reference are values for the corresponding exponents a_n , b_n , and c_n characterizing analogous probabilities involving the positions of n Brownian particles [22]. The only known [31] sequence of exponents is $a_n = n(n-1)/4$.

n	Maxima			Positions		
	α_n	β_n	γ_n	a_n	b_n	c_n
2	1/4	1/4	1/4	1/2	1/2	1/2
3	0.653	0.432	0.334	3/2	3/4	3/8
4	1.13	0.570	0.376	3	0.91342	0.306
5	1.60	0.674	0.401	5	1.02	0.265
6	2.01	0.759	0.418	15/2	1.11	0.234

simulations along with the analogous exponents for the positions, rather than the maxima [22].

All of the exponents are directly related to eigenvalues of the Laplace operator in high-dimensional space with suitable boundary conditions. Even for the simpler case of ordered positions, such eigenvalues are generally unknown (Table I). We expect that $\alpha_n > \beta_n > \gamma_n$ and, furthermore, that all three exponents increase with n . Further, it is possible to justify the behavior $\beta_n \approx b_n$ and consequently [22,33] obtain the logarithmic growth $\beta_n \approx \frac{1}{4} \ln n$ when the number of particles is large, $n \rightarrow \infty$. Also, it is simple to show that $\gamma_n \rightarrow 1/2$ in the limit $n \rightarrow \infty$ [19]. Based on the numerical results, we conjecture that one of the exponents is rational, $\gamma_n = (n-1)/2n$; this form is consistent with $\gamma_1 = 0$ and $\gamma_2 = 1/4$.

Our results thus far concern diffusion in one spatial dimension, yet closely related questions can be asked of Brownian motion in arbitrary dimension d . Consider, for example, the maximum distance traveled by a Brownian particle. If the particle starts at the origin, this distance equals the radial coordinate in a spherical coordinate system. We expect that the probability U_d that the maximal radial coordinate of one particle always exceeds that of another particle decays algebraically with time, $U_d \sim t^{-\nu_d}$. Our numerical simulations show that exponent ν grows rather slowly with dimension d

$$\nu_1 = 0.563, \quad \nu_2 = 0.602, \quad \nu_3 = 0.630. \quad (10)$$

It would also be interesting to study planar Brownian excursions and, in particular, the probability that the convex hull generated by one particle always contains that of a second particle [34,35].

We also mention that the first-passage process studied in this Letter is equivalent to a ‘‘competition’’ between two records [36]. As a data analysis tool, the first-passage probability P is a straightforward measure and can be used in finance [37], climate [38,39], and earthquakes [40,41]. The notion of competing maxima could also describe the

span of colloidal particles undergoing simple or anomalous diffusion [10,12].

In summary, we studied maxima of Brownian particles in one dimension and found that the probability that such maxima remain ordered decays as a power law with time. The exponent characterizing this decay varies continuously with the diffusion coefficients governing the motion of the particles. When there are two particles, the problem reduces to diffusion in two dimensions with mixed boundary conditions. Recent studies show that the eigenvalues characterizing ordering of a very large number of Brownian trajectories obey scaling laws in the thermodynamic limit [22], and an interesting open challenge would be to use such scaling methods to elucidate extreme-value statistics of many Brownian trajectories.

We acknowledge the DOE Grant No. DE-AC52-06NA25396 for support (E. B.).

-
- [1] P. Mörders and Y. Peres, *Brownian Motion* (Cambridge University Press, Cambridge, England, 2010).
 - [2] E. Sparre Andersen, *Math. Scand.* **1**, 263 (1953); **2**, 195 (1954).
 - [3] W. Feller, *An Introduction to Probability Theory and Its Applications* (Wiley, New York, 1968).
 - [4] S. Redner, *A Guide to First-Passage Processes* (Cambridge University Press, Cambridge, England, 2001).
 - [5] P. Lévy, *Processus Stochastiques et Mouvement Brownien* (Gauthier-Villars, Paris, 1948).
 - [6] K. Itô and H. P. McKean, *Diffusion Processes and Their Sample Paths* (Springer, New York, 1965).
 - [7] S. N. Majumdar, C. Sire, A. J. Bray, and S. J. Cornell, *Phys. Rev. Lett.* **77**, 2867 (1996).
 - [8] B. Derrida, V. Hakim, and R. Zeitak, *Phys. Rev. Lett.* **77**, 2871 (1996).
 - [9] Y. Takikawa and H. Orihara, *Phys. Rev. E* **88**, 062111 (2013).
 - [10] G. Coupier, M. Saint Jean, and C. Guthmann, *Europhys. Lett.* **77**, 60001 (2007).
 - [11] E. Ben-Naim and P. L. Krapivsky, *Phys. Rev. Lett.* **102**, 190602 (2009).
 - [12] C. Lutz, M. Kollmann, and C. Bechinger, *Phys. Rev. Lett.* **93**, 026001 (2004).
 - [13] E. Barkai and R. Silbey, *Phys. Rev. Lett.* **102**, 050602 (2009).
 - [14] P. L. Krapivsky, S. Redner, and E. Ben-Naim, *A Kinetic View of Statistical Physics* (Cambridge University Press, Cambridge, England, 2010).
 - [15] A. J. Bray, S. N. Majumdar, and G. Schehr, *Adv. Phys.* **62**, 225 (2013).
 - [16] B. Derrida, J. L. Lebowitz, and E. R. Speer, *Phys. Rev. Lett.* **87**, 150601 (2001).
 - [17] G. Schehr, S. N. Majumdar, A. Comtet, and J. Randon-Furling, *Phys. Rev. Lett.* **101**, 150601 (2008).
 - [18] S. N. Majumdar and R. M. Ziff, *Phys. Rev. Lett.* **101**, 050601 (2008).
 - [19] E. Ben-Naim and P. L. Krapivsky, *J. Phys. A* **47**, 255002 (2014).

- [20] See Supplementary material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.113.030604> for derivation of the boundary conditions.
- [21] G. H. Weiss, *Aspects and Applications of the Random Walk* (North-Holland, Amsterdam, 1994).
- [22] E. Ben-Naim and P. L. Krapivsky, *J. Phys. A* **43**, 495007 (2010); **43**, 495008 (2010).
- [23] D. Considine and S. Redner, *J. Phys. A* **22**, 1621 (1989).
- [24] S. Redner and P. L. Krapivsky, *J. Stat. Phys.* **82**, 999 (1996).
- [25] A. J. Bray and P. Gonos, *J. Phys. A* **37**, L361 (2004).
- [26] H. P. McKean, *Journal of mathematics of Kyoto University* **2**, 227 (1963); M. Goldman, *Ann. Math. Stat.* **42**, 2150 (1971); A. Lachal, *Ann. Inst. Henri Poincaré* **33**, 1 (1997).
- [27] T. W. Burkhardt, *J. Phys. A* **26**, L1157 (1993).
- [28] The first-passage exponent (1) governs Brownian maxima, and the dual exponent $(1/\pi) \arctan \sqrt{D_1/D_2}$ characterizes the image problem involving Brownian minima.
- [29] P. Grassberger, *Comput. Phys. Commun.* **147**, 64 (2002).
- [30] T. Oettel, V. V. Bulatov, A. Donev, M. H. Kalos, G. H. Gilmer, and B. Sadigh, *Phys. Rev. E* **80**, 066701 (2009).
- [31] M. E. Fisher, *J. Stat. Phys.* **34**, 667 (1984); D. A. Huse and M. E. Fisher, *Phys. Rev. B* **29**, 239 (1984).
- [32] M. Bramson and D. Griffeath, in *Random Walks, Brownian Motion, and Interacting Particle Systems: A Festschrift in Honor of Frank Spitzer*, edited by R. Durrett and H. Kesten (Birkhäuser, Boston, 1991).
- [33] P. L. Krapivsky and S. Redner, *J. Phys. A* **29**, 5347 (1996); D. ben-Avraham, B. M. Johnson, C. A. Monaco, P. L. Krapivsky, and S. Redner, *J. Phys. A* **36**, 1789 (2003).
- [34] B. Duplantier, in *Proceedings of Symposia in Pure Mathematics*, edited by M. L. Lapidus and M. van Frankenhuysen (American Mathematical Society, Providence, 2004), Vol. 72, p. 365.
- [35] S. N. Majumdar, A. Comtet, and J. Randon-Furling, *J. Stat. Phys.* **138**, 955 (2010).
- [36] B. C. Arnold, N. Balakrishnan, and H. N. Nagaraja, *Records* (Wiley-Interscience, New York, 1998).
- [37] J.-P. Bouchaud and M. Potters, *Theory of Financial Risk and Derivative Pricing* (Cambridge University Press, Cambridge, England, 2003).
- [38] W. I. Newman, B. D. Malamud, and D. L. Turcotte, *Phys. Rev. E* **82**, 066111 (2010).
- [39] G. Wergen, A. Hense, and J. Krug, *Climate Dynamics* **42**, 1275 (2014).
- [40] R. Shcherbakov, J. Davidsen, and K. F. Tiampo, *Phys. Rev. E* **87**, 052811 (2013).
- [41] E. Ben-Naim and P. L. Krapivsky, *Phys. Rev. E* **88**, 022145 (2013); P. W. Miller and E. Ben-Naim, *J. Stat. Mech.* (2013) P10025.