

Resonances in Coupled πK - ηK Scattering from Quantum Chromodynamics

Jozef J. Dudek,^{1,2,*} Robert G. Edwards,¹ Christopher E. Thomas,³ and David J. Wilson²

(for the Hadron Spectrum Collaboration)

¹Theory Center, Jefferson Lab, 12000 Jefferson Avenue, Newport News, Virginia 23606, USA

²Department of Physics, Old Dominion University, Norfolk, Virginia 23529, USA

³DAMTP, University of Cambridge, Cambridge CB3 0WA, United Kingdom

(Received 17 June 2014; revised manuscript received 15 July 2014; published 30 October 2014)

Using a first-principles calculation within quantum chromodynamics, we are able to determine a pattern of strangeness = 1 resonances that appear as complex singularities within coupled πK - ηK scattering amplitudes. We make use of numerical computation in the lattice discretized approach to the quantum field theory with light quark masses corresponding to $m_\pi \sim 400$ MeV and at a single lattice spacing. The energy dependence of scattering amplitudes is extracted through their relationship to the discrete spectrum in a finite volume, which we map out in unprecedented detail.

DOI: 10.1103/PhysRevLett.113.182001

PACS numbers: 14.40.Df, 12.38.Gc, 13.25.Es

Introduction.—Understanding how excited hadrons are built up from the basic quark and gluon degrees of freedom of quantum chromodynamics (QCD) remains a challenging problem. QCD should be able to explain the apparent regularity of the experimental spectrum, but its structure suggests additional “exotic” quark-gluon configurations, such as those in which the gluonic field is excited, which to date have not been unambiguously observed in experiment. An important feature of the theoretical challenge is that excited hadrons appear in experiment as resonances, which ultimately decay into pseudoscalar mesons, themselves complex aggregations of confined quarks, antiquarks, and gluons. To study resonances within QCD, one must explore the behavior of the theory’s hadron scattering amplitudes at low energy, a field of study that is still in its infancy.

Lattice QCD offers us an *ab initio* numerical approach to QCD calculations. By discretizing the quark and gluon fields on a finite lattice and by sampling possible space-time configurations of the gluon field through a Monte Carlo method, we can compute correlation functions with the quantum numbers of hadrons. These correlators contain information about the spectrum and interactions of hadrons.

Following Lüscher [1], a formalism has been derived that relates infinite-volume hadron scattering amplitudes to the discrete spectrum of hadron states in a finite volume. Through the computation of statistically precise excited state spectra in lattice QCD, we can extract the energy dependence of hadron scattering amplitudes and examine their resonant content. In a recent example considering the elastic scattering of two pions [2], following computation of the appropriate lattice QCD spectra, the scattering amplitude was determined at 32 discrete energy values

in a 300 MeV range. This determination demonstrated unambiguously the presence of the ρ resonance, whose mass and width were extracted.

Unlike the ρ resonance, most known hadron resonances do not decay into only one final state, but instead are enhancements in coupled channels. The recently derived formalism to extract coupled-channel scattering amplitudes from finite-volume spectra is somewhat more involved than the elastic case [3,4]. In this Letter, we will present the first application of this formalism to QCD, for the particular case of πK - ηK coupled-channel scattering.

Experimentally, a number of low-lying resonances appear in isospin-1/2 πK scattering: the $J^P = 1^-$, $K^*(892)$, the strange analog of the ρ meson, is a narrow elastic resonance, the $J^P = 0^+$, $K_0^*(1430)$ appears as a relatively broad resonance, and the $J^P = 2^+$, $K_2^*(1430)$, at a similar mass value, is a much narrower resonance [5]. Furthermore, it is observed that in even- J partial-waves, resonances decay dominantly to πK and not to ηK .

Resonances can be rigorously defined as pole singularities in scattering amplitudes when they are considered to be functions of *complex* values of the scattering energy. Experimentally, the scattering amplitudes are determined for real values of the energy above kinematic thresholds, which may then be parametrized by analytic functions that can be continued into the complex plane. One important application of this procedure concerns the $J^P = 0^+$ πK scattering amplitude at low energy, where the use of dispersion relations provides a particularly strong constraint when describing the available experimental scattering data, leading to a pole far from the real axis known as the κ pole [6].

In this Letter, we will present scattering amplitudes extracted from lattice QCD determinations of the excited

hadron spectrum for the $J^P = 0^+, 1^-, 2^+$ coupled πK - ηK channels. In this first calculation, we work with light quark masses in the QCD Lagrangian somewhat heavier than those known physically, meaning that initially our results can only be compared qualitatively with the experimental situation. We find, as in experiment, an approximate decoupling of the πK - ηK channels and a rapid increase of the πK $J^P = 0^+$ scattering phase shift above threshold, followed by a slower rise through 90° at higher energies. With light quark masses somewhat heavier than the physical case, the lightest vector resonance becomes bound and the κ pole appears to manifest itself as a “virtual bound state.” Behavior consistent with a narrow resonance is observed in the $J^P = 2^+$ channel.

Previous lattice QCD studies limited to elastic πK scattering in isospin-1/2 appear in Refs. [7,8]. The authors of Ref. [8], in calculations in a small volume with lower light quark masses compared to this study and without dynamical strange quarks, found a K^* above the πK threshold and extracted resonance parameters.

Finite-volume spectrum from lattice QCD.—We make use of the dynamical anisotropic lattices initially described in Ref. [9], which feature two degenerate light quarks (u, d) plus a heavier flavor with its mass tuned to approximate the physical strange quark. In this study, the light quark mass parameter is increased with respect to its physical value such that the pion mass is 391 MeV, the kaon mass is 549 MeV, and the η mass is 589 MeV. Three spatial volumes are utilized: 16^3 (~ 2 fm), 20^3 (~ 2.5 fm), and 24^3 (~ 3 fm), with a spatial lattice spacing $a_s \sim 0.12$ fm.

The discrete spectrum of hadron states can be extracted from two-point correlation functions $\langle 0 | \mathcal{O}_i(t) \mathcal{O}_j^\dagger(0) | 0 \rangle$. We make use of two classes of interpolating field \mathcal{O}_i^\dagger ; the first are “single-meson” operators, those which resemble a $q\bar{q}$ construction $\bar{\psi}\Gamma\psi$, where Γ is one of a large number of operators in spin, color, and position space [10], while the second resembles a pair of mesons with definite relative and total momentum $(\bar{\psi}\Gamma_1\psi)_{\vec{p}_1}(\bar{\psi}\Gamma_2\psi)_{\vec{p}_2}$ [11]. In an $L \times L \times L$

spatial volume with periodic boundary conditions, the momenta that appear in these constructions are quantized in integer multiples of $2\pi/L$: $\vec{p} = (2\pi/L) \times (n_x, n_y, n_z)$. More details of the constructions and a demonstration of their efficacy can be found in Ref. [11].

For this study, we construct a basis of operators including “meson-meson” constructions with quark flavors corresponding to πK and ηK . A matrix of correlation functions is built [12] using a large number of single-meson operators along with several relative momentum constructions of both πK -like and ηK -like operators. A spectrum of states that is best in a variational sense can be extracted from this matrix by solving a generalized eigenvalue problem [13].

The lattice has a cubic symmetry, rather than that of the full rotational group, so instead of being characterized by integer spins, independent spectra lie in a finite number of irreducible representations, or “irreps” Λ , of the reduced symmetry. For a meson system with nonzero total momentum \vec{P} , the symmetry group is classified by irreps of rotations that leave \vec{P} invariant [10]. We compute the excited state spectrum in all relevant irreps for $|\vec{P}|^2 \leq 4(2\pi/L)^2$ and find we can determine them with a high degree of statistical precision in all cases. In Fig. 1 we show two examples. Note that owing to the use of the variational method of state extraction, there is a robust extraction of near-degenerate states, even high in the spectrum.

If mesons had no residual strong interactions, the discrete spectrum in a finite volume would be predictable using the relativistic dispersion relation $E = \sqrt{m_1^2 + |\vec{p}_1|^2} + \sqrt{m_2^2 + |\vec{p}_2|^2}$ with the allowed quantized momentum values given above. These volume-dependent “noninteracting” energy levels are shown by the continuous curves in Fig. 1, and the fact that the energy levels extracted from our lattice QCD calculations do not lie on these curves is an indication that there are interactions between mesons. If these interactions are strong enough, there may be resonant behavior. To explore this rigorously, we must determine the meson-meson scattering amplitudes.




FIG. 1 (color online). Discrete spectrum of c.m.-frame energies obtained from variational analysis of correlation matrices featuring single-meson and meson-meson operators at three spatial volumes. Red bands are πK noninteracting level positions, and green bands are ηK noninteracting level positions. (a) $\vec{P} = [000]$, $\Lambda = A_1^+$ spectrum (dominated by $J^P = 0^+$ with negligible contributions from $J \geq 4$). (b) $\vec{P} = [001]$, $\Lambda = A_1$ spectrum ($J^P = 0^+, 1^-, 2^+$ all contribute).

Scattering amplitudes.—For an interacting quantum field theory, the relationship between the discrete spectrum of states in an $L \times L \times L$ volume, in a frame moving with momentum \vec{P} , for irrep Λ , can be written in a simplified form as

$$\det[\delta_{ij}\delta_{JJ'} + i\rho_i t_{ij}^{(J)}(E_{c.m.})(\delta_{JJ'} + i\mathcal{M}_{JJ'}^{\vec{P}\Lambda}(p_i L))] = 0,$$

where the scattering t matrix for partial-wave J appears along with the phase space in channel i , $\rho_i(s) = 2p_i/E_{c.m.}$, and the known volume-dependent functions \mathcal{M} [3,4,14]. Given knowledge of the energy dependence of the scattering amplitudes $t_{ij}^{(J)}(E_{c.m.})$, one can solve this equation for a discrete spectrum of states $\{E_{c.m.}\}$. The practical problem at hand, however, is the reverse of this: to find the t matrix given a lattice QCD calculation of the spectrum. For any single energy level value $E_{c.m.}$, this is an underconstrained problem, as there are multiple elements of the t matrix to be determined from only one condition.

The approach we will take is to parametrize the energy dependence of the t matrix and describe the spectrum as a whole. Such an approach was explored in the context of a toy model of coupled-channel scattering in Ref. [3]. A flexible K -matrix parametrization of partial-wave J , in terms of the variable $s = E_{c.m.}^2$, can be constructed that ensures the unitarity of the S matrix,

$$t_{ij}^{-1}(s) = \frac{1}{(2p_i)^J} K_{ij}^{-1}(s) \frac{1}{(2p_j)^J} + I_{ij}(s),$$

$$K_{ij}(s) = \sum_p \frac{g_i^{(p)} g_j^{(p)}}{m_p^2 - s} + \sum_n \gamma_{ij}^{(n)} s^n,$$

where we may choose how many poles and what order polynomial to include in K , with real parameters $g_i^{(p)}$, m_p , $\gamma_{ij}^{(n)}$. The function $I(s)$ must be chosen such that

$\text{Im}I_{ij}(s) = -\delta_{ij}\rho_i(s)$ above threshold to ensure unitarity is preserved. There is some freedom in the choice of the real part; we choose an implementation of the Chew-Mandelstam form [3], which has smooth behavior across kinematic thresholds.

We use 80 levels from 20 irreps to constrain the 0^+ amplitudes from slightly below πK threshold up to 1650 MeV, 19 levels to constrain the 1^- amplitude in the region around the πK threshold, and a further 24 levels to constrain the 2^+ amplitudes between 1250 and 1700 MeV. The 0^+ , 2^+ partial waves are described by a single K -matrix pole coupled to both πK and ηK plus a constant matrix, while the πK threshold region in 1^- is described by a relativistic Breit-Wigner form. We assume that the influence of partial waves $J \geq 3$ is negligible in this energy region.

The resulting t matrices are plotted in Fig. 2. For 0^+ , 2^+ , πK and ηK phase shifts and an inelasticity, defined in $t_{ii} = (\eta e^{2i\delta_i} - 1)/(2i\rho_i)$, $t_{ij} = [\sqrt{1 - \eta^2} e^{i(\delta_i + \delta_j)}] / (2\sqrt{\rho_i \rho_j})$, for channels $i = \pi K, \eta K$, are shown, while for 1^- we plot the function $p^3 \cot \delta$, which is real and continuous across the πK threshold. In each case, we present the χ^2/N_{dof} for the parameterized description of the input spectrum, which we find to be quite acceptable.

The points shown in the center of Fig. 2(a), which cover the whole energy region plotted, indicate that we are strongly constraining the energy dependence of the amplitudes; in particular, note that the low-energy behavior of the 0^+ πK amplitude is constrained by points at or below threshold. Similarly, in Fig. 2(c), the energy dependence of the 2^+ amplitude is well sampled in the region of the rapid rise of the phase shift. This region is above the πK threshold, which can, in principle, couple to the 2^+ partial-wave—we have assumed here that there is negligible coupling to this channel.




FIG. 2 (color online). (a) $J^P = 0^+$ amplitudes. Open circles on axis show πK and ηK thresholds. Upper panel: πK and ηK phase shifts in degrees. Lower panel: inelasticity. Points in center show the energy levels on three volumes used to constrain the t -matrix extraction; larger solid points show $\vec{P} = 0$, and smaller open circles show $\vec{P} \neq 0$. (b) $J^P = 1^-$ around the πK threshold. Points determined directly without parametrization of the vector amplitude from three volumes: 16^3 (boxes), 20^3 (circles), and 24^3 (triangles). Curve shows the result of a relativistic Breit-Wigner parametrization $p^3 \cot \delta_1 = (m_R^2 - s)[(6\pi\sqrt{s})/g_R^2]$. (c) $J^P = 2^+$ amplitudes. Open circles on axis show πK , ηK , and $\pi\pi K$ thresholds.

We observe that the πK phase shifts in 0^+ , 2^+ rise through 90° suggesting possible resonant behavior, and while this rise is rather slow in the scalar channel, it is rapid in the tensor, indicating a likely narrow resonance. The extracted inelasticities do not deviate significantly from unity, corresponding to an approximate decoupling of the ηK channel from πK . The ηK amplitudes are found to be weak and repulsive.

The 1^- amplitude around the πK threshold, shown in Fig. 2(b), has the behavior expected of a bound state, with the parametrization crossing zero below threshold. The energy dependence of the amplitude can be well described by a Breit-Wigner form, with a resulting mass of $m_R = 933(1)$ MeV, constrained by the zero crossing, and coupling of $g_R = 5.93(26)$, constrained by the slope.

Resonance poles.—We may now analyze the singularity structure of our parametrized t matrices by analytically continuing them to complex values of s . A general feature of relativistic scattering is that the square-root branch cut present in the phase space leads to a multisheeted structure for scattering amplitudes. The usual approach is to place the branch cuts along the real axis beginning at kinematic thresholds and running out to positive infinity. Physical scattering occurs just above these cuts on the first Riemann sheet, sheet I, where in our two-channel case, $\text{Im}p_{\pi K} > 0$, $\text{Im}p_{\eta K} > 0$. Resonance poles lie on the “unphysical” sheets reached by passing through the cuts, at pole positions which may be expressed in terms of a “pole mass” and “pole width,” $\sqrt{s_0} = m - i\Gamma/2$. Bound states may appear as poles on the physical sheet on the real axis below threshold, while poles on the real axis located on unphysical sheets are interpreted as virtual bound states.

In Fig. 3 we present the pole singularities found for the parametrizations given previously. As expected from the relatively slow increase through 90° in the πK 0^+ phase shift, we find a K_0^* resonance pole with a large width. A pole is present in this vicinity for a range of parametrizations (not presented here) capable of describing the finite-volume spectrum, and the uncertainty on the pole position shown in Fig. 3 includes such variation. In the 2^+ case, the rapid rise in phase shift is due to a K_2^* resonance pole at a slightly higher mass with a much smaller width. Both resonances have a dominant coupling to πK extracted from the residue of the pole. The 1^- amplitude has a bound-state pole just below πK threshold. The 0^+ amplitude has another pole that lies on the real axis some way below πK threshold. The 1^- pole is the only one lying on the physical sheet; the others lie on both sheet II where $\text{Im}p_{\pi K} < 0$, $\text{Im}p_{\eta K} > 0$ and sheet III where $\text{Im}p_{\pi K} < 0$, $\text{Im}p_{\eta K} < 0$.

The extracted poles are comparable to those found in experiment; the broad scalar and narrow tensor resonances resemble the $K_0^*(1430)$ and $K_2^*(1430)$ experimental states respectively, although the scalar state appears to have a somewhat larger width. The heavy (u, d) quark masses result in a bound state with $J^P = 1^-$ within a few MeV of




FIG. 3 (color online). Pole singularities of partial-wave t matrices in the complex plane for $J^P = 0^+$ (red), 1^- (orange), and 2^+ (green). Squares correspond to poles found on unphysical sheets, and the circle is a physical sheet bound state. Uncertainties include variation under changes in parametrization form.

πK threshold, where experiment finds a narrow elastic scattering resonance. It is our expectation that the K^* vector bound state will become a resonance as the light quark mass is decreased, with the observed proximity to the πK threshold being an accident of the particular quark mass value used in this calculation. We note that the Breit-Wigner coupling extracted above, $g_R = 5.93(26)$, is in quite reasonable agreement with the value $g_R^{\text{phys}} = 5.52(16)$, corresponding to the experimental width [5], in line with the theoretical suggestion that vector meson couplings are largely quark-mass independent [15].

Our observation of a 0^+ pole below threshold on unphysical sheets agrees with the qualitative prediction of unitarized chiral perturbation theory that as the pion mass is increased above its physical value, the κ resonance pole becomes a virtual bound state [15].

Summary.—Through lattice QCD computation of discrete excited-state spectra in finite volumes we have been able to determine coupled-channel scattering amplitudes via parametrizations of their energy dependence. The singularity structure of these amplitudes was then explored. This calculation serves as a demonstration that using these techniques it is possible to extract information about not only narrow resonances and bound states but also broad resonances and nonresonant features such as virtual bound states.

To compare this quantitatively with experiment, we would need calculations at the physical light quark mass, but the energy region of interest would then be above three-hadron (and higher multiplicity) thresholds. The formalism to relate finite-volume spectra to scattering amplitudes is not currently mature enough to be applied in that situation, although significant progress is being made [16].

While in this case relatively little coupling was observed between the two scattering channels, there are hadron scattering situations in which strong coupling is anticipated, such as $(\pi\eta, \bar{K}K)$ in which the $a_0(980)$ resonance appears, and the isoscalar channel $(\pi\pi, \bar{K}K, \eta\eta)$ whose resonance structure and interpretation poses a number of

questions for QCD. Building on the successful application of the finite-volume formalism for coupled channels presented in this Letter, we will consider these more strongly coupled systems, move to lighter quark masses, and explore states in exotic partial waves.

CHROMA [17] and QUDA [18] were used to perform this work at Jefferson Laboratory. For supplemental material see Ref. [19].

We thank our colleagues within the Hadron Spectrum Collaboration and M. R. Pennington and A. P. Szczepaniak for useful discussions. We acknowledge the resources used at Oak Ridge Leadership Computing Facility, the Texas Advanced Computer Center, and the Pittsburgh Supercomputer Center. Support is provided by U.S. Department of Energy Contract No. DE-AC05-06OR23177 under which Jefferson Science Associates manages Jefferson Lab and Early Career Award Contract No. DE-SC0006765.

*dudek@jlab.org

- [1] M. Lüscher, *Nucl. Phys.* **B354**, 531 (1991).
- [2] J. J. Dudek, R. G. Edwards, and C. E. Thomas, *Phys. Rev. D* **87**, 034505 (2013).
- [3] P. Guo, J. J. Dudek, R. G. Edwards, and A. P. Szczepaniak, *Phys. Rev. D* **88**, 014501 (2013).
- [4] S. He, X. Feng, and C. Liu, *J. High Energy Phys.* **07** (2005) 011; M. T. Hansen and S. R. Sharpe, *Phys. Rev. D* **86**, 016007 (2012); R. A. Briceno and Z. Davoudi, *Phys. Rev. D* **88**, 094507 (2013).
- [5] J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86**, 010001 (2012).
- [6] S. Descotes-Genon and B. Moussallam, *Eur. Phys. J. C* **48**, 553 (2006).

- [7] Z. Fu, *Phys. Rev. D* **85**, 074501 (2012); Z. Fu and K. Fu, *Phys. Rev. D* **86**, 094507 (2012); K. Sasaki, N. Ishizuka, M. Oka, and T. Yamazaki (PACS-CS Collaboration), *Phys. Rev. D* **89**, 054502 (2014).
- [8] C. B. Lang, L. Leskovec, D. Mohler, and S. Prelovsek, *Phys. Rev. D* **86**, 054508 (2012); S. Prelovsek, L. Leskovec, C. B. Lang, and D. Mohler, *Phys. Rev. D* **88**, 054508 (2013).
- [9] H.-W. Lin *et al.* (Hadron Spectrum Collaboration), *Phys. Rev. D* **79**, 034502 (2009).
- [10] J. J. Dudek, R. G. Edwards, M. J. Peardon, D. G. Richards, and C. E. Thomas, *Phys. Rev. Lett.* **103**, 262001 (2009); *Phys. Rev. D* **82**, 034508 (2010); C. E. Thomas, R. G. Edwards, and J. J. Dudek, *Phys. Rev. D* **85**, 014507 (2012).
- [11] J. J. Dudek, R. G. Edwards, and C. E. Thomas, *Phys. Rev. D* **86**, 034031 (2012).
- [12] M. Peardon, J. Bulava, J. Foley, C. Morningstar, J. Dudek, R. Edwards, B. Joó, H.-W. Lin, D. Richards, and K. Juge (Hadron Spectrum Collaboration), *Phys. Rev. D* **80**, 054506 (2009).
- [13] C. Michael, *Nucl. Phys.* **B259**, 58 (1985); M. Luscher and U. Wolff, *Nucl. Phys.* **B339**, 222 (1990).
- [14] K. Rummukainen and S. A. Gottlieb, *Nucl. Phys.* **B450**, 397 (1995); L. Leskovec and S. Prelovsek, *Phys. Rev. D* **85**, 114507 (2012).
- [15] J. Nebreda and J. R. Pelaez, *Phys. Rev. D* **81**, 054035 (2010).
- [16] M. T. Hansen and S. R. Sharpe, [arXiv:1311.4848](https://arxiv.org/abs/1311.4848).
- [17] R. G. Edwards and B. Joo, *Nucl. Phys. B, Proc. Suppl.* **140**, 832 (2005).
- [18] R. Babich, M. A. Clark, and B. Joo, *Proc. Comput. Sci.* **4**, 841 (2011).
- [19] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.113.182001> for a brief discussion of the coupled channel quantization condition, some details of the Chew-Mandelstam phase space, and the construction of operators to interpolate the η -meson.