

## Excitation of Chirping Whistler Waves in a Laboratory Plasma

B. Van Compernelle,<sup>1,\*</sup> X. An,<sup>2</sup> J. Bortnik,<sup>2</sup> R. M. Thorne,<sup>2</sup> P. Pribyl,<sup>1</sup> and W. Gekelman<sup>1</sup>

<sup>1</sup>*Department of Physics, University of California, Los Angeles, California 90095, USA*

<sup>2</sup>*Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, California 90095, USA*

(Received 3 March 2015; published 17 June 2015)

Whistler mode chorus emissions with a characteristic frequency chirp are important magnetospheric waves, responsible for the acceleration of outer radiation belt electrons to relativistic energies and also for the scattering loss of these electrons into the atmosphere. Here, we report on the first laboratory experiment where whistler waves exhibiting fast frequency chirping have been artificially produced using a beam of energetic electrons launched into a cold plasma. Frequency chirps are only observed for a narrow range of plasma and beam parameters, and show a strong dependence on beam density, plasma density, and magnetic field gradient. Broadband whistler waves similar to magnetospheric hiss are also observed, and the parameter ranges for each emission are quantified.

DOI: 10.1103/PhysRevLett.114.245002

PACS numbers: 94.30.Tz, 42.65.Re, 52.35.Hr, 94.30.cq

Whistler mode chorus waves are excited in the low density plasma region outside of Earth's plasmasphere following the convective injection of low-energy ( $\sim 10$  keV) plasma sheet electrons into the inner magnetosphere during periods of enhanced geomagnetic activity [1,2]. These waves are typically found in two distinct frequency bands with a gap near one half the electron cyclotron frequency [3]. They typically exhibit discrete rising or falling tones thought to originate from nonlinear processes [4], but can also occur as wideband incoherent hiss [5]. These chorus waves play a critical role in the acceleration of low-energy trapped radiation belt electrons to relativistic energies, and can also lead to rapid scattering loss into the atmosphere [6–9].

Extensive theoretical work has been done in the past but none adequately describes the features of discrete, chirping chorus waves. For instance, linear theory [10] predicts the regions in the frequency domain that are unstable to wave growth but cannot predict the saturation amplitude of the wave or the discreteness or frequency chirp rate. Extensive numerical simulations have been performed [4,11,12] under fairly restrictive assumptions that reproduce some of the features of chirping chorus waves, and certain scaling laws have been developed as a result, but these laws require extensive testing against observations to ascertain their validity. Such experimental testing is difficult to perform in space, since spacecraft do not generally have access to the source distribution of electrons that originally generated the waves, but can be readily performed in a laboratory setting, which is the motivation of the present study. Similarly, previous attempts to remotely excite artificial whistler waves have been made in Earth's near-space environment and have yielded interesting results [13,14] but have been difficult to probe and control.

This Letter reports the first experiment where choruslike whistler waves, i.e., discrete whistler waves exhibiting rapid

frequency chirping, have been artificially excited in a laboratory plasma. Energetic electrons emitted from a beam source are launched into a cold plasma and generate both broadband hisslike whistler waves and discrete choruslike whistler waves for specific beam and plasma parameters. Fast frequency chirping has been observed before in laboratory settings, for the Alfvén wave range of frequencies in experiments with energetic ion beams [15–17].

The experiment is performed on the upgraded Large Plasma Device (LAPD) [18,19] at the Basic Plasma Science Facility at University of California, Los Angeles (UCLA). The LAPD is a long cylindrical device with axial magnetic field and an 18 m long, 60 cm diameter quiescent plasma column (1 Hz repetition rate, helium fill gas at  $3 \times 10^{-5}$  Torr,  $T_e \leq 0.5$  eV). A range of plasma parameters were explored for this study, listed in Table I. Typical absolute values in the laboratory are quite different from those found in the magnetosphere but the dominant scaled dimensionless quantities are similar.

A 10 cm diameter electron beam source ( $0.5$  kV  $\leq V_{\text{beam}} \leq 4$  kV) [23,24] is introduced into the machine

TABLE I. Plasma parameters in the laboratory and in the magnetosphere: plasma density [20], magnetic field strength, ratio of plasma frequency to cyclotron frequency [21], ratio of beam density to plasma density [22], ratio of whistler wave amplitude to background magnetic field strength [21,22], and ratio of electron thermal pressure to magnetic pressure.

	LAPD	Inner magnetosphere
$n_0$ (cm <sup>-3</sup> )	$5 \times 10^8$ – $5 \times 10^{10}$	$1$ – $10^3$
$B_0$ (G)	20–60	$5 \times 10^{-4}$ – $10^{-1}$
$\omega_{pe}/\Omega_e$	1–12	2–15
$n_b/n_0$	0.001–0.04	$10^{-4}$ –1
$\tilde{B}/B_0$	$10^{-5}$ – $10^{-4}$	$10^{-5}$ – $10^{-3}$
$\beta_e$	$10^{-6}$ – $10^{-4}$	$10^{-5}$ – $10^{-3}$



as in Fig. 2(c) but they generally do not occur at the same time; i.e., when a discrete tone is present it suppresses any other emission. The wave activity at other frequencies outside of the discrete tone remains at values close to the natural plasma noise level that is present before the beam pulse starts. This indicates that the fast electron distribution is significantly modified by the excitation of the rising tone such that the broadband hisslike features are suppressed.

Apart from rising tones such as those illustrated in Fig. 2(c), a rich variety of beam-generated wave activity is observed in the whistler range. Figure 3 shows a representative selection ranging from (a) hisslike emissions in a lower band below  $\Omega_e/2$  and an upper band above  $\Omega_e/2$ , (b) falling tone followed by broadband activity below  $\Omega_e/2$ , (c) multiple consecutive short chirps, (d) double hook chirp emanating from the preexisting low frequency wave activity, (e) long extended rising and falling chirp crossing the  $\Omega_e/2$  mark, and (f) multiple simultaneous chirps at different frequencies followed by broadband wave activity below  $\Omega_e/2$ . For all these cases the beam source voltage is 3 kV. The relevant plasma and beam parameters are listed in the caption of Fig. 3. The magnetic field profile for these cases is plotted in Fig. 1(b) with  $L_B = -4.8$  m, except for Figs. 3(b) and 3(f), which were obtained at a nearly uniform 60 G field.

The relative occurrence rates of broadband hisslike emissions and frequency chirping choruslike emissions were investigated as a function of beam density, plasma density, and magnetic field gradient scale length. Data are taken in a radial line at  $z = 0.96$  m through the region with strongest wave activity. A plasma shot with at least one discrete frequency chirp is counted as a discrete event. Broadband wave activity an order of magnitude above the noise is counted as a broadband event. A single plasma shot can have both discrete and broadband events.

A scan of beam density was done at  $\omega_{pe}/\Omega_e = 3.2$  in a magnetic field profile with  $L_B = -4.8$  m [Fig. 1(b)]. Figure 4(a) shows that no wave activity above the noise is detected at low beam densities. As the beam density is increased discrete whistlers are first observed. There is a clear optimum beam density for excitation of discrete chirping waves; curiously, chirping whistler waves are not seen at the highest beam densities either. Broadband waves are mostly seen for larger density ratios,  $n_b/n_0 > 1\%$ , at this ratio of  $\omega_{pe}/\Omega_e = 3.2$ . Similar trends were observed for chorus and hiss in space [22].

A second parameter scan shown in Fig. 4(b) is performed by varying the plasma density at fixed beam density and with a fixed magnetic field profile the same as above. This changes both the ratio of  $\omega_{pe}/\Omega_e$  and the ratio of  $n_b/n_0$ . Frequency chirping occurs in a narrow range of  $\omega_{pe}/\Omega_e \approx 2-4$ , similar to space observations [5]. It is not clear if the absence of chirping waves at larger  $\omega_{pe}/\Omega_e$  is due to the increasing density or due to the decreasing ratio  $n_b/n_0$  as in Fig. 4(a) since these vary simultaneously. It is not possible to keep  $n_b/n_0$  fixed throughout this  $n_0$  scan because the beam source cannot deliver enough beam current at the higher plasma densities. Broadband waves occur both at low values of  $\omega_{pe}/\Omega_e$  [high values of  $n_b/n_0$  as in Fig. 4(a)], and at large  $\omega_{pe}/\Omega_e > 4$ , similar to hiss emissions found in the magnetosphere [5]. Higher plasma densities favor the excitation of broadband emissions, which is evident from their excitation even at low beam densities and from the lower threshold for broadband waves at large  $n_b/n_0$ , i.e.,  $n_b/n_0 > 1\%$  at  $\omega_{pe}/\Omega_e = 3.2$  [Fig. 4(a)] compared to  $n_b/n_0 > 3\%$  at  $\omega_{pe}/\Omega_e < 2$  [Fig. 4(b)]. The broadband waves generated at lower beam densities occur in a narrower frequency band, typically  $\Delta\omega/\Omega_e < 0.1$ , whereas at the higher beam densities  $\Delta\omega/\Omega_e > 0.2$  is routinely seen.




FIG. 3 (color online). Spectrograms of additional types of beam generated wave emissions, exhibiting hisslike emissions, falling tones, multiple consecutive chirps, hooks and chirping at multiple frequencies. Plasma and beam parameters for (a)–(f), respectively:  $\omega_{pe}/\Omega_e = 11.2, 2.8, 3.6, 2.8, 2.8, 2.8$ ;  $n_b/n_0(\%) = 0.1, 1.6, 1.0, 1.5, 0.8, 1.6$ . The beam source voltage is 3 kV for all cases. Audification of each of these cases is available in the Supplemental Material [32].






FIG. 4 (color online). Occurrence rates for discrete and broadband emissions versus (a) beam density at fixed plasma density, (b) plasma density at fixed beam density, and (c) magnetic field gradient at fixed plasma density and fixed beam density.

A third scan, displayed in Fig. 4(c), was done at  $\omega_{pe}/\Omega_e = 3.2$  by varying the magnetic field profile, as shown in Fig. 1(b). Steep magnetic field gradients dramatically suppress chirping whistler waves. At gentler gradients a clear optimum is reached for the occurrence of chirping waves; even for nearly uniform fields a relatively high occurrence rate is observed. Broadband emissions are relatively insensitive to the magnetic field gradient. The observed variation is thought to be due to the ratio of  $n_b/n_0 \approx 1.6\%$  being near the threshold value for the occurrence of broadband emissions, see Fig. 4(a).

The broadband emissions at  $\omega_{pe}/\Omega_e > 4$  in Fig. 4(b) are well described by linear theory. The measured power spectra, plotted in Fig. 5, demonstrate the upshift in frequency as the plasma density is lowered. The observed frequency shift agrees well with predictions based on linear excitation through the Doppler shifted cyclotron resonance, i.e., by solving  $\omega - k_{\parallel}v_{\text{beam},\parallel} = \Omega_e$  and the whistler wave dispersion relation simultaneously. The normalized linear growth rate  $\gamma/\Omega_e$  [33] for parallel propagating whistler waves is fairly insensitive to  $\omega_{pe}/\Omega_e$  at fixed  $n_b$ , which may explain why broadband emissions are observed at the higher densities even though the density ratio  $n_b/n_0$  decreases. Similarly, the absence of hiss in Fig. 4(a) at




FIG. 5 (color online). Measured power spectra at different values of  $\omega_{pe}/\Omega_e$ , with linear theory predictions for the excited frequency indicated by the dashed lines.

low  $n_b/n_0$  can be understood from linear growth rate estimates, given the dependence on the density ratio  $(n_b/n_0)^\alpha$  ( $\alpha = 1/3$  for a beam,  $\alpha = 1$  for a bi-Maxwellian plasma). We should note that, although we launched a gyrating beam into the background plasma, the electron distribution function is redistributed to have a long tail up to the beam energy by other processes [23,34–36].

The parametric behavior of the chirping whistler waves shows several similarities with the Omura model [11,37]. The model conjectures that the interplay between trapped and untrapped electron populations in the presence of a gentle gradient in the background field leads to the formation of an electron phase space hole. This gives rise to a nonlinear resonant current, which causes wave growth and frequency chirping. This is prohibited if a large background field gradient exists, which may explain the measured suppression of frequency chirps at large field gradients. The model predicts the existence of an optimum wave amplitude for the amplification of chorus waves, of which the measured optimum beam density in Fig. 4(a) may be evidence. Initial estimates of the predicted sweeping rate are an order of magnitude lower than observed, but may be due to the oblique nature of the observed discrete whistlers, whereas the model assumes parallel propagation.

In this Letter, we have summarized results from the first laboratory experiment to observe whistler waves exhibiting fast frequency chirping, a phenomenon that has been observed in space for decades known as chorus waves. Broadband wave activity reminiscent of magnetospheric hiss is also observed. The occurrence rates have strong dependencies on fundamental parameters such as the driving electron beam density, plasma density, and magnetic field profile. The experiment allows us, for the first time, to test under controlled conditions the leading hypotheses and identify missing elements in our current understanding of nonlinear whistler wave excitation.

The authors wish to thank G. J. Morales and T. A. Carter for insightful discussions. The research was funded by the Department of Energy and the National Science

Foundation by Grant No. DE-SC0010578, which was awarded to UCLA through the NSF/DOE Plasma Partnership program. Work was done at the Basic Plasma Science Facility also funded by DOE/NSF.

\*bvcomper@physics.ucla.edu

- [1] B. T. Ssurutani and E. J. Smith, *J. Geophys. Res.* **79**, 118 (1974).
- [2] O. Santolík, D. A. Gurnett, J. S. Pickett, M. Parrot, and N. Cornilleau-Wehrin, *J. Geophys. Res.: Space Phys.* **108**, 1278 (2003).
- [3] W. J. Burtis and R. A. Helliwell, *J. Geophys. Res.* **74**, 3002 (1969).
- [4] Y. Katoh and Y. Omura, *Geophys. Res. Lett.* **34**, L03102 (2007).
- [5] W. Li, R. M. Thorne, J. Bortnik, X. Tao, and V. Angelopoulos, *Geophys. Res. Lett.* **39**, L18106 (2012).
- [6] R. M. Thorne, W. Li, B. Ni, Q. Ma, J. Bortnik, L. Chen, D. N. Baker, H. E. Spence, G. D. Reeves, M. G. Henderson, C. A. Kletzing, W. S. Kurth, G. B. Hospodarsky, J. B. Blake, J. F. Fennell, S. G. Claudepierre, and S. G. Kanekal, *Nature (London)* **504**, 411 (2013).
- [7] G. D. Reeves, H. E. Spence, M. G. Henderson, S. K. Morley, R. H. W. Friedel, H. O. Funsten, D. N. Baker, S. G. Kanekal, J. B. Blake, J. F. Fennel, S. G. Claudepierre, R. M. Thorne, D. L. Turner, C. A. Kletzing, W. S. Kurth, B. A. Larsen, and J. T. Niehof, *Science* **341**, 991 (2013).
- [8] R. M. Thorne, *Geophys. Res. Lett.* **37**, L22107 (2010).
- [9] C. Cattell, J. R. Wygant, K. Goetz, K. Kersten, P. J. Kellogg, T. von Rosenvinge, S. D. Bale, I. Roth, M. Temerin, M. K. Hudson, R. A. Mewaldt, M. Wiedenbeck, M. Maksimovic, R. Ergun, M. Acuna, and C. T. Russell, *Geophys. Res. Lett.* **35**, L01105 (2008).
- [10] C. Kennel, *Phys. Fluids* **9**, 2190 (1966).
- [11] Y. Omura, Y. Katoh, and D. Summers, *J. Geophys. Res.: Space Phys.* **113**, A04223 (2008).
- [12] Y. Omura, M. Hikishima, Y. Katoh, D. Summers, and S. Yagitani, *J. Geophys. Res.: Space Phys.* **114**, A07217 (2009).
- [13] R. A. Helliwell, *Radio Sci.* **18**, 801 (1983).
- [14] M. Golkowski, M. B. Cohen, D. L. Carpenter, and U. S. Inan, *J. Geophys. Res.: Space Phys.* **116**, A04208 (2011).
- [15] K. McGuire *et al.*, *Phys. Rev. Lett.* **50**, 891 (1983).
- [16] M. Nave, D. Campbell, E. Joffrin, F. Marcus, G. Sadler, P. Smeulders, and K. Thomsen, *Nucl. Fusion* **31**, 697 (1991).
- [17] W. W. Heidbrink, *Plasma Phys. Controlled Fusion* **37**, 937 (1995).
- [18] W. Gekelman, H. Pfister, Z. Lucky, J. Bamber, D. Leneman, and J. Maggs, *Rev. Sci. Instrum.* **62**, 2875 (1991).
- [19] D. Leneman, W. Gekelman, and J. Maggs, *Rev. Sci. Instrum.* **77**, 015108 (2006).
- [20] B. Sheeley, M. Moldwin, H. Rassoul, and R. Anderson, *J. Geophys. Res.: Space Phys.* **106**, 25631 (2001).
- [21] W. Li, R. M. Thorne, Y. Nishimura, J. Bortnik, V. Angelopoulos, J. P. McFadden, D. E. Larson, J. W. Bonnell, O. Le Contel, A. Roux, and U. Auster, *J. Geophys. Res.: Space Phys.* **115**, A00F11 (2010).
- [22] X. Gao, W. Li, R. M. Thorne, J. Bortnik, V. Angelopoulos, Q. Lu, X. Tao, and S. Wang, *Geophys. Res. Lett.* **41**, 4805 (2014).
- [23] B. Van Compernelle, J. Bortnik, P. Pribyl, W. Gekelman, M. Nakamoto, X. Tao, and R. M. Thorne, *Phys. Rev. Lett.* **112**, 145006 (2014).
- [24] B. Van Compernelle, W. Gekelman, P. Pribyl, and C. Cooper, *Phys. Plasmas* **18**, 123501 (2011).
- [25] R. L. Stenzel and G. Golubyatnikov, *Phys. Fluids B* **5**, 3789 (1993).
- [26] E. Macúsová, O. Santolík, P. Decreau, A. G. Demekhov, D. Nunn, D. A. Gurnett, J. S. Pickett, E. E. Titova, B. V. Kozelov, J.-L. Rauch, and J.-G. Trotignon, *J. Geophys. Res.: Space Phys.* **115**, A12257 (2010).
- [27] E. Titova, B. V. Kozelov, F. Jiriček, J. Smilauer, A. G. Demekhov, V. Yu. Trakhtengerts, *Ann. Geophys.* **21**, 1073 (2003).
- [28] C. M. Cully, V. Angelopoulos, U. Auster, J. Bonnell, and O. Le Contel, *Geophys. Res. Lett.* **38**, L01106 (2011).
- [29] X. Tao, W. Li, J. Bortnik, R. Thorne, and V. Angelopoulos, *Geophys. Res. Lett.* **39**, L08106 (2012).
- [30] J. D. Means, *J. Geophys. Res.* **77**, 5551 (1972).
- [31] J. Bortnik, J. W. Cutler, C. Dunson, and T. E. Bleier, *J. Geophys. Res.: Space Phys.* **112**, A04204 (2007).
- [32] See supplemental material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.114.245002> for brief description.
- [33] T. F. Bell and O. Buneman, *Phys. Rev.* **133**, A1300 (1964).
- [34] A. A. Ivanov and L. I. Rudakov, *Sov. Phys. JETP* **24**, 1027 (1967).
- [35] S. Kainer, J. Dawson, R. Shanny, and T. Coffey, *Phys. Fluids* **15**, 493 (1972).
- [36] B. Lefebvre, L.-J. Chen, W. Gekelman, P. Kintner, J. Pickett, P. Pribyl, and S. Vincena, *Nonlinear Proc. Geophys.* **18**, 41 (2011).
- [37] Y. Omura and D. Nunn, *J. Geophys. Res.: Space Phys.* **116**, A05205 (2011).