



Semivisible Jets: Dark Matter Undercover at the LHC

Timothy Cohen,^{1,2,3} Mariangela Lisanti,¹ and Hou Keong Lou¹

¹*Department of Physics, Princeton University, Princeton, New Jersey 08544, USA*

²*School of Natural Sciences, Institute for Advanced Study, Princeton, New Jersey 08540, USA*

³*Institute of Theoretical Science, University of Oregon, Eugene, Oregon 97403, USA*

(Received 30 March 2015; published 23 October 2015)

Dark matter may be a composite particle that is accessible via a weakly coupled portal. If these hidden-sector states are produced at the Large Hadron Collider (LHC), they would undergo a QCD-like shower. This would result in a spray of stable invisible dark matter along with unstable states that decay back to the standard model. Such “semivisible” jets arise, for example, when their production and decay are driven by a leptophobic Z' resonance; the resulting signature is characterized by significant missing energy aligned along the direction of one of the jets. These events are vetoed by the current suite of searches employed by the LHC, resulting in low acceptance. This Letter will demonstrate that the transverse mass—computed using the final-state jets and the missing energy—provides a powerful discriminator between the signal and the QCD background. Assuming that the Z' couples to the standard model quarks with the same strength as the Z^0 , the proposed search can discover (exclude) Z' masses up to 2.5 TeV (3.5 TeV) with 100 fb^{-1} of 14 TeV data at the LHC.

DOI: 10.1103/PhysRevLett.115.171804

PACS numbers: 13.85.Rm, 12.10.-g, 14.80.-j, 95.35.+d

The existence of dark matter provides one of the strongest motivations for physics beyond the standard model (SM), and its discovery is one of the core missions for the Large Hadron Collider (LHC) program. Under the assumption that the dark-matter particle is neutral and stable, it escapes the detector and manifests as missing transverse energy (\cancel{E}_T). The LHC collaborations have developed a comprehensive search strategy to look for signals with significant \cancel{E}_T , accompanied by jets and/or leptons (see, e.g., [1] for a review). These searches are typically cast in terms of a simplified model [2] for supersymmetry or an effective theory of dark-matter interactions [3,4]. Yet if one relaxes the assumption that the dark sector is weakly coupled, a new class of dark-matter signatures emerges that evades this entire suite of analyses. Namely, it is possible that the dark matter has been lurking undercover within hadronic jets. The purpose of this Letter is to propose a straightforward discovery strategy for these “semivisible” jets.

Semivisible jets may occur if the dark matter is the stable (or metastable) remnant of a more complicated dark sector. The dynamics of nontrivial dark sectors have been explored in many contexts, e.g., [5–19]. In these models, the dark sector contains a dark-matter candidate(s) and possibly new force carriers and/or matter fields. Note that we are agnostic about how much of the cosmological relic density is accounted for by this dark-matter candidate. Messenger states that couple the dark sector to the SM can exist. If the messenger is accessible at colliders, dark-sector states can be produced, leading to unique signatures such as large particle multiplicities, displaced vertices, multiple resonances, and lepton or photon jets [20–33].

Another possibility is that the final state resulting from strongly coupled hidden sectors may contain a new type of

jet object—a semivisible jet. In this case, the dark matter is produced in a QCD-like parton shower along with other light degrees of freedom that decay hadronically. The result is a multijet+ \cancel{E}_T signature where one of the jets is closely aligned with the $\vec{\cancel{E}}_T$. A cornerstone of the standard multijet + \cancel{E}_T searches is to require a minimum angular separation between the jets and $\vec{\cancel{E}}_T$ to remove QCD background contamination arising from jet-energy mismeasurement [34,35]. This implies that events containing semivisible jets have a low acceptance for the currently implemented suite of searches.

To further illustrate this point, Fig. 1 compares selected variables for QCD with those for example weakly coupled and strongly coupled dark-matter models. The weakly coupled model is derived from supersymmetric theories and results from pair production of 1.5 TeV scalar quark partners. Each squark decays to a jet and 1 GeV neutral dark-matter particle. The signal from the strongly coupled model, which will be described more fully later, comes from the production of a 3 TeV resonance which then decays to a pair of dark-sector particles that subsequently shower and hadronize, yielding semivisible jets. Both these examples yield topologies with jets and missing energy. As the left panel shows, the weakly coupled (labeled WIMP) and strongly coupled (labeled semivisible jet) dark-sector models produce considerable \cancel{E}_T , with tails that extend beyond the QCD distribution. However, $\Delta\phi \equiv \min\{\Delta\phi_{j_1\cancel{E}_T}, \Delta\phi_{j_2\cancel{E}_T}\}$, where $j_{1,2}$ are the two hardest jets, is different between these models, as illustrated in the right panel. The $\Delta\phi$ distribution falls relatively steeply for the strongly coupled case, while it remains relatively flat for the weakly coupled scenario. Typical LHC searches require $\Delta\phi \gtrsim 0.4$ [34,35]. After requiring $\cancel{E}_T > 500 \text{ GeV}$ and $\Delta\phi > 0.4$, the acceptance




FIG. 1 (color online). (Left) The distribution of transverse missing energy E_T for the QCD background (solid blue), as well as the semivisible jet (dashed red) and WIMP (dotted green) examples. (Right) The distribution of $\Delta\phi \equiv \min\{\Delta\phi_{j_1 E_T}, \Delta\phi_{j_2 E_T}\}$, where $j_{1,2}$ are the two hardest jets.

of the WIMP (semivisible) example is $\sim 70\%$ (7%). We also verified that the α_T underperforms on semivisible jet signals; a standard cut of $\alpha_T > 0.55$ gives $\sim 20\%$ (3%) efficiency for the WIMP (semivisible) scenario [36]. Razor analyses [37] may prove useful, but require optimization for semivisible jets—see Supplemental Material [38].

To regain sensitivity to final states containing semivisible jets, the cut on the angular separation $\Delta\phi$ must be removed. This comes at the expense of an unsuppressed QCD multijet background, which must be eliminated using other techniques. In this Letter, we focus on the case where the dark sector is accessed via a heavy resonance. In such scenarios, one can take advantage of structure in the transverse mass—calculated using the final-state jets and E_T —to distinguish the signal from QCD. The strategy employed is similar to others proposed for semivisible Higgs decays [39].

We now introduce an example hidden valley [7] model that will enable us to analyze the LHC sensitivity for semivisible jets. This model is presented for illustration and concreteness; semivisible jets will be among the LHC signatures for a vast class of dark-sector theories. The messenger sector is described by a simple phenomenological model for a TeV-scale $U(1)'$ gauge boson. The new leptophobic Z' gauge boson couples to the SM baryon current J_{SM}^μ :

$$\mathcal{L} \supset -\frac{1}{4} Z'^{\mu\nu} Z'_{\mu\nu} - \frac{1}{2} M_{Z'}^2 Z'_\mu Z'^\mu - g_{Z'}^{\text{SM}} Z'_\mu J_{SM}^\mu. \quad (1)$$

Note that the Z' is treated as a Stueckelberg field—the Higgs sector has been neglected as it is not relevant for the LHC phenomenology discussed below; the additional matter needed to render the $U(1)$ of baryon number anomaly free is also ignored.

The dark sector is an $SU(2)_d$ gauge theory with coupling α_d and two fermionic quark flavors $\chi_i = \chi_{1,2}$ with masses M_i . The dark quark coupling to the Z' is $g_{Z'}^d$. In general, the couplings $g_{Z'}^d$ and $g_{Z'}^{\text{SM}}$ do not have to be comparable; we focus on the case where $g_{Z'}^d$ is large so that the Z' decays frequently to the dark sector.

The $SU(2)_d$ confines at a scale $\Lambda_d \ll M_{Z'}$. A QCD-like dark shower occurs when $M_i^2 \sim \Lambda_d^2$ so that many dark gluons and quarks are produced, which subsequently

hadronize. Some of these dark hadrons are stable, while others decay back to the SM via an off-shell Z' . The detailed spectrum of the dark hadrons depends on non-perturbative physics. Nonetheless, some properties of the low-energy states can be inferred from symmetry arguments. There are two accidental symmetries: a dark-isospin number $U(1)_{1-2}$ and a dark-baryon number $U(1)_{1+2}$, where “1” and “2” refer to the χ_i flavor index. For example, the mesons $\chi_1^\dagger \chi_1$ and $\chi_2^\dagger \chi_2$ are not charged under either of these symmetries, and are thus unstable. The other mesons ($\chi_1 \chi_2^\dagger$, $\chi_1^\dagger \chi_2$) and baryons ($\chi_1 \chi_2$, $\chi_1^\dagger \chi_2^\dagger$) are charged under $U(1)_{1-2}$ and $U(1)_{1+2}$, respectively, and are stable.

The spin of the dark mesons is also important. Similar to the η_b (η_c) and Υ (J/ψ) of the bottom (charm) system, the pseudoscalar and vector mesons should be degenerate. By naive degree-of-freedom counting, the vector mesons are produced roughly three times as often as the pseudoscalar mesons [29]. This impacts the phenomenology as the pseudoscalar decay is suppressed by a mass insertion, unlike the vector case. Therefore, the pseudoscalar decays are dominated by b quarks and are, in general, more displaced than the vector decays. The search strategy discussed below does not rely on b tags or displaced tracks; it may be possible to take advantage of the pseudoscalar meson decays with a more sophisticated analysis.

Despite the myriad of possibilities for the dark sector, only certain parameters have a direct impact on the jet observables and missing transverse energy. The strength of the dark shower, parametrized by α_d , plays a critical role. The coupling α_d controls how many dark hadrons are emitted in the shower as well as their p_T distributions, which has a direct and measurable impact on the jet observables. In addition, the mass scale of the dark quarks affects the jet masses.

The number of dark-matter particles produced in the shower impacts E_T . This effect can be parametrized as

$$r_{\text{inv}} \equiv \left\langle \frac{\text{number of stable hadrons}}{\text{number of hadrons}} \right\rangle. \quad (2)$$

The value of r_{inv} depends on the details of the dark-sector model. For the model described above with $M_1^2 = M_2^2$, the

average proportion of the stable and unstable hadrons is equal, implying $r_{\text{inv}} \approx 0.5$. This assumes that the hadronization process is flavor blind and that the dark quark masses are degenerate, and ignores baryon production, which is suppressed by a factor of $1/N_c^2$, where N_c is the number of dark colors.

A mass splitting between the flavors can lead to variations in r_{inv} . Assuming $M_2 \geq M_1$, in the Lund string model [40], fragmentation into heavier dark quark pairs is suppressed by the factor

$$T = \exp\left(-\frac{4\pi|M_2^2 - M_1^2|}{\Lambda_d^2}\right). \quad (3)$$

Because of the exponential dependence of the fragmentation process, r_{inv} is very sensitive to small splittings of the mass parameters. As a result, fewer stable mesons are produced when $M_2^2 - M_1^2 > \Lambda_d^2$. This decreases the value of r_{inv} below 0.5. To increase r_{inv} above 0.5, one can increase the number of flavors N_f , thereby enlarging the number of stable mesons by $N_f(N_f - 1)$, while only increasing the number of unstable mesons by N_f . Clearly, r_{inv} can take on any value between (0,1).

Table I summarizes the five parameters that are most relevant for semivisible jet observables. Three are sensitive to the details of the dark sector: the running dark-sector gauge coupling $\alpha_d(1 \text{ TeV})$, r_{inv} , and the mass scale for the dark mesons M_d . Note that by only including one value of M_d , we are assuming that the LHC will be insensitive to the dark spectrum mass splittings, i.e., $M_{Z'} \gg \Lambda_d$. Additionally, there are two portal parameters: the production rate $\sigma \times \text{Br}$ and the Z' mass.

To perform a detailed collider study, $u\bar{u}, d\bar{d} \rightarrow Z' \rightarrow \chi^\dagger \chi$ events were simulated for the 14 TeV LHC using PYTHIA 8 [41] with the default CTEQ 6 parton distribution functions. The dark-sector shower was simulated using the hidden valley PYTHIA module [28,29], modified to include the running of α_d as was done for [33]. Each meson had a probability r_{inv} to be a dark-matter particle. The possible decays of dark baryons or mesons into each other were neglected. The resulting particles were processed through DELPHES 3, with the CMS settings [42].

Anti- k_T $R = 0.5$ jets [43] were constructed and then reclustered into two large jets [44] using the Cambridge/Aachen (CA) algorithm [45] with $R = 1.1$. In a standard resonance search, one would use the invariant mass $M_{jj}^2 = (p_1 + p_2)^2$, where $p_{1,2}$ are the momenta of the

two final large jets $j_{1,2}$. However, the M_{jj} variable is not useful when there are a significant number of dark-matter particles in the shower. A variable that incorporates the missing momentum is the transverse mass:

$$M_T^2 = M_{jj}^2 + 2\left(\sqrt{M_{jj}^2 + p_{Tjj}^2} E_T - \vec{p}_{Tjj} \cdot \vec{E}_T\right). \quad (4)$$

In a detector with perfect resolution, $M_{jj} \leq M_T \leq M_{Z'}$. Figure 2 shows the distribution of M_{jj} , M_T , and M_{mc} after event selection. M_{mc} is the reconstructed $M_{Z'}$ computed from all the reclustered jets and truth-level dark-matter four vectors. M_T in general yields a narrower, more prominent peak closer to M_{mc} . The top panels of Fig. 2 show sample events for the different signals. The dark-sector particle multiplicity decreases for smaller α_d . As r_{inv} is increased, the signal degrades because more stable mesons are produced and more information is lost.

To estimate the reach at the LHC, we simulated 60×10^6 QCD events, 5×10^6 $W^\pm/Z + jj$ events, and 5×10^6 $t\bar{t}$ events. All samples were binned in H_T in order to increase statistics in the high- M_T tails [46] using MADGRAPH 5 [47] at parton level and PYTHIA 8 for the shower and hadronization. The dominant background after event selection is QCD and $W^\pm/Z + jj$. For the signal, 25000 events were generated for each choice of $M_{Z'}$ in increments of 500 GeV, using the benchmark parameters in Table I. An 8 TeV sample was used to validate the QCD background and limit-setting procedure [48] against the CMS dijet resonance search [49]. The E_T distribution was also validated [50].

Each event was required to have at least two $R = 0.5$ anti- k_T jets with $p_T > 200$ GeV and $|\eta| < 2.5$, as well as $E_T > 100$ GeV. These preselection cuts model the impact of the trigger. Then, the following cut flow was applied: (i) Recluster jets into $R = 1.1$ CA jets (j_1, j_2); (ii) require $|\eta_{j_1} - \eta_{j_2}| < 1.1$; (iii) require $\Delta\phi < 1$, where $\Delta\phi$ is the minimum azimuthal angle between \vec{E}_T and $\vec{p}_{Tj_{1,2}}$; (iv) veto isolated e^\pm/μ^\pm with $p_T > 20$ GeV, $|\eta| < 2.4$; and (v) require $E_T/M_T > 0.15$.

The $R = 1.1$ jets capture the wider radiation pattern expected from dark-shower dynamics. The cut on the pseudorapidity difference removes t -channel QCD [49,51]. The lepton veto and $\Delta\phi$ requirements suppress electroweak backgrounds. Finally, the E_T/M_T cut effectively acts as a missing energy requirement; cutting on the dimensionless ratio avoids sculpting the M_T distribution.

After applying these cuts, a bump hunt was performed using M_T . Following the dijet resonance searches [49,51,52], the resulting background distribution was parametrized using a fitting function—see Supplemental Material [38]. Assuming the background exactly follows the fit obtained from simulation, the exclusion reach for the signal benchmark can be computed. Figure 3 shows the results for 100 fb^{-1} of 14 TeV LHC data as a function of $M_{Z'}$ for the benchmark parameters (Table I). We assume a 10% width for the Z' , as computed using the benchmark parameters. The production cross section times branching

TABLE I. Parametrization for semivisible jet search.

	Description	Benchmark
$\sigma \times \text{Br}$	Production rate	80 fb
$M_{Z'}$	Z' pole mass	3 TeV
M_d	Dark hadron mass scale	20 GeV
$\alpha_d(1 \text{ TeV})$	Running dark coupling	0.2
r_{inv}	Fraction of stable hadrons	0.3




FIG. 2 (color online). Mass distributions after event selection cuts for the benchmark model in Table I, for various α_d and r_{inv} . M_{jj} is the mass of the two large reclustered jets, M_T the transverse mass, and M_{mc} the reconstructed Z' mass using all the dark-matter particles in the Monte Carlo simulation. The $\eta - \phi$ lego plots show the corresponding energy deposition in the detector. Red circles indicate visible SM hadrons, while the gray circles indicate undetected stable mesons. The crosses indicate the position of anti- k_T $R = 0.5$ jets. The relative size of each circle and cross is set by the $\sqrt{p_T}$ of the object.

ratio for a Z' with the same coupling as the SM Z^0 is shown as a reference. A Z' with SM couplings can be discovered (excluded) up to masses of ~ 2.5 TeV (3.5 TeV).

We estimate that the dijet limit on $\sigma \times \text{Br}(Z' \rightarrow q\bar{q})$ is comparable to the limit obtained for the dark-sector decay mode. For $g_{Z'}^d \approx 1$, the branching ratio to the dark sector varies from 80% to 50% along the expected exclusion bound as the Z' mass increases. Thus, the model would be discovered in the semivisible jet channel *before* it would be observed by the irreducible dijet channel; this conclusion only gets stronger for more integrated luminosity.

We simulate prompt decays for the dark mesons. For a sufficiently heavy Z' and small couplings, the dark vector meson decays could yield displaced vertices. Requiring that the lab-frame decay length be $\lesssim \mathcal{O}(1)$ mm, a lower bound on the couplings can be obtained:

$$g_{Z'}^{\text{SM}} \gtrsim 10^{-2} \left(\frac{1}{g_{Z'}^d} \right) \sqrt{\frac{B}{10}} \left(\frac{M_{Z'}}{3 \text{ TeV}} \right)^2 \left(\frac{20 \text{ GeV}}{\Lambda_d} \right)^{5/2}, \quad (5)$$

where $B \sim 10$ is the average boost factor computed from the benchmark simulation. Equation (5) gives the lower purple region in Fig. 3. Modifications of the search strategy can still be effective in this region.

This Letter proposed a new search strategy for the discovery of hidden-sector physics in resonance searches. In particular, the focus was on dark-sector showers that result in novel semivisible jets—objects that are composed of SM hadrons and dark matter. We argued that this generic signature could arise from a large class of strongly coupled dark-matter models. Furthermore, we gave a simplified parametrization that allows for a systematic treatment of the signature space. Finally, we provided expected exclusion limits using a bump hunt in transverse mass. A Z' with SM-size couplings to quarks could be discovered (excluded) up to ~ 2.5 TeV (3.5 TeV).

There are two main extensions that can be explored. First, one can allow for leptons, photons, and/or heavy-flavor particles to be produced in the shower. Second, one can consider other production modes. In this case, the semivisible jets may not be aligned with the E_T and

additional variables using jet substructure, along the lines of [53], displaced vertices, and/or the presence of low-mass resonances may be necessary.

With the LHC Run II on the horizon, it is important to rethink the program of dark-matter searches to guarantee that a wide range of new-physics scenarios are covered. Nontrivial dynamics in the dark-matter sector is one of the many fantastic and unexpected ways that new physics can emerge. This Letter provides a simple approach in preparation for this possibility.

We are grateful to Pedro Schwaller, Daniel Stolarski, and Andreas Weiler for providing us with the code to run the dark-sector coupling as implemented for [33]. We also thank Andrew Larkoski, Matthew Low, Duff Neill, Jesse Thaler, Scott Thomas, Tien-Tien Yu, and Daniel Whiteson for useful discussions. T.C. is supported by an LHC Theory Initiative Postdoctoral Fellowship, under the National Science Foundation Grant No. PHY-0969510.




FIG. 3 (color online). Expected $\sigma \times \text{Br}$ for the signal benchmark in Table I. The shaded green (yellow) band corresponds to ± 1 (2) standard deviations. The dashed black line shows the $\sigma \times \text{Br}$ for a Z' with the same coupling to quarks as the SM Z^0 ; the dashed red line shows estimates 5σ discovery. The shaded purple region indicates where the vertices are displaced, assuming $g_{Z'}^d \approx 1$ and that the shower is dominated by vector mesons.

- [1] V. A. Mitsou, Shedding light on dark matter at colliders, *Int. J. Mod. Phys. A* **28**, 1330052 (2013).
- [2] D. Alves *et al.* (LHC New Physics Working Group Collaboration), Simplified models for LHC new physics searches, *J. Phys. G* **39**, 105005 (2012).
- [3] J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T. M. P. Tait, and H.-B. Yu, Constraints on light majorana dark matter from colliders, *Phys. Lett. B* **695**, 185 (2011).
- [4] P. J. Fox, R. Harnik, J. Kopp, and Y. Tsai, Missing energy signatures of dark matter at the LHC, *Phys. Rev. D* **85**, 056011 (2012).
- [5] Z. Chacko, H.-S. Goh, and R. Harnik, The Twin Higgs: Natural Electroweak Breaking from Mirror Symmetry, *Phys. Rev. Lett.* **96**, 231802 (2006).
- [6] R. Schabinger and J. D. Wells, A minimal spontaneously broken hidden sector and its impact on Higgs boson physics at the Large Hadron Collider, *Phys. Rev. D* **72**, 093007 (2005).
- [7] M. J. Strassler and K. M. Zurek, Echoes of a hidden valley at hadron colliders, *Phys. Lett. B* **651**, 374 (2007).
- [8] G. Burdman, Z. Chacko, H.-S. Goh, and R. Harnik, Folded supersymmetry and the LEP paradox, *J. High Energy Phys.* **02** (2007) 009.
- [9] J. Kang and M. A. Luty, Macroscopic strings and “quirks” at colliders, *J. High Energy Phys.* **11** (2009) 065.
- [10] J. March-Russell, S. M. West, D. Cumberbatch, and D. Hooper, Heavy dark matter through the Higgs portal, *J. High Energy Phys.* **07** (2008) 058.
- [11] N. Arkani-Hamed, D. P. Finkbeiner, T. R. Slatyer, and N. Weiner, A theory of dark matter, *Phys. Rev. D* **79**, 015014 (2009).
- [12] D. E. Kaplan, M. A. Luty, and K. M. Zurek, Asymmetric dark matter, *Phys. Rev. D* **79**, 115016 (2009).
- [13] G. D. Kribs, T. S. Roy, J. Terning, and K. M. Zurek, Quirky composite dark matter, *Phys. Rev. D* **81**, 095001 (2010).
- [14] J. McDonald, Thermally Generated Gauge Singlet Scalars as Self-Interacting Dark Matter, *Phys. Rev. Lett.* **88**, 091304 (2002).
- [15] V. Barger, P. Langacker, M. McCaskey, M. J. Ramsey-Musolf, and G. Shaughnessy, LHC phenomenology of an extended standard model with a real scalar singlet, *Phys. Rev. D* **77**, 035005 (2008).
- [16] R. A. Porto and A. Zee, The private Higgs, *Phys. Lett. B* **666**, 491 (2008).
- [17] O. Bertolami and R. Rosenfeld, The Higgs portal and an unified model for dark energy and dark matter, *Int. J. Mod. Phys. A* **23**, 4817 (2008).
- [18] Y. Hochberg, E. Kuflik, T. Volansky, and J. G. Wacker, Mechanism for Thermal Relic Dark Matter of Strongly Interacting Massive Particles, *Phys. Rev. Lett.* **113**, 171301 (2014).
- [19] A. Alves, S. Profumo, and F. S. Queiroz, The dark Z' portal: Direct, indirect and collider searches, *J. High Energy Phys.* **04** (2014) 063.
- [20] M. J. Strassler, Possible effects of a hidden valley on supersymmetric phenomenology, [arXiv:hep-ph/0607160](https://arxiv.org/abs/hep-ph/0607160).
- [21] M. J. Strassler and K. M. Zurek, Discovering the Higgs through highly-displaced vertices, *Phys. Lett. B* **661**, 263 (2008).
- [22] T. Han, Z. Si, K. M. Zurek, and M. J. Strassler, Phenomenology of hidden valleys at hadron colliders, *J. High Energy Phys.* **07** (2008) 008.
- [23] M. J. Strassler, On the phenomenology of hidden valleys with heavy flavor, [arXiv:0806.2385](https://arxiv.org/abs/0806.2385).
- [24] N. Arkani-Hamed and N. Weiner, LHC signals for a superunified theory of dark matter, *J. High Energy Phys.* **12** (2008) 104.
- [25] C. Kilic, T. Okui, and R. Sundrum, Vectorlike confinement at the LHC, *J. High Energy Phys.* **02** (2010) 018.
- [26] M. Baumgart, C. Cheung, J. T. Ruderman, L.-T. Wang, and I. Yavin, Non-Abelian dark sectors and their collider signatures, *J. High Energy Phys.* **04** (2009) 014.
- [27] S. Chang and M. A. Luty, Displaced dark matter at colliders, [arXiv:0906.5013](https://arxiv.org/abs/0906.5013).
- [28] L. Carloni and T. Sjostrand, Visible effects of invisible hidden valley radiation, *J. High Energy Phys.* **09** (2010) 105.
- [29] L. Carloni, J. Rathsmann, and T. Sjostrand, Discerning secluded sector gauge structures, *J. High Energy Phys.* **04** (2011) 091.
- [30] Y. F. Chan, M. Low, D. E. Morrissey, and A. P. Spray, LHC signatures of a minimal supersymmetric hidden valley, *J. High Energy Phys.* **05** (2012) 155.
- [31] S. Chatrchyan *et al.* (CMS Collaboration), Search for microscopic black holes in pp collisions at $\sqrt{s} = 8$ TeV, *J. High Energy Phys.* **07** (2013) 178.
- [32] G. Aad *et al.* (ATLAS Collaboration), Search for pair-produced third-generation squarks decaying via charm quarks or in compressed supersymmetric scenarios in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *Phys. Rev. D* **90**, 052008 (2014).
- [33] P. Schwaller, D. Stolarski, and A. Weiler, Emerging jets, [arXiv:1502.05409](https://arxiv.org/abs/1502.05409).
- [34] S. Chatrchyan *et al.* (CMS Collaboration), Search for new physics in the multijet and missing transverse momentum final state in proton-proton collisions at $\sqrt{s} = 8$ TeV, *J. High Energy Phys.* **06** (2014) 055.
- [35] G. Aad *et al.* (ATLAS Collaboration), Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using $\sqrt{s} = 8$ TeV proton-proton collision data, *J. High Energy Phys.* **09** (2014) 176.
- [36] V. Khachatryan *et al.* (CMS Collaboration), Search for supersymmetry using razor variables in events with b -tagged jets in pp collisions at $\sqrt{s} = 8$ TeV, *Phys. Rev. D* **91**, 052018 (2015).
- [37] V. Khachatryan *et al.* (CMS Collaboration), Search for dark matter direct production using razor variables in events with two or more jets in pp collisions at 8 TeV, Report No. CMS-PAS-EXO-14-004.
- [38] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.115.171804> for a discussion of relevant background distributions, as well as Razor variables for semivisible jet signals.
- [39] C. Englert, M. Spannowsky, and C. Wymant, Partially (in) visible Higgs decays at the LHC, *Phys. Lett. B* **718**, 538 (2012).
- [40] B. Andersson, G. Gustafson, G. Ingelman, and T. Sjostrand, Parton fragmentation and string dynamics, *Phys. Rep.* **97**, 31 (1983).

- [41] T. Sjostrand, S. Mrenna, and P.Z. Skands, A brief introduction to PYTHIA 8.1, *Comput. Phys. Commun.* **178**, 852 (2008).
- [42] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaitre, A. Mertens, and M. Selvaggi, DELPHES 3, a modular framework for fast simulation of a generic collider experiment, *J. High Energy Phys.* **02** (2014) 057.
- [43] M. Cacciari, G.P. Salam, and G. Soyez, The anti- k_r jet clustering algorithm, *J. High Energy Phys.* **04** (2008) 063.
- [44] B. Nachman, P. Nef, A. Schwartzman, M. Swiatkowski, and C. Wanotayaroj, Jets from jets: Re-clustering as a tool for large radius jet reconstruction and grooming at the LHC, *J. High Energy Phys.* **02** (2015) 075.
- [45] S. Bentvelsen and I. Meyer, The Cambridge jet algorithm: Features and applications, *Eur. Phys. J. C* **4**, 623 (1998).
- [46] A. Avetisyan, J.M. Campbell, T. Cohen, N. Dhingra, J. Hirschauer *et al.*, Methods and results for standard model event generation at $\sqrt{s} = 14$ TeV, 33 TeV and 100 TeV proton colliders (a Snowmass whitepaper), [arXiv:1308.1636](https://arxiv.org/abs/1308.1636).
- [47] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, *J. High Energy Phys.* **07** (2014) 079.
- [48] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, Asymptotic formulae for likelihood-based tests of new physics, *Eur. Phys. J. C* **71**, 1554 (2011).
- [49] V. Khachatryan *et al.* (CMS Collaboration), Search for narrow resonances using the dijet mass spectrum with 19.6 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV, CERN Technical Report No. CMS-PAS-EXO-12-059, 2013.
- [50] V. Khachatryan *et al.* (CMS Collaboration), Performance of the CMS missing transverse momentum reconstruction in pp data at $\sqrt{s} = 8$ TeV, *J. Instrum.* **10**, P02006 (2015).
- [51] G. Aad *et al.* (ATLAS Collaboration), Search for new phenomena in the dijet mass distribution using pp collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector, [arXiv:1407.1376](https://arxiv.org/abs/1407.1376).
- [52] T. Aaltonen *et al.* (CDF Collaboration), Search for new particles decaying into dijets in proton-antiproton collisions at $\sqrt{s} = 1.96$ TeV, *Phys. Rev. D* **79**, 112002 (2009).
- [53] J. Fan, D. Krohn, P. Mosteiro, A. M. Thalapillil, and L.-T. Wang, Heavy squarks at the LHC, *J. High Energy Phys.* **03** (2011) 077.