

Did LIGO Detect Dark Matter?

Simeon Bird,^{*} Ilias Cholis, Julian B. Muñoz, Yacine Ali-Haïmoud, Marc Kamionkowski,
Ely D. Kovetz, Alvise Raccanelli, and Adam G. Riess

*Department of Physics and Astronomy, Johns Hopkins University,
3400 North Charles Street, Baltimore, Maryland 21218, USA*

(Received 4 March 2016; published 19 May 2016)

We consider the possibility that the black-hole (BH) binary detected by LIGO may be a signature of dark matter. Interestingly enough, there remains a window for masses $20M_\odot \lesssim M_{\text{bh}} \lesssim 100M_\odot$ where primordial black holes (PBHs) may constitute the dark matter. If two BHs in a galactic halo pass sufficiently close, they radiate enough energy in gravitational waves to become gravitationally bound. The bound BHs will rapidly spiral inward due to the emission of gravitational radiation and ultimately will merge. Uncertainties in the rate for such events arise from our imprecise knowledge of the phase-space structure of galactic halos on the smallest scales. Still, reasonable estimates span a range that overlaps the $2\text{--}53 \text{ Gpc}^{-3} \text{ yr}^{-1}$ rate estimated from GW150914, thus raising the possibility that LIGO has detected PBH dark matter. PBH mergers are likely to be distributed spatially more like dark matter than luminous matter and have neither optical nor neutrino counterparts. They may be distinguished from mergers of BHs from more traditional astrophysical sources through the observed mass spectrum, their high ellipticities, or their stochastic gravitational wave background. Next-generation experiments will be invaluable in performing these tests.

DOI: 10.1103/PhysRevLett.116.201301

The nature of dark matter (DM) is one of the most long-standing and puzzling questions in physics. Cosmological measurements have now determined with exquisite precision the abundance of DM [1,2], and from both observations and numerical simulations we know quite a bit about its distribution in galactic halos. Still, the nature of DM remains a mystery. Given the efficacy with which weakly interacting massive particles—for many years the favored particle-theory explanation—have eluded detection, it may be warranted to consider other possibilities for DM. Primordial black holes (PBHs) are one such possibility [3–6].

Here, we consider whether the two $\sim 30M_\odot$ black holes detected by LIGO [7] could plausibly be PBHs. There is a window for PBHs to be DM if the black-hole (BH) mass is in the range $20M_\odot \lesssim M \lesssim 100M_\odot$ [8,9]. Lower masses are excluded by microlensing surveys [10–12]. Higher masses would disrupt wide binaries [9,13,14]. It has been argued that PBHs in this mass range are excluded by cosmic microwave background (CMB) constraints [15,16]. However, these constraints require modeling of several complex physical processes, including the accretion of gas onto a moving BH, the conversion of the accreted mass to a luminosity, the self-consistent feedback of the BH radiation on the accretion process, and the deposition of the radiated energy as heat in the photon-baryon plasma. A significant (and difficult to quantify) uncertainty should therefore be associated with this upper limit [17], and it seems worthwhile to examine whether PBHs in this mass range could have other observational consequences.

In this Letter, we show that if DM consists of $\sim 30M_\odot$ BHs, then the rate for mergers of such PBHs falls within the

merger rate inferred from GW150914. In any galactic halo, there is a chance that two BHs will undergo a hard scatter, lose energy to a soft gravitational wave (GW) burst and become gravitationally bound. This BH binary will merge via the emission of GWs in less than the Hubble time. (This differs from Ref. [18] who considered binaries which form in the radiation era and merge over a Hubble time.) Below, we first estimate roughly the rate of such mergers and then present the results of more detailed calculations. We discuss uncertainties in the calculation and some possible ways to distinguish PBHs from BH binaries from more traditional astrophysical sources.

Consider two PBHs approaching each other on a hyperbolic orbit with some impact parameter and relative velocity v_{pbh} . As the PBHs approach each other, they produce a time-varying quadrupole moment and thus GW emission. The PBH pair becomes gravitationally bound if the GW emission exceeds the initial kinetic energy. The cross section for this process is [19,20]

$$\begin{aligned} \sigma &= \pi \left(\frac{85\pi}{3} \right)^{2/7} R_s^2 \left(\frac{v_{\text{pbh}}}{c} \right)^{-18/7} \\ &= 1.37 \times 10^{-14} M_{30}^2 v_{\text{pbh}-200}^{-18/7} \text{ pc}^2, \end{aligned} \quad (1)$$

where M_{pbh} is the PBH mass and M_{30} the PBH mass in units of $30M_\odot$, $R_s = 2GM_{\text{pbh}}/c^2$ is its Schwarzschild radius, v_{pbh} is the relative velocity of two PBHs, and $v_{\text{pbh}-200}$ is this velocity in units of 200 km sec^{-1} .

We begin with a rough but simple and illustrative estimate of the rate per unit volume of such mergers. Suppose that all DM in the Universe resided in Milky Way-like halos of

mass $M = M_{12} 10^{12} M_\odot$ and uniform mass density $\rho = 0.002 \rho_{0.002} M_\odot \text{ pc}^{-3}$, with $\rho_{0.002} \sim 1$. Assuming a uniform-density halo of the volume $V = M/\rho$, the rate of mergers per halo would be

$$N \simeq (1/2) V (\rho/M_{\text{pbh}})^2 \sigma v \simeq 3.10 \times 10^{-12} M_{12} \rho_{0.002} v_{\text{pbh}-200}^{-11/7} \text{ yr}^{-1}. \quad (2)$$

The relative velocity $v_{\text{pbh}-200}$ is specified by a characteristic halo velocity. The mean cosmic DM mass density is $\rho_{\text{dm}} \simeq 3.6 \times 10^{10} M_\odot \text{ Mpc}^{-3}$, so the spatial density of halos is $n \simeq 0.036 M_{12}^{-1} \text{ Mpc}^{-3}$. The rate per unit comoving volume in the Universe is thus

$$\Gamma \simeq 1.1 \times 10^{-4} \rho_{0.002} v_{\text{pbh}-200}^{-11/7} \text{ Gpc}^{-3} \text{ yr}^{-1}. \quad (3)$$

The normalized halo mass M_{12} drops out, as it should. The merger rate per unit volume also does not depend on the PBH mass, as the capture cross section scales like M_{pbh}^2 .

This rate is small compared to the $2\text{--}53 \text{ Gpc}^{-3} \text{ yr}^{-1}$ estimated by LIGO for a population of $\sim 30 M_\odot - 30 M_\odot$ mergers [21], but this is a very conservative estimate. As Eq. (3) indicates, the merger rate is higher in higher-density regions and in regions of lower DM velocity dispersion. The DM in Milky Way-like halos is known from simulations [22] and analytic models [23] to have substructure, regions of higher density and lower velocity dispersion. DM halos also have a broad mass spectrum, extending to very low masses where the densities can become far higher—and the velocity dispersion far lower—than in the Milky Way. To get a very rough estimate of the conceivable increase in the PBH merger rate due to these smaller-scale structures, we can replace ρ and v in Eq. (3) by the values they would have had in the earliest generation of collapsed objects, where the DM densities were largest and the velocity dispersions smallest. If the primordial power spectrum is nearly scale invariant, then gravitationally bound halos of mass $M_c \sim 500 M_\odot$, for example, will form at redshift $z_c \simeq 28 - \log_{10}(M_c/500 M_\odot)$. These objects will have virial velocities $v \simeq 0.2 \text{ km sec}^{-1}$ and densities $\rho \simeq 0.24 M_\odot \text{ pc}^{-3}$ [24]. Using these values in Eq. (3) increases the merger rate per unit volume to

$$\Gamma \simeq 700 \text{ Gpc}^{-3} \text{ yr}^{-1}. \quad (4)$$

This would be the merger rate if *all* of the DM resided in the smallest halos. Clearly, this is not true in the present day; substructures are at least partially stripped as they merge to form larger objects, so Eq. (4) should be viewed as a conservative upper limit.

Having demonstrated that rough estimates contain the merger-rate range $2\text{--}53 \text{ Gpc}^{-3} \text{ yr}^{-1}$ suggested by LIGO, we now turn to more careful estimates of the PBH merger rate. As Eq. (3) suggests, the merger rate will depend on a density-weighted average, over the entire cosmic DM

distribution, of $\rho_{0.002} v_{\text{pbh}-200}^{-11/7}$. To perform this average, we will (a) assume that DM is distributed within galactic halos with a Navarro-Frenk-White (NFW) profile [25] with concentration parameters inferred from simulations, and (b) try several halo mass functions taken from the literature for the distribution of halos.

The PBH merger rate $\mathcal{R}$ within each halo can be computed using

$$\mathcal{R} = 4\pi \int_0^{R_{\text{vir}}} r^2 \frac{1}{2} \left(\frac{\rho_{\text{nfw}}(r)}{M_{\text{pbh}}} \right)^2 \langle \sigma v_{\text{pbh}} \rangle dr, \quad (5)$$

where $\rho_{\text{nfw}}(r) = \rho_s [(r/R_s)(1 + r/R_s)^2]^{-1}$ is the NFW density profile with characteristic radius r_s and characteristic density ρ_s . R_{vir} is the virial radius at which the NFW profile reaches a value 200 times the comoving mean cosmic density and is cut off. The angle brackets denote an average over the PBH relative velocity distribution in the halo. The merger cross section σ is given by Eq. (1). We define the concentration parameter $C = R_{\text{vir}}/R_s$. To determine the profile of each halo, we require C as a function of halo mass M . We will use the concentration-mass relations fit to DM N -body simulations by both Ref. [26] and Ref. [27].

We now turn to the average of the cross section times the relative velocity. The one-dimensional velocity dispersion of a halo is defined in terms of the escape velocity at radius $R_{\text{max}} = 2.1626 R_s$, the radius of the maximum circular velocity of the halo, i.e.,

$$v_{\text{dm}} = \sqrt{\frac{GM(r < r_{\text{max}})}{r_{\text{max}}}} = \frac{v_{\text{vir}}}{\sqrt{2}} \sqrt{\frac{C}{C_m} \frac{g(C_m)}{g(C)}}, \quad (6)$$

where $g(C) = \ln(1 + C) - C/(1 + C)$, and $C_m = 2.1626 = R_{\text{max}}/R_s$. We approximate the relative velocity distribution of PBHs within a halo as a Maxwell-Boltzmann (MB) distribution with a cutoff at the virial velocity, i.e.,

$$P(v_{\text{pbh}}) = F_0 \left[\exp\left(-\frac{v_{\text{pbh}}^2}{v_{\text{dm}}^2}\right) - \exp\left(-\frac{v_{\text{vir}}^2}{v_{\text{dm}}^2}\right) \right], \quad (7)$$

where F_0 is chosen so that $4\pi \int_0^{v_{\text{vir}}} P(v) v^2 dv = 1$. This model provides a reasonable match to N -body simulations, at least for the velocities substantially less than the virial velocity which dominate the merger rate (e.g., Ref. [28]). Since the cross section is independent of the radius, we can integrate the NFW profile to find the merger rate in any halo:

$$\mathcal{R} = \left(\frac{85\pi}{12\sqrt{2}} \right)^{2/7} \frac{9G^2 M_{\text{vir}}^2}{c R_s^3} \left(1 - \frac{1}{(1 + C)^3} \right) \frac{D(v_{\text{dm}})}{g(C)^2}, \quad (8)$$

where

$$D(v_{\text{dm}}) = \int_0^{v_{\text{vir}}} P(v, v_{\text{dm}}) \left(\frac{2v}{c}\right)^{3/7} dv, \quad (9)$$

comes from Eq. (7).

Equation (1) gives the cross section for two PBHs to form a binary. However, if the binary is to produce an observable GW signal, these two PBHs must orbit and inspiral; a direct collision, lacking an inspiral phase, is unlikely to be detectable by LIGO. This requirement imposes a minimum impact parameter of roughly the Schwarzschild radius. The fraction of BHs direct mergers is $\sim v^{2/7}$ and reaches a maximum of $\sim 3\%$ for $v_{\text{pbh}} = 2000 \text{ km s}^{-1}$. Thus, direct mergers are negligible. We also require that once the binary is formed, the time until it merges (which can be obtained from Ref. [29]) is less than the Hubble time. The characteristic time it takes for a binary BH to merge varies as a function of halo velocity dispersion. It can be hours for $M_{\text{vir}} \approx 10^{12} M_{\odot}$ or thousands of years for $M_{\text{vir}} \approx 10^6 M_{\odot}$, and it thus is instantaneous on cosmological time scales. Given the small size of the binary and rapid time to merge, we can neglect disruption of the binary by a third PBH once formed. BH binaries can also form through nondissipative three-body encounters. The rate of these binary captures is non-negligible in small halos [19,30], but they generically lead to the formation of wide binaries that will not be able to harden and merge within the Hubble time. This formation mechanism should not affect our LIGO rates. The merger rate is therefore equal to the rate of binary BH formation, Eq. (8).

Figure 1 shows the contribution to the merger rate, Eq. (8), for two concentration-mass relations. As can be seen, both concentration-mass relations give similar results. An increase in halo mass produces an increased PBH merger rate. However, less massive halos have a higher concentration (since they are more likely to have virialized

earlier), so that the merger rate per unit mass increases significantly as the halo mass is decreased.

To compute the expected LIGO event rate, we convolve the merger rate $\mathcal{R}$ per halo with the mass function dn/dM . Since the redshifts ($z \lesssim 0.3$) detectable by LIGO are relatively low we will neglect redshift evolution in the halo mass function. The total merger rate per unit volume is, then,

$$\mathcal{V} = \int (dn/dM)(M) \mathcal{R}(M) dM. \quad (10)$$

Given the exponential falloff of dn/dM at high masses, despite the increased merger rate per halo suggested in Fig. 1, the precise value of the upper limit of the integrand does not affect the final result.

At the lower limit, discreteness in the DM particles becomes important, and the NFW profile is no longer a good description of the halo profile. Furthermore, the smallest halos will evaporate due to the periodic ejection of objects by dynamical relaxation processes. The evaporation time scale is [31]

$$t_{\text{evap}} \approx (14\mathcal{N}/\ln \mathcal{N})[R_{\text{vir}}/(Cv_{\text{dm}})], \quad (11)$$

where $\mathcal{N}$ is the number of individual BHs in the halo, and we assumed that the PBH mass is $30M_{\odot}$. For a halo of mass $400M_{\odot}$, the velocity dispersion is 0.15 km sec^{-1} , and the evaporation time scale is $\sim 3 \times 10^9 \text{ yr}$. In practice, during matter domination, halos which have already formed will grow continuously through mergers or accretion. Evaporation will thus be compensated for by the addition of new material, and as halos grow new halos will form from mergers of smaller objects. However, during dark-energy domination at $z \lesssim 0.3$, $3 \times 10^9 \text{ yr}$ ago, this process slows down. Thus, we will neglect the signal from halos with an evaporation time scale less than $3 \times 10^9 \text{ yr}$, corresponding to $M < 400M_{\odot}$. This is, in any case, 13 PBHs, and close to the point where the NFW profile is no longer valid.

The halo mass function dn/dM is computed using both semianalytic fits to N -body simulations and with analytic approximations. Computing the merger rate in the small halos discussed above requires us to extrapolate both the halo mass function and the concentration-mass relation around 6 orders of magnitude in mass beyond the smallest halos present in the calibration simulations. High-resolution simulations of $10^{-4}M_{\odot}$ cold dark matter microhalos [32,33] suggest that our assumed concentration-mass relations underestimate the internal density of these halos, making our rates conservative.

The mass functions depend on the halo mass through the perturbation amplitude $\sigma(R_{\text{vir}})$ at the virial radius R_{vir} of a given halo. Because of the scale invariance of the window functions on small scales, $\sigma(R_{\text{vir}})$ varies only by a factor of 2 between $M_{\text{vir}} = 10^9 M_{\odot}$ and $M_{\text{vir}} = 10^3 M_{\odot}$. Thus, the extrapolation in the mass function is less severe than it looks. We also note that the scale-invariant nature of the




FIG. 1. The PBH merger rate per halo as a function of halo mass. The solid line shows the trend assuming the concentration-mass relation from Ref. [27], and the dashed line from Ref. [26]. To guide the eye, the dot-dashed line shows a constant BH merger rate per unit halo mass.

initial conditions suggests that the shape of the halo mass function should not evolve unduly until it reaches the scale of the PBH mass, or the evaporation cutoff.

To quantify the uncertainty induced by the dn/dM extrapolation, we obtained results with two different mass functions: the classic analytic Press-Schechter calculation [34] and one calibrated to numerical simulations from Tinker *et al.* [35]. The agreement between the two small-scale behaviors suggests that extrapolating the mass functions is not as blind as it might otherwise seem. We also include a third mass function, from Jenkins *et al.* [36], that includes an artificial small-scale mass cutoff at a halo mass $M_{\text{vir}} \sim 10^6 M_\odot$. This cutoff is inserted to roughly model the mass function arising if there is no power on scales smaller than those currently probed observationally. We include it to provide a very conservative lower limit to the merger rate if, for some reason, small-scale power were to be suppressed. We do not, however, consider it likely that this mass function accurately represents the distribution of halo masses in our Universe.

Figure 2 shows the merger rate per logarithmic interval in halo mass. In all cases, halos with $M_{\text{vir}} \lesssim 10^9 M_\odot$ dominate the signal, due to the increase in concentration and decrease in velocity dispersion with smaller halo masses. The Tinker mass function, which asymptotes to a constant number density for small masses, produces the most mergers. The Press-Schechter calculation has $\sim 50\%$ fewer events in small halos, while the Jenkins mass function results in merger rates nearly 4 orders of magnitude smaller [and in rough agreement with Eq. (3)].

We integrate the curves in Fig. 2 to compute the total merger rate $\mathcal{V}$. All mass functions give a similar result, $\sim (3 \pm 1) \times 10^{-4} \text{ Gpc}^{-3} \text{ yr}^{-1}$, from halos of masses $\gtrsim 10^9 M_\odot$, representing for the Tinker and Press-Schechter mass function a small fraction of the events. When we include all halos with $M_{\text{vir}} > 400 M_\odot$, the number of events

increases dramatically and depends strongly on the lower cutoff mass M_c for the halo mass. Both the Press-Schechter and Tinker mass functions are for small halos linear in the integrated perturbation amplitude $\propto 1/\sigma(R_{\text{vir}})$ at the virial radius R_{vir} of the collapsing halo. In small halos, $1/\sigma(R_{\text{vir}})$ is roughly constant. Thus, for a mass function $\text{MF}(\sigma)$, we have

$$(dn/dM) \sim (C \log \sigma/dM) [\text{MF}(\sigma)/M_{\text{vir}}] \sim M_{\text{vir}}^{-2}. \quad (12)$$

The concentration is also a function of $1/\sigma(R_{\text{vir}})$ and it, too, becomes roughly constant for small masses. Assuming a constant concentration, the merger rate per halo scales as $\mathcal{R} \sim M^{10/21}$. Thus, Eq. (10) suggests that $\mathcal{V} \sim M_c^{-11/21}$. This compares well to the numerical differentiation of Fig. 2, which yields $\mathcal{V} \sim M_c^{-0.51}$.

The integrated merger rate is thus

$$\mathcal{V} = 2f(M_c/400M_\odot)^{-11/21} \text{ Gpc}^{-3} \text{ yr}^{-1}, \quad (13)$$

with $f \approx 1$ for the Tinker mass function, and $f \approx 0.6$ for the Press-Schechter mass function (the Jenkins mass function results in an event rate $\mathcal{V} \approx 0.02 \text{ Gpc}^{-3} \text{ yr}^{-1}$, independent of $M_c \lesssim 10^6 M_\odot$).

A variety of astrophysical processes may alter the mass function in some halos, especially within the dwarf galaxy range, $10^9 - 10^{10} M_\odot$. However, halos with $M_{\text{vir}} \lesssim 10^9 M_\odot$ are too small to form stars against the thermal pressure of the ionized intergalactic medium [37] and are thus unlikely to be affected by these astrophysical processes. Inclusion of galactic substructure, which our calculation neglects, should boost the results. However, since the event rate is dominated by the smallest halos, which should have little substructure, we expect this to make a negligible difference in our final result.

There is also the issue of the NFW density profile assumed. The results are fairly insensitive to the detailed density profile as long as the slope of the density profile varies no more rapidly than r^{-1} as $r \rightarrow 0$. For example, suppose we replace the NFW profile with the Einasto profile [38],

$$\rho(R) = \rho_0 \exp \left\{ -\frac{2}{\alpha} \left[\left(\frac{R}{R_s} \right)^\alpha - 1 \right] \right\}, \quad (14)$$

with $\alpha = 0.18$, which has a core as $r \rightarrow 0$. The reduction in the merger rate as $r \rightarrow 0$ is more than compensated for by an increased merger rate at larger radii leading to a total merger rate that is raised by 50% relative to NFW, to $\sim 3 \text{ Gpc}^{-3} \text{ yr}^{-1}$.

Our assumption of an isotropic MB-like velocity distribution in the halo may also underestimate the correct answer, as any other velocity distribution would have lower entropy and thus larger averaged $v^{-11/7}$. Finally, the discreteness of PBH DM will provide some Poisson enhancement of power on $\sim 400 M_\odot$ scales.




FIG. 2. The total PBH merger rate as a function of halo mass. Dashed and dotted lines show different prescriptions for the concentration-mass relation and halo mass function.

More small-scale power would probably lead to an enhancement of the event rate beyond Eq. (13).

The recent LIGO detection of two merging $\sim 30M_\odot$ black holes suggests a 90% C.L. event rate [21] of $2\text{--}53 \text{ Gpc}^{-3} \text{ yr}^{-1}$ if all mergers have the masses and emitted energy of GW150914. It is interesting that—although there are theoretical uncertainties—our best estimates of the merger rate for $30M_\odot$ PBHs, obtained with canonical models for the DM distribution, fall in the LIGO window.

The possibility that LIGO has seen DM thus cannot be immediately excluded. Even if the predicted merger rates turn out, with more precise treatments of the small-scale galactic phase-space distribution, to be smaller, conservative lower estimates of the merger rate for PBH DM suggests that the LIGO-VIRGO network should see a considerable number of PBH mergers over its lifetime.

We have assumed a population of PBHs with the same mass. The basic results obtained here should, however, remain unaltered if there is some small spread of PBH masses, as expected from PBH-formation scenarios, around the nominal value of $30M_\odot$.

PBH mergers may also be interesting for LIGO-VIRGO even if PBHs make up only a fraction f_{pbh} of the DM, as implied by CMB limits from Refs. [15,16] or the limits in Ref. [8]. In this case, the number density of PBHs will be reduced by f_{pbh} . The cutoff mass will increase as $M_c \sim f_{\text{pbh}}^{-1}$ if we continue to require > 13 PBHs in each halo to avoid halo evaporation. The overall event rate will be $\mathcal{V} \sim 2f_{\text{pbh}}^{3/21} \text{ Gpc}^{-3} \text{ yr}^{-1}$. Advanced LIGO will reach design PBH sensitivity in 2019 [39,40] and will probe $z < 0.75$, an increase in volume to $\approx 50 \text{ Gpc}^3$ (comoving). Thus, over the six planned years of aLIGO operation, while we should expect to detect ~ 600 events with $f_{\text{pbh}} = 1$, we will expect at least one event if $f_{\text{pbh}} > 0.1$.

Distinguishing whether any individual GW event, or even some population of events, are from PBH DM or more traditional astrophysical sources will be daunting. Still, there are some prospects. Most apparently, PBH mergers will be distributed more like small-scale DM halos and are thus less likely to be found in or near luminous galaxies than BH mergers resulting from stellar evolution. Moreover, PBH mergers are expected to have no electromagnetic or neutrino counterparts whatsoever. A DM component could conceivably show up in the BH mass spectrum as an excess of events with BH masses near $30M_\odot$ over a more broadly distributed mass spectrum from astrophysical sources (e.g., Ref. [41]).

Since the binary is formed on a very elongated orbit, the GW waveforms will initially have high ellipticity, exhibited by higher frequency harmonics in the GW signal [29]. We have verified that the ellipticities become unobservably small by the time the inspiral enters the LIGO band, but they may be detectable in future experiments [42]. Multiply lensed quasars [43,44], pulsar timing arrays [45], and fast

radio burst lensing searches [46] may also allow probes of the $\sim 30M_\odot$ PBH mass range.

Another potential source of information is the stochastic GW background. Models for the stochastic background due to BH mergers usually entail a mass distribution that extends to smaller BH masses and a redshift distribution that is somehow related to the star-formation history. Given microlensing limits, the PBH mass function cannot extend much below $30M_\odot$. Moreover, the PBH merger rate per unit comoving volume is likely higher for PBHs than for traditional BHs at high redshifts. Together, these suggest a stochastic background for PBHs that has more weight at low frequencies and less at higher ones than that from traditional BH sources.

The results of this work provide additional motivation for more sensitive next-generation GW experiments such as the Einstein Telescope [47], DECIGO [48], and BBO [49], which will continuously extend the aLIGO frequency range downwards. These may enable the tests described above for excesses in the BH mass spectrum, high ellipticity, and low-frequency stochastic background that are required to determine whether LIGO has detected dark matter.

We thank Liang Dai for the useful discussions. S. B. was supported by NASA through Einstein Postdoctoral Fellowship Grant No. PF5-160133. This work was supported by NSF Grant No. 0244990, NASA Grant No. NNX15AB18G, the John Templeton Foundation, and the Simons Foundation.

*sbird4@jhu.edu

- [1] G. Hinshaw *et al.* (WMAP Collaboration), Nine-year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Cosmological parameter results, *Astrophys. J. Suppl. Ser.* **208**, 19 (2013).
- [2] P. A. R. Ade *et al.* (Planck Collaboration), Planck 2015 results. XIII. Cosmological parameters, [arXiv:1502.01589](https://arxiv.org/abs/1502.01589).
- [3] B. J. Carr and S. W. Hawking, Black holes in the early Universe, *Mon. Not. R. Astron. Soc.* **168**, 399 (1974).
- [4] P. Meszaros, The behaviour of point masses in an expanding cosmological substratum, *Astron. Astrophys.* **37**, 225 (1974).
- [5] B. J. Carr, The primordial black hole mass spectrum, *Astrophys. J.* **201**, 1 (1975).
- [6] S. Clesse and J. Garcia-Bellido, Massive primordial black holes from hybrid inflation as dark matter and the seeds of galaxies, *Phys. Rev. D* **92**, 023524 (2015).
- [7] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Observation of Gravitational Waves from a Binary Black Hole Merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
- [8] B. J. Carr, K. Kohri, Y. Sendouda, and J. Yokoyama, New cosmological constraints on primordial black holes, *Phys. Rev. D* **81**, 104019 (2010).
- [9] M. A. Monroy-Rodríguez and C. Allen, The end of the MACHO era, revisited: New limits on MACHO masses from halo wide binaries, *Astrophys. J.* **790**, 159 (2014).

- [10] R. A. Allsman *et al.* (Macho Collaboration), MACHO project limits on black hole dark matter in the 1-30 solar mass range, *Astrophys. J.* **550**, L169 (2001).
- [11] L. Wyrzykowski *et al.*, The OGLE view of microlensing towards the Magellanic Clouds. IV. OGLE-III SMC data and final conclusions on MACHOs, *Mon. Not. R. Astron. Soc.* **416**, 2949 (2011).
- [12] P. Tisserand *et al.* (EROS-2 Collaboration), Limits on the macho content of the galactic halo from the EROS-2 survey of the Magellanic Clouds, *Astron. Astrophys.* **469**, 387 (2007).
- [13] J. Yoo, J. Chaname, and A. Gould, The end of the MACHO era: Limits on halo dark matter from stellar halo wide binaries, *Astrophys. J.* **601**, 311 (2004).
- [14] D. P. Quinn, M. I. Wilkinson, M. J. Irwin, J. Marshall, A. Koch, and V. Belokurov, On the reported death of the MACHO era, *Mon. Not. R. Astron. Soc.* **396**, L11 (2009).
- [15] M. Ricotti, J. P. Ostriker, and K. J. Mack, Effect of primordial black holes on the cosmic microwave background and cosmological parameter estimates, *Astrophys. J.* **680**, 829 (2008).
- [16] M. Ricotti, Bondi accretion in the early Universe, *Astrophys. J.* **662**, 53 (2007).
- [17] Y. Ali-Haïmoud *et al.* (to be published).
- [18] T. Nakamura, M. Sasaki, T. Tanaka, and K. S. Thorne, Gravitational waves from coalescing black hole MACHO binaries, *Astrophys. J.* **487**, L139 (1997).
- [19] G. D. Quinlan and S. L. Shapiro, Dynamical evolution of dense clusters of compact stars, *Astrophys. J.* **343**, 725 (1989).
- [20] H. Mouri and Y. Taniguchi, Runaway merging of black holes: Analytical constraint on the timescale, *Astrophys. J.* **566**, L17 (2002).
- [21] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), The rate of binary black hole mergers inferred from Advanced LIGO observations surrounding GW150914, [arXiv:1602.03842](https://arxiv.org/abs/1602.03842).
- [22] B. Moore, S. Ghigna, F. Governato, G. Lake, T. Quinn, J. Stadel, and P. Tozzi, Dark matter substructure within galactic halos, *Astrophys. J.* **524**, L19 (1999).
- [23] M. Kamionkowski and S. M. Koushiappas, Galactic substructure and direct detection of dark matter, *Phys. Rev. D* **77**, 103509 (2008).
- [24] M. Kamionkowski, S. M. Koushiappas, and M. Kuhlen, Galactic substructure and dark matter annihilation in the Milky Way halo, *Phys. Rev. D* **81**, 043532 (2010).
- [25] J. F. Navarro, C. S. Frenk, and S. D. M. White, The structure of cold dark matter halos, *Astrophys. J.* **462**, 563 (1996).
- [26] F. Prada, A. A. Klypin, A. J. Cuesta, J. E. Betancort-Rijo, and J. Primack, Halo concentrations in the standard LCDM cosmology, *Mon. Not. R. Astron. Soc.* **423**, 3018 (2012).
- [27] A. D. Ludlow *et al.*, The mass-concentration-redshift relation of cold and warm dark matter halos, [arXiv:1601.02624](https://arxiv.org/abs/1601.02624).
- [28] Y. Y. Mao, L. E. Strigari, R. H. Wechsler, H.-Y. Wu, and O. Hahn, Halo-to-halo similarity and scatter in the velocity distribution of dark matter, *Astrophys. J.* **764**, 35 (2013).
- [29] R. M. O'Leary, B. Kocsis, and A. Loeb, Gravitational waves from scattering of stellar-mass black holes in galactic nuclei, *Mon. Not. R. Astron. Soc.* **395**, 2127 (2009).
- [30] M. H. Lee, N-body evolution of dense clusters of compact stars, *Astrophys. J.* **418**, 147 (1993).
- [31] J. Binney and S. Tremaine, *Galactic Dynamics* (Princeton University Press, Princeton, NJ, 1987), p. 747.
- [32] T. Ishiyama, J. Makino, and T. Ebisuzaki, Gamma-ray signal from Earth-mass dark matter microhalos, *Astrophys. J.* **723**, L195 (2010).
- [33] T. Ishiyama, Hierarchical formation of dark matter halos and the free streaming scale, *Astrophys. J.* **788**, 27 (2014).
- [34] W. H. Press and P. Schechter, Formation of galaxies and clusters of galaxies by self-similar gravitational condensation, *Astrophys. J.* **187**, 425 (1974).
- [35] J. Tinker, A. V. Kravtsov, A. Klypin, K. Abazajian, M. Warren, G. Yepes, S. Gottlöber, and D. E. Holz, Toward a halo mass function for precision cosmology: The limits of universality, *Astrophys. J.* **688**, 709 (2008).
- [36] A. Jenkins, C. S. Frenk, S. D. M. White, J. M. Colberg, S. Cole, A. E. Evrard, H. M. P. Couchman, and N. Yoshida, The mass function of dark matter halos, *Mon. Not. R. Astron. Soc.* **321**, 372 (2001).
- [37] G. Efstathiou, Suppressing the formation of dwarf galaxies via photoionization, *Mon. Not. R. Astron. Soc.* **256**, 43P (1992).
- [38] J. Einasto, On the Construction of a Composite Model for the Galaxy and on the Determination of the System of Galactic Parameters, *Tr. Astrofiz. Inst., Alma-Ata*, **5**, 87 (1965).
- [39] B. Abbott *et al.* (LIGO Scientific and VIRGO Collaborations), Prospects for observing and localizing gravitational-wave transients with Advanced LIGO and Advanced Virgo, *Living Rev. Relativity* **19**, 1 (2016).
- [40] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Characterization of transient noise in Advanced LIGO relevant to gravitational wave signal GW150914, [arXiv:1602.03844](https://arxiv.org/abs/1602.03844).
- [41] K. Belczynski, D. E. Holz, T. Bulik, and R. O'Shaughnessy, The origin and evolution of LIGO's first gravitational-wave source, [arXiv:1602.04531](https://arxiv.org/abs/1602.04531).
- [42] I. Cholis *et al.* (to be published).
- [43] D. Pooley, S. Rappaport, J. Blackburne, P. L. Schechter, J. Schwab, and J. Wambsganss, The dark-matter fraction in the elliptical galaxy lensing the quasar PG1115+080, *Astrophys. J.* **697**, 1892 (2009).
- [44] E. Mediavilla, J. A. Muñoz, E. Falco, V. Motta, E. Guerras, H. Canovas, C. Jean, A. Oscoz, and A. M. Mosquera, Microlensing-based estimate of the mass fraction in compact objects in lens, *Astrophys. J.* **706**, 1451 (2009).
- [45] E. Bugaev and P. Klimai, Constraints on the induced gravitational wave background from primordial black holes, *Phys. Rev. D* **83**, 083521 (2011).
- [46] J. B. Muñoz, E. D. Kovetz, L. Dai, and M. Kamionkowski, Lensing of fast radio bursts as a probe of compact dark matter, [arXiv:1605.00008](https://arxiv.org/abs/1605.00008).
- [47] See the Einstein Telescope design at <http://www.et-gw.eu/>.
- [48] N. Seto, S. Kawamura, and T. Nakamura, Possibility of Direct Measurement of the Acceleration of the Universe Using 0.1 Hz Band Laser Interferometer Gravitational Wave Antenna in Space, *Phys. Rev. Lett.* **87**, 221103 (2001).
- [49] E. S. Phinney *et al.*, The big bang observer: Direct detection of gravitational waves from the birth of the Universe to the present, NASA Mission Concept Study, 2003.