



Rashba Splitting of Cooper Pairs

R. I. Shekhter,¹ O. Entin-Wohlman,^{2,3,*} M. Jonson,^{1,4} and A. Aharony^{2,3}

¹*Department of Physics, University of Gothenburg, SE-412 96 Göteborg, Sweden*

²*Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel*

³*Physics Department, Ben Gurion University, Beer Sheva 84105, Israel*

⁴*SUPA, Institute of Photonics and Quantum Sciences, Heriot-Watt University, Edinburgh EH14 4AS, United Kingdom*

(Received 19 February 2016; published 26 May 2016)

We investigate theoretically the properties of a weak link between two superconducting leads, which has the form of a nonsuperconducting nanowire with a strong Rashba spin-orbit coupling caused by an electric field. In the Coulomb-blockade regime of single-electron tunneling, we find that such a weak link acts as a “spin splitter” of the spin states of Cooper pairs tunneling through the link, to an extent that depends on the direction of the electric field. We show that the Josephson current is sensitive to interference between the resulting two transmission channels, one where the spins of both members of a Cooper pair are preserved and one where they are both flipped. As a result, the current is a periodic function of the strength of the spin-orbit interaction and of the bending angle of the nanowire (when mechanically bent); an identical effect appears due to strain-induced spin-orbit coupling. In contrast, no spin-orbit induced interference effect can influence the current through a single weak link connecting two normal metals.

DOI: 10.1103/PhysRevLett.116.217001

Superconducting spintronics [1] is of great interest in modern research on nanophysics owing to its potential for offering both spin control and dissipationless spin transport. In this Letter we consider the Josephson current between two superconductors connected by a weak link. The link suppresses the pairing potential, thus allowing external manipulation of the spin structure of the Cooper pairs. An obvious way to achieve this is to use the Rashba spin-orbit (SO) coupling [2,3], which can be generated by an external electric field. The SO coupling rotates the electron spin around an axis fixed by the electron momentum and the electric field [4]. Quite a number of papers investigated the effect of this interaction on the Josephson current [5–14]. Most found that modifying the Josephson current by the SO interaction necessitates breaking time-reversal symmetry, e.g., by a magnetic-field induced Zeeman splitting [5–9,11] or by magnetic exchange interactions [9,10,14]. Here, we achieve the desired spin control without any magnetic field [15].

A mechanically bent one-dimensional (1D) nanowire subjected to a strong Rashba SO interaction and suspended between two normal bulk conductors was considered in Ref. [16], where it was shown that none of the three components of the electronic spin are a good quantum number. Hence, the transfer of electrons through this *nonsuperconducting* weak link results in a split of the spinor wave function with respect to the two possible spin projections of the incident electrons. Since the dynamic transformation of the spin caused by such a “spin splitter” is fully deterministic—in contrast to the stochastic spin flips due to magnetic impurities—one may expect consequences in the form of various interference phenomena. Here, we

show that the splitting of the spin state of the paired electrons that carry the Josephson current may transform the spin-singlet Cooper pairs into a coherent mixture of singlet and triplet spin states. This mixture gives rise to interference between the channel in which both electrons preserve their spins and the channel where they are both flipped. The resulting interference pattern, which appears in the Josephson current but does not show up in the normal-state transmission of the junction, allows for electrical and mechanical control of the Josephson current between spin-singlet superconductors, corresponding to a new type of “spin gating” [17] of superconducting “weak links”.

To illustrate our calculation, Fig. 1 uses a semiclassical analogue of the quantum evolution of the spin states of electrons which move between two bulk leads via a weak link, where they are subjected to the Rashba SO interaction. For simplicity we assume for now that the weak link is a straight 1D wire along the \hat{x} axis. The SO interaction in the wire is due to an electric field, which, for the moment, is assumed to point along \hat{z} and therefore corresponds to an effective SO-interaction-induced magnetic field directed along \hat{y} . Figure 1(a) illustrates a single-electron transfer from one normal metal to another. Without loss of generality, we choose the \hat{z} axis in spin space to be along the direction of the polarization of the electron in the first (left) metal. Semiclassically, the spin of the injected electron will rotate in the XZ plane as it passes through the wire. As a result, the spins of the electrons that enter the second metal from the wire are rotated around the \hat{y} axis by an angle proportional to the strength of the SO interaction and the length of the wire. This rotation depends on the direction of the “initial” electron’s polarization. It occurs only if the

polarization has a component in the XZ plane. Quantum mechanically, the electron's spinor in the left metal is an eigenfunction of the Pauli spin matrix σ_z , and the spinor of the outgoing electron is in a coherent superposition of spin-up and spin-down eigenstates of σ_z .

How can this picture be generalized to describe the transfer of the two electrons of a Cooper pair between two bulk superconductors? The simplest case to consider, which we focus upon below, is when single-electron tunneling is Coulomb blocked throughout the wire. While the blockade can be lifted for one electron, double electron occupancy of the wire is suppressed; i.e., a Cooper pair is mainly transferred sequentially, as shown in Fig. 1(b). Each electron transfer is now accompanied by the spin rotation shown in Fig. 1(a). However, since the two transferred electrons are in time-reversed quantum states, the time evolution of their spins are reversed with respect to one another, and their rotation angles have opposite signs [step 4 in Fig. 1(b)]. This final state [Fig. 1(c)] can be expressed as a coherent mixture of a spin-singlet and a spin-triplet state, but only the former can enter into the second superconductor. As we show below, this projection onto the singlet causes a reduction of the Josephson current.

We consider a model where a Cooper pair is transferred between superconducting source and drain leads via virtual




FIG. 1. Schematic illustrations of the lowest-order perturbation expansion steps for tunneling (in the Coulomb-blockade regime) through a straight nanowire weak link subjected to the Rashba spin-orbit (SO) interaction caused by an electric field along the \hat{z} direction. In a semiclassical picture, the spin of each electron (the arrows) is rotated in the XZ plane as it goes through the link. (a) Single-electron tunneling from one normal metal to another, via an intermediate (rotating) state (the dashed circle). When the electron enters the second normal metal, its spin has been rotated. (b) Sequential tunneling in four steps of a Cooper pair between two superconductors connected by the same weak link. Because the two electrons that form the Cooper pair are in time-reversed states, the SO interaction rotates their spins in opposite directions. (c) As they enter the second superconductor, the Cooper pairs are in a coherent mixture (the dash-dotted circle) of a spin-singlet and a spin-triplet state. Inside this superconductor, this state is then projected onto the singlet state (the full circle).

states localized in a weak-link wire [see Fig. 2(a)]. The corresponding tunneling process, which supports multiple tunneling channels, was analyzed in detail in Ref. [18]. A significant simplification occurs in the Coulomb-blockade regime, defined by the inequality $E_e = E_C(N+1) - E_C(N) \gg |\Delta|$, where $|\Delta|$ is the energy gap parameter in the superconducting leads [19], and $E_C(N)$ is the Coulomb energy of the wire when it contains N electrons. In this regime, tunneling channels requiring two electrons to be simultaneously localized in a virtual state in the wire can be neglected—hence the sequential transmission. Another simplification follows from our assumption that the length of the wire d is short compared to the superconducting coherence length $\xi_0 \equiv \hbar v_F / |\Delta|$ [19], so that the dependence of the matrix element for a single-electron transfer on the electron energy in the virtual states can be ignored. More details are given in the Supplemental Material [20]. A final simplification, facilitated by the device geometry, concerns the conservation of the electrons' longitudinal momenta as they tunnel between the two leads. In Fig. 2, the wire ends are placed on top of the metal leads and separated from them by thin but long tunneling barriers. Since the direction of tunneling is nearly perpendicular to the direction of the current along the wire, such a geometry is conducive to longitudinal momentum conservation [21].

These simplifying but realistic assumptions allow us to describe the transfer of a Cooper pair between the two superconductors in terms of single-electron tunneling, as given by the Hamiltonian $\mathcal{H} = \mathcal{H}_L + \mathcal{H}_R + \mathcal{H}_T$. Here,

$$\mathcal{H}_{L(R)} = \sum_{\mathbf{k}(\mathbf{p})} \xi_{\mathbf{k}(\mathbf{p})} c_{\mathbf{k}(\mathbf{p})\sigma}^\dagger c_{\mathbf{k}(\mathbf{p})\sigma} + \left(\Delta_{L(R)} \sum_{\mathbf{k}(\mathbf{p})} c_{\mathbf{k}(\mathbf{p})\uparrow}^\dagger c_{-k(-\mathbf{p})\downarrow}^\dagger + \text{H.c.} \right), \quad (1)$$

where $\xi_{\mathbf{k}(\mathbf{p})} = \epsilon_{\mathbf{k}(\mathbf{p})} - \mu$ is the quasiparticle energy in the left (right) bulk superconducting lead with order parameter $\Delta_{L(R)}$, and μ is the common chemical potential. Single-electron tunneling between the leads is described by the tunnel Hamiltonian

$$\mathcal{H}_T = \sum_{\mathbf{k}, \mathbf{p}} \sum_{\sigma, \sigma'} (c_{\mathbf{p}\sigma'}^\dagger [W_{\mathbf{p}, \mathbf{k}}]_{\sigma', \sigma} c_{\mathbf{k}\sigma} + \text{H.c.}), \quad (2)$$




FIG. 2. (a) Sketch and (b) simplified model of a device that would allow the effects predicted in the text to be studied.

where $[W_{\mathbf{p},\mathbf{k}}]_{\sigma,\sigma'} = ([W_{-\mathbf{p},-\mathbf{k}}]_{-\sigma,-\sigma'})^*$ are elements of a matrix in spin space which obey time-reversal symmetry [22]. The operator $c_{\mathbf{k}(\mathbf{p})\sigma}^\dagger$ creates an electron in the left (right) lead, with momentum $\mathbf{k}(\mathbf{p})$ and a spin index σ , which denotes the eigenvalue of the spin projection along an arbitrary axis. The Cooper pairs in both superconducting leads are formed by electrons whose spin projections on their common quantization axis have opposite signs, denoted by σ and $\bar{\sigma}$. The Hamiltonian (2) describes how single electrons are transferred through the nanowire weak link, while their spin states evolve under the SO interaction, which is geometrically confined to the wire. Such tunneling amplitudes, for a normal-state junction, were considered before [23,24].

We assume a weak link containing a bent wire [see Fig. 2(a)]. The actual calculations are done for the geometry shown in Fig. 2(b), where the weak link comprises two straight 1D wires, \mathbf{r}_L and \mathbf{r}_R , of equal length $d/2$, connected by a ‘‘bend.’’ The angles between these wires and the $\hat{\mathbf{x}}$ axis are θ and $-\theta$, respectively. As shown below, the resulting Josephson current depends on θ , allowing its mechanical manipulation. The Rashba Hamiltonian for a wire lying in the XY plane is

$$\mathcal{H}_{\text{wire}} = -\frac{1}{2m^*} \frac{d^2}{d\mathbf{r}^2} - i \frac{k_{\text{so}}}{m^*} \hat{\mathbf{n}} \cdot \boldsymbol{\sigma} \times \frac{d}{d\mathbf{r}}, \quad (3)$$

where \mathbf{r} is the coordinate along a straight wire segment, $\boldsymbol{\sigma}$ is the vector of the Pauli matrices, m^* is the effective electronic mass, and k_{so} is the strength of the SO interaction in momentum units (using $\hbar = 1$), related to an electric field along $\hat{\mathbf{n}}$ that generates the SO coupling [2–4]. For this configuration, the tunneling amplitude, a 2×2 matrix in spin space, is [20]

$$W_{\mathbf{k},\mathbf{p}}/W_0 \equiv \mathcal{W}, \quad \mathcal{W} = e^{-ik_{\text{so}}\boldsymbol{\sigma}\cdot\mathbf{r}_L \times \hat{\mathbf{n}}} e^{-ik_{\text{so}}\boldsymbol{\sigma}\cdot\mathbf{r}_R \times \hat{\mathbf{n}}}. \quad (4)$$

In Eq. (4), $W_0 = G_0(|\mathbf{r}_L|; E) \mathcal{T} G_0(|\mathbf{r}_R|; E)$ comprises all of the characteristics of the tunneling matrix that are independent of the spin dynamics, where \mathcal{T} is the transfer matrix through the bend in the wire, and $G_0(|\mathbf{r}|; E) = -i(\pi m/\tilde{\kappa}) \exp[i\tilde{\kappa}|\mathbf{r}|]$ is the free propagator modified by the SO interaction according to $\tilde{\kappa} = (2mE + \tilde{k}_{\text{so}}^2)^{1/2}$. Here, $\tilde{k}_{\text{so}} = k_{\text{so}} \sqrt{1 - (\hat{\mathbf{n}} \cdot \hat{\mathbf{u}})^2}$, where $\hat{\mathbf{u}}$ is a unit vector along the direction of the straight segment. The unitary matrix \mathcal{W} performs two consecutive spin rotations around the axes $\mathbf{r}_L \times \hat{\mathbf{n}}$ and $\mathbf{r}_R \times \hat{\mathbf{n}}$, as described (for a single segment) semiclassically in Fig. 1(a). Equation (4) is derived to the lowest possible order in the tunneling; the explicit dependence of W_0 on the momenta is omitted for brevity [20].

The SO interaction modifies significantly the amplitude of the Josephson equilibrium current, while leaving the transmission of the junction in its normal state as in the absence of this coupling. The matrix \mathcal{W} that determines these quantities depends crucially on the direction $\hat{\mathbf{n}}$ of the

electric field. In the ‘‘genuine’’ Rashba configuration, $\hat{\mathbf{n}}$ is normal to the plane of the junction; this is the case described semiclassically in Fig. 1. For $\hat{\mathbf{n}} \parallel \hat{\mathbf{z}}$,

$$\begin{aligned} \mathcal{W} = & [\cos^2(k_{\text{so}}d/2) - \sin^2(k_{\text{so}}d/2) \cos(2\theta)] \\ & + i\boldsymbol{\sigma} \cdot [\hat{\mathbf{y}} \sin(k_{\text{so}}d) \cos(\theta) + \hat{\mathbf{z}} \sin^2(k_{\text{so}}d/2) \sin(2\theta)]. \end{aligned} \quad (5)$$

In contrast, when the electric field is in the plane of the junction, e.g., $\hat{\mathbf{n}} = \hat{\mathbf{y}}$, we find

$$\mathcal{W} = \cos[k_{\text{so}}d \cos(\theta)] - i\boldsymbol{\sigma} \cdot \hat{\mathbf{z}} \sin[k_{\text{so}}d \cos(\theta)]. \quad (6)$$

Spin-orbit coupling may also be induced by *strain* [25]; in that case, the last term in Eq. (3) is replaced by $\mathcal{H}_{\text{so}} = (\Delta_{\text{so}}^{\text{strain}}/2) \hat{\mathbf{k}} \cdot \boldsymbol{\sigma}$, where $\hat{\mathbf{k}}$ is a unit vector along the momentum (i.e., along the straight segment) [26,27]. By defining $\Delta_{\text{so}}^{\text{strain}}/2 = \hbar v_F k_{\text{so}}^{\text{strain}}$, one finds that \mathcal{W} of the strain-induced case has the same form as Eq. (5), except that $\hat{\mathbf{y}}$ is replaced by $\hat{\mathbf{x}}$. The resulting expressions for the Josephson current and for the normal-state transmission turn out to be the same as for the SO interaction in the genuine Rashba configuration.

A calculation of the current emerging from the left lead, $I_L = -e \langle \dot{N}_L \rangle = -ie \langle [\mathcal{H}, \sum_{\mathbf{k},\sigma} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma}] \rangle$, up to second order in \mathcal{H}_T , is straightforward [20]. It differs from the standard procedure [28] only in that the spin dynamics caused by the SO interaction must be properly taken into account by allowing the tunneling of electrons to be spin dependent. The current comprises the supercurrent $J(\varphi)$ and the quasiparticle current.

In the absence of the SO interaction, the quasiparticle current scales as the transmission of the junction when in the normal state [28,29]. The SO coupling modifies this transmission by the factor $\text{Tr}\{\mathcal{W}\mathcal{W}^\dagger\}$, where the trace is in spin space [20]. This factor is simply 2, the spin degeneracy [see Eq. (4)]; i.e., the SO interaction does not affect the electric conductance (unless the junction allows for geometrically interfering processes [24]). The superconducting Josephson current is

$$\frac{J(\varphi)}{J_0(\varphi)} = \frac{1}{2} \sum_{\sigma} [|\mathcal{W}_{\sigma\sigma}|^2 - |\mathcal{W}_{\sigma\bar{\sigma}}|^2] = \sum_{\sigma} \left[\frac{1}{2} - |\mathcal{W}_{\sigma\bar{\sigma}}|^2 \right], \quad (7)$$

where $J_0(\varphi) \propto \sin(\varphi)$ is the equilibrium Josephson current in the absence of the SO interaction [29], and φ is the superconducting phase difference.

The matrix element $\mathcal{W}_{\sigma\bar{\sigma}}$ depends on the quantization axis of the spins. Choosing this axis to be along $\hat{\mathbf{z}}$, the genuine Rashba configuration implies that $\hat{\mathbf{n}} \parallel \hat{\mathbf{z}}$, and Eq. (5) yields (cf. Fig. 3)

$$1 - 2|\mathcal{W}_{\sigma\bar{\sigma}}|^2 = 1 - 2\cos^2(\theta) \sin^2(k_{\text{so}}d). \quad (8)$$




FIG. 3. The Josephson current $J(\varphi)$ divided by its value without the SO interaction, $J_0(\varphi)$, for the genuine Rashba configuration [Eqs. (7) and (8)] as a function of $k_{\text{so}}d/(2\pi)$. The largest amplitude is for the zero bending angle, $\theta = 0$, decreasing gradually for $\theta = \pi/6, \pi/5, \pi/4, \pi/3, \pi/2.5$ [Fig. 2(b)]. Relevant values of k_{so} are estimated in the text.

The Josephson current is thus significantly modified. In contrast, when the electric field is in the plane of the junction, $\hat{\mathbf{n}} = \hat{\mathbf{y}}$, the matrix \mathcal{W} is diagonal [Eq. (6)], $J(\varphi) = J_0(\varphi)$, and the superconducting current is not affected by the spin dynamics. Similar qualitative results are found [20] for all of the directions of the spin quantization axis. The splitting of the Cooper-pair spin state by the SO interaction reduces the Josephson current through the superconducting weak link under consideration.

Two features determine the magnitude of the effect for a given spin quantization axis in the leads (in addition to the strength k_{so} of the SO interaction and the length d over which it acts). One is the extent to which the nanowire is bent [θ in Fig. 2(b)], and the other is the orientation $\hat{\mathbf{n}}$ of the electric field responsible for the SO coupling relative to the spin quantization axis. Both break spin conservation, which results in Rabi oscillations between the singlet and triplet spin states of the (originally spin-singlet) Cooper pairs passing through the spin-orbit-active weak link. The consequence is a spin splitting of the Cooper pairs that reach the second superconducting lead, where their spin state is projected onto the singlet state. This splitting can result in a Josephson current that is an oscillatory function of the “action” $k_{\text{so}}d$ of the SO interaction (Fig. 3); the effect may be absent for special directions of the electric field. Both results can be understood in terms of a semiclassical picture (Fig. 1).

As seen in Eq. (7), the Josephson current can be written as a sum of two contributions. One, $|\mathcal{W}_{\sigma\sigma}|^2$, comes from a channel where the spin projections of the Cooper-pair electrons, when leaving and entering the weak link, are identical; the other, $|\mathcal{W}_{\sigma\bar{\sigma}}|^2$, arises from another channel, where both electron spins are flipped during the passage. It is remarkable that the two contributions have opposite signs. This is due to a Josephson tunneling “ π shift” caused by electronic spin flips (and is similar to the effect predicted

for tunneling through a Kondo impurity [30]). In particular, a total cancellation of the Josephson current is possible when, e.g., $\theta = 0$ and $k_{\text{so}}d = \pi/4$; in the limit $\theta = 0$ and $k_{\text{so}}d = \pi/2$, the Josephson current even changes its sign. This spin-orbit induced interference effect on the Josephson current is specific to a weak link subjected to SO interaction between superconductors. There is no such effect on the current through a single weak link connecting two normal metals.

According to Eq. (7), none or both of the Cooper pair electrons must have flipped their spins as they leave the weak link in order to contribute to the Josephson current. This is because only spin-singlet Cooper pairs can enter the receiving s -type bulk superconductor. However, single-flip processes, where only one of the two tunneling electrons flips its spin, are also possible results of injecting Cooper pairs into a Rashba weak link. Those processes correspond to a triplet component of the spin state of the transferred pair, and they can be viewed as evidence for spin polarization of injected Cooper pairs. The triplet component could be responsible for a spin-triplet proximity effect [15] and would presumably contribute a spin supercurrent if higher-order tunneling processes were taken into account.

In conclusion, we have shown that the Josephson current through a weak-link nanowire, locally subjected to the Rashba SO interaction, is sensitive to both the amount of bending and to the orientation of the electric field generating the SO coupling. This allows for a tuning of the supercurrent by mechanical and electrical manipulations of the spin polarization of the Cooper pairs. Specifically, the Josephson current through an electrostatically gated device becomes an oscillatory function of the gate voltage. We emphasize that these results follow from the interference of two transmission channels, one where the spins of both members of a Cooper pair are preserved and one where they are both flipped, and that this interference does not require any external magnetic field. It is important, however, that those parts of the device where the superconducting pairing potential is nonzero and where the SO coupling is finite are spatially separated. To lowest order in the tunneling, this separation prevents the superconductivity in the leads to have any effect on the dynamical spin evolution in the wire.

Carbon nanotubes and semiconductor wires seem particularly suitable to be used as spin splitters. Measured Rashba spin-orbit-coupling induced energy gaps in InGaAs/InAlAs ($\Delta_{\text{so}} = 2\hbar v_F k_{\text{so}} \approx 5$ meV) [31] and InAs/AlSb ($\Delta_{\text{so}} \approx 4$ meV) [32] quantum wells correspond to $k_{\text{so}} \approx 4 \times 10^6$ m $^{-1}$. The strain-induced SO energy gap for a carbon nanotube is $\Delta_{\text{so}}^{\text{strain}} = 2\hbar v_F k_{\text{so}}^{\text{strain}} \approx 0.4$ meV, corresponding to $k_{\text{so}}^{\text{strain}} \approx 0.4 \times 10^6$ m $^{-1}$ for $v_F \approx 0.5 \times 10^6$ m/sec [33]. For d of the order of μm , $k_{\text{so}}^{\text{(strain)}}d$ can therefore be of the order 1–5.

This work was partially supported by the Swedish Research Council (VR), by the Israel Science

Foundation (ISF), and by the infrastructure program of the Israel Ministry of Science and Technology under Contract No. 3-11173. O. E. W. and A. A. thank the Department of Physics at the University of Gothenburg for its hospitality.

*oraentin@bgu.ac.il

- [1] J. Linder and J. W. A. Robinson, Superconducting spintronics, *Nat. Phys.* **11**, 307 (2015).
- [2] E. I. Rashba, *Fiz. Tverd. Tela (Leningrad)* **2**, 1224 (1960) [*Sov. Phys. Solid State* **2**, 1109 (1960)]; Y. A. Bychkov and E. I. Rashba, Oscillatory effects and the magnetic susceptibility of carriers in inversion layers, *J. Phys. C* **17**, 6039 (1984).
- [3] R. Winkler, *Spin-Orbit Coupling Effects in Two-Dimensional Electron and Hole Systems* (Springer-Verlag, Berlin, 2003).
- [4] S. Datta and B. Das, Electronic analog of the electro-optic modulator, *Appl. Phys. Lett.* **56**, 665 (1990).
- [5] E. V. Bezuglyi, A. S. Rozhavsky, I. D. Vagner, and P. Wyder, Combined effect of Zeeman splitting and spin-orbit interaction on the Josephson current in a superconductor-two-dimensional electron gas-superconductor structure, *Phys. Rev. B* **66**, 052508 (2002).
- [6] I. V. Krive, L. Y. Gorelik, R. I. Shekhter, and M. Jonson, Chiral symmetry breaking and the Josephson current in a ballistic superconductor-quantum wire-superconductor junction, *Low Temp. Phys.* **30**, 398 (2004); I. V. Krive, A. M. Kadigrobov, R. I. Shekhter, and M. Jonson, Influence of the Rashba effect on the Josephson current through superconductor/Luttinger liquid/superconductor tunnel junction, *Phys. Rev. B* **71**, 214516 (2005).
- [7] L. Dell'Anna, A. Zazunov, R. Egger, and T. Martin, Josephson current through a quantum dot with spin-orbit coupling, *Phys. Rev. B* **75**, 085305 (2007); A. Zazunov, R. Egger, T. Jonckheere, and T. Martin, Anomalous Josephson Current through a Spin-Orbit Coupled Quantum Dot, *Phys. Rev. Lett.* **103**, 147004 (2009); A. Brinetti, A. Zazunov, A. Kundu, and R. Egger, Anomalous Josephson current, incipient time reversed symmetry breaking, and Majorana bound states in interacting multilevel dots, *Phys. Rev. B* **88**, 144515 (2013); T. Yokoyama, M. Eto, and Yu. V. Nazarov, Anomalous Josephson effect induced by spin-orbit interaction and Zeeman effect in semiconductor nanowires, *Phys. Rev. B* **89**, 195407 (2014).
- [8] A. A. Reynoso, Gonzalo Usaj, C. A. Balseiro, D. Feinberg, and M. Avignon, Anomalous Josephson Current in Junctions with Spin Polarizing Quantum Point Contacts, *Phys. Rev. Lett.* **101**, 107001 (2008).
- [9] A. Buzdin, Direct Coupling between Magnetism and Superconducting Current in the Josephson φ_0 Junction, *Phys. Rev. Lett.* **101**, 107005 (2008); S. V. Mironov, A. S. Mel'nikov, and A. I. Buzdin, Double Path Interference and Magnetic Oscillations in Cooper Pair Transport through a Single Nanowire, *Phys. Rev. Lett.* **114**, 227001 (2015).
- [10] S. H. Jacobsen and J. Linder, *Phys. Rev. B* **92**, 024501 (2015).
- [11] G. Campagnano, P. Lucignano, D. Giuliano, and A. Tagliacozzo, Spin-orbit coupling and anomalous Josephson effect in nanowires, *J. Phys. Condens. Matter* **27**, 205301 (2015).
- [12] B. Béri, J. H. Bardarson, and C. W. J. Beenakker, Splitting of Andreev levels in a Josephson junction by spin-orbit coupling, *Phys. Rev. B* **77**, 045311 (2008).
- [13] O. V. Dimitrova and M. V. Feigel'man, Two-dimensional S-N-S junction with Rashba spin-orbit coupling, *J. Exp. Theor. Phys.* **102**, 652 (2006).
- [14] A. V. Samokhvalov, R. I. Shekhter, and A. I. Buzdin, Stimulation of a singlet superconductivity in SFS weak links by spin-exchange scattering of Cooper pairs, *Sci. Rep.* **4**, 5671 (2014).
- [15] Spin-triplet pairing correlations in a conductor with SO interactions in the proximity of an *s*-wave superconductor were found in the absence of a magnetic field by C. P. Reeg and D. L. Maslov, Proximity-induced triplet superconductivity in Rashba materials, *Phys. Rev. B* **92**, 134512 (2015).
- [16] R. I. Shekhter, O. Entin-Wohlman, and A. Aharony, Suspended Nanowires as a Mechanically Controlled Rashba Spin Splitters, *Phys. Rev. Lett.* **111**, 176602 (2013); Mechanically controlled spin-selective transport, *Phys. Rev. B* **90**, 045401 (2014).
- [17] R. I. Shekhter and M. Jonson, Spin gating of mesoscopic devices, *Synth. Met.* **216**, 2 (2016).
- [18] M. Gissel-fält, Parity effects in Josephson tunneling through a metallic island, *Phys. Scr.* **54**, 397 (1996); see, in particular, Eqs. (18) and (39).
- [19] In order to keep the estimates simple, it is assumed that both the Fermi velocities, v_F , and the superconducting gaps of the two leads are the same.
- [20] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.116.217001> for a detailed derivation.
- [21] Since electron tunneling is confined to a restricted interval Δx at one or the other end of the wire, where the overlap between the wave functions in the lead and the wire is finite, there is an uncertainty $\Delta k \sim 1/\Delta x$ in the longitudinal momentum of the tunneling electrons. We view this uncertainty as a splitting of the incoming well-defined momentum state into several channels k within a range Δk . This enables cotunneling processes, where the momentum of the electron entering the wire from the left electrode and the momentum of the electron leaving the wire for the right electrode are different. However, for the geometry in Fig. 2, the direction of tunneling is nearly orthogonal to that of the current. This is why Δx can be chosen to be large enough to suppress quantum fluctuations of the longitudinal electron momentum (and therefore cotunneling). A resonant tunneling effect exploiting this longitudinal momentum conservation was recently observed by L. Britnell, R. V. Gorbachev, A. K. Geim, L. A. Ponomarenko, A. Mishchenko, M. T. Greenaway, T. M. Fromhold, K. S. Novoselov, and L. Eaves, Resonant tunneling and negative differential conductance in graphene transistors, *Nat. Commun.* **4**, 1794 (2013).
- [22] P. W. Anderson, Special effects in superconductivity, in *Lectures on the Many-Body Problem*, Vol. 2, edited by E. R. Caianiello (Academic Press, New York, 1964), p. 113.
- [23] T. V. Shahbazyan and M. E. Raikh, Low-Field Anomaly in 2D Hopping Magnetoresistance Caused by Spin-Orbit

- Term in the Energy Spectrum, *Phys. Rev. Lett.* **73**, 1408 (1994).
- [24] O. Entin-Wohlman, A. Aharony, Y.M. Galperin, V.I. Kozub, and V. Vinokur, Orbital ac Spin-Hall Effect in the Hopping Regime, *Phys. Rev. Lett.* **95**, 086603 (2005); A. Aharony, Y. Tokura, G.Z. Cohen, O. Entin-Wohlman, and S. Katsumoto, Filtering and analyzing mobile qubit information via Rashba-Dresselhaus-Aharonov-Bohm interferometers, *Phys. Rev. B* **84**, 035323 (2011).
- [25] We thank one of the referees for pointing this out.
- [26] M. S. Rudner and E. I. Rashba, *Phys. Rev. B* **81**, 125426 (2010); K. Flensberg and C. M. Marcus, *Phys. Rev. B* **81**, 195418 (2010).
- [27] In this case, the electric field is a host crystal field; see L. Chico, M. P. López-Sancho, and M. C. Muñoz, *Phys. Rev. B* **79**, 235423 (2009).
- [28] I. O. Kulik and I. K. Janson, *The Josephson Effect in Superconductive Tunneling Structures* (Israel Program for Scientific Translations, Jerusalem, 1972).
- [29] V. Ambegaokar and A. Baratoff, Tunneling between Superconductors, *Phys. Rev. Lett.* **10**, 486 (1963); **11**, 104(E) (1963).
- [30] L. I. Glazman and K. A. Matveev, Resonant Josephson current through Kondo impurities in a tunnel barrier, *Pis'ma Zh. Eksp. Teor. Fiz.* **49**, 570 (1989) [*JETP Lett.* **49**, 660 (1989)]; B. I. Spivak and S. A. Kivelson, Negative local superfluid densities: The difference between superconductors and dirty Bose liquids, *Phys. Rev. B* **43**, 3740 (1991).
- [31] J. Nitta, T. Akazaki, H. Takayanagi, and T. Enoki, Gate Control of Spin-Orbit Interaction in an Inverted $\text{In}_{0.53}\text{Ga}_{0.47}\text{As}/\text{In}_{0.52}\text{Al}_{0.48}\text{As}$ Heterostructure, *Phys. Rev. Lett.* **78**, 1335 (1997).
- [32] J. P. Heida, B. J. van Wees, J. J. Kuipers, T. M. Klapwijk, and G. Borghs, Spin-orbit interaction in a two-dimensional electron gas in a InAs/AlSb quantum well with gate-controlled electron density, *Phys. Rev. B* **57**, 11911 (1998).
- [33] F. Kuemmeth, S. Ilani, D. C. Ralph, and P. L. McEuen, Coupling of spin and orbital motion of electrons in carbon nanotubes, *Nature (London)* **452**, 448 (2008).