Isotensor Axial Polarizability and Lattice QCD Input for Nuclear Double- β Decay Phenomenology

Phiala E. Shanahan,¹ Brian C. Tiburzi,^{2,3} Michael L. Wagman,^{4,5} Frank Winter,⁶ Emmanuel Chang,⁵ Zohreh Davoudi,¹ William Detmold,¹ Kostas Orginos,^{7,6} and Martin J. Savage⁵

(NPLQCD Collaboration)

¹Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

²Department of Physics, The City College of New York, New York, New York 10031, USA

³Graduate School and University Center, The City University of New York,

New York, New York 10016, USA

⁴Department of Physics, University of Washington, Box 351560, Seattle, Washington 98195, USA

⁵Institute for Nuclear Theory, University of Washington, Seattle, Washington 98195-1550, USA

⁶Jefferson Laboratory, 12000 Jefferson Avenue, Newport News, Virginia 23606, USA

¹Department of Physics, College of William and Mary, Williamsburg, Virginia 23187-8795, USA

(Received 27 February 2017; published 10 August 2017)

The potential importance of short-distance nuclear effects in double- β decay is assessed using a lattice QCD calculation of the $nn \rightarrow pp$ transition and effective field theory methods. At the unphysical quark masses used in the numerical computation, these effects, encoded in the isotensor axial polarizability, are found to be of similar magnitude to the nuclear modification of the single axial current, which phenomenologically is the quenching of the axial charge used in nuclear many-body calculations. This finding suggests that nuclear models for neutrinoful and neutrinoless double- β decays should incorporate this previously neglected contribution if they are to provide reliable guidance for next-generation neutrinoless double- β decay searches. The prospects of constraining the isotensor axial polarizabilities of nuclei using lattice QCD input into nuclear many-body calculations are discussed.

DOI: 10.1103/PhysRevLett.119.062003

Double- β ($\beta\beta$) decays of nuclei are of significant phenomenological interest; they probe fundamental symmetries of nature and admit both tests of the standard model (SM) and investigations of physics beyond it [1]. Consequently, these decays are the subject of intense experimental study, and next-generation $\beta\beta$ -decay experiments are currently being planned [2-4]. At present, both the robust prediction of the efficacy of different detector materials, necessary for optimal design sensitivity, and the robust interpretation of the highly sought-after neutrinoless $\beta\beta$ -decay ($0\nu\beta\beta$) mode are impeded by the lack of knowledge of second-order weak-interaction nuclear matrix elements. These quantities bear uncertainties from nuclear modeling that are both significant and difficult to quantify [5]. Controlling the nuclear uncertainties in $\beta\beta$ -decay matrix elements by connecting the nuclear many-body methods to the underlying parameters of the SM is a critical task for nuclear theory.

In this Letter, lattice QCD and pionless effective field theory $[EFT(\pi))]$ are used to investigate the strong-interaction uncertainties in the second-order weak transition of the two-nucleon system in the SM by determining the threshold transition matrix element for $nn \rightarrow pp$. This matrix element receives long-distance contributions from the deuteron intermediate state whose size is governed by the squared magnitude of the $\langle pp | \tilde{J}^+_{\mu} | d \rangle$ matrix element of the axial current that has been recently calculated using lattice quantum chromodynamics (LQCD) [6]. In that work, the two-body contribution to the matrix element (i.e., that beyond the coupling of the axial current to a single nucleon) was constrained, quantifying the effective modification (quenching) of the axial charge of the nucleon from two-body effects. Here, it is highlighted that the $nn \rightarrow$ *pp* matrix element receives additional short-distance contributions beyond those in $|\langle pp | \tilde{J}^+_{\mu} | d \rangle|^2$ arising from the two axial currents being separated by $r < \Lambda^{-1} \sim m_{\pi}^{-1}$ [where Λ is the cutoff scale of EFT(π)], referred to herein as the isotensor axial polarizability. Using $EFT(\pi)$ to analyze both the second-order weak transition calculated here for the first time and the first-order $\langle pp | \tilde{J}_{\mu}^{+} | d \rangle$ transition [6], the short- and long-distance contributions to the $nn \rightarrow pp$ matrix element are separately determined. Interestingly, the short-distance contribution to the total matrix element from the axial polarizability is found to be of comparable size (within the uncertainties of the present calculation) to the two-nucleon current contribution to $|\langle pp|\tilde{J}^+_{\mu}|d\rangle|^2$. Described in phenomenological terms, the polarizability thus appears to be as important as the effective quenching of the axial charge in the two-nucleon system.

The numerical calculations in this work are performed at unphysical values of the quark masses and for a disallowed decay. While there is no immediate phenomenological impact of the numerical values of the matrix elements that are extracted, the observed behavior does provide an important lesson for many-body calculations. In typical calculations of two-neutrino $(2\nu\beta\beta)$ decay, the nuclear matrix elements are calculated using two insertions of the axial current in a truncated model space, with a quenched value of g_A tuned to reproduce experiment. If the findings presented here persist at the physical values of the quark masses, they would imply that a potentially significant contribution has been ignored in standard $2\nu\beta\beta$ calculations, resulting in a source of uncertainty in the nuclear matrix elements that remains to be quantified. Importantly, this uncertainty can only be constrained using $\beta\beta$ -decay measurements or numerical calculations. In $0\nu\beta\beta$ decays, the situation becomes even less certain, in part due to dependence on possible scenarios of physics beyond the SM. With a light Majorana neutrino, generalizations of the axial polarizability will also likely be relevant.

In what follows, the lattice QCD and EFT(π) calculations and the analysis of the axial polarizability are summarized, with complete details presented in a subsequent paper [7]. The potential for future lattice QCD calculations to provide the necessary input to constrain many-body calculations of $2\nu\beta\beta$ and $0\nu\beta\beta$ matrix elements, and thereby reduce the uncertainties in calculated $\beta\beta$ -decay rates, is also discussed.

Two-neutrino $\beta\beta$ *decay.*—The focus of this Letter is on $2\nu\beta\beta$ decay of the dinucleon system. The decay width is given by

$$[T_{1/2}^{2\nu}]^{-1} = G_{2\nu}(Q) |M_{\rm GT}^{2\nu}|^2, \tag{1}$$

where $Q = E_{nn} - E_{pp}$, $G_{2\nu}(Q)$ is a known phase-space factor [8,9], and the Gamow-Teller matrix element in the two-nucleon system is

$$M_{\rm GT}^{2\nu} = 6\left(\frac{1}{2}\int d^4x d^4y \langle pp|T[J_3^+(x)J_3^+(y)]|nn\rangle\right)$$
$$= 6\sum_{\mathfrak{l}'} \frac{\langle pp|\tilde{J}_3^+|\mathfrak{l}'\rangle \langle \mathfrak{l}'|\tilde{J}_3^+|nn\rangle}{E_{\mathfrak{l}'} - (E_{nn} + E_{pp})/2}.$$
(2)

Here, $J_3^+ = (J_3^1 + iJ_3^2)/\sqrt{2}$ is the third component of the $\Delta I_3 = 1$ axial-vector current, $J_{\mu}^a(x) = \bar{q}(x)\gamma_{\mu}\gamma_5(\tau^a/2)q(x)$, and \mathfrak{l}' indexes a complete set of zero-momentum hadronic states with the quantum numbers of the deuteron. The factors of 6 in Eq. (2) are due to rotational symmetry and our normalization of the currents. We employ $\tilde{J}_3^+ =$

 $\int d\mathbf{x} J_3^+(\mathbf{x}, t=0)$ to denote the zero-momentum current at t=0.

As with forward Compton scattering, the amplitude can be written in terms of a Born term, corresponding to an intermediate deuteron state, and the isotensor axial polarizability which absorbs the contributions from the remaining states in the above summation. By isospin symmetry, this polarizability is most cleanly identified as the forward matrix element of the I = 2, $I_3 = 0$ component of the timeordered product of two axial-vector currents in the 1S_0 npground state with the deuteron pole (the Born term) omitted. For use below, isospin relations allow this matrix element to be written as

$$\langle pp|J_{3}^{+}(x)J_{3}^{+}(y)|nn\rangle = \langle np|J_{3}^{(u)}(x)J_{3}^{(u)}(y)|np\rangle -\frac{1}{2}\langle nn|J_{3}^{(u)}(x)J_{3}^{(u)}(y)|nn\rangle -\frac{1}{2}\langle nn|J_{3}^{(d)}(x)J_{3}^{(d)}(y)|nn\rangle, \quad (3)$$

where $J_{3}^{(q)}(x) = \bar{q}(x)\gamma_{3}\gamma_{5}q(x)$.

Pionless effective field theory.—EFT(π) [10–15] efficiently describes two-nucleon systems in the regime where momenta are small compared to the pion mass. This is an appropriate tool with which to address $2\nu\beta\beta$ decays at heavier quark masses, but the inclusion of explicit pion degrees of freedom will likely be required at the physical quark masses $(0\nu\beta\beta)$ decay probes higher momenta, $k \sim 100$ MeV, in large nuclei and likely also requires an EFT with explicit pion degrees of freedom). In what follows, the dibaryon formalism of $EFT(\pi)$ is utilized, using the conventions for the strong-interaction sector described in Ref. [15]. The nucleon degrees of freedom are encoded in the field N, and the two-nucleon degrees of freedom enter as the isosinglet t_i and isotriplet s_a dibaryon fields while y_t and y_s describe the couplings between two nucleons and the corresponding dibaryon fields. In this formalism, the single axial-current interactions enter through the Lagrangian [16–19]

$$\mathcal{L}^{(1)} = -\frac{g_A}{2} N^{\dagger} W_3^a \sigma_3 \tau^a N + \left(g_A - \frac{\tilde{l}_{1,A}}{2M\sqrt{r_s r_t}} \right) (W_3^a t_3^{\dagger} s^a + \text{H.c.}), \quad (4)$$

where $r_{s(t)}$ is the effective range in the ${}^{1}S_{0}$ (${}^{3}S_{1}$) twonucleon channel, σ_{i} (τ^{a}) are Pauli matrices in spin (flavor) space, g_{A} and $\tilde{l}_{1,A}$ are the one- and two-nucleon axial couplings, and W_{3}^{a} is an axial isovector field aligned in the j = 3 spatial direction. The second term is constructed so that $\tilde{l}_{1,A}$ corresponds to a purely two-body current effect. The second-order isotensor axial interaction in the ${}^{1}S_{0}$ channel enters as

$$\mathcal{L}^{(2)} = -\left(\frac{Mg_A^2}{4\gamma_s^2} + \frac{\tilde{h}_{2,S}}{2Mr_s}\right) \mathcal{W}^{ab} s^{a\dagger} s^b, \tag{5}$$

where $W_3^{ab} = W_3^{\{a\}} W_3^{b\}}$ is the traceless symmetric combination of two axial fields at the same location, $\tilde{h}_{2,S}$ is the scalar isotensor weak two-nucleon coupling, and $\gamma_s = \sqrt{MB_{nn}}$ with the binding energy of the 1S_0 system being B_{nn} (at the unphysical masses used herein, the 1S_0 system is bound [20]).

Calculation of the $nn \rightarrow pp$ amplitude, presented in detail in Ref. [7], shows that

$$M_{nn \to pp} = -\frac{|\langle pp | \tilde{J}_3^+ | d \rangle|^2}{\Delta} + \frac{Mg_A^2}{4\gamma_s^2} - \mathbb{H}_{2,S}, \qquad (6)$$

where $\Delta = E_{nn} - E_d$ is the difference of the ground-state ${}^{1}S_{0}$ and ${}^{3}S_{1}$ energies. The parameter $\mathbb{H}_{2,S}$ is the shortdistance two-nucleon, two-axial current coupling $\tilde{h}_{2,S}$ of Eq. (5), redefined and rescaled to capture the effects beyond the deuteron pole and two-nucleon states at energies below Λ [7]. This expression depends on both the long-distance contribution from the deuteron pole (the first term) and the short-distance contributions encapsulated in the second and third terms. (where "short-distance" here means the "nonadiabatic" contribution; that is, every process other than those proceeding via the bound ${}^{3}S_{1}$ ground state). The deuteron-pole contribution includes the effective quenching of the axial charge through $l_{1,A}$. A determination of the $nn \rightarrow pp$ transition matrix element, along with an extraction of the $pp \rightarrow d$ amplitude allows for the isolation of the unknown short-distance contribution, $\mathbb{H}_{2,S}$. Once this counterterm is determined, few-body methods based on $EFT(\pi)$ or matched to them (for example, see Ref. [21]) can incorporate the axial polarizability in computations of decay rates of larger nuclei.

Lattice QCD calculations.—The present lattice QCD calculations extend those of the *pp*-fusion cross section and tritium β decay in Ref. [6]. The same hadronic correlators calculated in the presence of external axial fields are analyzed further to access second-order weak responses to the external field. Recent calculations by the RBC/UKQCD Collaboration in the kaon sector [22–25] have demonstrated that long-distance second-order weak effects can be constrained using lattice QCD. These methods are extended to determine the second-order weak matrix elements of the two-nucleon system.

As discussed in Ref. [6], calculations are performed on one ensemble of gauge-field configurations generated using a Lüscher-Weisz gauge action [26] and a cloverimproved fermion action [27] with $N_f = 3$ degenerate flavors of quarks. The quark masses are tuned to the physical strange-quark mass, producing a pion of mass $m_{\pi} \approx 806$ MeV. The ensemble has a spacetime volume of $L^3 \times T = 32^3 \times 48$ and a gauge coupling that corresponds to a lattice spacing of $a \sim 0.145$ fm. For these calculations, 437 configurations spaced by 10 hybrid Monte Carlo trajectories are used and seven different sets of compound propagators are generated from 16 smeared sources on each configuration with both smeared (SS) and point (SP) sinks. The compound propagators are produced with a single insertion of $J_3^{(u,d)}$ with couplings $\lambda_{u,d} = \{0, \pm 0.05, \pm 0.1, \pm 0.2\}$. These propagators are used to produce correlation functions,

$$C_{\lambda_{u};\lambda_{d}}^{(h)}(t) = \sum_{\mathbf{x}} \langle 0|\chi_{h}(\mathbf{x},t)\chi_{h}^{\dagger}(0)|0\rangle_{\lambda_{u};\lambda_{d}}, \qquad (7)$$

for all the allowed spin states of the one- and two-nucleon systems, $h \in \{p, np({}^{3}S_{1}), nn, np({}^{1}S_{0}), pp\}$. Results for all source locations on each configuration are averaged before subsequent analysis.

The calculations use a lattice axial current with the finite renormalization factor $Z_A = 0.867(43)$ [6]. Because of the isotensor nature of the bilinear insertions, mixing with other operator structures is highly suppressed. As the dinucleon and deuteron states are both compact bound states at this value of the quark masses [20], only exponentially small volume effects are anticipated in the extracted matrix element. This will become a more subtle issue for future calculations with quark masses near the physical values, as discussed in Ref. [7].

The second-order axial responses of the dinucleon system are the primary focus of the current work. For an up-quark axial current, the relevant background-field correlators have the form

$$C_{\lambda_{u};\lambda_{d}=0}^{(h)}(t) = \sum_{\mathbf{x}} \langle 0|\chi_{h}(\mathbf{x},t)\chi_{h}^{\dagger}(0)|0\rangle + \lambda_{u}\sum_{\mathbf{x},\mathbf{y}}\sum_{t_{1}=0}^{t} \langle 0|\chi_{h}(\mathbf{x},t)J_{3}^{(u)}(\mathbf{y},t_{1})\chi_{h}^{\dagger}(0)|0\rangle + \frac{\lambda_{u}^{2}}{2}\sum_{\mathbf{x},\mathbf{y},\mathbf{z}}\sum_{t_{1,2}=0}^{t} \langle 0|\chi_{h}(\mathbf{x},t)J_{3}^{(u)}(\mathbf{y},t_{1})J_{3}^{(u)}(\mathbf{z},t_{2})\chi_{h}^{\dagger}(0)|0\rangle + \mathcal{O}(\lambda_{u}^{3}), \qquad (8)$$

from which the second-order term in the field strength λ_u can be extracted from determinations at multiple values of λ_u . Combining these correlators, and those for the downquark axial current, as specified in Eq. (3), leads to the isotensor matrix element

$$C(t) = 2C_{\lambda_{u};0}^{(np({}^{1}S_{0}))}(t)|_{\lambda_{u}^{2}} - C_{\lambda_{u};0}^{(nn)}(t)|_{\lambda_{u}^{2}} - C_{0;\lambda_{d}}^{(nn)}(t)|_{\lambda_{d}^{2}}, \quad (9)$$

where each term on the right-hand side is the component of the correlation functions that is second order in λ_u or λ_d , as denoted. Using Eq. (3) and isospin symmetry, Eq. (9) can be written as

Up to discretization effects, insertion of appropriate complete sets of states allows this expression to be written as

$$C(t) = \frac{2}{a^2} \sum_{\mathbf{n},\mathbf{m},\mathbf{l}'} Z_{\mathbf{n}} Z_{\mathbf{m}}^{\dagger} e^{-E_{\mathbf{n}}t} \frac{\langle \mathbf{n} | \hat{J}_3^+ | \mathbf{l}' \rangle \langle \mathbf{l}' | \hat{J}_3^+ | \mathbf{m} \rangle}{E_{\mathbf{l}'} - E_{\mathbf{m}}} \times \left(\frac{e^{-(E_{\mathbf{l}'} - E_{\mathbf{n}})t} - 1}{E_{\mathbf{l}'} - E_{\mathbf{n}}} + \frac{e^{(E_{\mathbf{n}} - E_{\mathbf{m}})t} - 1}{E_{\mathbf{n}} - E_{\mathbf{m}}} \right),$$
(11)

where $|\mathbf{n}\rangle$, $|\mathbf{m}\rangle$, and $|\mathbf{l}'\rangle$ are zero-momentum energy eigenstates with the quantum numbers of the pp, nn, and deuteron systems, respectively. Here, $Z_{\mathbf{n}} = \sqrt{V}\langle 0|\chi_{pp}|\mathbf{n}\rangle$ and $Z_{\mathbf{m}} = \sqrt{V}\langle 0|\chi_{nn}|\mathbf{m}\rangle$ are overlap factors, and $E_{\mathbf{l}'} = E_{nn} + \delta_{\mathbf{l}'}$ and $E_{\mathbf{n}} = E_{nn} + \delta_{\mathbf{n}}$ are the energies of the \mathbf{l}' th and \mathbf{n} th excited states in the ${}^{3}S_{1}$ and ${}^{1}S_{0}$ channels, respectively.

Forming a ratio of Eq. (11) to the zero-field two-point function,

$$\mathcal{R}(t) = \frac{C(t)}{2C_{0:0}^{(nn)}(t)},\tag{12}$$

it is straightforward, utilizing the isospin symmetry of the calculation, to show that [7]

$$a^{2}\hat{\mathcal{R}}(t) = a^{2}\mathcal{R}(t) - \frac{|\langle pp|\tilde{J}_{3}^{+}|d\rangle|^{2}}{\Delta} \left(\frac{e^{\Delta t} - 1}{\Delta} - t\right)$$
$$= t\sum_{\mathfrak{l}'\neq d} \frac{\langle pp|\tilde{J}_{3}^{+}|\mathfrak{l}'\rangle\langle\mathfrak{l}'|\tilde{J}_{3}^{+}|nn\rangle}{E_{\mathfrak{l}'} - E_{nn}}$$
$$+ c + de^{\Delta t} + \mathcal{O}(e^{-\hat{\delta}t}), \tag{13}$$

where *c* and *d* involve complicated combinations of ground- and excited-state transition amplitudes, and $\hat{\delta}$ is the minimum energy gap to the first excited state in either channel, and, for these calculations, $\hat{\delta} \gg \Delta$. Importantly, the coefficient of the linear term determines the axial polarizability and can be extracted from

$$\mathcal{R}^{(\mathrm{lin})}(t) = \frac{(e^{a\Delta} + 1)\hat{\mathcal{R}}(t+a) - \hat{\mathcal{R}}(t+2a) - e^{a\Delta}\hat{\mathcal{R}}(t)}{e^{a\Delta} - 1} \quad (14)$$

at late times. Finally, this result can be combined with the deuteron-pole contribution to give a quantity that asymptotes to the bare Gamow-Teller matrix element at late times:

$$\mathcal{R}^{(\text{full})}(t) = \mathcal{R}^{(\text{lin})}(t) - \frac{|\langle pp | \tilde{J}_3^+ | d \rangle|^2}{a\Delta} \stackrel{t \to \infty}{\to} \frac{M_{\text{GT}}^{2\nu}}{6aZ_A^2}.$$
 (15)

The four ratios used to determine $M_{GT}^{2\nu}$ are shown in Fig. 1 for both SS and SP source-sink combinations. Fits are performed to the statistically more precise SP correlators, and the values of the total matrix element and the





FIG. 1. Ratios from Eqs. (12)–(15) used in the analysis. In each panel, the orange diamonds (blue circles) correspond to the SS (SP) data. The green bands show fits to the SP data in the lower two panels. The SS data are slightly offset in the horizontal direction for clarity. The difference between the SS and SP ratios in the upper two panels is due to contamination that is removed in constructing the subsequent quantities in the lower panels.

short-distance contribution, normalized by the naive deuteron-pole matrix element g_A^2/Δ , are given by

$$\frac{\Delta}{g_A^2} \sum_{\mathfrak{l}' \neq d} \frac{\langle pp | \tilde{J}_3^+ | \mathfrak{l}' \rangle \langle \mathfrak{l}' | \tilde{J}_3^+ | nn \rangle}{E_{\mathfrak{l}'} - E_{nn}} = -0.04(2)(1), \quad (16)$$

$$6\frac{\Delta}{g_A^2}M_{\rm GT}^{2\nu} = -1.04(4)(4). \tag{17}$$

In these expressions, the first uncertainties arise from statistical sampling and from systematic effects from fitting choices and deviations from Wigner symmetry [7]. The second uncertainties encompass differences between analysis methods. The leading discretization effects, which are potentially large on the numerically smaller polarizability term, are removed by normalizing to the square of the proton axial charge computed using the same lattice axial current on the same ensemble.

Discussion.—The computed value of $M_{GT}^{2\nu}$ above can be used to determine the unknown EFT(π) low-energy constant $\mathbb{H}_{2,S}$. Taking the values of g_A and the two-body single-current matrix element from Ref. [6], and using the calculated binding energies and effective ranges of the two-nucleon systems [20,28], the result is $\mathbb{H}_{2,S} = 4.7(1.3)(1.8)$ fm. The dominant contribution to $M_{GT}^{2\nu}$ comes from the deuteron pole with coupling g_A^2 . This is modified by two-body effects in the axial current $(\tilde{l}_{1,A})$; this contribution shifts the leading-order result from g_A^2/Δ to $|\langle pp|\tilde{J}_3^+|d\rangle|^2/\Delta$, which is approximately a 5% shift. Interestingly, the calculated result suggests that the additional axial polarizability contribution is of similar size. The existence of this short-distance contribution precludes accurately predicting $\beta\beta$ -decay matrix elements in a nuclear many-body calculation by simply rescaling (quenching) g_A .

The present results are obtained at an unphysical quark mass without the inclusion of electromagnetism and isospin breaking effects, and at a single lattice spacing and volume. All these caveats may be important, particularly given that the short-distance two-nucleon effects are only few-percent contributions to the matrix elements. Such effects require further investigation but are not expected to qualitatively alter the conclusions of this work. Despite these qualifications, the potential for a relatively large contribution from the isotensor axial polarizability is important, as terms of this form are not included in current phenomenological analyses of $\beta\beta$ decay. This observation, supported and motivated by the numerical calculations, is the central point of this Letter. In order to accurately predict $2\nu\beta\beta$ decay rates with fully quantified uncertainties, the isotensor axial polarizabilities of nuclei must be determined. Future lattice QCD calculations in few-nucleon systems and light nuclei, matched to nuclear many-body methods such as $EFT(\pi)$, offer the possibility of determining these contributions which are difficult to access experimentally. However, in order to undertake such calculations at the physical quark masses, a number of difficulties related to the bilocal nature of such weak processes must be overcome [7]. Additional complications arise for $0\nu\beta\beta$ where these polarizability contributions are also likely to be important in the case of light Majorana neutrinos. Furthermore, the short-distance strong-interaction contributions encapsulated in the isotensor axial polarizability provide an inherent background to extracting contributions from short-distance leptonnumber violating operators that are possible beyond the light Majorana scenario [29–33].

This research was supported in part by the National Science Foundation under Grant No. NSF PHY11-25915, and Z. D., W. D., M. J. S., P. E. S., B. C. T., and M. L. W.

acknowledge the Kavli Institute for Theoretical Physics for hospitality during completion of this work. Calculations were performed using computational resources provided by NERSC (supported by U.S. Department of Energy Grant No. DE-AC02-05CH11231), and by the USQCD Collaboration. This research used resources of the Oak Ridge Leadership Computing Facility at the Oak Ridge National Laboratory, which is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC05-00OR22725. The PRACE Research Infrastructure resources at the Très Grand Centre de Calcul and Barcelona Supercomputing Center were also used. Parts of the calculations used the CHROMA software suite [34] and the QUDA library [35,36]. E. C. was supported in part by the USQCD SciDAC project, the U.S. Department of Energy through Grant No. DE-SC00-10337, and by U.S. Department of Energy Grant No. DE-FG02-00ER41132. Z. D., W. D., and P. E. S. were partially supported by U.S. Department of Energy Early Career Research Grant No. DE-SC0010495 and Grant No. DE-SC0011090. The work of W. D. is supported in part by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, within the framework of the TMD Topical Collaboration. K.O. was partially supported by the U.S. Department of Energy through Grant No. DE-FG02-04ER41302 and through Contract No. DE-AC05-06OR23177 under which JSA operates the Thomas Jefferson National Accelerator Facility. M. J. S. was supported by DOE Grant No. DE-FG02-00ER41132, and in part by the USQCD SciDAC project, the U.S. Department of Energy through Grant No. DE-SC00-10337. B. C. T. was supported in part by a joint City College of New York-RIKEN/Brookhaven Research Center fellowship, and by the U.S. National Science Foundation, under Grant No. PHY15-15738. M.L.W. was supported in part by DOE Grant No. DE-FG02-00ER41132. F. W. was partially supported through the USQCD Scientific Discovery through Advanced Computing (SciDAC) project funded by U.S. Department of Energy, Office of Science, Offices of Advanced Scientific Computing Research, Nuclear Physics and High Energy Physics and by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under Contract No. DE-AC05-06OR23177.

- [1] R. N. Mohapatra et al., Rep. Prog. Phys. 70, 1757 (2007).
- [2] F. T. Avignone, III, S. R. Elliott, and J. Engel, Rev. Mod. Phys. 80, 481 (2008).
- [3] S. M. Bilenky and C. Giunti, Mod. Phys. Lett. A 27, 1230015 (2012).
- [4] S. Dell'Oro, S. Marcocci, M. Viel, and F. Vissani, Adv. High Energy Phys. (2016) 2162659.
- [5] J. Engel and J. Menendez, Rep. Prog. Phys. 80, 046301 (2017).

- [6] M. J. Savage, P. E. Shanahan, B. C. Tiburzi, M. L. Wagman, F. Winter, S. R. Beane, E. Chang, Z. Davoudi, W. Detmold, and K. Orginos, preceding Letter, Phys. Rev. Lett. 119, 062002 (2017).
- [7] B. C. Tiburzi, M. L. Wagman, F. Winter, E. Chang, Z. Davoudi, W. Detmold, K. Orginos, M. J. Savage, and P. E. Shanahan, arXiv:1702.02929.
- [8] J. Kotila and F. Iachello, Phys. Rev. C 85, 034316 (2012).
- [9] S. Stoica and M. Mirea, Phys. Rev. C 88, 037303 (2013).
- [10] D. B. Kaplan, Nucl. Phys. B494, 471 (1997).
- [11] D. B. Kaplan, M. J. Savage, and M. B. Wise, Phys. Lett. B 424, 390 (1998).
- [12] D. B. Kaplan, M. J. Savage, and M. B. Wise, Nucl. Phys. B534, 329 (1998).
- [13] U. van Kolck, Nucl. Phys. A645, 273 (1999).
- [14] J.-W. Chen, G. Rupak, and M. J. Savage, Nucl. Phys. A653, 386 (1999).
- [15] S. R. Beane and M. J. Savage, Nucl. Phys. A694, 511 (2001).
- [16] M. Butler and J.-W. Chen, Nucl. Phys. A675, 575 (2000).
- [17] M. Butler, J.-W. Chen, and X. Kong, Phys. Rev. C 63, 035501 (2001).
- [18] X. Kong and F. Ravndal, Phys. Rev. C 64, 044002 (2001).
- [19] M. Butler and J.-W. Chen, Phys. Lett. B 520, 87 (2001).
- [20] S. R. Beane, E. Chang, S. D. Cohen, W. Detmold, H. W. Lin, T. C. Luu, K. Orginos, A. Parreño, M. J. Savage, and A. Walker-Loud (NPLQCD Collaboration), Phys. Rev. D 87, 034506 (2013).
- [21] N. Barnea, L. Contessi, D. Gazit, F. Pederiva, and U. van Kolck, Phys. Rev. Lett. 114, 052501 (2015).
- [22] N. H. Christ, T. Izubuchi, C. T. Sachrajda, A. Soni, and J. Yu (RBC and UKQCD Collaborations), Phys. Rev. D 88, 014508 (2013).

- [23] N. H. Christ, X. Feng, G. Martinelli, and C. T. Sachrajda, Phys. Rev. D 91, 114510 (2015).
- [24] N. H. Christ, X. Feng, A. Portelli, and C. T. Sachrajda (RBC and UKQCD Collaborations), Phys. Rev. D 93, 114517 (2016).
- [25] N. H. Christ, X. Feng, A. Jüttner, A. Lawson, A. Portelli, and C. T. Sachrajda, Phys. Rev. D 94, 114516 (2016).
- [26] M. Lüscher and P. Weisz, Commun. Math. Phys. 97, 59 (1985).
- [27] B. Sheikholeslami and R. Wohlert, Nucl. Phys. B259, 572 (1985).
- [28] S. R. Beane *et al.* (NPLQCD Collaboration), Phys. Rev. C 88, 024003 (2013).
- [29] M. J. Savage, Phys. Rev. C 59, 2293 (1999).
- [30] G. Prezeau, M. Ramsey-Musolf, and P. Vogel, Phys. Rev. D 68, 034016 (2003).
- [31] M. L. Graesser, arXiv:1606.04549.
- [32] A. Nicholson, E. Berkowitz, C. C. Chang, M. A. Clark, B. Joó, T. Kurth, E. Rinaldi, B. Tiburzi, P. Vranas, and A. Walker-Loud, *Proc. Sci.*, LATTICE2016 (2016) 017.
- [33] V. Cirigliano, W. Dekens, M. Graesser, and E. Mereghetti, Phys. Lett. B **769**, 460 (2017).
- [34] R. G. Edwards and B. Joó (SciDAC Collaboration, LHPC Collaboration, UKQCD Collaboration), Nucl. Phys. B, Proc. Suppl. 140, 832 (2005).
- [35] M. A. Clark, R. Babich, K. Barros, R. C. Brower, and C. Rebbi, Comput. Phys. Commun. 181, 1517 (2010).
- [36] R. Babich, M. A. Clark, B. Joó, G. Shi, R. C. Brower, and S. Gottlieb, in *Proceedings of the SC'11 International Conference for High Performance Computing, Networking, Storage and Analysis, Seattle, 2011* (Association for Computing Machinery, New York, 2011).