

Determination of the Density and Temperature Dependence of the Shear Viscosity of a Unitary Fermi Gas Based on Hydrodynamic Flow

Marcus Bluhm,^{1,2} Jiaxun Hou,¹ and Thomas Schäfer¹

¹*Department of Physics, North Carolina State University, Raleigh, North Carolina 27695, USA*

²*Institute of Theoretical Physics, University of Wrocław, 50204 Wrocław, Poland*

(Received 16 April 2017; revised manuscript received 19 June 2017; published 10 August 2017)

We determine the shear viscosity of the ultracold Fermi gas at unitarity in the normal phase using hydrodynamic expansion data. The analysis is based on a generalized fluid dynamic framework which ensures a smooth transition between the fluid dynamic core of the cloud and the ballistic corona. We use expansion data taken by Joseph, Elliott, and Thomas [Shear Viscosity of a Universal Fermi Gas Near the Superfluid Phase Transition, *Phys. Rev. Lett.* **115**, 020401 (2015).] and measurements of the equation of state by Ku *et al.* [Revealing the superfluid lambda transition in the universal thermodynamics of a unitary Fermi gas, *Science* **335**, 563 (2012).]. We find that the shear viscosity to particle density ratio just above the critical temperature is $\eta/n|_{T_c} = 0.41 \pm 0.11$. We also obtain evidence that the shear viscosity to entropy density ratio has a minimum slightly above T_c with $\eta/s|_{\min} = 0.50 \pm 0.10$.

DOI: 10.1103/PhysRevLett.119.065302

Introduction.—The dilute Fermi gas at unitarity is a very attractive physical system for studying the transport properties of strongly correlated quantum fluids [1–3]. From a theoretical point of view, the unitary Fermi gas is a parameter-free, scale invariant, and intrinsically quantum-mechanical many-body system. A lot of interest has centered on the question of how close the viscosity to entropy density ratio of this system comes to the proposed string theory bound $\eta/s = \hbar/(4\pi k_B)$ [4]. Experimentally, the unitary Fermi gas can be realized in dilute atomic gases using Feshbach resonances [5,6]. The experimental control provided by Feshbach resonances implies that we can study the transition from the strongly correlated unitary Fermi gas to weakly coupled Bose and Fermi gases.

In this Letter we focus on the problem of extracting the shear viscosity of the unitary Fermi gas from experiments with trapped ultracold gases [7–15]. Our main interest is in the low-temperature regime, where the density dependence of the shear viscosity is relevant, and the minimum of η/s is likely to be achieved. There are two main types of experiments that are relevant to this problem. The first class involves measuring the damping rate of collective excitations, and the second focuses on the expansion of the cloud after removing the trapping potential. From a theoretical perspective the damping experiments would appear to be more attractive, because even a very small viscosity leads to a clear signature in the exponential decay of the collective mode. In practice, however, the expansion experiments take place in a cleaner environment and have achieved greater accuracy. In an expansion experiment what is observed is the time evolution of the aspect ratio of the cloud. Hydrodynamic pressure gradients accelerate the cloud along the short direction, so that the aspect ratio increases as a function of time. Viscosity counteracts the pressure

gradients, and slows the growth of the aspect ratio. These flow experiments are very similar to elliptic flow experiments in relativistic heavy ion physics [16–18].

The main difficulty in analyzing these experiments is that the viscosity $\eta(n, T)$ is a local quantity that varies with the density n and temperature T of the cloud, while the observed aspect ratio is a global property of the trapped gas. This means that the dependence of the data on initial cloud energy, particle number, and expansion time has to be unfolded to determine $\eta(n, T)$. An even more significant problem is that the viscosity is a parameter that appears in the fluid dynamic description of the cloud. However, fluid dynamics breaks down in the dilute, dissipative corona of the gas.

We have recently made significant progress in dealing with the physics of the dilute corona. We have introduced a new method, anisotropic fluid dynamics [19–21], that takes into account the effects of nonhydrodynamic modes. These modes quickly relax in the dense part of the cloud so that Navier-Stokes fluid dynamics is recovered. In the dilute corona nonhydrodynamic modes ensure a smooth transition to a free-streaming, ballistic expansion. We have checked numerically that anisotropic fluid dynamics reproduces the Navier-Stokes equation in the dense limit [19] as well as numerical solutions of the Boltzmann equation in the dilute regime [22,23]. We have also shown that the anisotropic fluid dynamics, combined with the kinetic theory prediction for the shear viscosity $\eta = 15/(32\sqrt{\pi})(mT)^{3/2}$ [24], reproduces the high-temperature expansion data obtained in [12]. Note that here and in the remainder of the Letter we set \hbar and k_B equal to unity.

In this Letter we extend our studies to lower temperature. For this purpose we fit the expansion data to a systematic expansion of the viscosity in powers of the density.

We show that the data clearly demand that the shear viscosity has nontrivial density dependence. We also show that the density dependence in the normal phase is quite smooth, and that the existing data place strong constraints on η/n near T_c . This study requires several refinements of our previous work. We extend the fluid dynamic analysis to three-dimensional systems with no axial symmetry. We include an accurate parametrization of the measured equation of state, and a more general functional form of the shear viscosity.

Anisotropic fluid dynamics.—In this section we briefly summarize the anisotropic fluid dynamics method [19]. The fluid dynamical variables that characterize a non-relativistic fluid in the normal phase are the mass density ρ , the momentum density $\vec{\pi} = \rho\vec{u}$, and the energy density \mathcal{E} . The equations of motion follow from the conservation laws

$$D_0\rho = -\rho\vec{\nabla}\cdot\vec{u}, \quad (1)$$

$$D_0u_i = -\frac{1}{\rho}\nabla_j(\delta_{ij}P + \delta\Pi_{ij}), \quad (2)$$

$$D_0\epsilon = -\frac{1}{\rho}\nabla_i(u_iP + \delta J_i^\mathcal{E}). \quad (3)$$

Here, we defined the comoving time derivative $D_0 = \partial_0 + \vec{u}\cdot\vec{\nabla}$, the energy per mass $\epsilon = \mathcal{E}/\rho$, and the pressure P . We also introduce the energy density in the rest frame of the fluid, $\mathcal{E}^0 = \mathcal{E} - \frac{1}{2}\rho\vec{u}^2$. In order for the equations to close we have to provide an equation of state $P = P(\mathcal{E}^0, \rho)$, and constitutive equations for the dissipative stresses $\delta\Pi_{ij}$ and the dissipative energy current $\delta J_i^\mathcal{E}$. For the unitary Fermi gas scale invariance implies that $P = \frac{2}{3}\mathcal{E}^0$.

In the Navier-Stokes approximation the dissipative stresses are expanded to first order in gradients of the thermodynamic variables. We get $\delta\Pi_{ij} = -\eta\sigma_{ij}$ with

$$\sigma_{ij} = \nabla_i u_j + \nabla_j u_i - \frac{2}{3}\delta_{ij}\vec{\nabla}\cdot\vec{u} \quad (4)$$

and $\delta J_i^\mathcal{E} = u_j\delta\Pi_{ij}$. Scale invariance implies that the bulk viscosity vanishes. We have also used the fact that in expansion experiments the effects of heat conduction are of higher order in the gradient expansion. This is related to the fact that the initial temperature is constant, and that the expansion of an ideal gas preserves the isothermal nature of the temperature profile [25].

In anisotropic fluid dynamics we treat the components of the dissipative stress tensor as independent fluid dynamical variables. The symmetries of the trap imply that the stresses are diagonal. We define anisotropic components of the pressure, P_a for $a = 1, 2, 3$, and define

$$\delta\Pi_{ij} = \text{diag}(\Delta P_1, \Delta P_2, \Delta P_3), \quad (5)$$

where $\Delta P_a = P_a - P$. We also define anisotropic components of the energy density \mathcal{E}_a such that $\mathcal{E} = \sum_a \mathcal{E}_a$. The

anisotropic components of the energy per mass satisfy the equation of motion [19]

$$D_0\epsilon_a = -\frac{1}{\rho}\nabla_i[\delta_{ia}u_iP + (\delta J_a^\mathcal{E})_i] - \frac{P}{2\eta\rho}\Delta P_a, \quad (6)$$

where $\epsilon_a = \mathcal{E}_a/\rho$ and $(\delta J_a^\mathcal{E})_i = \delta_{ia}u_j\delta\Pi_{ij}$. The anisotropic pressure is related to the anisotropic energy density by an equation of state. In the case of a scale-invariant fluid we have $P_a(\mathcal{E}_a^0) = 2\mathcal{E}_a^0$ with $\mathcal{E}_a^0 = \mathcal{E}_a - \frac{1}{2}\rho u_a^2$. Then $P = \frac{1}{3}\sum_a P_a$ satisfies the isotropic equation of state, and Eq. (6) gives the isotropic equation of energy conservation Eq. (3) when summed over a . In our previous work we have described a three-dimensional fluid dynamics code that solves Eqs. (1)–(3) and Eq. (6) [19,25]. This code is based on the piecewise parabolic method of Colella and Woodward [26,27].

We have shown that in the limit of small viscosity, $\eta(\vec{\nabla}\cdot\vec{u}) \ll P$, the anisotropic pressure terms relax to the viscous stress tensor in Navier-Stokes theory, $\Delta P_a = -\eta\sigma_{aa}$. We observe that in the opposite limit, that of very large viscosity, Eq. (6) becomes a conservation law. This conservation law ensures that anisotropic fluid dynamics reproduces the free-streaming limit. Finally, we have checked that anisotropic fluid dynamics provides a very accurate representation of numerical solutions of the Boltzmann equation in the limit that two-body scattering dominates [23].

In general the viscosity is a function of density and temperature. In the unitary limit scale invariance implies that $\eta(n, T) = (mT)^{3/2}f(n\lambda^3)$, where $\lambda = [(2\pi)/(mT)]^{1/2}$ is the de Broglie wavelength. In this Letter we will expand the function $f(x)$ in powers of the diluteness of the gas

$$\eta(n, T) = \eta_0(mT)^{3/2}\{1 + \eta_2(n\lambda^3) + \eta_3(n\lambda^3)^2 + \dots\}. \quad (7)$$

We note that the leading term is purely a function of temperature, the first correction is solely a function of density, and higher-order terms depend on increasing powers of the density. In general this expansion is not expected to be useful near T_c , but we will show that terms that scale as $(n\lambda^3)^2$ and higher are surprisingly small.

Experimental parameters.—We will analyze the expansion data reported in [7]. This work represents the most complete set of elliptic flow measurements for the unitary Fermi gas over a wide range of temperatures currently available. The gas is released from a harmonic trap $V_{\text{ext}} = \frac{1}{2}m\omega_x^2 x^2$ with trap frequencies $(\omega_x, \omega_y, \omega_z) = (2\pi)(2210, 830, 64.3)$ Hz. After the optical trap is turned off there is a residual magnetic bowl characterized by $\omega_{\text{mag}} = 2\pi \times 21.5$ Hz. The total energy per particle of the gas varies between $E/(NE_F) = (0.56\text{--}1.91)$. Here, N is the number of particles and $E_F \equiv (3N)^{1/3}\bar{\omega}$, where $\bar{\omega}$ is the geometric mean of the trap frequencies. The energy and temperature of the cloud are extracted using absorption

images and an equation of state $\mathcal{E}^0(n, T)$. We describe a parametrization of the equation of state measured by the MIT group [28] in the Supplemental Material [29], see also [25,30,31]. Based on this equation of state we find that the critical energy where superfluidity occurs at the center of the trap is $E/(NE_F) = 0.70$. In the high-temperature limit many relations simplify. For example, the total cloud energy is given by $E = 3NT$. We will characterize the initial temperature using the dimensionless ratio T/T_F , where $T_F = E_F$.

Scaling of the aspect ratio with the initial energy.—Expansion experiments measure the time evolution of the aspect ratio $A_R(t)$ for different initial energies and particle numbers. The experiment of Joseph *et al.* [7] focuses on the ratio σ_x/σ_y , which reaches its asymptotic behavior more quickly than σ_x/σ_z or σ_y/σ_z . The radii σ_i are determined from a Gaussian fit to two-dimensional absorption images. As noted in [22] it is important to follow this definition when analyzing the data using transport theory. In particular, there is a significant difference between the ratio of rms radii, $\sqrt{\langle x^2 \rangle}/\sqrt{\langle y^2 \rangle}$, and the ratio of Gaussian fit radii, σ_x/σ_y . This is the case even if the initial density distribution is a Gaussian.

Joseph *et al.* [7] observed that the main information about the density and temperature dependence of $\eta(n, T)$ is not carried by the time dependence of $A_R(t)$ for fixed initial energy, but by the dependence of $A_R(t^*)$ at a fixed time t^* on the initial energy. In Fig. 1 we show $A_R(t^*) = \sigma_x/\sigma_y$ as a function of $E/(NE_F)$ at $t^* = 1.2$ msec. Note that the plot covers a fairly narrow range in A_R . Individual data points are more accurate than previously published data, which spanned a much larger range in aspect ratio.

A difficulty in interpreting the results is that the data points correspond to a range of particle numbers. The data




FIG. 1. Aspect ratio $A_R = \sigma_x/\sigma_y$ at $t^* = 1.2$ msec as a function of the energy $E/(NE_F)$ of the cloud. Data (gray points) compared to hydrodynamic fits based on the equation of state of a free gas. The solid red line corresponds to the shear viscosity $\eta = \eta_0(mT)^{3/2}$ predicted by kinetic theory, and the dashed and dotted line show the $\pm 25\%$ and $\pm 50\%$ range in η_0 . The thick green line is the best fit to the high-energy data, corresponding to $\eta_0 = 0.301$.

are clustered around a mean $\bar{N} = 1.94 \times 10^5$, and the variance in $N^{1/3}$, which is relevant to the effective viscosity, is about 7%. We show all the data points on the same plot, but when performing hydrodynamic fits we use the correct number of particles for each individual data point.

Figure 1 shows a fit to the data based on the high-temperature theory only. This means that we use the free gas equation of state, and only the first coefficient, η_0 , in the virial expansion of the shear viscosity. The best fit to the high-temperature data gives $\eta_0 = 0.301$ which is somewhat higher than the value $\eta_0 = 0.264$ predicted by kinetic theory. The best-fit value shifts slightly if the full equation of state is used, but the shape of $A_R(t^*)$ as a function of $E/(NE_F)$ does not change. We observe that the data at lower energy clearly demand a more complicated functional form of the shear viscosity.

Figure 2 shows a fit to the data above the superfluid transition based on the full equation of state and an expansion of the shear viscosity up to second order in density. The best fit is

$$\eta_0 = 0.265 \pm 0.02, \quad \eta_2 = 0.060 \pm 0.02, \quad (8)$$

and $\eta_3 = -(2 \pm 8) \times 10^{-4}$. We observe that the n^2 coefficient is consistent with zero within error bars. We also find that the fit is stable with respect to including higher-order terms in n . The χ^2/N_{dof} of the fit is of order unity, indicating that this simple model provides a very good representation of all the data in the entire regime above the superfluid phase transition. We note that η_0 agrees to better than 1% with the kinetic theory prediction $\eta_0 = 0.264$.

Conclusions.—Our determination of η/n for the homogeneous Fermi gas is shown in Figs. 3 and 4. The result is shown as a function of T/T_F^{loc} , where $T_F^{\text{loc}} = k_F^2/(2m)$ is the local Fermi temperature of the gas. The best fit to the data, shown as the thick red line in Fig. 3, is




FIG. 2. Aspect ratio $A_R = \sigma_x/\sigma_y$ at $t^* = 1.2$ msec as a function of the energy $E/(NE_F)$ of the cloud. Data (gray points) compared to hydrodynamic fits based on the measured equation of state. The red short-dashed line shows the high temperature fit, the blue dashed line includes density corrections, and the green solid line contains a density squared term.




FIG. 3. The reconstructed ratio η/n as a function of T/T_F^{loc} for a homogeneous gas. The local Fermi temperature is defined as $T_F^{\text{loc}} = k_F^2/(2m)$ where k_F is defined via the density of the gas, $n = k_F^3/(3\pi^2)$. The red line shows the density expansion together with the error band described in the text. The curves terminate at T_c . The gray dots show the reconstruction obtained in [7], and the dashed line shows the T -matrix calculation of Enss *et al.* [32].

$$\eta/n = 2.773x^{3/2} + 0.251 - 0.0013x^{-3/2}, \quad (9)$$

where $x = T/T_F^{\text{loc}}$. The coefficients in Eq. (9) are given by the central values of η_0, η_2, η_3 normalized by the density n . The local Fermi momentum k_F is defined in terms of the density of the gas, $n = k_F^3/(3\pi^2)$. We show the reconstruction for temperatures above the critical temperature $T_c = 0.167(13)T_F^{\text{loc}}$ [28]. We find that the value of the viscosity at T_c is $\eta/n|_{T_c} = 0.41 \pm 0.11$. We have not attempted to reconstruct the shear viscosity below T_c , since a proper treatment of this regime requires superfluid hydrodynamics.

For comparison the gray data points show the reconstructed values of η/n obtained in the experimental work of Joseph *et al.* [7]. These results are based on the same expansion data, but involve a number of assumptions [34]. The main assumption is that there is a critical radius R_i^{crit} so




FIG. 4. Same as Fig. 3, zooming in on the low-temperature regime. Our analysis (red band) is compared to the results (gray points) obtained in [7], the T -matrix calculation (dashed line) of Enss *et al.* [32], and the lattice calculation (green band) of Wlazlowski *et al.* [33].

that the atomic cloud inside this radius can be described as a viscous fluid, and the particles outside the radius are a noninteracting gas. The critical radius is assumed to be a constant fraction of the cloud size. The overall constant is adjusted to reproduce the expected behavior of the high-temperature viscosity, $\eta \sim \eta_0(mT)^{3/2}$. This implies that the agreement of the data points with kinetic theory for large T/T_F^{loc} is not a result, but an input. In contrast, the agreement of our reconstruction with kinetic theory is a nontrivial result. There is some discrepancy between the two reconstructions in the regime $T = (0.2-1.0)T_F^{\text{loc}}$. In this regime our result for η/n is systematically lower. This makes sense if one assumes that as the temperature is lowered and the viscosity drops the effective fluid radius increases. This implies that assuming a constant radius of the fluid core leads to an overestimate of the viscosity. It is interesting that directly at T_c the two reconstructions agree.

We also show the T -matrix calculation of Enss *et al.* [32], which agrees quite well with our reconstructed viscosity near T_c . It will be interesting to study the physical consequences of this result, for example, possible implications for quasiparticle models. We also show the lattice calculation of Wlazlowski *et al.* [33]. The calculation does not match the shape of our reconstruction, and has a substantially smaller $\eta/n|_{T_c}$.

Finally, Fig. 5 shows the ratio of shear viscosity to entropy density, based on our reconstruction of η/n and the measurement of s/n by the MIT group [28]. The result is compared to high and low temperature predictions for η/s in kinetic theory [24,35]. We find a shallow minimum of $\eta/s|_{\text{min}} = 0.50 \pm 0.10$ slightly above T_c . The minimum is related to the fact that the entropy per particle drops significantly as T_c is approached from above, whereas no structure is seen in η/n . We note that at present we can only weakly exclude (at about 1σ) a minimum in η/s at or below T_c . A minimum in η/s above T_c was predicted in [36], but is in tension with the Monte Carlo data in [33].




FIG. 5. Low-temperature behavior of the shear viscosity to entropy density ratio η/s as a function of T/T_F^{loc} . Our analysis (red band) is compared to the high and low temperature predictions from kinetic theory, see [24,35].

This work was supported in part by the U.S. Department of Energy Grant No. DE-FG02-03ER41260. The work of M. Bluhm is funded by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska Curie Grant No. 665778 via the National Science Center, Poland, under Grant No. Polonez UMO-2016/21/P/ST2/04035. We would like to thank James Joseph and John Thomas for many useful discussions, and for providing us with the data in [7]. We thank Martin Zwierlein for providing us with the data in [28].

-
- [1] T. Schäfer and D. Teaney, Nearly perfect fluidity: From cold atomic gases to hot quark gluon plasmas, *Rep. Prog. Phys.* **72**, 126001 (2009).
- [2] A. Adams, L. D. Carr, T. Schäfer, P. Steinberg, and J. E. Thomas, Strongly correlated quantum fluids: Ultracold quantum gases, quantum chromodynamic plasmas, and holographic duality, *New J. Phys.* **14**, 115009 (2012).
- [3] T. Schäfer, Fluid dynamics and viscosity in strongly correlated fluids, *Annu. Rev. Nucl. Part. Sci.* **64**, 125 (2014).
- [4] P. K. Kovtun, D. T. Son, and A. O. Starinets, Viscosity in Strongly Interacting Quantum Field Theories from Black Hole Physics, *Phys. Rev. Lett.* **94**, 111601 (2005).
- [5] I. Bloch, J. Dalibard, and W. Zwerger, Many-body physics with ultracold gases, *Rev. Mod. Phys.* **80**, 885 (2008).
- [6] S. Giorgini, L. P. Pitaevskii, and S. Stringari, Theory of ultracold atomic Fermi gases, *Rev. Mod. Phys.* **80**, 1215 (2008).
- [7] J. A. Joseph, E. Elliott, and J. E. Thomas, Shear Viscosity of a Universal Fermi Gas Near the Superfluid Phase Transition, *Phys. Rev. Lett.* **115**, 020401 (2015).
- [8] J. Kinast, A. Turlapov, and J. E. Thomas, Breakdown of hydrodynamics in the radial breathing mode of a strongly interacting Fermi gas, *Phys. Rev. A* **70**, 051401(R) (2004).
- [9] T. Schäfer, The shear viscosity to entropy density ratio of trapped fermions in the unitarity limit, *Phys. Rev. A* **76**, 063618 (2007).
- [10] A. Turlapov, J. Kinast, B. Clancy, L. Luo, J. Joseph, and J. E. Thomas, Is a gas of strongly interacting atomic fermions a nearly perfect fluid, *J. Low Temp. Phys.* **150**, 567 (2008).
- [11] G. M. Bruun and H. Smith, Frequency and damping of the scissors mode of a Fermi gas, *Phys. Rev. A* **76**, 045602 (2007).
- [12] C. Cao, E. Elliott, J. Joseph, H. Wu, J. Petricka, T. Schäfer, and J. E. Thomas, Universal quantum viscosity in a unitary Fermi gas, *Science* **331**, 58 (2011).
- [13] E. Elliott, J. A. Joseph, and J. E. Thomas, Anomalous Minimum in the Shear Viscosity of a Fermi Gas, *Phys. Rev. Lett.* **113**, 020406 (2014).
- [14] E. Elliott, J. A. Joseph, and J. E. Thomas, Observation of Conformal Symmetry Breaking and Scale Invariance in Expanding Fermi Gases, *Phys. Rev. Lett.* **112**, 040405 (2014).
- [15] J. Brewer, M. Mendoza, R. E. Young, and P. Romatschke, Lattice Boltzmann simulations of a strongly interacting two-dimensional Fermi gas, *Phys. Rev. A* **93**, 013618 (2016).
- [16] K. H. Ackermann *et al.* (STAR Collaboration), Elliptic Flow in Au + Au Collisions at $(s_{NN})^{1/2} = 130$ GeV, *Phys. Rev. Lett.* **86**, 402 (2001).
- [17] S. S. Adler *et al.* (PHENIX Collaboration), Elliptic Flow of Identified Hadrons in Au + Au Collisions at $(s_{NN})^{1/2} = 200$ GeV, *Phys. Rev. Lett.* **91**, 182301 (2003).
- [18] K. Aamodt *et al.* (ALICE Collaboration), Elliptic Flow of Charged Particles in Pb-Pb Collisions at 2.76 TeV, *Phys. Rev. Lett.* **105**, 252302 (2010).
- [19] M. Bluhm and T. Schäfer, Dissipative fluid dynamics for the dilute Fermi gas at unitarity: Anisotropic fluid dynamics, *Phys. Rev. A* **92**, 043602 (2015).
- [20] W. Florkowski and R. Ryblewski, Highly anisotropic and strongly dissipative hydrodynamics for early stages of relativistic heavy-ion collisions, *Phys. Rev. C* **83**, 034907 (2011).
- [21] M. Martinez and M. Strickland, Dissipative dynamics of highly anisotropic systems, *Nucl. Phys.* **A848**, 183 (2010).
- [22] P. A. Pantel, D. Davesne, and M. Urban, Numerical solution of the Boltzmann equation for trapped Fermi gases with in-medium effects, *Phys. Rev. A* **91**, 013627 (2015).
- [23] M. Bluhm and T. Schäfer, Model-Independent Determination of the Shear Viscosity of a Trapped Unitary Fermi Gas: Application to High-Temperature Data, *Phys. Rev. Lett.* **116**, 115301 (2016).
- [24] G. M. Bruun and H. Smith, Viscosity and thermal relaxation for a resonantly interacting Fermi gas, *Phys. Rev. A* **72**, 043605 (2005).
- [25] T. Schäfer, Dissipative fluid dynamics for the dilute Fermi gas at unitarity: Free expansion and rotation, *Phys. Rev. A* **82**, 063629 (2010).
- [26] P. Colella and P. R. Woodward, The piecewise parabolic method (PPM) for gas-dynamical simulations, *J. Comput. Phys.* **54**, 174 (1984).
- [27] J. M. Blondin and E. A. Lufkin, The piecewise-parabolic method in curvilinear coordinates, *Astrophys. J. Suppl. Ser.* **88**, 589 (1993).
- [28] M. J. H. Ku, A. T. Sommer, L. W. Cheuk, and M. W. Zwierlein, Revealing the superfluid lambda transition in the universal thermodynamics of a unitary Fermi gas, *Science* **335**, 563 (2012).
- [29] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.119.065302> for a description of the equation of state, and for a definition of a set of units that can be used to scale out the dependence on the number of particles.
- [30] S. Nascimbene, N. Navon, K. Jiang, F. Chevy, and C. Salomon, Exploring the thermodynamics of a universal Fermi gas, *Nature (London)* **463**, 1057 (2010).
- [31] P. Romatschke and R. E. Young, Implications of hydrodynamic fluctuations for the minimum shear viscosity of the dilute Fermi gas at unitarity, *Phys. Rev. A* **87**, 053606 (2013).
- [32] T. Enss, R. Haussmann, and W. Zwerger, Viscosity and scale invariance in the unitary Fermi gas, *Ann. Phys. (Amsterdam)* **326**, 770 (2011).
- [33] G. Wlazlowski, P. Magierski, A. Bulgac, and K. J. Roche, The temperature evolution of the shear viscosity in a unitary Fermi gas, *Phys. Rev. A* **88**, 013639 (2013).
- [34] The χ^2/N_{dof} of our fit is close to 1, whereas the reconstruction of [7], used as an input to our hydrodynamic

- model, has $\chi^2/N_{\text{dof}} \approx 4.5$. More importantly, we stress that the result in [7] has an unknown overall normalization, which was fixed using kinetic theory.
- [35] G. Rupak and T. Schäfer, Shear viscosity of a superfluid Fermi gas in the unitarity limit, *Phys. Rev. A* **76**, 053607 (2007).
- [36] W. Zwerger, in *Quantum Matter at Ultralow Temperatures*, Proceedings of the International School of Physics “Enrico Fermi”, Course CXCI, edited by M. Inguscio, W. Ketterle, S. Stringari, and G. Roati (IOS Press, Amsterdam, 2016), pp. 63–142, ISBN .