

Universal Scaling Laws in the Dynamics of a Homogeneous Unitary Bose Gas

Christoph Eigen,¹ Jake A. P. Glidden,¹ Raphael Lopes,¹ Nir Navon,^{1,2} Zoran Hadzibabic,¹ and Robert P. Smith^{1,*}

¹*Cavendish Laboratory, University of Cambridge, J. J. Thomson Avenue, Cambridge CB3 0HE, United Kingdom*

²*Department of Physics, Yale University, New Haven, Connecticut 06511, USA*

(Received 19 September 2017; published 22 December 2017)

We study the dynamics of an initially degenerate homogeneous Bose gas after an interaction quench to the unitary regime at a magnetic Feshbach resonance. As the cloud decays and heats, it exhibits a crossover from degenerate- to thermal-gas behavior, both of which are characterized by universal scaling laws linking the particle-loss rate to the total atom number N . In the degenerate and thermal regimes, the per-particle loss rate is $\propto N^{2/3}$ and $N^{26/9}$, respectively. The crossover occurs at a universal kinetic energy per particle and at a universal time after the quench, in units of energy and time set by the gas density. By slowly sweeping the magnetic field away from the resonance and creating a mixture of atoms and molecules, we also map out the dynamics of correlations in the unitary gas, which display a universal temporal scaling with the gas density, and reach a steady state while the gas is still degenerate.

DOI: 10.1103/PhysRevLett.119.250404

Strong interactions and correlations are at the heart of the most interesting many-body quantum phenomena. The possibility to control the interaction strength via Feshbach resonances [1] makes ultracold atomic gases an excellent setting for studies of strongly correlated behavior. On resonance, the s -wave scattering length a , which characterizes two-body contact interactions, diverges. In this so-called unitary regime, the interactions are as strong as allowed by quantum mechanics, and the physics cannot explicitly depend on a , leading to the possibility of new types of universal behavior [2–6].

Of particular interest are the interaction-dominated degenerate unitary gases. Within the “universality hypothesis,” they have only one relevant length scale—the average interparticle spacing, given by the density n , which also sets the natural energy and time scales [2]:

$$E_n = \frac{\hbar^2}{2m} (6\pi^2 n)^{2/3} \quad \text{and} \quad t_n = \hbar/E_n, \quad (1)$$

where m is the atom mass. These “Fermi energy” and “Fermi time” scales are applicable to both Fermi and Bose gases. In Bose gases, however, the universality can be broken by Efimov physics [7–18]. The Feshbach dimer molecular state, responsible for the resonance, is of size a and becomes unbound as $a \rightarrow \infty$, but the infinite series of Efimov trimer states, each of a size 22.7 times larger than the previous one, can introduce new length scales into the problem.

While unitary Fermi gases have been extensively explored [3–5], the experimental [15,16,19–22] and theoretical [23–36] studies of unitary Bose gases are only recently emerging. An experimental challenge is that they exhibit rapid three-body loss and heating, which also raises fundamental questions about the extent to which they have

well-defined equilibrium properties [21], but the loss dynamics also offer a valuable probe of the unitary behavior [16,19–22]. While coherent three-body correlations [15] and Efimov trimers [16] have been observed, the decay dynamics [16,19–22] have been consistent with universal scalings (see also [37]). All experiments so far were performed with harmonically trapped gases, and their interpretation relies on knowledge of the inhomogeneous density profiles. For a degenerate gas, the density profile is known prior to a quench to unitarity, and hence right after it, but the subsequent evolution is complicated by strong interactions and inhomogeneous losses and heating.

In this Letter, we study the dynamics of a *homogeneous*, initially degenerate Bose gas quenched to unitarity. In our gas, produced in an optical box trap [39], E_n and t_n are global variables simply set by the total atom number N [40]. We can thus quantitatively study the full evolution of a cloud as it decays and heats. In both degenerate and thermal regimes, we observe (different) universal atom-loss scaling laws in agreement with theoretical predictions, and we characterize the universal features of the crossover between the two regimes. By slowly ramping the magnetic field away from the resonance and creating an atom-molecule mixture, we also study the dynamics of correlations in the unitary gas. These correlations show t_{n_0} (where n_0 is the initial density) as the only relevant time scale and indicate that the system reaches a strongly correlated quasiequilibrium state while it is still degenerate.

Our clouds are produced in the lowest hyperfine state of ³⁹K, which has a background scattering length of $-29a_0$ (where a_0 is the Bohr radius) and a 52-G-wide Feshbach resonance centered at $B_0 = 402.70(3)$ G [15,41]. The atoms are confined in a cylindrical box trap of radius $15(1)$ μm and length $50(2)$ μm [39,42], and we vary the initial atom number N_0 in the range $(48\text{--}214) \times 10^3$. This




FIG. 1. Measurement of atom-loss and correlation dynamics. (a) A BEC is quenched to unitarity, and the cloud is held there for a variable time t_{hold} and then ramped away from unitarity at a variable rate $R = -dB/dt$. An infinitely fast ramp-out (solid line) would project the resonantly interacting cloud onto free-atom states, while a finite-rate ramp-out (dashed line) creates a mixture of atoms and molecules. (b) Open symbols: The observed atom number versus $1/R$, for $N_0 = 98 \times 10^3$ and $t_{\text{hold}} = 80 \mu\text{s}$. Solid line: An exponential fit, which gives an exponential constant of $2.2(3) \mu\text{s}/\text{G}$. Solid symbols: N_{obs} if the molecules are dissociated by a second magnetic-field pulse to resonance. (c) Evolution of N_{obs} with t_{hold} for our fastest ramp-out ($0.3 \mu\text{s}/\text{G}$) and a much slower one ($6 \mu\text{s}/\text{G}$). The fast ramp-out data show the on-resonance atom loss, and the difference between the two curves reveals the correlation dynamics in the unitary gas.

corresponds to t_{n_0} values of a few tens of microseconds, and the range of fields near B_0 where $n_0 a^3 > 1$ is $\Delta B \sim 0.1 \text{ G}$ [43].

We start with a weakly interacting quasipure Bose-Einstein condensate (BEC) more than $10\Delta B$ away from resonance, where $n_0 a^3 < 10^{-3}$. As outlined in Fig. 1(a), we then rapidly (within $2 \mu\text{s}$) quench the magnetic field to B_0 [44], wait for a time t_{hold} , and then ramp the field away from unitarity. We finally image the atoms after 8–32 ms of time-of-flight (TOF) expansion. The number of atoms that we observe, N_{obs} , can reduce with t_{hold} for two reasons: (i) due to losses at unitarity and (ii) because the ramp-out from unitarity results in a mixture of atoms and (Feshbach and/or Efimov) molecules [16], and our imaging detects only free atoms. The molecular fraction after the ramp-out depends on both the correlations in the unitary gas and the ramp-out rate [16,45–47], $R = -dB/dt$ [48]. Independently of the many-body state at unitarity, an infinitely rapid ramp-out should essentially project the resonantly interacting cloud onto free-atom states, so that $N_{\text{obs}} = N$.




FIG. 2. Growth of the kinetic energy, for $N_0 = 214 \times 10^3$. (a) Absorption images taken for various t_{hold} and after 12 ms of TOF expansion at weak interactions. (b) Kinetic energy per particle, E , versus t_{hold} . Note that the interaction energy per particle after the ramp-out is $< 20 \text{ nK}$. The solid line shows the $2/13$ power law predicted for a thermal gas at long t_{hold} , where $E \propto T$.

To disentangle the two sources of the reduction of N_{obs} , we vary the ramp-out rate, as illustrated in Fig. 1(b). Here the open symbols show N_{obs} versus $1/R$ for $N_0 = 98 \times 10^3$ and a fixed $t_{\text{hold}} = 80 \mu\text{s}$. The data are fitted well by an exponential, characteristic of a Landau-Zener process [45,47]. Extrapolating to $1/R = 0$, we assess that, with our technically limited fastest ramp-out, $1/R = 0.3 \mu\text{s}/\text{G}$, we get $N_{\text{obs}} = N$ to within $< 10\%$; we verified this for our full range of N_0 and at several t_{hold} . For slower ramp-outs, N_{obs} is reduced by up to 40% (for this N_0 and t_{hold}). To verify that this reduction occurs due to the creation of an atom-molecule mixture, after the first trip to the resonance (with a slow ramp-out) we apply a brief ($8 \mu\text{s}$) second magnetic-field pulse to B_0 to break up the molecules and find that most of the missing atoms reappear—see the solid symbols in Fig. 1(b) [49].

In Fig. 1(c), we show, for $N_0 = 98 \times 10^3$, the evolution of N_{obs} with t_{hold} for both our fastest ramp-out and a much slower one, $1/R = 6 \mu\text{s}/\text{G}$. Note that there is very little difference between the two curves for very short t_{hold} , showing that it takes time for the system to develop the correlations that lead to the creation of an atom-molecule mixture.

From here on, we separately study the atom-loss and correlation dynamics. We first focus on the atom loss and assume that for our largest R to a good approximation $N_{\text{obs}} = N$. From the TOF images (see Fig. 2), we also extract the kinetic energy per particle, E , which monotonically grows with t_{hold} .

The per-particle loss rate is $\dot{N}/N = -L_3 \langle n^2 \rangle$, where L_3 is the three-body loss coefficient. For a homogeneous gas, we simply have $\langle n^2 \rangle = n^2 = N^2/V^2$ (where V is the volume), and one can generally predict a scaling-law behavior:

$$\dot{N}/N = -\text{const} \times N^\gamma. \quad (2)$$

Away from unitarity $L_3 \propto a^4$ [51,52] and in a degenerate unitary gas (assuming universal behavior), a is replaced by




FIG. 3. Atom-loss scaling laws, for $N_0 = 214 \times 10^3$. We observe both the $\gamma = 2/3$ and the $\gamma = 26/9$ law, predicted (respectively) for a degenerate and a thermal unitary Bose gas. The crossover between the two regimes occurs at a well-defined atom number.

$n^{-1/3} \propto N^{-1/3}$, so $\gamma = 2/3$; that is, the instantaneous t_n is the only time scale, so at all times $\dot{N}/N \propto -1/t_n$. Such scaling was recently seen in the initial loss rate in a harmonically trapped ^{85}Rb gas quenched to unitarity [16].

In a thermal unitary Bose gas, $\dot{N}/N \propto -\langle n^2 \rangle / T^2$ [19,20,22], where T is the temperature, so in our homogeneous case $\dot{N}/N \propto -N^2/T^2$. Here a is replaced by the thermal wavelength $\lambda \propto 1/\sqrt{T}$, which is a statistical measure of the inverse relative atomic momenta. We now have two length scales, $n^{-1/3}$ and λ , but their dynamics are coupled and we can still predict a simple scaling law in the form of Eq. (2). Heating occurs because atoms with lower relative momenta have a higher unitarity-limited loss rate and are preferentially lost from the system. Following Ref. [19], for a homogeneous gas we get $dT/T = -(4/9)dN/N$, so $T \propto N^{-4/9}$, and we thus predict $\gamma = 2 + 2 \times 4/9 = 26/9$. Note that this theory also implies that at long times $T \propto t_{\text{hold}}^{2/13}$, in agreement with our measurements shown in Fig. 2(b).

To experimentally study the instantaneous loss rate, we numerically differentiate our atom-loss curves, $N(t_{\text{hold}})$. In Fig. 3, we plot \dot{N} versus N for $N_0 = 214 \times 10^3$. We clearly observe both the degenerate-gas $\gamma = 2/3$ behavior, for large N (short t_{hold}), and the thermal-gas $\gamma = 26/9$ behavior, for small N (long t_{hold}).

In Fig. 4, we explore the universal features of the crossover between the two distinct unitary-gas regimes. In Fig. 4(a), we plot \dot{N} versus N for different N_0 and see that all the degenerate-gas data (large N/N_0) follow the same $\gamma = 2/3$ law (solid line). Writing $\dot{N}/N = -A/t_n$, we get $A = 0.28(3)$; we assess that due to the $< 10\%$ difference between N_{obs} and the true N we might overestimate A by up to 0.04. For comparison, from the ^{85}Rb data [16] we extract a slightly lower $A \approx 0.18$, which is consistent [26]




FIG. 4. Universal crossover from a degenerate to a thermal unitary gas. (a) The same $2/3$ law, $\dot{N}/N = -0.28/t_n$ (solid line), describes degenerate gases with different N_0 . The dashed lines show the $\gamma = 26/9$ scaling. (b) For all N_0 the crossover occurs at almost the same N/N_0 and t_{hold}/t_{n_0} ; averaging gives $N_c = 0.43(4)N_0$ and $t_c = 4.0(4)t_{n_0}$. (c) Plotting the dimensionless loss-rate Γ , defined in Eq. (3), versus E/E_n yields a single universal curve, with the crossover at $E_c = 1.7(2)E_n$. The solid line is $\Gamma = 0.28$, and the dashed one shows the expected $\Gamma \propto (E_n/E)^2$.

with the difference in the Efimov width parameters, $\eta^* \approx 0.09$ for ^{39}K [20] and ≈ 0.06 for ^{85}Rb [12].

For all N_0 , the small N/N_0 data agree with $\gamma = 26/9$ [dashed lines in Fig. 4(a)]. Within errors, the crossover to this regime always occurs at the same fraction of the initial population and the same t_{hold} expressed in units of t_{n_0} ; see Fig. 4(b). We get $N_c = 0.43(4)N_0$ and $t_c = 4.0(4)t_{n_0}$ [53].

In Fig. 4(c), we relate the change in γ to the growth of the dimensionless E/E_n . We define the dimensionless per-particle loss rate

$$\Gamma = -t_n \dot{N}/N, \quad (3)$$

such that in a degenerate gas $\Gamma = A$ and in a thermal gas $\Gamma \propto N^{4/3}/T^2 \propto (E_n/E)^2$ (using $\dot{N}/N \propto -N^2/T^2$ and $E \propto T$). Plotting Γ versus E/E_n , the data for different N_0 collapse onto a single universal curve, with the crossover at $E_c = 1.7(2)E_n$. For an ideal Bose gas in equilibrium, this energy would be quite high, corresponding to $T \approx 3T_c$, where T_c is the BEC critical temperature. However, in a unitary gas, we expect E/E_n to be of the order of unity even at $T = 0$ [23,27,28,31]. An important challenge for future work is to disentangle the contributions to the initial growth of E/E_n (at $t_{\text{hold}} < t_c$) due to heating and due to the development of the interaction-induced correlations that coherently broaden the momentum distribution [21,23,32,35,36].




FIG. 5. Correlation dynamics. (a) ΔN versus t_{hold} for different N_0 and the same slow ramp-out, $1/R = 6 \mu\text{s}/\text{G}$. (b) $\Delta N/N$ versus the reduced time t_{hold}/t_{n_0} , for different N_0 and $1/R$; see the legend in (a). The shaded region marks the crossover to the thermal regime at $t_c = 4.0(4)t_{n_0}$. Inset: Zoom-in on the degenerate regime shows that the correlations reach a (quasi)steady state well before t_c .

We now turn to the dynamics of correlations at unitarity. Here we denote by ΔN the reduction in N_{obs} due to a slow ramp-out (still setting $N_{\text{obs}} = N$ for our largest R). In Fig. 5(a), we show ΔN versus t_{hold} for different N_0 and a same slow ramp-out. In Fig. 5(b), we plot the fractional $\Delta N/N$ versus the dimensionless t_{hold}/t_{n_0} and also show a measurement with a different R . Plotted this way, apart from their heights all the curves look essentially the same, showing that the correlation dynamics are universally determined by the initial density.

Qualitatively, the nonmonotonic behavior of $\Delta N/N$ arises due to the competition of two effects. It takes time for the correlations to develop after the interaction quench, but, on the other hand, at very long times the system is again uncorrelated, because it heats up and the phase space density drops significantly (see also [47,54]).

In the inset in Fig. 5(b), we highlight (on a linear time scale) that $\Delta N/N$ becomes essentially flat well before t_c , meaning that at least in some sense the system reaches a quasiequilibrium while it is still degenerate. The same conclusion was drawn in Ref. [21] based on momentum-distribution dynamics in a harmonically trapped gas; the

short-range correlations reflected in the molecule formation here go hand in hand with the high-momentum tails observed in Ref. [21], and in both cases these correlations are seen to grow and saturate on a time scale of about $1t_{n_0}$. Importantly, in our homogeneous system this implies a *global* (quasi)equilibrium.

Quantitatively understanding the apparently universal shape of the $\Delta N/N$ curves, and their heights, is an interesting and challenging problem for future work. Based on the recent observation with ^{85}Rb [16], our atom-molecule mixture likely contains trimers in the first excited Efimov state, which could lead to a nontrivial dependence of the peak value of $\Delta N/N$ on the initial density. This state has an on-resonance size of the order of $1 \mu\text{m}$ [9,10,14,55] and could set a scale that separates “small” and “large” densities.

In conclusion, we have performed a comprehensive study of the particle-loss, energy, and correlation dynamics in an initially degenerate homogeneous Bose gas quenched to unitarity. We have demonstrated the anticipated scaling laws characterizing both a degenerate and a nondegenerate gas, observed universal features of both the dynamical crossover to the thermal regime and of the correlation dynamics, and found that the cloud attains a quasiequilibrium state while it is still degenerate. In the future, it would be interesting to also study the composition of the molecular cloud created by the magnetic-field sweep [16] and distinguish the two- and three-body correlation dynamics at unitarity.

Data supporting this publication are available for download at [56].

We acknowledge inspiring discussions with Alex Gaunt, Eric Cornell, Debbie Jin, and Francesca Ferlaino and thank Jean-Loup Ville and Jinyi Zhang for experimental assistance. This work was supported by the Royal Society, EPSRC [Grants No. EP/N011759/1 and No. EP/P009565/1], ERC (QBox), AFOSR, and ARO. R. L. acknowledges support from the EU Marie-Curie program [Grant No. MSCA-IF-2015 704832] and Churchill College, Cambridge. N. N. acknowledges support from Trinity College, Cambridge.

* rps24@cam.ac.uk

- [1] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Feshbach resonances in ultracold gases, *Rev. Mod. Phys.* **82**, 1225 (2010).
- [2] T.-L. Ho, Universal Thermodynamics of Degenerate Quantum Gases in the Unitarity Limit, *Phys. Rev. Lett.* **92**, 090402 (2004).
- [3] M. Inguscio, W. Ketterle, and C. Salomon, Ultra-cold Fermi gases, in *Proceedings of the International School of Physics “Enrico Fermi,” Course CLXIV, Varenna, Italy* (IOS Press, Amsterdam, 2007).

- [4] *BCS-BEC Crossover and the Unitary Fermi Gas*, edited by W. Zwerger (Springer, Berlin, 2012).
- [5] M. W. Zwierlein, in *Novel Superfluids*, edited by K.-H. Bennemann and J. B. Ketterson (Oxford University Press, Oxford, 2014), Vol. 2, Sec. 18.
- [6] F. Chevy and C. Salomon, Strongly correlated Bose gases, *J. Phys. B* **49**, 192001 (2016).
- [7] V. Efimov, Energy levels arising from resonant two-body forces in a three-body system, *Phys. Lett.* **33B**, 563 (1970).
- [8] T. Kraemer, M. Mark, P. Waldburger, J. G. Danzl, C. Chin, B. Engeser, A. D. Lange, K. Pilch, A. Jaakkola, H.-C. Nägerl, and R. Grimm, Evidence for Efimov quantum states in an ultracold gas of caesium atoms *Nature (London)* **440**, 315 (2006).
- [9] E. Braaten and H.-W. Hammer, Efimov physics in cold atoms, *Ann. Phys. (Amsterdam)* **322**, 120 (2007).
- [10] M. Zaccanti, B. Deissler, C. D’Errico, M. Fattori, M. Jona-Lasinio, S. Mueller, G. Roati, M. Inguscio, and G. Modugno, Observation of an Efimov spectrum in an atomic system, *Nat. Phys.* **5**, 586 (2009).
- [11] F. Ferlaino, A. Zenesini, M. Berninger, B. Huang, H. C. Nägerl, and R. Grimm, Efimov resonances in ultracold quantum gases, *Few-Body Syst.* **51**, 113 (2011).
- [12] R. J. Wild, P. Makotyn, J. M. Pino, E. A. Cornell, and D. S. Jin, Measurements of Tan’s Contact in an Atomic Bose-Einstein Condensate, *Phys. Rev. Lett.* **108**, 145305 (2012).
- [13] O. Machtey, Z. Shotan, N. Gross, and L. Khaykovich, Association of Efimov Trimers from a Three-Atom Continuum, *Phys. Rev. Lett.* **108**, 210406 (2012).
- [14] S. Roy, M. Landini, A. Trenkwalder, G. Semeghini, G. Spagnolli, A. Simoni, M. Fattori, M. Inguscio, and G. Modugno, Test of the Universality of the Three-Body Efimov Parameter at Narrow Feshbach Resonances, *Phys. Rev. Lett.* **111**, 053202 (2013).
- [15] R. J. Fletcher, R. Lopes, J. Man, N. Navon, R. P. Smith, M. W. Zwierlein, and Z. Hadzibabic, Two- and three-body contacts in the unitary Bose gas, *Science* **355**, 377 (2017).
- [16] C. E. Klauss, X. Xie, C. Lopez-Abadia, J. P. D’Incao, Z. Hadzibabic, D. S. Jin, and E. A. Cornell, Observation of Efimov Molecules Created from a Resonantly Interacting Bose Gas, *Phys. Rev. Lett.* **119**, 143401 (2017).
- [17] J. P. D’Incao, Few-body physics in resonantly interacting ultracold quantum gases, [arXiv:1705.10860](https://arxiv.org/abs/1705.10860).
- [18] C. Langmack, R. Schmidt, and W. Zwerger, Efimov states near a Feshbach resonance and the limits of van der Waals universality at finite background scattering length, [arXiv:1709.00749](https://arxiv.org/abs/1709.00749).
- [19] B. S. Rem, A. T. Grier, I. Ferrier-Barbut, U. Eismann, T. Langen, N. Navon, L. Khaykovich, F. Werner, D. S. Petrov, F. Chevy, and C. Salomon, Lifetime of the Bose Gas with Resonant Interactions, *Phys. Rev. Lett.* **110**, 163202 (2013).
- [20] R. J. Fletcher, A. L. Gaunt, N. Navon, R. P. Smith, and Z. Hadzibabic, Stability of a Unitary Bose Gas, *Phys. Rev. Lett.* **111**, 125303 (2013).
- [21] P. Makotyn, C. E. Klauss, D. L. Goldberger, E. A. Cornell, and D. S. Jin, Universal dynamics of a degenerate unitary Bose gas, *Nat. Phys.* **10**, 116 (2014).
- [22] U. Eismann, L. Khaykovich, S. Laurent, I. Ferrier-Barbut, B. S. Rem, A. T. Grier, M. Delehaye, F. Chevy, C. Salomon, L.-C. Ha, and C. Chin, Universal Loss Dynamics in a Unitary Bose Gas, *Phys. Rev. X* **6**, 021025 (2016).
- [23] J. M. Diederix, T. C. F. van Heijst, and H. T. C. Stoof, Ground state of a resonantly interacting Bose gas, *Phys. Rev. A* **84**, 033618 (2011).
- [24] W. Li and T.-L. Ho, Bose Gases near Unitarity, *Phys. Rev. Lett.* **108**, 195301 (2012).
- [25] X. Yin and L. Radzihovsky, Quench dynamics of a strongly interacting resonant Bose gas, *Phys. Rev. A* **88**, 063611 (2013).
- [26] A. G. Sykes, J. P. Corson, J. P. D’Incao, A. P. Koller, C. H. Greene, A. M. Rey, K. R. A. Hazzard, and J. L. Bohn, Quenching to unitarity: Quantum dynamics in a three-dimensional Bose gas, *Phys. Rev. A* **89**, 021601 (2014).
- [27] H. T. C. Stoof and J. J. R. M. van Heugten, Resummation of infrared divergencies in the theory of atomic Bose gases, *J. Low Temp. Phys.* **174**, 159 (2014).
- [28] S.-J. Jiang, W.-M. Liu, G. W. Semenoff, and F. Zhou, Universal Bose gases near resonance: A rigorous solution, *Phys. Rev. A* **89**, 033614 (2014).
- [29] A. Rançon and K. Levin, Equilibrating dynamics in quenched Bose gases: Characterizing multiple time regimes, *Phys. Rev. A* **90**, 021602 (2014).
- [30] S. Laurent, X. Leyronas, and F. Chevy, Momentum Distribution of a Dilute Unitary Bose Gas with Three-Body Losses, *Phys. Rev. Lett.* **113**, 220601 (2014).
- [31] M. Rossi, L. Salasnich, F. Ancilotto, and F. Toigo, Monte Carlo simulations of the unitary Bose gas, *Phys. Rev. A* **89**, 041602 (2014).
- [32] D. H. Smith, E. Braaten, D. Kang, and L. Platter, Two-Body and Three-Body Contacts for Identical Bosons Near Unitarity, *Phys. Rev. Lett.* **112**, 110402 (2014).
- [33] S. Piatecki and W. Krauth, Efimov-driven phase transitions of the unitary Bose gas, *Nat. Commun.* **5**, 3503 (2014).
- [34] S.-J. Jiang, J. Maki, and F. Zhou, Long-lived universal resonant Bose gases, *Phys. Rev. A* **93**, 043605 (2016).
- [35] T. Comparin and W. Krauth, Momentum Distribution in the Unitary Bose Gas from First Principles, *Phys. Rev. Lett.* **117**, 225301 (2016).
- [36] X. Yin and L. Radzihovsky, Postquench dynamics and prethermalization in a resonant Bose gas, *Phys. Rev. A* **93**, 033653 (2016).
- [37] Related loss studies have also been performed with Bose-Fermi mixtures [38].
- [38] S. Laurent, M. Pierce, M. Delehaye, T. Yefsah, F. Chevy, and C. Salomon, Connecting Few-Body Inelastic Decay to Quantum Correlations in a Many-Body System: A Weakly Coupled Impurity in a Resonant Fermi Gas, *Phys. Rev. Lett.* **118**, 103403 (2017).
- [39] A. L. Gaunt, T. F. Schmidutz, I. Gotlibovych, R. P. Smith, and Z. Hadzibabic, Bose-Einstein Condensation of Atoms in a Uniform Potential, *Phys. Rev. Lett.* **110**, 200406 (2013).
- [40] In experiments the density is never perfectly uniform due to edge effects, but for our box trap we assess that for more than 80% of the atoms the local E_n is within 10% of the trap-average value.
- [41] C. D’Errico, M. Zaccanti, M. Fattori, G. Roati, M. Inguscio, G. Modugno, and A. Simoni, Feshbach resonances in ultracold ^{39}K , *New J. Phys.* **9**, 223 (2007).
- [42] C. Eigen, A. L. Gaunt, A. Suleymanzade, N. Navon, Z. Hadzibabic, and R. P. Smith, Observation of Weak Collapse

- in a Bose-Einstein Condensate, *Phys. Rev. X* **6**, 041058 (2016).
- [43] Note that we use a magnetic-field gradient to cancel out the gravitational force on the atoms [39], and this results in a variation of the field across the cloud. However, this variation is only ≈ 25 mG, much smaller than ΔB .
- [44] We prepare the quasipure BEC in a 50 nK deep trap, but for experiments at unitarity we raise the trap depth to $2 \mu\text{K}$, to avoid any atom loss due to spilling or evaporation; by varying the trap depth, we have verified that the $2 \mu\text{K}$ depth is sufficient to suppress these effects.
- [45] C. A. Regal, C. Ticknor, J. L. Bohn, and D. S. Jin, Creation of ultracold molecules from a Fermi gas of atoms, *Nature (London)* **424**, 47 (2003).
- [46] E. Altman and A. Vishwanath, Dynamic Projection on Feshbach Molecules: A Probe of Pairing and Phase Fluctuations, *Phys. Rev. Lett.* **95**, 110404 (2005).
- [47] E. Hodby, S. T. Thompson, C. A. Regal, M. Greiner, A. C. Wilson, D. S. Jin, E. A. Cornell, and C. E. Wieman, Production Efficiency of Ultracold Feshbach Molecules in Bosonic and Fermionic Systems, *Phys. Rev. Lett.* **94**, 120402 (2005).
- [48] We exponentially ramp B to $> 10\Delta B$ away from B_0 , so the ramp is essentially linear in and close to the unitary regime.
- [49] Reminiscent of Ref. [50], we observe an oscillation of N_{obs} as a function of the time spent at low a between the two magnetic-field pulses and optimize this time to maximize the atom-number recovery.
- [50] E. A. Donley, N. R. Claussen, S. T. Thompson, and C. E. Wieman, Atom-molecule coherence in a Bose-Einstein condensate, *Nature (London)* **417**, 529 (2002).
- [51] P. O. Fedichev, M. W. Reynolds, and G. V. Shlyapnikov, Three-Body Recombination of Ultracold Atoms to a Weakly Bound s Level, *Phys. Rev. Lett.* **77**, 2921 (1996).
- [52] T. Weber, J. Herbig, M. Mark, H.-C. Nägerl, and R. Grimm, Three-Body Recombination at Large Scattering Lengths in an Ultracold Atomic Gas, *Phys. Rev. Lett.* **91**, 123201 (2003).
- [53] The small (not statistically conclusive) trend in the central values of t_c/t_{n_0} can arise due to the experimental imperfection of the optical-box trapping potential [39].
- [54] F. Werner and Y. Castin, General relations for quantum gases in two and three dimensions. II. Bosons and mixtures, *Phys. Rev. A* **86**, 053633 (2012).
- [55] P. M. A. Mestrom, J. Wang, C. H. Greene, and J. P. D’Incao, Efimov–van der Waals universality for ultracold atoms with positive scattering lengths, *Phys. Rev. A* **95**, 032707 (2017).
- [56] <https://doi.org/10.17863/CAM.16741>.