## Logarithmically Slow Relaxation in Quasiperiodically Driven Random Spin Chains

Philipp T. Dumitrescu,<sup>1,\*</sup> Romain Vasseur,<sup>2,3,4</sup> and Andrew C. Potter<sup>1</sup>

<sup>1</sup>Department of Physics, University of Texas at Austin, Austin, Texas 78712, USA

<sup>2</sup>Department of Physics, University of California, Berkeley, Berkeley, California 94720, USA

<sup>3</sup>Materials Science Division, Lawrence Berkeley National Laboratories, Berkeley, California 94720, USA

<sup>4</sup>Department of Physics, University of Massachusetts, Amherst, Massachusetts 01003, USA

(Received 11 August 2017; revised manuscript received 20 November 2017; published 16 February 2018)

We simulate the dynamics of a disordered interacting spin chain subject to a quasiperiodic timedependent drive, corresponding to a stroboscopic Fibonacci sequence of two distinct Hamiltonians. Exploiting the recursive drive structure, we can efficiently simulate exponentially long times. After an initial transient, the system exhibits a long-lived glassy regime characterized by a logarithmically slow growth of entanglement and decay of correlations analogous to the dynamics at the many-body delocalization transition. Ultimately, at long time scales, which diverge exponentially for weak or rapid drives, the system thermalizes to infinite temperature. The slow relaxation enables metastable dynamical phases, exemplified by a "time quasicrystal" in which spins exhibit persistent oscillations with a distinct quasiperiodic pattern from that of the drive. We show that in contrast with Floquet systems, a highfrequency expansion strictly breaks down above fourth order, and fails to produce an effective static Hamiltonian that would capture the prethermal glassy relaxation.

DOI: 10.1103/PhysRevLett.120.070602

Introduction.—Interacting quantum many-body systems often exhibit chaotic dynamics that rapidly scramble quantum information and lead to highly entangled states whose local properties are thermal and classical [1,2]. A dramatic exception occurs in isolated and disordered systems where many-body localization (MBL) can arrest thermalization, resulting in quantum coherent dynamics at arbitrarily high energy density [3–5]. This dichotomy naturally raises fundamental questions about when and how a system thermalizes. What are the universal features governing the dynamical approach to the final—thermal or nonthermal—state? More practically, what classes of protocols allow one to manipulate a many-body system without rapidly scrambling its stored quantum information?

Given their large bandwidth and dense spectrum, one might naively expect that *any* persistent dynamical manipulation of an isolated, interacting quantum many-body system leads to runaway heating to a featureless infinite-temperature state. Indeed, random time-dependent manipulations have recently been shown to cause rapid growth of entanglement, accompanied by universal hydrodynamic features [6–8]. However, this expectation is violated in time-periodically driven (Floquet) systems with strong disorder, in which sufficiently rapid driving maintains MBL and indefinitely avoids heating [9–11]. Even in the absence of disorder, rapid periodic driving leads to long-lived prethermal phenomena [12–21]. Floquet-MBL systems have been shown to exhibit remarkable dynamic phenomena from spontaneous time-translation symmetry

breaking [22–27] to dynamical topological phases with no equilibrium analog [22,28–37].

The stark contrast between the behaviors under random and periodic driving can be understood by a simple argument: local time-dependent Hamiltonians can only make local rearrangements. In strongly disordered systems, such rearrangements have a nonzero energy cost and are generically nonresonant with harmonics of the driving frequency. This heuristic forms the basis for more sophisticated considerations for the stability of Floquet-MBL systems [11], which are supported by numerical simulations [9,10], and cold-atom experiments [38]. Using similar arguments, one can rule out the stability of MBL to random time-dependent drives, which have continuous frequency spectra capable of resonantly inducing arbitrary local transitions leading to thermalization.

In this Letter, we consider an intermediate case between periodic and random driving by subjecting a strongly disordered quantum many-body system to a drive with quasiperiodic time dependence. The quasiperiodic drive has a dense, but sharply discontinuous frequency spectrum that occupies a set of measure zero. *A priori*, it is not clear whether the density of spectral content will drive heating and thermalization or whether its sparsity will preserve MBL. We find that quasiperiodic driving does eventually lead to thermalization to a featureless infinite temperature state, but only after a long time  $t_{th}$  that grows exponentially in the inverse driving strength and the rate of driving. While reminiscent of prethermalization in delocalized Floquet systems [12–16], the dynamics before  $t_{th}$  are not described by an effective finite temperature equilibrium. Instead, this regime shows a logarithmically slow relaxation of correlations and growth of entanglement, which we will call glassy dynamics. This glassy behavior is analogous to the critical dynamics at the transition between MBL and thermal systems in nondriven settings [39–41]. We explore to what extent the quasiperiodic evolution can be reduced to an effective static Hamiltonian, connecting our study to the question of reducibility of differential equations with quasiperiodic coefficients [42,43]. The glassy relaxation regime can host new metastable dynamical phases, which we illustrate with a quasiperiodic analog of time-translation symmetry breaking—a "time quasicrystal."

*Model.*—To address the fate of a quantum many-body system under quasiperiodic driving, we numerically simulate spin-1/2 chains, subjected to a stroboscopic drive consisting of a Fibonacci sequence of unitary evolutions:

$$U_n = U_{n-2}U_{n-1},$$
 (1)

for  $n \ge 2$ . The sequence is initialized by two elementary unitaries formed from two different static Hamiltonian evolutions:  $U_0 = \exp(-i\lambda H_+)$  and  $U_1 = \exp(-i\lambda H_-)$ , where

$$H_{\pm} = \sum_{i=1}^{L} h_i S_i^z + \sum_{i=1}^{L-1} (J_0 \pm \delta J) \mathbf{S}_i \cdot \mathbf{S}_{i+1}.$$
 (2)

The  $h_i$  are random fields drawn independently for each site from a uniform distribution  $h \in [-2\pi, 2\pi)$ ,  $J_0$  is a static interaction,  $\delta J$  represents the strength of the quasiperiodic driving, and  $\lambda \in [0, 1]$  is the characteristic driving time scale. We will focus on the regime  $|J_0 \pm \delta J| \lesssim 1.7$ , where  $H_{\pm}$  as static Hamiltonians would be MBL [44]. As such, they are separately described by emergent local integrals of motion (LIOM) with definite  $S^z$  value [45]. Unless otherwise noted, we will take  $J_0 = 0$ . An appealing feature of the recursive nature of the drive is that it enables simulation of exponentially long Fibonacci times  $t_n = F_{n+1} \sim \varphi^{n+1}$ with only *n* unitary multiplications; here  $\varphi = (1 + \sqrt{5})/2$ is the golden ratio. This enables us to simulate the longtime physics, limited only by machine precision.

*Results.*—We focus on three observables: the *z* component of spin  $C^{zz}(t) = 4\langle S_i^z(t)S_i^z(0)\rangle$ , whose total value is conserved by the evolution, and whose local dynamics are related to spin transport; the transverse spin fluctuations  $C^{+-}(t) = 4|\langle S_i^+(t)S_i^-(0)\rangle|$ , which encodes the dephasing of quantum superpositions of up and down spins; and the bi-partite (half-system) entanglement entropy  $S_{bp}(t)$ .

Before discussing the results, we summarize the behavior of these quantities in static MBL, periodically driven (Floquet) MBL, and thermalizing systems. In a static or Floquet-MBL system,  $C^{zz}(t)$  tends to a nonzero constant at long times, indicating the absence of spin transport and emergent conservation laws that produce infinite memory of the initial spin configuration [9,10,45,46]. The transverse fluctuations  $C^{+-}(t)$  decay as a power law in time from dephasing due to classical interactions among the local conserved quantities [47]. This dephasing also produces a logarithmically slow growth of entanglement  $S_{bp}(t) \sim \log t$ [48–50]. On the other hand, in strongly thermal or randomly driven systems, the nonzero spin conductivity and chaotic scrambling leads to an exponential decay of correlation functions  $C^{zz}$ ,  $C^{+-} \sim e^{-t/t_{\rm th}}$  and a linear growth in  $S_{\rm hp}(t) \sim$ t [51,52]. Finally, a clean delocalized system subject to rapid periodic driving exhibits a prethermalization regime, in which the system initially equilibrates with respect to an effective Hamiltonian at finite temperature. Prethermalization persists up to a time exponentially long in the driving frequency [12-15], after which the system heats to a featureless infinite temperature state.

Figure 1 shows  $C^{zz}$ ,  $C^{+-}$ , and  $S_{bp}$  for quasiperiodic driving, in a quench from an initial product state. These



FIG. 1. Quasiperiodically driven spin chain. Time evolution under the quasiperiodic driving sequence, with  $J_0 = 0$ ,  $\delta J = \pi/30$ ,  $\lambda = 1$ and varying L [markers defined in (c)]. All quantities are averaged over states in the global  $S^z = 0$  sector of the spin chain and averaged over at least 3000 disorder realizations. (a) The bipartite entanglement  $S_{bp}(t)$ . Inset: The normalized entanglement  $S_{bp}/L$ . (b) Onsite correlation function  $C^{zz}(t)$  on site i = L/2. This plot additionally shows (dashed line) the case of driving an L = 150 chain in the noninteracting limit of Eq. (2); see the Supplemental Material [53]. Inset: Comparison of driving with periodic (P), quasiperiodic (QP), and random (R) sequences of the elementary unitaries, with L = 8. The random case is averaged over 20 different random sequences, each with 100 disorder realizations. (c) Correlation function  $C^{+-}(t)$  on site i = L/2.

observables are averaged over initial states and disorder realizations. We observe three distinct regimes: First, there is a short-time transient regime in which there is no distinction between periodic, quasiperiodic and random driving [Fig. 1(b), inset]. Next, there is a long-lived glassy relaxation regime where  $S_{bp}$  grows and  $C^{zz}$  decays logarithmically slowly. Finally, after a time-scale  $t_{th}$  that is exponentially long for weak or rapid driving, the system ultimately heats up to infinite temperature with a nonzero rate, signaled by linear growth of entanglement and rapid decay of correlations. Ultimately,  $S_{bp}$  will saturate to its thermal value and  $C^{zz}$ ,  $C^{+-}$  decay to zero.

The behavior of this quasiperiodic system is markedly distinct from the other scenarios mentioned above, as contrasted in the inset of Fig. 1(b). Similar to an MBL system,  $C^{+-}$  shows aperiodic oscillations that decay slowly. Unlike an MBL system, however,  $C^{zz}$  does not saturate to a nonzero value. Taken together, these imply that the glassy relaxation regime does not possess LIOM. Nonetheless, it does not exhibit the rapid decay characteristic of a thermal system.

There are two ways we can identify the thermalization time  $t_{\rm th}$ : as the time where  $C^{zz}$  curves of different *L* separate from each other after the logarithmic decay or as the time where the normalized entanglement curves  $S_{\rm bp}/L$  cross at a single point as a function of *Lt* [Fig. 1(a), inset]. These two ways of extracting  $t_{\rm th}$  follow each other closely and allow us to extract the parametric dependence of  $t_{\rm th}$  on  $\delta J$  and  $\lambda$ (Fig. 2) [55]. At small  $\lambda$  and  $\delta J$  we find an asymptotic dependance which is consistent with  $t_{\rm th} \sim e^{1/\lambda}$ ,  $t_{\rm th} \sim e^{1/\delta J}$ , implying an anomalously slow dephasing and decay over extremely long time scales. At larger  $\lambda$ ,  $\delta J$  there may be deviations from this form. In this respect, the logarithmic decay is reminiscent of the long-lived prethermal regime of non-MBL Floquet systems [12–16]. However, the



FIG. 2. Thermalization time  $t_{\text{th}}$ . Thermalization time extracted from the crossing of  $S_{\text{bp}}/L$ , between pairs of L (6,8),(8,10), (10,12). (a) As a function of  $1/\delta J$  for  $\lambda = 1/2\pi$ , 1 and (b) as a function of  $\lambda$  for  $\delta J = \pi/20$ ,  $\pi/5$ . Error bars are linear estimates in Fibonacci time; dashed lines are fits of form log  $t_n \sim 1/\lambda$ ,  $1/\delta J$ to the (10,12) crossing.

entanglement growth in this region is slower than linear and consistent with logarithmic growth, which would not be the case of a system equilibrating to an effective finite temperature and prethermal Hamiltonian. We note that such logarithmic decay is observed at the phase transition between MBL and thermal phases [39–41]; here, we see this critical-like behavior without fine-tuning.

It is interesting to compare these results to those of a noninteracting analog of Eq. (2) (dashed line in Fig. 1(b), for detailed comparison see [53]). The noninteracting system also exhibits a slow decay regime, but in this case there is no crossover to fast thermalization ( $t_{th} = +\infty$ ). This suggests that, despite the absence of local conserved quantities, the long lived glassy relaxation regime in the interacting case is nonetheless governed by the dynamics of emergent single-particle-like degrees of freedom.

(*Ir*)reducibility of the quasiperiodic drive.—Highfrequency expansions provide a useful tool for understanding prethermalization behavior in Floquet systems. They enable the computation of an effective static prethermal Hamiltonian and the expansion breakdown at long times indicates the onset of thermalization. Here, we attempt to develop a generic expansion of the many-body time-evolution operator organized in powers of  $\lambda$ —effectively a Magnus expansion—taking advantage of the special selfsimilar structure of the Fibonacci drive. Technical details are given in the Supplemental Material [53].

We can analytically construct a recursive Magnus expansion for  $\Omega_n = \log U_n$ , using the local deflation rule structure of quasiperiodic sequences [56,57]. We can generate  $U_{n+1}$  from  $U_n$  by replacing  $U_0 \rightarrow U_1$  and  $U_1 \rightarrow$  $U_0U_1$  in the product defining  $U_n$ . We expand  $\Omega_n$  onto a basis of nested commutators and construct and solve difference equations for the coefficients in this expansion, order by order in the degree k of the commutator basis. Up to degree two,

$$\Omega_n = F_{n-1}\Omega_0 + F_n\Omega_1 + \frac{1}{2}\{(-1)^n + F_{n-2}\}[\Omega_0, \Omega_1].$$

Explicit expressions for degrees k = 3, 4 are given in the Supplemental Material [53]. In order to assign an effective static Hamiltonian interpretation, the asymptotic form for all coefficients need to be  $\sim \varphi^n$ , as above. However, for  $k \ge 4$ , the asymptotic behavior is  $\sim \varphi^{(k-2)n}$ . Therefore, the time where the non-Hamiltonian evolution dominates becomes increasingly short  $t_n \sim \lambda^{-(k-1)/(k-3)}$ . We note that this breakdown is fundamentally different from the breakdown of thermalization in the Floquet-Magnus case for periodic driving, which is due to a lack of convergence of the expansion.

Despite this, we find that truncating the expansion at k = 3 gives a Hamiltonian evolution which reproduces the data at small  $\lambda$  remarkably well, with the exception of rare anomalous disorder configurations. Indeed, the time where this expansion deviates from the data scales with  $\lambda^{-5}$ , much



FIG. 3. Magnus expansion. (a) Onsite correlation function  $C^{zz}(t)$  for L = 8,  $J_0 = 0$ ,  $\delta J = \pi/5$  and different  $\lambda$  compared to that obtained by the Magnus expansion Hamiltonian at third order. (b) Initial time at which  $C^{zz}(t)$  of the Magnus expansion deviates by more than  $10^{-4}$  from the data.

later than the expected  $\lambda^{-3}$  (Fig. 3). In no case, however, does the Magnus expansion capture the anomalous logarithmic decay of  $C^{zz}$  or growth of  $S_{bp}$  for  $t < t_{th}$ , suggesting these are inherently dynamical phenomena not governed by a static Hamiltonian, i.e., not governed by an effective conserved (quasi)-energy.

*Fibonacci time quasicrystal.*—The existence of an exponentially long-lived quasi-MBL regime, with only logarithmically slow decay, raises the prospect of transient phases unique to quasiperiodically driven systems. These are analogous to metastable phases in prethermal Floquet settings, but with the important distinction that the quasiperiodically driven system does not require cooling to observe quantum coherent behavior. To illustrate this possibility, we now construct a model that exhibits the quasiperiodic analog of discrete time-translation symmetry-breaking [22–27]—a "time quasicrystal" (TQC). The model uses the Fibonacci sequence of Eq. (1), but with elementary unitaries

$$U_0 = e^{-i\theta \sum_i S_i^x}, \qquad U_1 = e^{-i\lambda \sum_i (J_i S_i^z S_{i+1}^z + h_i^z S_i^z + h_i^x S_i^x)}.$$
 (3)

This model is closely inspired by the periodic version introduced in Refs. [22,24].

Consider the ideal case of Eq. (3), where  $\theta = \pi$ ,  $h_i^x = 0$ and random  $J_i$ ,  $h_i^z$ . Then  $U_0 \sim \prod_i S_i^x$  applies a perfect, global spin flip, while  $U_1$  is made of only  $S^z$  operators. A simple  $S^z$ -product state would merely acquire a phase under  $U_1$  and flip under  $U_0$ . The time evolution of a specific spin  $\langle S_i^z(t) S_i^z(0) \rangle$  exhibits an oscillating quasiperiodic pattern that is sharply distinct from the driving pattern. An elegant way to capture this difference is to view the quasiperiodic sequence as a projection of a 1D strip cutting through a regular 2D square lattice at an irrational angle (see Ref. [53]). The TQC spin response corresponds to a projection from a 2D lattice having a doubled unit cell compared to that for the drive.

Alternatively, we can directly compare the Fourier spectrum of the spin response to that of the drive [56,57]. For this, it is convenient to interpret  $U_0$  in Eq. (1) as arising from an instantaneous pulse, so that we can write the evolution in terms of a Hamiltonian with quasiperiodic delta-function "kicking":  $H(t) = H_1 + \sum_{m=1}^M \delta(t - t_m) H_0$ , where  $t_m =$  $\lfloor \varphi m \rfloor$  and M is the largest integer such that  $t_M \leq t$ . In the ideal limit  $\theta = \pi$ ,  $h_i^x = 0$ , the correlation function would satisfy  $dC^{zz}(t)/dt = 2\sum_{m=1}^{M} (-1)^m \delta(t-t_m)$ . The spectrum of the spin response is shifted compared to the drive (see Fig. 4 and the Supplemental Material [53]). The distinction between the spin response and drive patterns is even simpler if we consider stroboscopically measuring  $C^{zz}(t)$ at Fibonacci times  $t_n = F_n$ . At these times, the initial spins have been flipped  $F_{n-1} \mod 2$  times from their initial state. Since  $F_k \mod 2$  form a repeating pattern with period 3; the TQC is characterized by persistent period-3 oscillations in Fibonacci time.

These aspects also generalize straightforwardly to other time quasicrystal phases. For example, we may replace the Ising spins ( $\mathbb{Z}_2$ ) by *N*-state clock spins ( $\mathbb{Z}_N$ ) in  $U_1$  and replace  $S^x$  by the operator that increments the clock spins in  $U_0$  of Eq. (3). In Fibonacci time, the spins would oscillate



FIG. 4. Time quasicrystal. (a) TEBD data of a single spin in a spin-1/2 chain subjected to drive Eq. (3) with  $U_0$  occurring instantaneously. Parameters are L = 60,  $\lambda = 1$ , and  $\theta = \pi - 0.1$ , as well as random variables drawn from uniform distributions  $J \in [2, 8]$ ,  $h_i^z \in [0, 2]$ ,  $h_i^x \in [0, 0.6]$ . We show a single disorder realization. (b) Fourier spectra of the quasiperiodic (QP) drive pattern, of the ideal TQC pattern and of the TEBD data. (c) Magnetization at Fibonacci times, for an ideal ( $\theta = \pi$ ) and nonideal ( $\theta = \pi - 0.1$ ) pulse, shows period-3 oscillations characteristic of the TQC.

with the Pisano period  $\pi(N)$ ; for  $N = 2, 3, 4, 5, \pi(N) = 3$ , 8, 6, 20. While the emergence of quasiperiodic correlations that have a different pattern from the drive can occur in ideally driven single spins [58], this is special to fine-tuned drivings. In the many-body set-up (3), the interactions give phase rigidity even away from the ideal limit  $\theta = \pi$ , as for a Floquet time crystal [25].

For  $\theta \neq \pi$  or  $h_x \neq 0$ , the model becomes nonintegrable and we lose analytic control. Figure 4 shows  $C^{zz}(t)$  from time-evolving block decimation (TEBD) [59-61] for system size L = 60 starting from a product state. The TEBD calculations were done with Trotter step  $0.01\lambda$ , keeping the discarded weight below  $10^{-7}$  throughout the time evolution. Away from the ideal limit, the results largely track the ideal oscillations, but we clearly see the overall logarithmic decay in the quasiperiodic oscillations due to the quasi-MBL nature as discussed in the previous sections. In the Heisenberg chain [Eq. (2)] discussed above, the glassy relaxation was smoothly connected to the noninteracting limit. It is intriguing that this behavior is again observed in a system that is unconnected to any free fermion limit due to the longitudinal fields. This again suggests a possible description in terms of an emergent set of effectively singleparticle, though nonconserved, degrees of freedom.

Despite that the system eventually thermalizes, for moderately small  $\lambda$  the decay is sufficiently slow to permit many period-3 oscillations in Fibonacci time. This is a fundamentally different type of approximate nonequilibrium order than previously discussed for the cases of prethermal order in Floquet systems [12–16], which require cooling to an effective prethermal ground state.

Beyond this quasiperiodic generalization of a Floquet time crystal, the slow relaxation in the long-lived regime of glassy relaxation opens the door to more exotic quantum dynamical behavior such as long-lived quasiperiodic topological phenomena. Investigating this intriguing possibility, and developing a systematic theoretical framework to characterize such metastable quantum phases will be an important challenge for future work.

We thank M. Kolodrubetz and N. Y. Yao for insightful discussions. We especially thank S. Gopalakrishnan for pointing out that the glassy relaxation also occurs in the noninteracting limit. Numerical simulations were performed at the Texas Advanced Computing Center (TACC) at the University of Texas at Austin. This work was supported by NSF DMR-1653007 (A. C. P.) and the LDRD program at LBNL (R. V.). This work was performed in part at Aspen Center for Physics, which is supported by National Science Foundation Grant No. PHY-1607611 (P. T. D. & A. C. P).

<sup>\*</sup>philippd@utexas.edu

- [3] R. Nandkishore and D. A. Huse, Annu. Rev. Condens. Matter Phys. 6, 15 (2015).
- [4] E. Altman and R. Vosk, Annu. Rev. Condens. Matter Phys. 6, 383 (2015).
- [5] R. Vasseur and J. E. Moore, J. Stat. Mech. (2016) 064010.
- [6] A. Nahum, J. Ruhman, S. Vijay, and J. Haah, Phys. Rev. X 7, 031016 (2017).
- [7] C. von Keyserlingk, T. Rakovszky, F. Pollmann, and S. Sondhi, arXiv:1705.08910.
- [8] A. Nahum, S. Vijay, and J. Haah, arXiv:1705.08975.
- [9] A. Lazarides, A. Das, and R. Moessner, Phys. Rev. Lett. 115, 030402 (2015).
- [10] P. Ponte, Z. Papić, F. Huveneers, and D. A. Abanin, Phys. Rev. Lett. **114**, 140401 (2015).
- [11] D. A. Abanin, W. D. Roeck, and F. Huveneers, Ann. Phys. (Amsterdam) 372, 1 (2016).
- [12] D. A. Abanin, W. De Roeck, W. W. Ho, and F. Huveneers, Phys. Rev. B 95, 014112 (2017).
- [13] D. A. Abanin, W. De Roeck, and F. Huveneers, Phys. Rev. Lett. 115, 256803 (2015).
- [14] T. Kuwahara, T. Mori, and K. Saito, Ann. Phys. (Amsterdam) 367, 96 (2016).
- [15] T. Mori, T. Kuwahara, and K. Saito, Phys. Rev. Lett. 116, 120401 (2016).
- [16] D. V. Else, B. Bauer, and C. Nayak, Phys. Rev. X 7, 011026 (2017).
- [17] S. A. Weidinger and M. Knap, Sci. Rep. 7, 45382 (2017).
- [18] A. Chandran and S. L. Sondhi, Phys. Rev. B 93, 174305 (2016).
- [19] E. Canovi, M. Kollar, and M. Eckstein, Phys. Rev. E 93, 012130 (2016).
- [20] M. Bukov, S. Gopalakrishnan, M. Knap, and E. Demler, Phys. Rev. Lett. 115, 205301 (2015).
- [21] S. Gopalakrishnan, K. R. Islam, and M. Knap, Phys. Rev. Lett. 119, 046601 (2017).
- [22] V. Khemani, A. Lazarides, R. Moessner, and S. L. Sondhi, Phys. Rev. Lett. 116, 250401 (2016).
- [23] C. W. von Keyserlingk, V. Khemani, and S. L. Sondhi, Phys. Rev. B 94, 085112 (2016).
- [24] D. V. Else, B. Bauer, and C. Nayak, Phys. Rev. Lett. 117, 090402 (2016).
- [25] N. Y. Yao, A. C. Potter, I.-D. Potirniche, and A. Vishwanath, Phys. Rev. Lett. **118**, 030401 (2017).
- [26] J. Zhang, P. W. Hess, A. Kyprianidis, P. Becker, A. Lee, J. Smith, G. Pagano, I. D. Potirniche, A. C. Potter, A. Vishwanath, N. Y. Yao, and C. Monroe, Nature (London) 543, 217 (2017).
- [27] S. Choi, J. Choi, R. Landig, G. Kucsko, H. Zhou, J. Isoya, F. Jelezko, S. Onoda, H. Sumiya, V. Khemani, C. von Keyserlingk, N. Y. Yao, E. Demler, and M. D. Lukin, Nature (London) 543, 221 (2017).
- [28] T. Kitagawa, E. Berg, M. Rudner, and E. Demler, Phys. Rev. B 82, 235114 (2010).
- [29] L. Jiang, T. Kitagawa, J. Alicea, A. R. Akhmerov, D. Pekker, G. Refael, J. I. Cirac, E. Demler, M. D. Lukin, and P. Zoller, Phys. Rev. Lett. 106, 220402 (2011).
- [30] M. S. Rudner, N. H. Lindner, E. Berg, and M. Levin, Phys. Rev. X 3, 031005 (2013).
- [31] C. W. von Keyserlingk and S. L. Sondhi, Phys. Rev. B 93, 245145 (2016).

<sup>[1]</sup> J. M. Deutsch, Phys. Rev. A 43, 2046 (1991).

<sup>[2]</sup> M. Srednicki, Phys. Rev. E 50, 888 (1994).

- [32] D. V. Else and C. Nayak, Phys. Rev. B 93, 201103 (2016).
- [33] A. C. Potter, T. Morimoto, and A. Vishwanath, Phys. Rev. X 6, 041001 (2016).
- [34] R. Roy and F. Harper, Phys. Rev. B 94, 125105 (2016).
- [35] H. C. Po, L. Fidkowski, T. Morimoto, A. C. Potter, and A. Vishwanath, Phys. Rev. X 6, 041070 (2016).
- [36] F. Harper and R. Roy, Phys. Rev. Lett. 118, 115301 (2017).
- [37] H. C. Po, L. Fidkowski, A. Vishwanath, and A. C. Potter, Phys. Rev. B 96, 245116 (2017).
- [38] P. Bordia, H. Luschen, U. Schneider, M. Knap, and I. Bloch, Nat. Phys. 13, 460 (2017).
- [39] R. Vosk, D. A. Huse, and E. Altman, Phys. Rev. X 5, 031032 (2015).
- [40] A. C. Potter, R. Vasseur, and S. A. Parameswaran, Phys. Rev. X 5, 031033 (2015).
- [41] M. Serbyn, Z. Papić, and D. A. Abanin, Phys. Rev. X 5, 041047 (2015).
- [42] H. Jauslin and J. Lebowitz, Chaos 1, 114 (1991).
- [43] P. Blekher, H. Jauslin, and J. Lebowitz, J. Stat. Phys. 68, 271 (1992).
- [44] D. J. Luitz, N. Laflorencie, and F. Alet, Phys. Rev. B 91, 081103 (2015).
- [45] M. Serbyn, Z. Papić, and D. A. Abanin, Phys. Rev. Lett. 111, 127201 (2013).
- [46] D. A. Huse, R. Nandkishore, and V. Oganesyan, Phys. Rev. B 90, 174202 (2014).
- [47] M. Serbyn, Z. Papić, and D. A. Abanin, Phys. Rev. B 90, 174302 (2014).

- [48] M. Žnidarič, T. Prosen, and P. Prelovšek, Phys. Rev. B 77, 064426 (2008).
- [49] J. H. Bardarson, F. Pollmann, and J. E. Moore, Phys. Rev. Lett. 109, 017202 (2012).
- [50] M. Serbyn, Z. Papić, and D. A. Abanin, Phys. Rev. Lett. 110, 260601 (2013).
- [51] P. Calabrese and J. Cardy, J. Stat. Mech. (2005) P04010.
- [52] H. Kim and D. A. Huse, Phys. Rev. Lett. 111, 127205 (2013).
- [53] See Supplemental Material at http://link.aps.org/ supplemental/10.1103/PhysRevLett.120.070602 for details regarding the quasiperiodic drive in the noninteracting limit, the construction and breakdown of the recursive Magnus expansion, and Fourier response of the time quasicrystal, which includes Ref. [54].
- [54] C. Reutenauer, *Free Lie Algebras*, London Mathematical Society Vol. 7 (Clarendon Press, Oxford, 1993).
- [55] Since  $t_{\text{th}}$  is a cross-over scale, some observables show deviations earlier than others.
- [56] D. Levine and P.J. Steinhardt, Phys. Rev. B 34, 596 (1986).
- [57] J. E. S. Socolar and P. J. Steinhardt, Phys. Rev. B 34, 617 (1986).
- [58] U. Feudel, A. S. Pikovsky, and M. A. Zaks, Phys. Rev. E 51, 1762 (1995).
- [59] G. Vidal, Phys. Rev. Lett. 91, 147902 (2003).
- [60] G. Vidal, Phys. Rev. Lett. 93, 040502 (2004).
- [61] U. Schollwöck, Ann. Phys. (Amsterdam) **326**, 96 (2011), january 2011 Special Issue.