

Generalized Entanglement Entropies of Quantum Designs

Zi-Wen Liu,^{1,2,*} Seth Lloyd,^{3,2} Elton Yechao Zhu,^{1,2} and Huangjun Zhu^{4,5,6,7,8}

¹*Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

²*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

³*Department of Mechanical Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

⁴*Institute for Theoretical Physics, University of Cologne, 50937 Cologne, Germany*

⁵*Department of Physics and Center for Field Theory and Particle Physics, Fudan University, Shanghai 200433, China*

⁶*Institute for Nanoelectronic Devices and Quantum Computing, Fudan University, Shanghai 200433, China*

⁷*State Key Laboratory of Surface Physics, Fudan University, Shanghai 200433, China*

⁸*Collaborative Innovation Center of Advanced Microstructures, Nanjing 210093, China*



(Received 11 September 2017; published 28 March 2018)

The entanglement properties of random quantum states or dynamics are important to the study of a broad spectrum of disciplines of physics, ranging from quantum information to high energy and many-body physics. This Letter investigates the interplay between the degrees of entanglement and randomness in pure states and unitary channels. We reveal strong connections between designs (distributions of states or unitaries that match certain moments of the uniform Haar measure) and generalized entropies (entropic functions that depend on certain powers of the density operator), by showing that Rényi entanglement entropies averaged over designs of the same order are almost maximal. This strengthens the celebrated Page's theorem. Moreover, we find that designs of an order that is logarithmic in the dimension maximize all Rényi entanglement entropies and so are completely random in terms of the entanglement spectrum. Our results relate the behaviors of Rényi entanglement entropies to the complexity of scrambling and quantum chaos in terms of the degree of randomness, and suggest a generalization of the fast scrambling conjecture.

DOI: [10.1103/PhysRevLett.120.130502](https://doi.org/10.1103/PhysRevLett.120.130502)

Introduction.—The interplay between entanglement and randomness plays important roles in many areas of physics. A particular notion of wide interest is “scrambling,” which describes the phenomenon that initially localized quantum information spreads over the entire system via global entanglement, so that the information is lost from the perspective of any local observer, or the state of the system is effectively randomized. The concept of scrambling originates from the study of black holes and quantum gravity [1–4], and similar mechanisms also underlie many other key concepts in physics, such as quantum chaos [5–7], quantum thermalization [8,9], and quantum data hiding [10,11]. The entanglement properties of random or pseudorandom quantum states and channels can illuminate such phenomena and are fundamental to relevant studies.

It has long been noted that a random state is typically highly entangled [12,13]. This observation is formalized by Page's theorem [1,14–16], which states that the expected von Neumann entropy of small subsystems of a completely random state (drawn from the Haar measure) is very close to the maximum. Similar observations for the entanglement in random unitary channels have been made recently in [7]. However, such results are not “tight” from the perspective of complexity. On the one hand, the complexity of Haar randomness is high: the number of local gates needed to even approximate the Haar distribution grows

exponentially in the number of qubits [17]. On the other hand, however, pseudorandom distributions with low complexity [18–21] are sufficient to acquire the Page-like entanglement property. That is, there is a significant complexity gap between complete randomness and large entanglement entropy. In dynamical scenarios, this gap corresponds to a substantial but poorly understood regime beyond scrambling, chaos and thermalization, where the randomness and complexity of the system can keep growing. Indeed, the common characteristics of information scrambling, such as global entanglement [2,7], remote signaling [3], and local indistinguishability [22], do not need nor imply complete randomization, and there is little knowledge about the physics of later times.

To fill this gap, we consider more stringent entanglement measures. The study is also extended to unitary channels via the Choi isomorphism. More specifically, we employ various techniques from representation theory, random matrix theory, combinatorics, and Weingarten calculus to analytically study the generalized entanglement entropies (which depend on higher powers of the reduced density operator) of random and pseudorandom states and unitaries. A key collective finding is that the Rényi- α entanglement entropy averaged over α -designs is almost maximal, where α -designs stand for evenly distributed ensembles of states or unitaries that mimic the first α

moments of the Haar measure, in analogy to α -wise independent distributions which have wide applications in classical computer science and combinatorics. In other words, designs represent finite-degree approximations of the truly random distribution on states or unitaries, which are of great interest in quantum information. This result links the order of entanglement entropies and that of designs and closes the complexity gap in Page-like theorems. It also suggests Rényi entanglement entropies as diagnostics of the randomness complexity of corresponding designs in scrambling, as well as (truly quantum) witnesses of quantum pseudorandomness. The infinite order limit of Rényi entropy, which only depends on the largest eigenvalue, is known as the min entropy. We further show that the min entanglement entropy (and therefore all Rényi entanglement entropies) becomes almost maximal, which we also call “max-scrambling,” for designs of an order that is only logarithmic in the dimension of the system. So designs of higher orders are essentially completely random in terms of entanglement. This leads to a strong estimate of the time needed to achieve max-scrambling based on the fast scrambling [3] and design [21] conjectures. Finally, we are able to construct state 2-designs such that all Rényi entanglement entropies of orders greater than 2 are bounded away from the maximum, which establishes an explicit separation between the complexities diagnosed by Rényi entanglement entropies.

This Letter distills the key quantum information results of [23], which is written from the perspective of scrambling complexity. Please refer to [23] for technical details and more discussions.

Preliminaries.—Here we recall the formal definitions of designs and generalized entropies, the central mathematical concepts of this study.

Designs are ensembles of quantum states (unitaries) that are evenly distributed on the complex unit sphere (unitary group). They are efficient to implement [18–21] and useful in many important quantum information processing tasks such as randomized benchmarking [24,25] and decoupling [26]. There are several ways to characterize exact or approximate designs (see, e.g., [27]), among which the one based on polynomials is the most relevant to this work. Let $\text{Hom}_{(t,t)}(\mathbb{C}^d)$ be the space of polynomials homogeneous of degree t both in the coordinates of vectors in \mathbb{C}^d and in their complex conjugates. An ensemble ν of pure state vectors in dimension d is a (complex projective) t -design if

$$\mathbb{E}_\nu p(\psi) = \int d\psi p(\psi) \quad \forall p \in \text{Hom}_{(t,t)}(\mathbb{C}^d),$$

where \mathbb{E}_ν denotes the expectation value over ν . The integral is taken with respect to the (normalized) uniform measure on the complex unit sphere in \mathbb{C}^d . Designs of unitary channels can be defined analogously. Let $\text{Hom}_{(t,t)}[\text{U}(d)]$ be the space of polynomials homogeneous of degree t both in the matrix elements of $U \in \text{U}(d)$ and in their complex conjugates. An

ensemble μ of unitary operators in dimension d is a “unitary t -design” if

$$\mathbb{E}_\mu p(U) = \int dU p(U) \quad \forall p \in \text{Hom}_{(t,t)}[\text{U}(d)],$$

where the integral is taken over the normalized Haar measure on $\text{U}(d)$.

Order- α entropies of a density operator ρ are entropic functions (which we call characteristic functions) of $\text{tr}\{\rho^\alpha\}$. A unified definition of such entropies is given by $S_s^{(\alpha)}(\rho) = \{1/[s(1-\alpha)]\}[(\text{tr}\{\rho^\alpha\})^s - 1]$, where s is a parameter that identifies the characteristic function and the family of entropies. The most representative families are Rényi (the limiting case $s \rightarrow 0$) and Tsallis ($s = 1$) entropies. In this Letter, we mostly focus on Rényi entropies

$$S_R^{(\alpha)}(\rho) = \frac{1}{1-\alpha} \log \text{tr}\{\rho^\alpha\},$$

with orders α being positive integers. In contrast to other generalized entropies, Rényi entropies have the following desirable properties, which make this family most relevant: (i) they are convex in $\text{tr}\{\rho^\alpha\}$, which makes it possible to use Jensen’s inequality to lower bound the design-averaged values by Haar integrals; (ii) they have the same roof value n for uniform spectrum for systems of n qubits, which allows meaningful comparisons with the maximum and between different orders; (iii) they are additive on product states (otherwise, it is not natural to define generalized quantities such as mutual information and tripartite information). When the order α increases, $S_R^{(\alpha)}$ becomes more and more sensitive to the nonuniformity in the spectrum: $S_R^{(\alpha_1)} \geq S_R^{(\alpha_2)}$, when $\alpha_1 < \alpha_2$. Taking the $\alpha \rightarrow \infty$ limit yields the min entropy

$$S_{\min}(\rho) = -\log \|\rho\| = -\log \lambda_{\max}(\rho),$$

where $\|\cdot\|$ denotes the operator norm and $\lambda_{\max}(\cdot)$ denotes the largest eigenvalue. The min entropy lower bounds all Rényi entropies.

Random states.—We first introduce results on random pure states. Consider a bipartite system with Hilbert space $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$, where \mathcal{H}_A and \mathcal{H}_B have dimensions d_A and d_B , respectively. The entanglement entropy between partitions A and B of a pure state $|\psi\rangle$ is given by the entropy of the reduced density operator $\rho_A = \text{tr}_B(|\psi\rangle\langle\psi|)$.

A key observation is that, given an α -design ν_α , we have $\mathbb{E}_{\nu_\alpha} \text{tr}\{\rho_A^\alpha\} = \int d\psi \text{tr}\{\rho_A^\alpha\}$, since $\text{tr}\{\rho_A^\alpha\}$ only involves $\text{Hom}_{(\alpha,\alpha)}$ terms of the entries of $|\psi\rangle$. Since the characteristic function for the Rényi- α entropy is convex, $\mathbb{E}_{\nu_\alpha} S_R^{(\alpha)}(\rho_A)$ is lower bounded by the characteristic function of the Haar integral $\int d\psi \text{tr}\{\rho_A^\alpha\}$ by Jensen’s inequality. Calculation shows that

$$\int d\psi \text{tr}\{\rho_A^\alpha\} = \frac{1}{\alpha! D_{[\alpha]}} \sum_{\sigma \in S_\alpha} d_A^{\xi(\sigma\tau)} d_B^{\xi(\sigma)}, \quad (1)$$

where $D_{[\alpha]} = \binom{d_A d_B + \alpha - 1}{\alpha}$ is the dimension of the symmetric subspace of $\mathcal{H}^{\otimes \alpha}$, S_α is the symmetric group of α symbols, $\xi(\sigma)$ is the number of disjoint cycles associated with σ [28], and $\tau := (1\ 2\ \dots\ \alpha)$ is the 1-shift (canonical full cycle). We noticed that similar results have been derived and rederived several times [12,29–32]. A simple derivation was presented in [23].

First, consider equal partitions $d_A = d_B$ and the limit of large dimension. Here we introduce the following cycle lemma (proof in [23], cf. [33]): for all $\sigma \in S_\alpha$, $\xi(\sigma\tau) + \xi(\sigma) \leq \alpha + 1$. Then Eq. (1) reduces to

$$\int d\psi \text{tr}\{\rho_A^\alpha\} = \text{Cat}_\alpha d_A^{-\alpha+1} + O(d_A^{-(\alpha+1)}), \quad (2)$$

where Cat_α is the α th Catalan number, satisfying $[(\log \text{Cat}_\alpha)/(\alpha - 1)] \leq 2$ for all $\alpha \geq 2$. So we obtain the following Theorem.

Theorem 1.—Let ν_α be a projective α -design. Consider equal partitions $d_A = d_B$. As $d_A \rightarrow \infty$,

$$\mathbb{E}_{\nu_\alpha} S_R^{(\alpha)}(\rho_A) \geq \log d_A - \frac{\log \text{Cat}_\alpha}{\alpha - 1} + O(d_A^{-2}). \quad (3)$$

So,

$$\mathbb{E}_{\nu_\alpha} S_R^{(\alpha)}(\rho_A) \geq \log d_A - O(1). \quad (4)$$

That is, the Rényi- α entanglement entropy across any cut averaged over an α -design is very close to (at most a constant away from) the maximum.

In fact, we are able to derive explicit bounds for finite dimensions and nonequal partitions.

Theorem 2.—Let ν_α be a projective α -design. Let $q := \alpha^3/(32d_B^2) < 1$, $h(q) := 1 + 2q/[3(1 - q)]$. For any $d_A \leq d_B$ and all α ,

$$\begin{aligned} \mathbb{E}_{\nu_\alpha} S_R^{(\alpha)}(\rho_A) &\geq \log d_A - \frac{2\alpha - \frac{3}{2}\log \alpha + \log h(q) - \frac{1}{2}\log \pi}{\alpha - 1} \\ &\geq \log d_A - 2. \end{aligned} \quad (5)$$

We also obtain the following bound, which improves the above result when d_A is small:

$$\begin{aligned} \mathbb{E}_{\nu_\alpha} S_R^{(\alpha)}(\rho_A) &\geq \log d_A - 2 \log \left(1 + \sqrt{\frac{d_A}{d_B}} \right) - \log c \\ &\geq \log d_A - 2\sqrt{\frac{d_A}{d_B}} - \log c, \end{aligned} \quad (6)$$

where $c = 1$ if \mathcal{H} is real and $c = 2$ if \mathcal{H} is complex.

Error bounds indicating that the above results are highly robust against small deviations from exact designs can be found in [23]. These results can be regarded as improved Page’s theorems that are tight in terms of the complexity.

Now we focus on the min entropy, given by $\alpha \rightarrow \infty$. Large min entropy implies that the spectrum is almost completely uniform. Are designs of infinite orders needed

to achieve almost maximal min entanglement entropy? The following result answers the question in the negative.

Theorem 3.—Let ν_α be a projective α -design, where $\alpha = \lceil (\log d_A)/a \rceil \leq (16d_B^2)^{1/3}$ with $0 < a \leq 1$. Then

$$\mathbb{E}_{\nu_\alpha} S_{\min}(\rho_A) \geq \log d_A - 2 - a. \quad (7)$$

In particular, $\mathbb{E}_{\nu_\alpha} S_{\min}(\rho_A) \geq \log d_A - 3$ if $\alpha = \lceil \log d_A \rceil$.

That is, $\Omega(\log d_A)$ -designs maximize all Rényi entanglement entropies and so are essentially indistinguishable from the Haar measure by the entanglement spectrum.

Conversely, one may wonder whether there exist α -designs such that Rényi entanglement entropies of orders larger than α are bounded away from the maximum, which we call “gap α -designs.” This indicates that they do not behave like designs of higher orders in a strong sense. Here we present an explicit example of gap 2-designs. Let $G = U_A \otimes U_B$, where U_A, U_B are the unitary groups on $\mathcal{H}_A, \mathcal{H}_B$, respectively. Calculation shows that the orbit of $|\psi\rangle$ under the action of G forms a 2-design if and only if $\text{tr}\{\rho_A^2\}$, with $\rho_A = \text{tr}_B(|\psi\rangle\langle\psi|)$, is equal to the average over the uniform ensemble, that is,

$$\text{tr}\{\rho_A^2\} = \frac{d_A + d_B}{d_A d_B + 1}. \quad (8)$$

The same conclusion still holds if U_A, U_B are replaced by subgroups that form unitary 2-designs on $\mathcal{H}_A, \mathcal{H}_B$, respectively. Equation (8) holds if ρ_A has the following spectrum

$$\begin{aligned} \lambda_1 &= \frac{d_A d_B + 1 + (d_A - 1)\sqrt{(d_A + 1)(d_A d_B + 1)}}{d_A(d_A d_B + 1)}, \\ \lambda_2 = \dots = \lambda_{d_A} &= \frac{d_A d_B + 1 - \sqrt{(d_A + 1)(d_A d_B + 1)}}{d_A(d_A d_B + 1)}. \end{aligned}$$

Suppose $d_B/d_A \leq r$, where r is a constant, then $\lambda_1 \geq (rd_A)^{-1/2}$, and so

$$S_R^{(\alpha)}(\rho_A) \leq \frac{1}{1 - \alpha} \log \lambda_1^\alpha \leq \frac{\alpha}{2(\alpha - 1)} (\log d_A + \log r). \quad (9)$$

As d_A increases, the gap of $S_R^{(\alpha)}(\rho_A)$ from the maximum is unbounded for all $\alpha > 2$.

Random unitary channels.—Now we extend the above analysis of pure states to the intrinsic entanglement properties of random unitary channels. The key results are similar in spirit to those for states, although the derivations are considerably more involved.

The Choi isomorphism (more generally, the channel-state duality) is widely used in quantum information theory to study quantum channels as states, by which a unitary operator U acting on a d -dimensional Hilbert space $U = \sum_{i,j=0}^{d-1} U_{ij}|i\rangle\langle j|$ is dual to the pure state

$$|U\rangle = \frac{1}{\sqrt{d}} \sum_{i,j=0}^{d-1} U_{ji}|i\rangle_{\text{in}} \otimes |j\rangle_{\text{out}},$$

which is called the Choi state of U . Consider bipartitions of the input register into A and B and the output register into C and D . Let d_A, d_B, d_C, d_D be the dimensions of subregions A, B, C, D , respectively ($d_A d_B = d_C d_D = d$). We study the entropy of ρ_{AC} with $\rho_{AC} = \text{tr}_{BD}(|U\rangle\langle U|)$. Consider the negative tripartite information

$$-I_3(A:C:D) := I(A:CD) - I(A:C) - I(A:D),$$

which is suggested in [7] to diagnose information scrambling, since it intuitively measures the delocalization of local information. Here $I(A:C) = S(A) + S(C) - S(AC)$ is the mutual information, which measures the total correlation between A and C . Since the input and output are maximally mixed due to unitarity, the four subregions are all maximally mixed. As a result, $-I_3$ is determined by the entanglement entropy $S(AC)$. Indeed, $-I_3$ essentially measures the ability of a channel to generate global entanglement that “hides” the delocalized information. Note that $-I_3$ can be reduced to the conditional mutual information $I(A:B|C)$ [34], which is of great interest in quantum information theory.

Let μ_α be a unitary α -design. By similar arguments involving the definition of unitary designs and the convexity of the Rényi characteristic function, the problem of bounding $\mathbb{E}_{\mu_\alpha} S_R^{(\alpha)}(\rho_{AC})$ boils down to computing the Haar integral $\int dU \text{tr}\{\rho_{AC}^\alpha\}$. In general, we find that

$$\int dU \text{tr}\{\rho_{AC}^\alpha\} = \frac{1}{d^\alpha} \sum_{\sigma, \gamma \in S_\alpha} d_A^{\xi(\sigma\tau)} d_B^{\xi(\sigma)} d_C^{\xi(\gamma\tau)} d_D^{\xi(\gamma)} \text{Wg}(d, \sigma\gamma^{-1}), \quad (10)$$

where

$$\text{Wg}(d, \sigma) = \frac{1}{(\alpha!)^2} \sum_{\lambda \vdash \alpha} \frac{\chi^\lambda(1)^2 \chi^\lambda(\sigma)}{s_{\lambda, d}(1, \dots, 1)}$$

are Weingarten functions of $U(d)$. Here $\lambda \vdash \alpha$ means λ is a partition of α , χ^λ is the corresponding character of S_α , and s_λ is the corresponding Schur function (polynomial). Notice that $s_{\lambda, d}(1, \dots, 1)$ is simply the dimension of the irreducible representation of $U(d)$ associated with λ . The Weingarten function can be derived by various tools in representation theory, such as Schur-Weyl duality [35,36] and Jucys-Murphy elements [37].

For equal partitions, in the limit of large dimension, we obtain the following analogous result by applying the cycle lemma.

Theorem 4.—Let μ_α be a unitary α -design. Consider equal partitions of the input and output registers, $d_A = d_B = d_C = d_D$. As $d \rightarrow \infty$,

$$\mathbb{E}_{\mu_\alpha} S_R^{(\alpha)}(\rho_{AC}) \geq \log d - \frac{\log \text{Cat}_\alpha}{\alpha - 1} + O(d^{-1}). \quad (11)$$

So,

$$\mathbb{E}_{\mu_\alpha} S_R^{(\alpha)}(\rho_{AC}) \geq \log d - O(1). \quad (12)$$

Therefore, the Rényi- α entanglement entropy between AC and BD (and the corresponding negative tripartite information based on the Rényi- α entropy) averaged over unitary α -designs is almost maximal.

We also provide more explicit bounds for finite dimensions and non-equal partitions.

Theorem 5.—Let μ_α be a unitary α -design. Suppose $d > \sqrt{6}\alpha^{7/4}$ and $d_A \leq d_B$. Then

$$\mathbb{E}_{\mu_\alpha} S_R^{(\alpha)}(\rho_{AC}) \geq \log d - \frac{\log \text{Cat}_\alpha}{\alpha - 1} - \frac{\log \left[\frac{a_\alpha h(q)}{8} (7 + \cosh \frac{2\alpha(\alpha-1)}{d}) \right]}{\alpha - 1}, \quad (13)$$

where $a_\alpha := \{1 - [(6\alpha^{7/2})/d^2]\}^{-1}$.

Similarly, these results do not deviate much for approximate unitary designs (see [23] for detailed error analysis).

The result on the min entropy is also similar.

Theorem 6.—Let μ_α be a unitary α -design, where $1 \leq \alpha = \lceil \log d/a \rceil \leq \sqrt{d}/2$ and $a > 0$; then

$$\mathbb{E}_{\mu_\alpha} S_{\min}(\rho_{AC}) \geq \log d - 2 - a. \quad (14)$$

In particular, $\mathbb{E}_{\mu_\alpha} S_{\min}(\rho_{AC}) \geq \log d - 3$ if $\alpha \geq \lceil \log d \rceil$.

Therefore, unitary $\Omega(\log d)$ -designs maximize all Rényi entanglement entropies.

Design complexities by Rényi.—In the above, we presented kinematic results revealing fundamental correspondences between Rényi entanglement entropies and quantum designs, which imply that states or unitaries sampled from α -designs typically exhibit nearly maximal Rényi- α entanglement entropy. This also suggests Rényi- α entanglement entropy as potential diagnostics of the randomness complexity of α -designs beyond information scrambling, in dynamical scenarios. Note that a recent work [38] generalizes the out-of-time-order correlators (which are widely used in the study of scrambling; see, e.g., [39–45]) to higher points and establishes similar connections to unitary designs via frame potentials.

In particular, the saturation of the min entanglement entropy indicates that the system looks completely random (and the local information is completely lost) to any local observer, which is the strongest form of scrambling that we call max-scrambling. How fast can physical systems achieve max-scrambling? The recent design Hamiltonian conjecture [21] argues, based on the original fast scrambling conjecture [3], that there exist physical dynamics (represented by local [46] and time-independent random Hamiltonians) that achieve approximate unitary α -designs in $O(\alpha \log n)$ time, where n is the number of qubits. Our result on logarithmic designs indicates that $O(n)$ -designs

are sufficient for max-scrambling and therefore suggests the following fast max-scrambling conjecture: max-scrambling can be achieved by physical dynamics in $O(n \log n)$ time.

Outlook.—The mathematical results of this Letter concern the average Rényi entanglement entropies of state and unitary designs. Some technical problems are left open. For example, we are not yet able to construct gap α -designs for $\alpha > 2$ and for unitaries. Moreover, due to the lack of subadditivity, the negative tripartite information $-I_3$ in terms of Rényi entropies is not necessarily positive. It is worth looking into when this situation occurs and further considering the meanings of such derived quantities. Also, the results here are about expected values. It would be nice to further analyze the variances or derive probabilistic bounds on concentration to talk about “typical” behaviors in a more rigorous sense.

Our results suggest Rényi entanglement entropies as powerful tools to further advance the study of quantum randomness and pseudorandomness. For example, a particularly interesting insight is that Rényi entropies of noninteger orders are naturally defined, which indicates that they can be helpful in understanding the mysterious but potentially important notion of noninteger designs. The results may find more applications in relevant areas in quantum information, such as entanglement theory, quantum complexity theory, quantum computing, and quantum cryptography.

The physical aspects are certainly worth further exploration. For example, it would be interesting to study the dynamical behaviors of Rényi entanglement entropies and randomness in specific many-body or holographic systems to learn about the physics in the postscrambling regime and extend existing studies of entanglement growth (e.g., “entanglement tsunami” [47,48]). A recent study [49] on (a 1D variant of) the strongly chaotic Sachdev-Ye-Kitaev model [50,51] (which has drawn considerable interest as a solvable toy model of quantum black holes and holography) shows that, after a quench, there is a “prethermal” regime where light modes rapidly scramble, but the Rényi entanglement entropies do not reach thermal values, which confirms our expectation that the randomness complexity of the system is still low; however, the late-time behaviors remain open. Moreover, it would be nice to extend the techniques and results of this work to the finite temperature regime or systems with conserved quantities, so as to apply our ideas in such physical scenarios and in general the study of quantum thermalization and many-body localization more carefully. We also hope to establish more solid connections between the randomness complexities and the conventional ones, such as computational, gate, and Kolmogorov complexities, which play active roles in recent studies of holographic duality and black holes [52–54] and are of independent interest. In general, the study of randomness complexities may also shed new light on the

fruitful idea of modeling complex systems (especially black holes [2]) by random states or dynamics. Further research along these lines are essential to our understanding of quantum chaos, quantum statistical mechanics, quantum many-body physics, and quantum gravity.

Z. W. L. and S. L. are supported by AFOSR and ARO. E. Y. Z. is supported by the National Science Foundation under Grant Contract No. CCF-1525130. Research at MIT CTP is supported by the DOE. H. Z. is supported by the Excellence Initiative of the German Federal and State Governments (ZUK 81) and the DFG.

*zwliu@mit.edu

- [1] D. N. Page, *Phys. Rev. Lett.* **71**, 1291 (1993).
- [2] P. Hayden and J. Preskill, *J. High Energy Phys.* **09** (2007) 120.
- [3] Y. Sekino and L. Susskind, *J. High Energy Phys.* **10** (2008) 065.
- [4] L. Susskind, [arXiv:1101.6048](https://arxiv.org/abs/1101.6048).
- [5] K. Furuya, M. C. Nemes, and G. Q. Pellegrino, *Phys. Rev. Lett.* **80**, 5524 (1998).
- [6] A. Lakshminarayan, *Phys. Rev. E* **64**, 036207 (2001).
- [7] P. Hosur, X.-L. Qi, D. A. Roberts, and B. Yoshida, *J. High Energy Phys.* **02** (2016) 4.
- [8] R. Nandkishore and D. A. Huse, *Annu. Rev. Condens. Matter Phys.* **6**, 15 (2015).
- [9] S. Popescu, A. J. Short, and A. Winter, *Nat. Phys.* **2**, 754 (2006).
- [10] B. M. Terhal, D. P. DiVincenzo, and D. W. Leung, *Phys. Rev. Lett.* **86**, 5807 (2001).
- [11] D. P. DiVincenzo, D. W. Leung, and B. M. Terhal, *IEEE Trans. Inf. Theory* **48**, 580 (2002).
- [12] E. Lubkin, *J. Math. Phys.* **19**, 1028 (1978).
- [13] S. Lloyd and H. Pagels, *Ann. Phys. (N.Y.)* **188**, 186 (1988).
- [14] S. K. Foong and S. Kanno, *Phys. Rev. Lett.* **72**, 1148 (1994).
- [15] J. Sánchez-Ruiz, *Phys. Rev. E* **52**, 5653 (1995).
- [16] S. Sen, *Phys. Rev. Lett.* **77**, 1 (1996).
- [17] E. Knill, [arXiv:quant-ph/9508006](https://arxiv.org/abs/quant-ph/9508006).
- [18] F. G. S. L. Brandao, A. W. Harrow, and M. Horodecki, *Commun. Math. Phys.* **346**, 397 (2016).
- [19] F. G. S. L. Brandão, A. W. Harrow, and M. Horodecki, *Phys. Rev. Lett.* **116**, 170502 (2016).
- [20] G. Tóth and J. J. García-Ripoll, *Phys. Rev. A* **75**, 042311 (2007).
- [21] Y. Nakata, C. Hirche, M. Koashi, and A. Winter, *Phys. Rev. X* **7**, 021006 (2017).
- [22] N. Lashkari, D. Stanford, M. Hastings, T. Osborne, and P. Hayden, *J. High Energy Phys.* **04** (2013) 22.
- [23] Z.-W. Liu, S. Lloyd, E. Y. Zhu, and H. Zhu, [arXiv:1703.08104](https://arxiv.org/abs/1703.08104).
- [24] E. Magesan, J. M. Gambetta, and J. Emerson, *Phys. Rev. Lett.* **106**, 180504 (2011).
- [25] J. J. Wallman and S. T. Flammia, *New J. Phys.* **16**, 103032 (2014).
- [26] O. Szehr, F. Dupuis, M. Tomamichel, and R. Renner, *New J. Phys.* **15**, 053022 (2013).

- [27] H. Zhu, R. Kueng, M. Grassl, and D. Gross, [arXiv:1609.08172](#).
- [28] Every element of the symmetric group can be uniquely decomposed into a product of disjoint cycles (up to relabeling).
- [29] K. Życzkowski and H.-J. Sommers, *J. Phys. A* **34**, 7111 (2001).
- [30] L. C. Malacarne, R. S. Mendes, and E. K. Lenzi, *Phys. Rev. E* **65**, 046131 (2002).
- [31] B. Collins and I. Nechita, *Commun. Math. Phys.* **297**, 345 (2010).
- [32] B. Collins and I. Nechita, *Ann. Appl. Probab.* **21**, 1136 (2011).
- [33] A. Nica and R. Speicher, *Lectures on the Combinatorics of Free Probability* (Cambridge University Press, Cambridge, England, 2006).
- [34] D. Ding, P. Hayden, and M. Walter, *J. High Energy Phys.* **12** (2016) 145.
- [35] B. Collins, *Int. Math. Res. Not.* **2003**, 953 (2003).
- [36] B. Collins and S. Matsumoto, *J. Math. Phys.* **50**, 113516 (2009).
- [37] P. Zinn-Justin, *Lett. Math. Phys.* **91**, 119 (2010).
- [38] D. A. Roberts and B. Yoshida, *J. High Energy Phys.* **04** (2017) 121.
- [39] S. H. Shenker and D. Stanford, *J. High Energy Phys.* **03** (2014) 67.
- [40] S. H. Shenker and D. Stanford, *J. High Energy Phys.* **12** (2014) 46.
- [41] S. H. Shenker and D. Stanford, *J. High Energy Phys.* **05** (2015) 132.
- [42] D. A. Roberts, D. Stanford, and L. Susskind, *J. High Energy Phys.* **03** (2015) 51.
- [43] D. A. Roberts and D. Stanford, *Phys. Rev. Lett.* **115**, 131603 (2015).
- [44] J. Maldacena, S. H. Shenker, and D. Stanford, *J. High Energy Phys.* **08** (2016) 106.
- [45] I. Kukuljan, S. Grozdanov, and T. Prosen, *Phys. Rev. B* **96**, 060301 (2017).
- [46] Here “local” should mean all interaction terms involve at most k qubits where k is some constant (k -local), rather than spatially local on a finite-dimensional graph. This was not made very clear in [21].
- [47] H. Liu and S. J. Suh, *Phys. Rev. Lett.* **112**, 011601 (2014).
- [48] H. Liu and S. J. Suh, *Phys. Rev. D* **89**, 066012 (2014).
- [49] Y. Gu, A. Lucas, and X.-L. Qi, *J. High Energy Phys.* **09** (2017) 120.
- [50] S. Sachdev and J. Ye, *Phys. Rev. Lett.* **70**, 3339 (1993).
- [51] A. Kitaev (2015), <http://online.kitp.ucsb.edu/online/entangled15/kitaev/>; <http://online.kitp.ucsb.edu/online/entangled15/kitaev2/>.
- [52] A. R. Brown, D. A. Roberts, L. Susskind, B. Swingle, and Y. Zhao, *Phys. Rev. D* **93**, 086006 (2016).
- [53] A. R. Brown, D. A. Roberts, L. Susskind, B. Swingle, and Y. Zhao, *Phys. Rev. Lett.* **116**, 191301 (2016).
- [54] A. R. Brown and L. Susskind, [arXiv:1701.01107](#).