

Nontopological Majorana Zero Modes in Inhomogeneous Spin Ladders

Neil J. Robinson,^{1,2,*} Alexander Altland,³ Reinhold Egger,⁴ Niklas M. Gergs,⁵ Wei Li,^{6,7} Dirk Schuricht,⁵
Alexei M. Tsvelik,² Andreas Weichselbaum,^{2,7} and Robert M. Konik^{2,†}

¹*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

²*CMPMS Division, Brookhaven National Laboratory, Upton, New York 11973, USA*

³*Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Strasse 77, D-50937 Köln, Germany*

⁴*Institut für Theoretische Physik, Heinrich-Heine-Universität, D-40225 Düsseldorf, Germany*

⁵*Institute for Theoretical Physics, Center for Extreme Matter and Emergent Phenomena, Utrecht University, Leuvenlaan 4, 3584 CE Utrecht, The Netherlands*

⁶*Department of Physics, International Research Institute of Multidisciplinary Science, Beihang University, Beijing 100191, China*

⁷*Physics Department, Arnold Sommerfeld Center for Theoretical Physics, and Center for NanoScience, Ludwig-Maximilians-Universität, 80333 Munich, Germany*



(Received 26 June 2018; revised manuscript received 10 October 2018; published 15 January 2019)

We show that the coupling of homogeneous Heisenberg spin-1/2 ladders in different phases leads to the formation of interfacial zero energy Majorana bound states. Unlike Majorana bound states at the interfaces of topological quantum wires, these states are void of topological protection and generally susceptible to local perturbations of the host spin system. However, a key message of our Letter is that, in practice, they show a high degree of resilience over wide parameter ranges which may make them interesting candidates for applications.

DOI: [10.1103/PhysRevLett.122.027201](https://doi.org/10.1103/PhysRevLett.122.027201)

Introduction.—The Majorana fermion has become one of the most important fundamental quasiparticles of condensed matter physics. Besides its key role as a building block in correlated quantum matter, much of this interest is motivated by perspectives in quantum information [1–3]. Majorana qubits have unique properties which make them ideal candidates for applications in, e.g., stabilizer code quantum computation [4]. Current experimental attempts to isolate and manipulate Majorana bound states (MBSs) focus on interfaces between distinct phases of symmetry protected topological (SPT) quantum matter. These material platforms have the appealing property that MBSs are protected against local perturbations by principles of topology. In practice, however, topological protection may play a lesser role than one might hope, and various obtrusive aspects of realistic quantum materials appear to challenge the isolation and manipulation of MBSs. Specifically, in topological quantum wires based on the hybrid semiconductor-superconductor platform [5] or on coupled ferromagnetic atoms [6], all relevant scales are confined to narrow windows in energy. In this regard, proposals to realize MBSs in topological insulator nanowires [7] may offer superior solutions. However, these realizations require a high level of tuning of external parameters, notably of magnetic fields, and may be met with their own difficulties.

In this Letter, we suggest an alternative hardware platform for the isolation of zero-energy MBSs. Our proposal does not engage topology. Specifically, local perturbations

of the microscopic Hamiltonian may induce nonlocal correlations between the emergent Majorana quantum particles. However, we argue below that in practice this problem is less drastic than one might fear and that the current architecture may grant a high level of effective protection. The numerical evidence provided below certainly points in this direction.

The material platform we suggest is based on spin ladder materials. Their phases can be classified by combining standard Landau-Ginzburg symmetry breaking with the presence of SPT order [8,9]. Here, we show that combining ladders in different phases provides a systematic means to generating interface MBSs. The formal bridge between the physics of spin ladders and that of Majorana fermions is provided by a two-step mapping, first, representing the spin degrees of freedom by bosons, followed by refermionization of the latter into an effective Majorana theory [10]. We will discuss how numerous spin ladder properties that are difficult to access in the spin language are made simple and transparent in the Majorana representation. In particular, SU(2) invariant spin ladders with two legs are described by a theory of four massive Majorana fermions, comprising a triplet and a singlet of different masses, together with a global parity constraint. The ground state degeneracies of the spin systems are then encoded entirely in zero-energy MBSs localized on the boundaries of the system.

Two surprising findings arise from this Majorana representation. The first is that additional ground state degeneracies can appear in *inhomogeneous* ladders, where the

spin-spin interactions vary spatially along the ladder. In the fermionic language, these degeneracies manifest themselves in new MBSs appearing at the phase boundaries via the Jackiw-Rebbi mechanism [11], according to which a sign change in the fermion mass creates a zero mode. This may happen even if all of the bulk phases composing the ladder do not support MBSs on their own. The second finding is that zero-energy MBSs exist only if the spatial variation of spin couplings about the boundary is sufficiently gentle (a few lattice sites, in practice). The spatial smoothness across the interface is required to stabilize the mapping onto a continuum description and to prevent the coupling of distant MBSs via higher-energy states [12]. This condition manifests the lack of topological protection. (For other zero energy modes in topologically trivial phases, see Refs. [13–17].) However, we present numerical evidence that these MBSs are, nonetheless, close to zero energy over parametrically wide regions.

Spin ladders.—Ladder geometries provide an important viewpoint on the physics of strongly correlated electron systems [18] and are a research focus of condensed matter physics in their own regard. They are close enough to being one dimensional (1D) that powerful theoretical techniques can be deployed in their understanding, running the gamut from field theory [10,19–21] and Bethe ansatz [22] to density matrix renormalization group (DMRG) [23–28]. However, they are also far enough removed from 1D that they capture the physics of two-dimensional systems. Here, we focus on ladders where the fluctuations of spin-1/2 degrees of freedom are dominant (e.g., SrCu₂O₃ [29]) over ladders where charge degrees are mobile (e.g., Sr_{14-x}Ca_xCu₂₄O₄₁ [30]). For concreteness, we consider the two-leg ladder Hamiltonian

$$\begin{aligned}
 H = & J \sum_{\ell=1,2} \sum_{r=1}^{N-1} \mathbf{S}_{\ell,r} \cdot \mathbf{S}_{\ell,r+1} + J_{\perp} \sum_{r=1}^N \mathbf{S}_{1,r} \cdot \mathbf{S}_{2,r} \\
 & + J_{\times} \sum_{r=1}^{N-1} (\mathbf{S}_{1,r} \cdot \mathbf{S}_{1,r+1})(\mathbf{S}_{2,r} \cdot \mathbf{S}_{2,r+1}), \quad (1)
 \end{aligned}$$

where $S_{\ell,r}^a$ is the $a = x, y, z$ spin-1/2 operator located on leg ℓ and rung r of the ladder. The exchange parameters $J := 1, J_{\perp}, J_{\times}$ characterize leg, rung, and plaquette interactions, respectively. For uncoupled Heisenberg chains, the total spin of each leg would be conserved, and we could work in a representation where $S_{\ell}^z = \sum_r S_{\ell,r}^z$ are good quantum numbers. Assuming an even number N of sites per chain, both $S_{\ell}^z \in \mathbb{Z}$ are integer valued. The coupling J_{\perp} exchanges spin in integer units, $S_1^z \rightarrow S_1^z \pm 1, S_2^z \rightarrow S_2^z \mp 1$, violating the conservation of the individual S_{ℓ}^z , but still constraining the even and odd combinations, $S_{\pm}^z = S_1^z \pm S_2^z$, to have identical parity,

$$S_{+}^z \equiv S_{-}^z \pmod{2}. \quad (2)$$




FIG. 1. Phase diagram for the Hamiltonian (1) obtained from SU(2) DMRG simulations of a 100×2 site ladder with bond dimension $\chi = 1500$ states. The red (blue) phase boundary shows the critical line with Majorana fermion mass $m_t = 0$ ($m_s = 0$). Inset figures show schematic representations of singlet (blue) and triplet (red) bond order within each phase, and the corresponding signs of m_t and m_s .

Thus, we expect an effective fermionized theory of the system to display a U(1) symmetry reflecting the conservation of S_{\pm}^z plus a \mathbb{Z}_2 parity condition implementing (2). The latter introduces correlations between the S_{\pm}^z and the S_{\mp}^z sectors and will play a key role throughout.

Phase diagram.—Depending on the couplings J_{\perp}, J_{\times} , the Hamiltonian (1) supports different phases. For strong positive rung interaction J_{\perp} and weak plaquette interaction J_{\times} , the formation of rung singlets (RS) is favored, cf. the lower right part of Fig. 1. For strong negative couplings J_{\perp} , rung triplets are formed instead, and effectively implement an $S = 1$ Haldane-Heisenberg chain (Haldane phase, H). For strong J_{\times} , one may anticipate “valence bond solids” (VBSs) distinguished by different types of periodically repeated intrachain dimerization, VBS₊ and VBS₋ (see Fig. 1). While the existence of different dimerization patterns is relatively easy to anticipate, it takes more effort to determine the symmetries characterizing them, the respective order parameters, the ground state degeneracies, and the phase boundaries. For example, the Haldane phase is an SPT phase without a local order parameter. It exhibits a fourfold ground state degeneracy due to two spin-1/2 degrees of freedom dangling at the boundaries. In particular, the identification of the symmetries of the VBS phases is a nontrivial matter [31,32]. The boundaries between the phases as well as the ensuing ground state degeneracies can be established via DMRG simulations [12]: in Fig. 1, we present the phase diagram and, in Table I, the ground state degeneracies.

The presence of distinct dimerization patterns also provides a first clue as to the formation of zero energy degrees of freedom if chains of competing order are coupled by interfaces of sufficiently smooth variation. As an example, consider the RS–VBS₋–RS setup in Fig. 2. The VBS₋ chain breaks a translational \mathbb{Z}_2 symmetry

TABLE I. Phases of the spin model, the signs of their fermion masses, m_t/m_s , the ground state degeneracies, d_{\pm} , of their even or odd sectors (S_{\pm}^z) before the parity restriction (2) is applied, and, finally, their overall actual ground state degeneracies from SU(2) DMRG.

Phase	m_t/m_s	d_+/d_-	Ground state degeneracies
H	+/-	4/2	4
RS	-/+	1/2	1
VBS ₊	+/+	4/4	8
VBS ₋	-/-	1/1	1

via the choice of the links harboring singlet configurations (indicated as blue ovals). If the interface is sharp, one such configuration is rigidly pinned between two RS phases, and the ground state is unique. However, for a smooth interface, dimerization patterns of either parity can be put at no difference in energy (cf. the bottom part of the figure). This leads to a \mathbb{Z}_2 ground state degeneracy between phases whose ground states are individually nondegenerate.

Majorana representation.—All the structures and phenomena alluded to above afford a simple and surprisingly quantitative description in a language of Majorana fermions. The passage to this representation involves the Abelian bosonization [33] of the spin ladder as an intermediate step. In a second step, the bosonic degrees of freedom are mapped to an equivalent system of Majorana fermions [10]. Within the bosonized framework, smooth and rapid changes of the spin magnetization in the interaction terms are represented as gradient (“current-current”) and transcendental (“massive”) perturbations of the boson fields, respectively [12]. Within the fermion language, these, in turn, correspond to interaction terms and bilinear fermion operators, where, crucially, the former turn out to be irrelevant in a renormalization group sense. This means that, perhaps counterintuitively, the spin ladder is represented by a system of two noninteracting fermion fields, representing the sum and the difference S_{\pm} of the magnetization, respectively. The fermion bilinears describe scattering between left and right moving fermions together with effectively superconducting correlations in the S_{\pm}^z sector reflecting the absence of U(1) symmetry. Much as for the case of topological superconducting wires [2], it then pays off to switch to a Majorana fermion representation. As a result, one arrives at the low-energy continuum Hamiltonian

$$H = \int dx \left(-\frac{iv}{2} (\xi_R^0 \partial_x \xi_R^0 - \xi_L^0 \partial_x \xi_L^0) - im_s \xi_R^0 \xi_L^0 - \frac{iv}{2} (\xi_R \partial_x \xi_R - \xi_L \partial_x \xi_L) - im_t \xi_R \cdot \xi_L \right), \quad (3)$$

where $\xi^{0,1,2,3}$ are Majorana fields arranged into a singlet, ξ^0 , and a triplet, $\xi = (\xi^1, \xi^2, \xi^3)$, subject to masses [10]

$$m_t \propto 9J_{\times}/\pi^2 - J_{\perp}, \quad m_s \propto 9J_{\times}/\pi^2 + 3J_{\perp}. \quad (4)$$

The doublets (ξ^1, ξ^2) and (ξ^0, ξ^3) represent the S_{+}^z and S_{-}^z sectors, respectively. In the Majorana language, the U(1) \simeq O(2) symmetry of the S_{+}^z sector is realized as a continuous rotation symmetry between the mass-degenerate fields (ξ^1, ξ^2) , and the \mathbb{Z}_2 symmetry of the S_{-}^z sector via sign inversion of $\xi^{0,3}$. Importantly, these Majorana fields are not independent but are correlated via the spin parity relation (2). In the present language, the global S_{\pm}^z quantum numbers assume the form $S_{+}^z = i \sum_a \xi_a^2 \xi_a^1 / 2$ and $S_{-}^z = i \sum_b \xi_b^3 \xi_b^0 / 2$, where $\sum_{a,b}$ is a formal sum over all eigenmodes of the system. (In translational invariant cases, these are momentum modes. However, for systems with boundaries or interfaces, the situation becomes more interesting.) Thus, the constraint (2) translates to

$$\exp\left(\pi \sum_a \xi_a^1 \xi_a^2 / 2\right) = \exp\left(\pi \sum_b \xi_b^3 \xi_b^0 / 2\right), \quad (5)$$

introducing entanglement between the four Majorana sectors [12].

Interfacial Majorana states.—In the Majorana representation, the ground state degeneracy of a phase is diagnosed via the appearance of MBSs localized at the system’s boundaries. Here, the vacuum can be represented as a fictitious Majorana system with an infinitely large negative mass [34]. A vacuum interface of a system with a bulk positive mass then amounts to the zero crossing of a spatially dependent mass function $m(x)$, where the Jackiw-Rebbi mechanism implies the presence of a zero-energy MBS at each end. Since two MBSs define a fermion Hilbert space of dimension two, prior to imposing the parity constraint (5), the ground state degeneracy of a system of definite (J_{\times}, J_{\perp}) is given by $d = d_+ d_-$, $d_+ = 2^{2\Theta(m_t)}$, $d_- = 2^{\Theta(m_t) + \Theta(m_s)}$, where Θ is the Heaviside function, and we use Eq. (4). For $d > 1$, (5) then implies a factor of 2 reduction in the actually realized ground state degeneracy, $d \rightarrow d/2$. This integer agrees exactly with the DMRG results listed in Table I. The same ground state degeneracies also follow from the bosonized formulation (Sec. II of Ref. [12]) from a truncated conformal space approach [35,36] for sine-Gordon-like models [37–51].

What happens at interfaces between ladders of different symmetry can now be understood in equally straightforward ways. Let us, then, return to the RS–VBS₋–RS hybrid, see Fig. 2. Provided the interface varies smoothly enough, the system is described by the Majorana theory with $m_t < 0$ but with m_s changing from positive values to negative and back. Thus, we have MBSs at both interfaces with spatial extension determined by the width of the interface region. Naively, one might think that the same principle secures the existence of MBSs in the complementary case of VBS₋–RS–VBS₋ hybrids as well.




FIG. 2. Bond formation patterns when parameters (J_{\times}, J_{\perp}) are varied (in green) in order to form a RS–VBS_––RS ladder. (a) For sudden parameter changes at interfaces. (b) For smooth parameter changes, Jackiw-Rebbi zero modes emerge when the singlet mass, m_s , changes sign (lowest sketch).

However, there is a catch: The above argument does not make reference to the parity constraint (5). In the RS–VBS_––RS case, since $m_s > 0$ in the outer RS segments, MBSs will not only exist at the internal interfaces but also at the outer vacuum boundaries, cf. Fig. 2(b). This implies that changes in the occupation of the internal MBS system can be compensated by the outer MBS system, which may act as a “parity sink” to restore the condition (5). In concrete terms, the + sector of the RS–VBS_––RS ladder is even parity and has a unique ground state as $d_+ = 1$ for the RS and VBS_– segments. On the other hand, the – sector is nominally fourfold degenerate (as $d_- = 2$ for each RS segment), but only two of the four states have even parity, thus, leaving only two allowed states once we combine the \pm sectors.

In Fig. 3, we present DMRG results showing that the RS–VBS_––RS ladder, indeed, has a doubly degenerate ground state for smooth interfaces. If J_{\perp} and J_{\times} defining these phases vary too sharply, the ground state remains unique. We explain why this is so field theoretically in Sec. III of [12]. However, once the scale of variation extends over just a few lattice sites, one rapidly approaches a twofold degenerate ground state. We also note that the energy gap protecting the ground state degeneracy is rather large for the example in Fig. 3. It is remarkable that MBSs are generated in the RS–VBS_––RS example, where none of the individual parts, VBS_– or RS, support such states. These MBSs also provide a means to distinguish two different SPT-trivial phases, cf. Refs. [31,32]. The situation is rather different for the VBS_––RS–VBS_– system. Since one of the two fermion states formed from the central MBS pair is parity blocked, MBSs are effectively removed from the zero energy Hilbert space [52]. See Sec. IV. B. 2 of [12] for verification of this via DMRG. In this way, the parity constraint trumps the Jackiw-Rebbi principle.

Interfaces between phases of enriched symmetry define higher-dimensional MBS systems. As an example, consider




FIG. 3. Finite size scaling of low-energy SU(2) DMRG eigenstates in RS–VBS_––RS ladders of total length L . Blue and red indicate singlet and triplet states ($S = 0/1$) (the ground state is not shown). We use $J = 1$, with the other couplings varied as $J_{\perp}(x) = \frac{4}{3}[1 - w(x)]$ and $J_{\times}(x) = -\frac{4}{3}w(x)$, with $w(x) = f(x - x_+) - f(x - x_-)$, $f(x) = [1 + \exp(x/\delta x)]^{-1}$, $L_c \equiv x_+ - x_-$, and $x_{\pm} = (L \pm L_c)/2$. The width δx controls the interface smoothness. The ground state degeneracy develops quickly with increasing δx and is only marginally affected by the length L_c of the VBS_– region (see Sec. IV of [12]).

the RS– H –RS hybrid. Although the ground state degeneracy of the central H segment (the outer RS phases) is only fourfold (unique), the interfaces harbor a potential 32D zero-energy space, with four MBSs at either side of the H segment since four masses change sign upon crossing from one phase into the other. Parity, as in the RS–VBS_––RS ladder, reduces this by one-half (see Sec. IV. B. 3 of [12]).

Reality check.—The above constructions demonstrate that spin ladder materials provide a remarkably rich platform for the isolation of zero energy MBSs, with sizeable energy gaps to higher-lying states. In view of the general interest in MBSs, it is imperative to ask how our non-topological MBSs fare in comparison to topologically protected MBSs. At first sight, the absence of topological protection appears to be a crucial setback. However, at present, the probably most obtrusive effect hampering Majorana device functionality is the buildup of long-range MBS hybridizations. In topological devices, the hybridization exponentially approaches zero with increasing distance but can, nonetheless, be large in practice. For example, in hybrid semiconductor wires, topological protection crucially relies on the rather tiny superconducting proximity gaps [2,53–56]. In the present setup, the lack of topological protection manifests itself in long-range correlations between MBSs when short range correlations of the underlying spin chains are changed (in particular, the interface roughness). However, the degrees of freedom behind such changes are highly inert in realistic systems since they require energy scales comparable to the exchange couplings. Even though these energy scales do not grow with system size, they can be sufficiently large to provide efficient MBS protection at low temperatures.

Outlook.—A promising aspect of our approach is that it brings a plethora of material platforms into play. While we

have focused on spin ladders, similar considerations apply to many quasi-1D materials, in particular, those that admit a bosonization treatment, e.g., N -leg Heisenberg ladders with $SU(2)$ spin symmetry [18,27,57] or a more general $SU(M)$ symmetry [19,28,58], coupled chains of itinerant electrons [21,59–61], or coupled Luttinger liquid systems [62,63]. In addition, our setup directly comes with an intrinsic source of strong entanglement. Indeed, the Majorana parity constraint (5) plays a similar role to the strong Coulomb charging energy [64–67] in mesoscopic MBS systems, where a related parity constraint implies qubit functionality [68,69]. The question of how this entanglement mechanism may be turned into an operational resource, and how the MBSs discussed here can be probed and/or manipulated, is an interesting subject for future study.

N. J. R. thanks F. Burnell, F. Harper, and D. Schimmel for discussions. Work at BNL (N. J. R., A. M. T., A. W., R. M. K.) was supported by the CMPMS Division funded by the U.S. Department of Energy, Office of Basic Energy Sciences, under Contract No. DE-SC0012704. N. J. R. was supported by the EU Horizon 2020 program, Grant Agreement No. 745944. A. W. acknowledges support in the initial states of this project from the Deutsche Forschungsgemeinschaft (DFG), Grants No. WE4819/2-1 and No. WE4819/3-1. W. L. was funded by NSFC under No. 11504014 and No. 11834014. D. S. is member of the D-ITP consortium of the Netherlands Organisation for Scientific Research. A. A. and R. E. acknowledge DFG support via Grants No. EG 96/11-1 and No. CRC TR 183 (Project C4). R. M. K. and A. M. T. acknowledge the hospitality of LMU Munich and of HHU Düsseldorf where parts of this work have been done.

*n.j.robinson@uva.nl

†rmk@bnl.gov

- [1] S. R. Elliott and M. Franz, Colloquium: Majorana fermions in nuclear, particle, and solid-state physics, *Rev. Mod. Phys.* **87**, 137 (2015).
- [2] J. Alicea, New directions in the pursuit of Majorana fermions in solid state systems, *Rep. Prog. Phys.* **75**, 076501 (2012).
- [3] J. Alicea, Y. Oreg, G. Refael, F. von Oppen, and M. P. A. Fisher, Non-Abelian statistics and topological quantum information processing in 1D wire networks, *Nat. Phys.* **7**, 412 (2011).
- [4] S. Das Sarma, M. Freedman, and C. Nayak, Majorana zero modes and topological quantum computation, *npj Quantum Inf.* **1**, 15001 (2015).
- [5] R. M. Lutchyn, E. P. A. M. Bakkers, L. P. Kouwenhoven, P. Krogstrup, C. M. Marcus, and Y. Oreg, Majorana zero modes in superconductor semiconductor heterostructures, *Nat. Rev. Mater.* **3**, 52 (2018).
- [6] S. Nadj-Perge, I. K. Drozdov, J. Li, H. Chen, S. Jeon, J. Seo, A. H. MacDonald, B. A. Bernevig, and A. Yazdani, Observation of Majorana fermions in ferromagnetic atom chains on a superconductor, *Science* **346**, 602 (2014).
- [7] A. Cook and M. Franz, Majorana fermions in a topological-insulator nanowire proximity-coupled to an s -wave superconductor, *Phys. Rev. B* **84**, 201105 (2011).
- [8] Z.-C. Gu and X.-G. Wen, Tensor-entanglement-filtering renormalization approach and symmetry-protected topological order, *Phys. Rev. B* **80**, 155131 (2009); X. Chen, Z.-C. Gu, and X.-G. Wen, Classification of gapped symmetric phases in one-dimensional spin systems, *Phys. Rev. B* **83**, 035107 (2011); Complete classification of one-dimensional gapped quantum phases in interacting spin systems, *Phys. Rev. B* **84**, 235128 (2011); F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa, Symmetry protection of topological phases in one-dimensional quantum spin systems, *Phys. Rev. B* **85**, 075125 (2012).
- [9] N. Schuch, D. Pérez-García, and I. Cirac, Classifying quantum phases using matrix product states and projected entangled pair states, *Phys. Rev. B* **84**, 165139 (2011).
- [10] D. G. Shelton, A. A. Nersisyan, and A. M. Tsvelik, Antiferromagnetic spin ladders: Crossover between spin $S = 1/2$ and $S = 1$ chains, *Phys. Rev. B* **53**, 8521 (1996).
- [11] R. Jackiw and C. Rebbi, Solitons with fermion number $1/2$, *Phys. Rev. D* **13**, 3398 (1976).
- [12] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.122.027201> for additional details on the bosonized theory and on DMRG simulations, an in-depth explanation of why sharp interfaces will gap out MBSs, and alternative computations of the ground state degeneracies within the bosonized formalism.
- [13] Z. Yan, R. Bi, and Z. Wang, Majorana Zero Modes Protected by a Hopf Invariant in Topologically Trivial Superconductors, *Phys. Rev. Lett.* **118**, 147003 (2017).
- [14] C. Chan, L. Zhang, T. F. J. Poon, Y.-P. He, Y.-Q. Wang, and X.-J. Liu, Generic Theory for Majorana Zero Modes in 2D Superconductors, *Phys. Rev. Lett.* **119**, 047001 (2017).
- [15] Z. Yan, F. Song, and Z. Wang, Majorana Corner Modes in a High-Temperature Platform, *Phys. Rev. Lett.* **121**, 096803 (2018).
- [16] T. H. Hsieh, H. Ishizuka, L. Balents, and T. L. Hughes, Bulk Topological Proximity Effect, *Phys. Rev. Lett.* **116**, 086802 (2016); T. H. Hsieh, Y.-M. Lu, and A. W. W. Ludwig, Topological bootstrap: Fractionalization from Kondo coupling, *Sci. Adv.* **3**, e1700729 (2017).
- [17] V. Kaladzhyan, C. Bena, and P. Simon, Topology from triviality, *Phys. Rev. B* **97**, 104512 (2018).
- [18] E. Dagotto and T. M. Rice, Surprises on the way from one- to two-dimensional quantum magnets: The ladder materials, *Science* **271**, 618 (1996).
- [19] P. Lecheminant and A. M. Tsvelik, Two-leg $SU(2n)$ spin ladder: A low-energy effective field theory approach, *Phys. Rev. B* **91**, 174407 (2015).
- [20] R. Konik and A. W. W. Ludwig, Exact zero-temperature correlation functions for two-leg Hubbard ladders and carbon nanotubes, *Phys. Rev. B* **64**, 155112 (2001).
- [21] R. M. Konik, T. M. Rice, and A. M. Tsvelik, Doped Spin Liquid: Luttinger Sum Rule and Low Temperature Order, *Phys. Rev. Lett.* **96**, 086407 (2006).
- [22] Y. Wang, Exact solution of a spin-ladder model, *Phys. Rev. B* **60**, 9236 (1999); M. T. Batchelor and M. Maslen, Exactly solvable quantum spin tubes and ladders, *J. Phys. A* **32**, L377 (1999).

- [23] S. R. White, Density Matrix Formulation for Quantum Renormalization Groups, *Phys. Rev. Lett.* **69**, 2863 (1992).
- [24] U. Schollwöck, The density-matrix renormalization group, *Rev. Mod. Phys.* **77**, 259 (2005).
- [25] U. Schollwöck, The density-matrix renormalization group in the age of matrix product states, *Ann. Phys. (Amsterdam)* **326**, 96 (2011).
- [26] R. M. Noack, S. R. White, and D. J. Scalapino, The ground state of the two-leg Hubbard ladder a density-matrix renormalization group study, *Physica (Amsterdam)* **270C**, 281 (1996).
- [27] F. B. Ramos and J. C. Xavier, N -leg spin- S Heisenberg ladders: A density-matrix renormalization group study, *Phys. Rev. B* **89**, 094424 (2014).
- [28] A. Weichselbaum, S. Capponi, P. Lecheminant, A. M. Tsvelik, and A. M. Läuchli, Unified phase diagram of antiferromagnetic SU(N) spin ladders, *Phys. Rev. B* **98**, 085104 (2018).
- [29] M. Azuma, Z. Hiroi, M. Takano, K. Ishida, and Y. Kitaoka, Observation of a Spin Gap in SrCu₂O₃ Comprising Spin-1/2 Quasi-1D Two-Leg Ladders, *Phys. Rev. Lett.* **73**, 3463 (1994).
- [30] T. Vuletić, B. Korin-Hamzić, T. Ivek, S. Tomić, B. Gorshunov, M. Dressel, and J. Akimitsu, The spin-ladder and spin-chain system (La, Y, Sr, Ca)₁₄Cu₂₄O₄₁: Electronic phases, charge and spin dynamics, *Phys. Rep.* **428**, 169 (2006).
- [31] Y. Fuji, F. Pollmann, and M. Oshikawa, Distinct Trivial Phases Protected by a Point-Group Symmetry in Quantum Spin Chains, *Phys. Rev. Lett.* **114**, 177204 (2015).
- [32] Y. Fuji, Effective field theory for one-dimensional valence-bond-solid phases and their symmetry protection, *Phys. Rev. B* **93**, 104425 (2016).
- [33] S. Eggert and I. Affleck, Magnetic impurities in half-integer-spin Heisenberg antiferromagnetic chains, *Phys. Rev. B* **46**, 10866 (1992).
- [34] P. Lecheminant and E. Orignac, Magnetization and dimerization profiles of the cut two-leg spin ladder, *Phys. Rev. B* **65**, 174406 (2002).
- [35] V. P. Yurov and A. I. B. Zamolodchikov, Truncated conformal space approach to scaling Lee-Yang model, *Int. J. Mod. Phys. A* **05**, 3221 (1990); Truncated-fermionic-space approach to the critical 2D Ising model with magnetic field, *Int. J. Mod. Phys. A* **06**, 4557 (1991).
- [36] A. J. A. James, R. M. Konik, P. Lecheminant, N. J. Robinson, and A. M. Tsvelik, Non-perturbative methodologies for low-dimensional strongly-correlated systems: From non-Abelian bosonization to truncated spectrum methods, *Rep. Prog. Phys.* **81**, 046002 (2018).
- [37] G. Feverati, F. Ravanini, and G. Takács, Truncated conformal space at $c = 1$, nonlinear integral equation and quantization rules for multi-soliton states, *Phys. Lett. B* **430**, 264 (1998).
- [38] G. Feverati, F. Ravanini, and G. Takács, Scaling functions in the odd charge sector of sine-Gordon/massive Thirring theory, *Phys. Lett. B* **444**, 442 (1998).
- [39] G. Feverati, F. Ravanini, and G. Takács, Non-linear integral equation and finite volume spectrum of sine-Gordon theory, *Nucl. Phys.* **B540**, 543 (1999).
- [40] Z. Bajnok, C. Dunning, L. Palla, G. Takács, and F. Wágner, SUSY sine-Gordon theory as a perturbed conformal field theory and finite size effects, *Nucl. Phys.* **B679**, 521 (2004).
- [41] Z. Bajnok, L. Palla, and G. Takács, Finite size effects in boundary sine-Gordon theory, *Nucl. Phys.* **B622**, 565 (2002).
- [42] Z. Bajnok, L. Palla, and G. Takács, The spectrum of boundary sine-Gordon theory, in *Statistical Field Theories* (Springer, Dordrecht, 2002), pp. 195–204.
- [43] Z. Bajnok, L. Palla, G. Takács, and F. Wágner, Non-perturbative study of the two-frequency sine-Gordon model, *Nucl. Phys.* **B601**, 503 (2001).
- [44] Z. Bajnok, L. Palla, G. Takács, and F. Wágner, The k -folded sine-Gordon model in finite volume, *Nucl. Phys.* **B587**, 585 (2000).
- [45] G. Takács and F. Wágner, Double sine-Gordon model revisited, *Nucl. Phys.* **B741**, 353 (2006).
- [46] G. Zs. Tóth, A nonperturbative study of phase transitions in the multi-frequency sine-Gordon model, *J. Phys. A* **37**, 9631 (2004).
- [47] T. Pálmai and G. Takács, Diagonal multisoliton matrix elements in finite volume, *Phys. Rev. D* **87**, 045010 (2013).
- [48] R. M. Konik, Exciton Hierarchies in Gapped Carbon Nanotubes, *Phys. Rev. Lett.* **106**, 136805 (2011).
- [49] R. M. Konik, M. Y. Sfeir, and J. A. Misewich, Predicting excitonic gaps of semiconducting single-walled carbon nanotubes from a field theoretic analysis, *Phys. Rev. B* **91**, 075417 (2015).
- [50] Z. Bajnok, L. Palla, and G. Takacs, Boundary states and finite size effects in sine-Gordon model with Neumann boundary condition, *Nucl. Phys.* **B614**, 405 (2001).
- [51] Z. Bajnok, L. Palla, G. Takacs, and G. Zs. Toth, The spectrum of boundary states in sine-Gordon model with integrable boundary conditions, *Nucl. Phys.* **B622**, 548 (2002).
- [52] One may change the occupation of the MBS pair only at the expense of populating high-energy states via a mechanism similar to the “quasiparticle poisoning” [2] of topological Majorana wires.
- [53] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven, Signatures of Majorana fermions in hybrid superconductor-semiconductor nanowire devices, *Science* **336**, 1003 (2012).
- [54] W. Chang, S. M. Albrecht, T. S. Jespersen, F. Kuemmeth, P. Krogstrup, J. Nygård, and C. M. Marcus, Hard gap in epitaxial semiconductor-superconductor nanowires, *Nat. Nanotechnol.* **10**, 232 (2015).
- [55] A. P. Higginbotham, S. M. Albrecht, G. Kiršanskas, W. Chang, F. Kuemmeth, P. Krogstrup, T. S. Jespersen, J. Nygård, K. Flensberg, and C. M. Marcus, Parity lifetime of bound states in a proximitized semiconductor nanowire, *Nat. Phys.* **11**, 1017 (2015).
- [56] S. M. Albrecht, A. P. Higginbotham, M. Madsen, F. Kuemmeth, T. S. Jespersen, J. Nygård, P. Krogstrup, and C. M. Marcus, Exponential protection of zero modes in Majorana islands, *Nature (London)* **531**, 206 (2016).
- [57] D. C. Cabra, A. Honecker, and P. Pujol, Magnetization plateaux in N -leg spin ladders, *Phys. Rev. B* **58**, 6241 (1998).
- [58] V. Bois, P. Fromholz, and P. Lecheminant, One-dimensional two-orbital SU(N) ultracold fermionic quantum gases at

- incommensurate filling: A low-energy approach, *Phys. Rev. B* **93**, 134415 (2016).
- [59] F. H. L. Essler and A. M. Tsvelik, Weakly coupled one-dimensional Mott insulators, *Phys. Rev. B* **65**, 115117 (2002).
- [60] F. L. Pedrocchi, S. Chesi, S. Gangadharaiah, and D. Loss, Majorana states in inhomogeneous spin ladders, *Phys. Rev. B* **86**, 205412 (2012).
- [61] W. DeGottardi, D. Sen, and S. Vishveshwara, Topological phases, Majorana modes and quench dynamics in a spin ladder system, *New J. Phys.* **13**, 065028 (2011).
- [62] J. C. Y. Teo and C. L. Kane, From Luttinger liquid to non-Abelian quantum Hall states, *Phys. Rev. B* **89**, 085101 (2014); Y. Fuji and P. Lecheminant, Non-Abelian $SU(N-1)$ -singlet fractional quantum Hall states from coupled wires, *Phys. Rev. B* **95**, 125130 (2017).
- [63] H.-C. Jiang, Z.-X. Li, A. Seidel, and D.-H. Lee, Symmetry protected topological Luttinger liquids and the phase transition between them, *Sci. Bull.* **63**, 753 (2018).
- [64] L. Fu, Electron Teleportation via Majorana Bound States in a Mesoscopic Superconductor, *Phys. Rev. Lett.* **104**, 056402 (2010).
- [65] B. Béri and N. R. Cooper, Topological Kondo Effect with Majorana Fermions, *Phys. Rev. Lett.* **109**, 156803 (2012).
- [66] A. Altland and R. Egger, Multiterminal Coulomb-Majorana Junction, *Phys. Rev. Lett.* **110**, 196401 (2013).
- [67] B. Béri, Majorana-Klein Hybridization in Topological Superconductor Junctions, *Phys. Rev. Lett.* **110**, 216803 (2013).
- [68] S. Plugge, A. Rasmussen, R. Egger, and K. Flensberg, Majorana box qubits, *New J. Phys.* **19**, 012001 (2017).
- [69] T. Karzig, C. Knapp, R. M. Lutchyn, P. Bonderson, M. B. Hastings, C. Nayak, J. Alicea, K. Flensberg, S. Plugge, Y. Oreg, C. M. Marcus, and M. H. Freedman, Scalable designs for quasiparticle-poisoning-protected topological quantum computation with Majorana zero modes, *Phys. Rev. B* **95**, 235305 (2017).