

Unique Access to u -Channel Physics: Exclusive Backward-Angle Omega Meson Electroproduction

W. B. Li,^{1,2} G. M. Huber,¹ H. P. Blok,^{3,4} D. Gaskell,⁵ T. Horn,⁶ K. Semenov-Tian-Shansky,^{7,8} B. Pire,⁹ L. Szymanowski,¹⁰
 J.-M. Laget,⁵ K. Aniol,¹¹ J. Arrington,¹² E. J. Beise,¹³ W. Boeglin,¹⁴ E. J. Brash,¹⁵ H. Breuer,¹³ C. C. Chang,¹³
 M. E. Christy,¹⁶ R. Ent,⁵ E. F. Gibson,¹⁷ R. J. Holt,¹⁸ S. Jin,¹⁹ M. K. Jones,⁵ C. E. Keppel,^{16,5} W. Kim,¹⁹ P. M. King,²⁰
 V. Kovaltchouk,²¹ J. Liu,²² G. J. Lolos,¹ D. J. Mack,⁵ D. J. Margaziotis,¹¹ P. Markowitz,¹⁴ A. Matsumura,²³ D. Meekins,⁵
 T. Miyoshi,²³ H. Mkrtchyan,²⁴ I. Niculescu,²⁵ Y. Okayasu,²³ L. Pentchev,⁵ C. Perdrisat,²⁶ D. Potterveld,¹² V. Punjabi,²⁷
 P. E. Reimer,¹² J. Reinhold,¹⁴ J. Roche,²⁰ P. G. Roos,¹³ A. Sarty,²⁸ G. R. Smith,⁵ V. Tadevosyan,²⁴ L. G. Tang,^{16,5}
 V. Tvaskis,^{29,4} J. Volmer,^{29,30} W. Vulcan,⁵ G. Warren,³¹ S. A. Wood,⁵ C. Xu,¹ and X. Zheng³²

(Jefferson Lab F_π Collaboration)

¹University of Regina, Regina, Saskatchewan S4S 0A2, Canada

²College of William and Mary, Williamsburg, Virginia 23185, USA

³VU University, NL-1081 HV Amsterdam, Netherlands

⁴NIKHEF, Postbus 41882, NL-1009 DB Amsterdam, Netherlands

⁵Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

⁶Catholic University of America, Washington, D.C. 20064, USA

⁷National Research Centre Kurchatov Institute: Petersburg Nuclear Physics Institute, RU-188300 Gatchina, Russia

⁸Saint Petersburg National Research Academic University of the Russian Academy of Sciences, RU-194021 St. Petersburg, Russia

⁹CPHT, CNRS, École Polytechnique, IP Paris, 91128-Palaiseau, France

¹⁰National Centre for Nuclear Research (NCBJ), 02-093 Warsaw, Poland

¹¹California State University Los Angeles, Los Angeles, California 90032, USA

¹²Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA

¹³University of Maryland, College Park, Maryland 20742, USA

¹⁴Florida International University, Miami, Florida 33119, USA

¹⁵Christopher Newport University, Newport News, Virginia 23606, USA

¹⁶Hampton University, Hampton, Virginia 23668, USA

¹⁷California State University, Sacramento, California 95819, USA

¹⁸Caltech, Pasadena, California 91125, USA

¹⁹Kyungpook National University, Daegu, 702-701, Republic of Korea

²⁰Ohio University, Athens, Ohio 45701, USA

²¹Ontario Tech University, Oshawa, Ontario L1G 0C5, Canada

²²Shanghai Jiao Tong University, Shanghai 200240, China

²³Tohoku University, Sendai, Miyagi 980-8578, Japan

²⁴A.I. Alikhanyan National Science Laboratory, Yerevan 0036, Armenia

²⁵James Madison University, Harrisonburg, Virginia 22807, USA

²⁶College of William and Mary, Williamsburg, Virginia 23187, USA

²⁷Norfolk State University, Norfolk, Virginia 23504, USA

²⁸Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada

²⁹VU University, NL-1081 HV Amsterdam, Netherlands

³⁰DESY, Hamburg 22607, Germany

³¹Pacific Northwest National Laboratory, Richland, Washington 99352, USA

³²University of Virginia, Charlottesville, Virginia 22904, USA

 (Received 27 June 2019; revised manuscript received 1 October 2019; published 31 October 2019)

Backward-angle meson electroproduction above the resonance region, which was previously ignored, is anticipated to offer unique access to the three quark plus sea component of the nucleon wave function. In this Letter, we present the first complete separation of the four electromagnetic structure functions above the resonance region in exclusive ω electroproduction off the proton, $ep \rightarrow e'p\omega$, at central Q^2 values of 1.60, 2.45 GeV², at $W = 2.21$ GeV. The results of our pioneering $-u \approx -u_{\min}$ study demonstrate the existence of a unanticipated backward-angle cross section peak and the feasibility of full $L/T/LT/TT$ separations in this never explored kinematic territory. At $Q^2 = 2.45$ GeV², the observed dominance of σ_T over σ_L , is qualitatively consistent with the collinear QCD description in the near-backward

regime, in which the scattering amplitude factorizes into a hard subprocess amplitude and baryon to meson transition distribution amplitudes: universal nonperturbative objects only accessible through backward-angle kinematics.

DOI: 10.1103/PhysRevLett.123.182501

Deep exclusive reactions have recently gained much attention, as they provide direct access to the internal structure of hadrons. Measurements of such reactions at different squared four momenta of the exchanged virtual photon (γ^*), Q^2 , and at different hadron four-momentum transfer, Mandelstam variable t and u (defined in Fig. 1), are used to probe QCD's transition from hadronic degrees of freedom at the long distance scale to quark-gluon degrees of freedom at the short distance scale.

The standard experimental configuration to probe deep exclusive reactions involves accelerated charged lepton collisions with a hydrogen target. While most experiments detect the scattered leptons and forward going final state particles (in the laboratory reference frame), the reaction of interest of this Letter concerns final state particles produced at the backward angle. The visualization of the backward-angle interaction gives rise to a unique physical picture: a target proton absorbs most of the momentum transfer (by γ^*), and recoils forward, whereas the produced meson remains close to the target nearly at rest. This type of reaction is sometimes referred to as a “knocking a proton out of a proton” process. The backward-angle exclusive observables accessed by the methodology presented in this Letter opens up new opportunities to extend the current knowledge on the nucleon structure to an unexplored kinematic region.

In the Bjorken limit (sufficiently large Q^2 and invariant mass W , and $-t/Q^2 \ll 1$), the longitudinal scattering amplitude factorizes into a hard scattering perturbative contribution, and soft generalized parton distributions (GPDs) of the nucleon and distribution amplitudes (DAs) of the meson. The vector meson (ω) production through the GPD in the near-forward kinematics is demonstrated in Fig. 1(a). GPDs are light-cone matrix elements




FIG. 1. QCD collinear factorization mechanisms for exclusive ω electroproduction off a proton (p) probed by γ^* at large Q^2 and W : (a) Forward regime (small $-t$), GPDs (bottom oval), and the ω -DA (top-right oval); (b) backward regime (small $-u$), ωN TDAs (bottom oval) and the proton N -DA (top-right oval).

of nonlocal bilinear quark and gluon operators that describe the three-dimensional structure of hadrons, by correlating the internal transverse position of partons to their longitudinal momentum. For a review, see Refs. [1–11].

Analogous universal structure functions accessible in “near-backward” kinematics are known as baryon-to-meson transition distribution amplitudes (TDAs) [12–16], see Fig. 1(b), which are light-cone matrix elements of nonlocal three quark operators. In the TDA picture, the backward-angle meson is produced as the γ^* probes the meson cloud structure of the nucleon.

The TDA collinear factorization regime for hard meson production has two key marking signs in near-backward kinematics which can be tested experimentally [12–16]:

(i) The dominance of the transverse polarization of the virtual photon results in the suppression of the longitudinal cross section (σ_L) at large Q^2 by at least a factor of $1/Q^2$: $\sigma_L/\sigma_T < \mu^2/Q^2$ and $\sigma_T \gg \sigma_L$, where μ is a typical hadronic scale.

(ii) The characteristic $1/Q^8$ scaling behavior of the transverse cross section (σ_T) for fixed Bjorken x : $x_B = (Q^2/2p_p q)$, where p_p and q are the four momenta of the virtual proton and γ^* , respectively.

In a recent publication [17], the CLAS Collaboration reported the first measurement of the cross sections for exclusive π^+ electroproduction off the proton in near-backward kinematics. The result gives promising signs of the predicted $1/Q^8$ scaling of the cross section by TDA; however, the critical evidence for σ_T dominance remains missing.

In this Letter, we present a pioneering study of backward-angle ω cross sections from exclusive electroproduction: $e p \rightarrow e' p \omega$ using the missing-mass reconstruction technique. The extracted cross sections are separated into the transverse (T), longitudinal (L), and LT , TT interference terms. This allows for comparing the individual σ_L and σ_T contributions to the TDA calculations and verifying the predicted σ_T dominance.

The general form of twofold virtual-photon differential cross section in terms of the structure functions is given as

$$2\pi \frac{d^2\sigma}{dt d\phi} = \frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} + \sqrt{2\epsilon(1+\epsilon)} \frac{d\sigma_{LT}}{dt} \cos\phi + \epsilon \frac{d\sigma_{TT}}{dt} \cos 2\phi, \quad (1)$$

where ϵ is the γ^* longitudinal polarization $\epsilon = [1 + 2(|\vec{q}|^2/Q^2)\tan^2(\theta_e/2)]^{-1}$; θ_e is the scattered electron

polar angle; ϕ is the azimuthal angle between the electron scattering plane and the proton target reaction plane. For brevity, differential cross sections such as $d\sigma_T/dt$ will be expressed as σ_T . Separating σ_L from σ_T , and extracting the interference terms relies on an experimental technique known as Rosenbluth separation. This technique requires two measurements at different ϵ (dependent upon the beam energy and electron scattering angle), while other Lorentz invariant quantities are kept constant. The interference terms σ_{LT} and σ_{TT} dictate the azimuthal modulation for a given opening angle θ between the proton recoil momentum and the γ^* momentum.

The analyzed data were part of experiment E01-004 (F_π -2), which used 2.6–5.2 GeV electron beams on a liquid hydrogen target and the high precision particle spectrometers in Jefferson Lab Hall C [18,19]. The dataset has two central Q^2 values: $Q^2 = 1.60$ and 2.45 GeV², at common central $W = 2.21$ GeV. The primary objective of the experiment was to detect coincidence e - π at a forward angle, but backward-angle ω (e - p) were fortuitously acquired.

The recoil protons were detected in the high momentum spectrometer (HMS), while the scattered electrons were detected in the short orbit spectrometer (SOS). Both spectrometers include two sets of drift chambers for tracking and scintillator arrays for triggering. A detailed description of the experimental configuration is documented in Refs. [19].

In order to select e^- in the SOS, a gas Cherenkov detector containing Freon-12 at 1 atm was used in combination with a lead-glass calorimeter. The positively charged π^+ were rejected in the HMS using an aerogel Cherenkov detector with a refractive index of 1.03. The rare e^+ were rejected using a gas Cherenkov detector filled with C_4F_{10} at 0.47 atm. Most remaining contamination of the e - p events was rejected by a coincidence time cut of ± 1 ns. Background originated from the aluminum target cell and random coincidence events, $< 5\%$ contribution to the total yield, was subtracted from the charge normalized yield. Proton loss due to multiple scattering inside the HMS was estimated as 7–10% [20].

Unlike the exclusive π^+ channel [18,19], the ω events sit on a broad background, as shown in the reconstructed missing-mass spectrum for $ep \rightarrow e'pX$ in Fig. 2. The final state particle X could include ω , ρ , or two- π production ($\pi\pi$). For each Q^2 - ϵ - u - ϕ bin, extracting ω is a two step process. First, simulations were used to determine the contribution of each final state particle to the M_X distribution. Here, the shape of the distribution for each particle is dictated by the detector acceptance and the particle decay width, while the normalization (scale) factor of the simulated distribution is determined by the fit to the data (simultaneously). In the second step, the background (scaled ρ and $\pi\pi$ simulations) is subtracted from the data to obtain the ω experimental yield.




FIG. 2. Reconstructed missing mass (M_X) for $ep \rightarrow e'pX$ at $Q^2 = 2.45$ GeV² (blue crosses). The simulated distributions for ρ (blue), ω (red), and $\pi\pi$ (green) are used to describe the measured reaction.

Two quality control criteria were introduced to validate the background subtraction procedure: (1) The χ^2 per-degree-of-freedom ($\chi^2/\text{d.o.f.}$) comparison between the experimental and simulated ω yields, defined as $Y_{\omega \text{ exp}} = Y_{\text{data}} - Y_{\rho \text{ sim}} - Y_{\pi\pi \text{ sim}}$, and $Y_{\omega \text{ sim}}$; (2) $\chi^2/\text{d.o.f.}$ comparison between the experimental and simulated background yields, defined as $Y_{\text{BG exp}} = Y_{\text{data}} - Y_{\omega \text{ sim}}$ and $Y_{\text{BG sim}} = Y_{\rho \text{ sim}} + Y_{\pi\pi \text{ sim}}$. Both $\chi^2/\text{d.o.f.}$ distributions obey Poisson statistics with center values: 0.94, 1.3, and widths: 0.77, 0.97, respectively. The detailed analysis procedure is documented in Ref. [20].

For each Q^2 setting, two datasets with different ϵ values were acquired: $Q^2 = 1.6$ GeV², $\epsilon = 0.32, 0.59$; at $Q^2 = 2.45$ GeV², $\epsilon = 0.27, 0.55$. Data at each Q^2 - ϵ setting were divided into three u bins and eight ϕ bins. Figure 3 shows the unseparated experimental cross section at $Q^2 = 1.6$ GeV² as functions of ϕ at three $-u$ bins. The separated cross section is obtained from fitting the data at both ϵ settings simultaneously using Eq. (1).

The experimental acceptance covers a range of Q^2 , W values; thus the measured experimental yields represent an average over the covered range. As a result, each $-u$ bin has a slightly different average value $\overline{Q^2}$ and \overline{W} . In order to minimize errors that resulted from the averaging, the experimental cross sections were calculated by comparing the experimental yields to a Monte Carlo simulation of the experiment. The Monte Carlo simulation includes a detailed description of the spectrometer acceptance, multiple scattering, energy loss due to ionization, decay, and radiative process.

The shape of the simulated ω tail is influenced by radiative effects describing the emission of real or virtual photons, and multiple scattering. For more information on the simulation, see Ref. [19]. The matching between the




FIG. 3. Unseparated cross section as a function of ϕ at $-u = 0.057, 0.135,$ and 0.245 GeV^2 (from bottom to top) at $Q^2 = 1.6 \text{ GeV}^2$. The higher $\epsilon = 0.59$ and lower $\epsilon = 0.32$ data are shown in red circles and black crosses, respectively. Red dashed (higher ϵ) and black solid (lower ϵ) lines are the fitting results used in Eq. (1). Note that the fitting performed takes into account data at both ϵ settings simultaneously.

simulated and the experimental resolutions was verified with ep elastic scattering data and the relatively small observed effect is included in the point-to-point systematic uncertainty. Additionally, to avoid sensitivity to some kinematic regions (at larger θ) with large contributions from the radiative events, bins with simulated ω tail contribution $> 60\%$ of the overall distribution are excluded from the analysis (9% of the bins).

The uncertainty in the separated cross sections includes both statistical and systematic contributions. The statistical contribution consists of the error in determining “good” ω from the background subtraction procedure (fitting error included), the uncertainties in detector performance (efficiencies and tracking) and beam characteristics on a run-by-run basis. A comprehensive study was carried out to obtain the total systematic uncertainties for the separated cross section. It includes three parts: (1) correlated scale error of the unseparated cross section (2.6%); (2) point-to-point variations due to the cross section model dependence in simulation; (3) effects of the error amplification (by a factor of $1/\Delta\epsilon$) of the ϵ uncorrelated u correlated systematic error (1.7–2.0%). The effects of all three parts are added in quadrature as the total systematic error and are reported separately for each u bin.

To investigate the Q^2 dependence, σ_L and σ_T for the smallest $-u$ bin ($u - u_{\min} = 0$) from the two Q^2 settings are plotted on the left panel of Fig. 4, whereas the σ_L/σ_T ratio




FIG. 4. Left: $\sigma_L(u = u_{\min})$ and $\sigma_T(u = u_{\min})$ as function of Q^2 for the lowest $-u$ bin. Right: $\sigma_L(u = u_{\min})/\sigma_T(u = u_{\min})$ ratio as function of Q^2 . Fitted lines are for visualization purpose only.

is plotted on the right. σ_T shows a flat Q^2 dependence, whereas σ_L decreases significantly as Q^2 rises. The drop in the σ_L/σ_T ratio as a function of Q^2 is qualitatively consistent with the prediction of TDA collinear factorization.

The extracted σ_L and σ_T as a function of $-u$ at $Q^2 = 1.6$ and 2.45 GeV^2 are shown in Fig. 5. The two sets of TDA predictions for σ_T each assume different nucleon DAs as input. The predictions were calculated at the specific $\overline{Q}^2, \overline{W}$ values of each u bin. The predictions at three u bins are joined by straight lines. At $Q^2 = 2.45 \text{ GeV}^2$, TDA predictions are within the same order of magnitude as the data, whereas at $Q^2 = 1.6 \text{ GeV}^2$, the TDA model overpredicts the data by a factor of ~ 10 . This is very similar to the recent backward-angle π^+ data from the CLAS Collaboration [17], where the TDA prediction is within 50% of the data at $Q^2 = 2.5 \text{ GeV}^2$, but far higher than the unseparated data at $Q^2 = 1.7 \text{ GeV}^2$. Together, datasets suggest that the boundary where the TDA factorization applies may begin around $Q^2 = 2.5 \text{ GeV}^2$.




FIG. 5. σ_T (triangles), σ_L (circles) as function of $-u$, at $Q^2 = 1.6 \text{ GeV}^2$ (left), 2.45 GeV^2 (right). For the lowest $-u$ bin. TDA predictions for σ_T : Chernyak-Ogloblin-Zhitnitsky (COZ) [21] (blue dashed lines) and KS [22] (red solid lines). The green bands indicate correlated systematic uncertainties for σ_T , the uncertainties for σ_L have similar magnitude.




FIG. 6. Exclusive ω electroproduction cross section as a function of $-t$ at $Q^2 = 1.75$ (left panel) and $Q^2 = 2.35$ GeV^2 (right panel). The CLAS Collaboration data are the black dots in the near-forward kinematics region ($-t < 2.5$ GeV^2), and the $F_{\pi-2}$ are the red crosses in the backward region ($-t > 5$ GeV^2), scaled to the kinematics of the CLAS Collaboration data, as described in the text. The blue and magenta dashed thick lines are Regge trajectory-based JML04 and JML18 predictions, respectively. The short curves above the $F_{\pi-2}$ data are TDA predictions based on the COZ model [21] (blue solid) and KS [22] (red solid) DAs.

The behavior of σ_L differs greatly at the two Q^2 settings. At $Q^2 = 1.6$ GeV^2 , σ_L falls almost exponentially as a function of $-u$; at $Q^2 = 2.45$ GeV^2 , σ_L is constant near zero (within one standard deviation) and this is consistent with the leading-twist TDA prediction: $\sigma_L \approx 0$.

The combined data from CLAS Collaboration [23] and $F_{\pi-2}$ cover both forward and backward-angle kinematics, and jointly form a complete $-t$ evolution picture for the $ep \rightarrow e'p\omega$ reaction. The CLAS Collaboration data, at $W \sim 2.48$ GeV^2 , $Q^2 = 1.75$ and 2.35 GeV^2 , are shown in the left and right panels of Fig. 6, respectively. Because of the similarities in the kinematics, the $F_{\pi-2}$ data (this work) are scaled to those of the CLAS Collaboration data. The W dependence of the backward-angle cross section is unknown; therefore the scaling procedure: $(W^2 - m_p^2)^{-2}$, based on the forward-angle phenomenology studies, is applied [24]. The Q^2 scaling is based on the empirical fit used to extract the separated cross sections of this work. This empirical model is documented in Ref. [20]. In addition to the scaling, the extracted $-u$ dependent cross section from $F_{\pi-2}$ is translated to the $-t$ space of the CLAS Collaboration data.

Figure 6 indicates strong evidence of the existence of the backward-angle peak at $-t > 5$ GeV^2 for both Q^2 settings, with strength $\sim 1/10$ of the forward-angle cross section. Previously, the “forward-backward” peak phenomenon was only observed in π^+ photoproduction data [25–29]. This was successfully interpreted using the Regge trajectory-based VGL model [25,26].

The results presented in this Letter have demonstrated that the missing-mass reconstruction technique, in combination with the high precision spectrometers in coincidence mode at Hall C of Jefferson Lab, is able to reliably perform a full L/T separation of the backward-angle exclusive reaction $ep \rightarrow e'p\omega$. Since the missing mass reconstruction method does not require the detection of the produced meson, this allows the possibility to extend experimental kinematic coverage that was considered to be inaccessible through the standard direct detection method. If used in combination with a large acceptance detector, such as CLAS-12, one could systematically study the complete t evolution of a given interaction, thus unveiling new aspects of nucleon structure. The separated cross sections show indications of a regime where $\sigma_T \gg \sigma_L$ for $ep \rightarrow e'p\omega$, qualitatively consistent with the TDA factorization approach in backward-angle kinematics. However, the approach relying on the QCD partonic picture applying at large enough Q^2 involves different mechanisms for the forward and backward peaks and could not provide a unique description in the whole range in $-t$.

An alternative description for the ω -meson electroproduction cross section is given by the Regge-based JML model. It describes the JLab π electroproduction cross sections over a wide kinematic range without destroying good agreement at $Q^2 = 0$ [30,31]. Two L.-M. Laget model predictions are plotted in Fig. 6: JML04 [32] (prior to $F_{\pi-2}$ data) and JML18. JML04 includes the near-forward Regge contribution at $-t < 1$ GeV^2 and N exchange in the u channel with a t -dependent cutoff mass. It significantly underpredicts the backward-angle cross section. In JML18 [33], the principle of the u -channel treatment is the same as in the t -channel neutral pion electroproduction [31]. It includes, in addition, an estimation of the contribution of the ρ - N and ρ - Δ unitarity rescattering (Regge) cuts, allowing an excellent description of the combined data within a unique framework. In particular, the $-u$ dependence and the strength of the backward-angle peak are described well at both Q^2 settings. The inelastic exchange diagrams are the main sources to the observed backward-angle peak, with one-third of the contribution coming from the ρ^0 - ω transition, and the rest coming from ρ^+ - N and Δ resonance. However, JML18 lacks the prediction of the Q^2 dependence of the σ_L/σ_T ratio.

In conclusion, the presented experimental data hint on the early onset of the QCD-based factorized description of electroproduction of ω in the backward kinematics regime for Q^2 in the few GeV^2 range. This opens a way to the experimental access of nucleon-to-meson TDAs and provides a new window on the quark-gluon structure of nucleons. These data also supply a new interesting testing bench for Regge-based hadronic models.

We acknowledge the excellent efforts provided by the staff of the Accelerator and the Physics Divisions at

Jefferson Lab. This work is supported by NSERC (Canada) FRN: SPAPIN-2016-00031, DOE and NSF (U.S.), FOM (Netherlands), North Atlantic Treaty Organization (NATO), and NRF (Republic of Korea). Additional support from Jefferson Science Associates and the University of Regina is gratefully acknowledged. This material is based upon work supported by the U.S. Department of Energy under Contracts No. DE-AC05-06OR23177 and No. DE-AC02-06CH11357. L. S. is supported by Grant No. 2017/26/M/ST2/01074 of the National Science Center in Poland. He thanks the French Laboratories of Excellence (LABEX) P2IO and the French Groupement de Recherche Chromodynamique Quantique et Physique des Hadrons (GDR QCD) for support.

-
- [1] D. Mueller, D. Robaschik, B. Geyer, F. M. Dittes, and J. Horejsi, *Fortschr. Phys.* **42**, 101 (1994).
- [2] A. V. Radyushkin, *Phys. Lett. B* **380**, 417 (1996).
- [3] X. Ji, *Phys. Rev. Lett.* **78**, 610 (1997).
- [4] X. Ji, *Phys. Rev. D* **55**, 7114 (1997).
- [5] K. Goeke, M. V. Polyakov, and M. Vanderhaeghen, *Prog. Part. Nucl. Phys.* **47**, 401 (2001).
- [6] M. Diehl, *Phys. Rep.* **388**, 41 (2003).
- [7] A. Belitsky and A. Radyushkin, *Phys. Rep.* **418**, 1 (2005).
- [8] S. Boffi and B. Pasquini, *Riv. Nuovo Cimento* **30**, 387 (2007).
- [9] M. Guidal, H. Moutarde, and M. Vanderhaeghen, *Rep. Prog. Phys.* **76**, 066202 (2013).
- [10] K. Kumericki, S. Liuti, and H. Moutarde, *Eur. Phys. J. A* **52**, 157 (2016).
- [11] L. L. Frankfurt, P. V. Pobylitsa, M. V. Polyakov, and M. Strikman, *Phys. Rev. D* **60**, 014010 (1999).
- [12] B. Pire and L. Szymanowski, *Phys. Rev. D* **71**, 111501(R) (2005).
- [13] B. Pire and L. Szymanowski, *Phys. Lett. B* **622**, 83 (2005).
- [14] J. P. Lansberg, B. Pire, K. Semenov-Tian-Shansky, and L. Szymanowski, *Phys. Rev. D* **85**, 054021 (2012).
- [15] B. Pire, K. Semenov-Tian-Shansky, and L. Szymanowski, *Phys. Rev. D* **91**, 094006 (2015).
- [16] K. M. Semenov-Tian-Shansky, JLab Presentation (2015), https://userweb.jlab.org/~kenjo/EMP/1400_Semenov-Tian-Shansky.pdf.
- [17] K. Park *et al.*, *Phys. Lett. B* **780**, 340 (2018).
- [18] T. Horn *et al.*, *Phys. Rev. Lett.* **97**, 192001 (2006).
- [19] H. P. Blok *et al.*, *Phys. Rev. C* **78**, 045202 (2008).
- [20] W. Li, [arXiv:1712.03214](https://arxiv.org/abs/1712.03214).
- [21] V. Chernyak, A. A. Ogloblin, and I. R. Zhitnitsky, *Z. Phys. C* **42**, 583 (1989).
- [22] I. D. King and C. T. Sachrajda, *Nucl. Phys.* **B279**, 785 (1987).
- [23] L. Morand *et al.* (CLAS Collaboration), *Eur. Phys. J. A* **24**, 445 (2005).
- [24] P. Brauel, T. Canzler, D. Cords, R. Felst, G. Grindhammer, M. Helm, W.-D. Kollmann, H. Krehbiel, and M. Schädlich, *Z. Phys. C* **3**, 101 (1979).
- [25] M. Guidal, J. M. Laget, and M. Vanderhaeghen, *Proceedings of ELFE Workshop on Hadronic Physics* (1996), FR9806080, <https://www.osti.gov/etdweb/servlets/purl/20058898>.
- [26] M. Guidal, J. M. Laget, and M. Vanderhaeghen, *Phys. Lett. B* **400**, 6 (1997).
- [27] R. L. Anderson, D. Gustavson, J. Johnson, I. Overman, D. Ritson, and B. H. Wiik, *Phys. Rev. Lett.* **23**, 721 (1969).
- [28] J. L. Anderson and J. W. Ryon, *Phys. Rev.* **181**, 1765 (1969).
- [29] A. M. Boyarski, F. Bulos, W. Busza, R. Diebold, S. D. Ecklund, G. E. Fischer, J. R. Rees, and B. Richter *Phys. Rev. Lett.* **20**, 300 (1968).
- [30] J.-M. Laget, *Phys. Lett. B* **685**, 146 (2010).
- [31] J.-M. Laget, *Phys. Lett. B* **695**, 199 (2011).
- [32] J.-M. Laget, *Phys. Rev. D* **70**, 054023 (2004).
- [33] J.-M. Laget (to be published).