

Supernova-Produced ^{53}Mn on EarthG. Korschinek^{1,*}, T. Faestermann¹, M. Poutivtsev^{1,†}, A. Arazi^{2,3}, K. Knie^{1,‡}, G. Rugel^{1,4} and A. Wallner^{1,4}¹Physik-Department, Technische Universität München, 85748 Garching, Germany²Laboratorio TANDAR, Comisión Nacional de Energía Atómica, Av. Gral. Paz 1499, B1650KNA San Martín, Argentina³Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Av. Rivadavia 1917, C1033AAJ Buenos Aires, Argentina⁴Helmholtz-Zentrum Dresden-Rossendorf, 01328 Dresden, Germany

(Received 16 December 2019; revised 13 April 2020; accepted 18 June 2020; published 17 July 2020)

For the time period from 1.5 to 4 Myr before the present we found in deep ocean ferromanganese crusts a ^{53}Mn excess concentration in terms of $^{53}\text{Mn}/\text{Mn}$ of about 4×10^{-14} over that expected for cosmogenic production. We conclude that this ^{53}Mn is of supernova origin because it is detected in the same time window, about 2.5 Myr ago, where ^{60}Fe has been found earlier. This overabundance confirms the supernova origin of that ^{60}Fe . For the first time, supernova-formed ^{53}Mn has been detected and it is the second positively identified radioisotope from the same supernova. The ratio $^{53}\text{Mn}/^{60}\text{Fe}$ of about 14 is consistent with that expected for a SN with a 11–25 M_{\odot} progenitor mass and solar metallicity.

DOI: [10.1103/PhysRevLett.125.031101](https://doi.org/10.1103/PhysRevLett.125.031101)

Stars heavier than about 10 solar masses (M_{\odot}) end their life in a supernova explosion (SN). At the same time they eject large amounts of stable and radio nuclides into the interstellar medium. This happened close to the solar system several times during the last few millions of years (Myr) [1]. One of the expelled nuclides is the long-lived nuclide ^{60}Fe ($T_{1/2} = 2.61 \pm 0.04$ Myr) [2,3]. If the explosion is close enough to the solar system, some material can enter the solar system [4,5]. Traces of ^{60}Fe have already been found in Earth's reservoirs like ferromanganese crusts and ocean sediments, and even on the Moon [6–11]. Several indications support the SN origin of the ^{60}Fe [1,12]. However, there is also the possibility of ^{60}Fe being formed in asymptotic giant branch (AGB) stars [13]. Despite having lower nucleosynthesis yields, they could be the origin of the observed ^{60}Fe , as well. An unambiguous, only SN formed radionuclide, such as ^{53}Mn , detected in the same samples as the ^{60}Fe , can solve this open question. Therefore the concomitant finding of ^{53}Mn ($T_{1/2} = 3.7 \pm 0.4$ Myr) [14] would be a compelling support for the SN origin of this ^{60}Fe . The proton-rich nuclide ^{53}Mn is generated primarily during so-called silicon burning before the supernova explosion. During this process also ^{53}Fe is formed [15] that decays in a few minutes to ^{53}Mn and both end up eventually as stable ^{53}Cr . Until now, ^{53}Mn , formed by nucleosynthesis, has not been detected in the interstellar space because ^{53}Mn decays by electron capture directly to the ground state of ^{53}Cr , hence only low-energy x rays are emitted, not detectable by space borne detectors. This is different to isotopes like ^{60}Fe [16] or ^{26}Al [17] whose 1173 and 1333 keV (^{60}Fe), or 1809 keV (^{26}Al) gamma lines following β decay have been observed in the Galaxy.

A feasible way to detect ^{53}Mn in Earth's reservoirs is, as in the case of the finding of ^{60}Fe , direct atom counting by accelerator mass spectrometry (AMS). Unfortunately, and in contrast to ^{60}Fe , the dominate fraction of the ^{53}Mn in the solar system is produced in dust that originates from asteroid collisions or comets. Individual contributions [18], and the absolute amounts deposited on Earth are still under debate. In these dust particles, ^{53}Mn is mainly formed by cosmic ray reactions mostly on iron via the nuclear reactions $^{\text{nat}}\text{Fe}(p, x)^{53}\text{Mn}$ or $^{\text{nat}}\text{Fe}(n, x)^{53}\text{Mn}$ (with neutrons from the secondary component of the cosmic rays) and on nickel. However, due to its lower abundance, nickel contributes only less than 5% [19,20]. Compared to interplanetary ^{53}Mn influx on Earth the ^{53}Mn influx via interstellar dust [21] represents only a very small contribution, and makes its detection challenging considering that the overwhelming bulk of ^{53}Mn is cosmogenically produced within the Solar System.

Reservoirs collecting the ^{53}Mn from the dust, and preserving time information of the flux over millions of years, are ferromanganese crusts [22] and seafloor sediments [23]. As the terrestrial ^{55}Mn abundance remains constant over time in these reservoirs, knowledge of $^{53}\text{Mn}/^{55}\text{Mn}$ ratios in dated layers of such archives reveals the dust flux over very long periods. Here, we have focused on ferromanganese crusts. Applying AMS we measured time profiles of $^{53}\text{Mn}/\text{Mn}$ isotope ratios over time periods of more than 10 Myr. The dating of the different layers in crusts was performed by means of ^{53}Mn and ^{10}Be [24]. ^{10}Be is formed by cosmic-ray induced nuclear reactions on nitrogen and oxygen in Earth's atmosphere at approximately constant rates over Myr time periods and becomes eventually incorporated into deep-ocean crusts. It has a half-life of

1.387 Myr [25,26]. In contrast to ^{53}Mn , ^{10}Be is routinely measured with AMS with high precision. There have been three previous attempts to measure extraterrestrial ^{53}Mn [23,27,28], but none measured a time dependence.

At present, only AMS systems with high-energy accelerators in conjunction with a gas-filled magnet have the ability to suppress the interfering isobar ^{53}Cr sufficiently to determine $^{53}\text{Mn}/\text{Mn}$ ratios at levels down to about 10^{-14} in milligram amounts of sample material. MnO^- beams are extracted from an ion source. The ions are mass separated and accelerated to energies of $\sim 150\text{--}180$ MeV and ^{53}Mn finally detected as single events in an energy-sensitive particle detector. Details of the measurements can be found elsewhere [29].

We measured four different crusts for their ^{53}Mn content, all of them of hydrogenetic origin, and from different locations in the Pacific Ocean. Two, 29DR-32 and 29DR-45, were from the Midway atoll ($28^\circ 13'\text{N}$, $177^\circ 22'\text{W}$) from depths of 2938 and 1589 m [30], respectively. One, 4DR [31], from the Donizetti ridge ($32^\circ 14.99'\text{N}$, $159^\circ 56.99'\text{W}$ from a depth of 5120 m) and the fourth, (237KD), from the Central Pacific ($9^\circ 18'\text{N}$, $146^\circ 03'\text{W}$) from a depth of 4830 m [32].

Typical mass concentrations in the crusts were about 20% for Mn and in the range of 10–15 ppm for Cr. The Mn fraction was extracted using chemical treatments as described in Ref. [29]. Thus, around 100 mg of crust material yielded $\sim 15\text{--}20$ mg MnO_2 sample material available for the subsequent AMS measurements. Around 15 samples (each about 100 mg crust material) representing 2 mm depth intervals (and representing ~ 1 Myr time integral) were taken from each crust, which resulted in more than 60 samples totally. The measured isotope ratios $^{53}\text{Mn}/\text{Mn}$ are plotted in Fig. 1 and listed in the supplement [33]. In addition, also iron had been separated for two crusts for the determination of $^{60}\text{Fe}/\text{Fe}$ ratios as reported in Refs. [7,11].

The crust 237KD, where the enhanced $^{60}\text{Fe}/\text{Fe}$ ratio about 2.5 Myr ago was reported [7], was dated with ^{10}Be [35]. We used these data but the improved value of the ^{10}Be half-life of 1.387 Myr [25,26] for the timescale given in Fig. 1. The $^{53}\text{Mn}/\text{Mn}$ ratios of all crusts were fitted to the function

$$C = C_0 \exp(-d \ln 2 / (gr T_{1/2})).$$

Here, C is the ratio at depth d , C_0 the result of the fit at the surface, gr is the growth rate, and $T_{1/2}$ the half-life of ^{53}Mn . Then, the age of a sample equals d/gr and the uncertainty of the exponent is subsumed into the uncertainty of gr . The results of the least-squares fits for the two adjusted parameters are listed in Table I.

Although there are considerable fluctuations, especially at ages older than 7 Myr, all crusts show an exponential




FIG. 1. Shows the measured isotope ratios $^{53}\text{Mn}/\text{Mn}$ at different depths/ages in the crusts. Top to bottom are for the four crusts from the Pacific: the two from Midway atoll, (29DR-32 and 29DR-45), one from the Donizetti ridge (4DR) and one from Central Pacific (237KD). Error bars for small event numbers were calculated following the prescription of Feldman and Cousins [34]. The 1σ limits are drawn.

TABLE I. Results for growth rate and initial concentration of the four crusts. The values of the normalized χ^2 are also given.

Crust	29DR-32	29DR-45	4DR	237KD
Growth rate [mm/Myr]	2.04(25)	2.00(18)	1.52(27)	2.55(22)
C_0 [10^{-13}]	1.64(19)	1.72(14)	0.98(16)	1.50(11)
χ^2/dof	1.51	0.52	1.78	1.14

behavior of the $^{53}\text{Mn}/\text{Mn}$ ratio as a function of depth due to the half-life of ^{53}Mn . In one case the value of χ^2/dof is considerably larger than unity.

To check the validity of dating by ^{53}Mn , we compare the dating by ^{10}Be with ^{53}Mn for the case of the crust 237 KD. The growth rate for ^{10}Be dating, was 2.32 mm/Myr in Ref. [35], but if we fit these ^{10}Be data between 2 and 12 mm, where the growth rate seems constant and use the most precise half-life of ^{10}Be , namely, 1.387 Myr [25,26], we obtain a growth rate of 2.56(13) mm/Myr. Within the uncertainties the ^{53}Mn data yield the same values for the growth rate, 2.55(22) mm/Myr, meaning that the adopted half-life value for ^{53}Mn is supported. Therefore, the dating of the other three crusts was based on ^{53}Mn data only.

In order to deduce the contribution of interstellar, SN-produced ^{53}Mn , we select the time range between 1.5 and 4 Myr where the highest $^{60}\text{Fe}/\text{Fe}$ ratios have been found [7,9,10]. These data were compared with the concentrations obtained for younger and older time periods. The rather wide time window was chosen to allow for uncertainties in the dating of the individual samples which is mainly caused by the inhomogeneous structure of the different crusts. In a first step we calculated all $^{53}\text{Mn}/\text{Mn}$ ratios at the time of incorporation by means of their age (i.e., corrected for radioactive decay). In a second step we merged the data from the four crusts as shown in Fig. 2.

In the next step, we formed the weighted average of the data in the relevant time interval T_2 , where the influx of ^{60}Fe has been detected [6,7,9–11], from 1.5 until 4 Myr, and the adjacent intervals, T_1 from 0 until 1.5 Myr and T_3 from 4 until 5.5 Myr. The fact that the decay-corrected concentrations at T_1 and T_3 are smaller than 1 is the result of fitting to a pure exponential and underlines the excess at T_2 . The ratio of the concentration at T_2 and the average at T_1 and T_3 yields $2C(T_2)/[C(T_1) + C(T_3)] = 1.34 \pm 0.09$. This is more than 3σ larger than unity.

As an alternative, we fitted the data in Fig. 2 (from 0 to 13 Myr) to a constant b and a Gaussian

$$\frac{C}{C_0} = b + a \exp(-(t - t_p)^2 / (2\sigma^2)),$$

with fixed width $\sigma = 0.8$ Myr on the basis of the ^{60}Fe data [9,10], and varied the position t_p and height a as well as the base value b . The result is plotted in Fig. 2 and the parameters are




FIG. 2. Merged $^{53}\text{Mn}/\text{Mn}$ ratios (C/C_0) from the different crusts at the time of incorporation. The red curve is the result of a fit of a Gaussian with fixed width $\sigma = 0.8$ Myr.

$$b = 0.928 (39),$$

$$a = 0.29 (10),$$

$$t_p = 2.56 (33) \text{ Myr.}$$

The resulting $\chi^2 = 58.9$ is larger than the 50 degrees of freedom. However, the Gaussian is probably not the correct shape of the distribution. Again, the peak has a significance of about 3σ .

Now, we can compare our measured values with nucleosynthesis models, in order to get a hint on the SN progenitor. We assume that the ^{53}Mn ejected from a SN follows the same path as the ^{60}Fe until it becomes incorporated into the ferromanganese crusts. We have measured a ratio $^{60}\text{Fe}/\text{Fe} \sim 3.4 \times 10^{-15}$ in the crusts [7] for the time range (2.58 ± 0.43) Myr, calculated back to the time of deposition. The additional $^{53}\text{Mn}/\text{Mn}$ ratio above the cosmogenically formed ratio is $0.95abC_0 = (3.8 \pm 1.5) \times 10^{-14}$, the factor 0.95 represents the average value of the fitted Gaussian in the same range ± 0.43 Myr as evaluated for the ^{60}Fe data. As the weighted average of the four C_0 values from Table I we obtain $C_0 = (1.47 \pm 0.15) \times 10^{-13}$, where the error is inflated by the square root of the normalized χ^2 . If we consider a Fe/Mn ratio in the crusts of ~ 0.8 (it varies between 0.6 and 0.9 [22]), we get for the $^{53}\text{Mn}/^{60}\text{Fe}$ ratio a value of about 14. However, the uptake factors into the crust for Mn and Fe can be quite different. In Ref. [7] the uptake factor for Mn was estimated 7 times greater than for Fe. If this were the case, the relative mass yields of ^{53}Mn to ^{60}Fe in the interstellar dust would be 2:1. As well, the ratio of the mass yields $^{53}\text{Mn}/^{60}\text{Fe}$, calculated by Woosley and Weaver [15], varies between about 1 and 20 for a SN with a mass of the progenitor star between 11 and 25 M_\odot and solar metallicity. For the time being, we only can state that the

observed ratio of $^{53}\text{Mn}/^{60}\text{Fe}$ in the crusts is in the expected range.

Since the excess of ^{53}Mn is detected in the same samples and time range in which ^{60}Fe has been identified, it confirms the SN origin of that ^{60}Fe . Thus, ^{53}Mn is the second radioisotope from the same SN where ^{60}Fe has been detected, and it is for the first time that ^{53}Mn , formed by nucleosynthesis during a SN, has been found. This finding might initiate further searches in deep sea sediments or manganese crusts for long-lived radionuclides, like the proton-rich nuclei ^{92}Nb ($T_{1/2} = 35$ Myr) or ^{146}Sm ($T_{1/2} = 68$ Myr), to corroborate nucleosynthesis calculations. Up to now, the search for SN-produced ^{26}Al only resulted in an upper limit [36].

This work was supported by the DFG cluster of excellence “Origin and Structure of the Universe.”

*Corresponding author.

korschin@tum.de

[†]Present address: DAW SE, 64372 Ober-Ramstadt, Germany.

[‡]Present address: GSI Helmholtz-Zentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany.

- [1] D. Breitschwerdt, J. Feige, M. M. Schulreich, M. A. de Avelles, C. Dettbarn, and B. Fuchs, *Nature (London)* **532**, 73 (2016).
- [2] G. Rugel, T. Faestermann, K. Knie, G. Korschinek, M. Poutivtsev, D. Schumann, N. Kivel, I. Günther-Leopold, R. Weinreich, and M. Wohlmuther, *Phys. Rev. Lett.* **103**, 072502 (2009).
- [3] A. Wallner, M. Bichler, K. Buczak, R. Dressler, L. K. Fifield, D. Schumann, J. H. Sterba, S. G. Tims, G. Wallner, and W. Kutschera, *Phys. Rev. Lett.* **114**, 041101 (2015).
- [4] G. Korschinek, T. Faestermann, K. Knie, and C. Schmidt, *Radiocarbon* **38**, 68 (1996).
- [5] J. R. Ellis, B. D. Fields, and D. N. Schramm, *Astrophys. J.* **470**, 1227 (1996).
- [6] K. Knie, G. Korschinek, T. Faestermann, C. Wallner, J. Scholten, and W. Hillebrandt, *Phys. Rev. Lett.* **83**, 18 (1999).
- [7] K. Knie, G. Korschinek, T. Faestermann, E. A. Dorfi, G. Rugel, and A. Wallner, *Phys. Rev. Lett.* **93**, 171103 (2004).
- [8] L. Fimiani, D. L. Cook, T. Faestermann, J. M. Gomez-Guzman, K. Hain, G. Herzog, K. Knie, G. Korschinek, P. Ludwig, J. Park, R. C. Reedy, and G. Rugel, *Phys. Rev. Lett.* **116**, 151104 (2016).
- [9] P. Ludwig, S. Bishop, R. Egli, V. Chernenko, B. Deneva, T. Faestermann, N. Famulok, L. Fimiani, J. M. Gomez-Guzman, K. Hain, G. Korschinek, M. Hanzlik, S. Merchel, and G. Rugel, *Proc. Natl. Acad. Sci. U.S.A.* **113**, 9232 (2016).
- [10] A. Wallner, J. Feige, N. Kinoshita, M. Paul, L. K. Fifield, R. Golser, M. Honda, U. Linnemann, H. Matsuzaki, S. Merchel, G. Rugel, S. G. Timms, P. Steier, T. Yamagata, and S. R. Winkler, *Nature (London)* **532**, 69 (2016).
- [11] G. Korschinek and T. Faestermann, *Nucl. Instrum. Methods Phys. Res., Sect. B* **438**, 148 (2019).
- [12] M. Kachelriess, A. Neronov, and D. V. Semikoz, *Phys. Rev. Lett.* **115**, 181103 (2015).
- [13] G. J. Wasserburg, M. Busso, R. Gallino, and K. M. Nolett, *Nucl. Phys.* **777**, 5 (2006).
- [14] M. Honda and M. Imamura, *Phys. Rev. C* **4**, 1182 (1971).
- [15] S. E. Woosley and T. A. Weaver, *Astrophys. J. Suppl. Ser.* **101**, 181 (1995).
- [16] W. Wang, M. J. Harris, R. Diehl, H. Halloin, B. Cordier, K. Kretschmer, P. Jean, G. G. Lichti, J. P. Roques, S. Schanne *et al.*, *Astron. Astrophys.* **469**, 1005 (2007).
- [17] R. Diehl, H. Halloin, K. Kretschmer, A. W. Strong, W. Wang, P. Jean, G. G. Lichti, J. Knödlseher, J. P. Roques, S. Schanne *et al.*, *Nature (London)* **439**, 45 (2006).
- [18] D. Nesvory, P. Jenniskens, H. F. Levison, W. F. Bottke, D. Vokrouhlicy, and M. Gounelle, *Astrophys. J.* **713**, 816 (2010).
- [19] R. Michel, P. Dragovitsch, P. Cloth, G. Dagge, and D. Filges, *Meteoritics*, **26**, 221 (1991).
- [20] R. Michel, I. Leya, and L. Borges, *Nucl. Instrum. Methods Phys. Res., Sect. B* **113**, 434 (1996).
- [21] N. Altobelli, F. Postberg, K. Fiege, M. Trieloff, H. Kimura, V. J. Sterken, H.-W. Hsu, J. Hillier, N. Khawaja, G. Moragas-Klostermeyer, J. Blum, M. Burton, R. Srama, S. Kempf, and E. Gruen, *Science* **352**, 312 (2016).
- [22] M. Poutivtsev, Ph.D. thesis, Technische Universität München, 2007; https://webarchiv.typo3.tum.de/PH/gams/fileadmin/w00bjs/www/publications/2007_Poutivtsev.pdf.
- [23] M. Imamura, T. Inoue, K. Nisiizumi, and S. Tanaka, ^{53}Mn in Deep-sea sediment cores - an indicator of past solar activity, in *16th International Cosmic Ray Conference*, Vol. 2. OG 12-21 (Institute for Cosmic Ray Research, University of Tokyo, Tokyo, 1979), p. 304.
- [24] M. Segl, A. Mangini, G. Bonani, H. J. Hofmann, M. Nessi, M. Suter, W. Wölfli, G. Friedrich, W. L. Plüger, A. Wiechowski, and J. Beer, *Nature (London)* **309**, 540 (1984).
- [25] J. Chmeleff, F. V. Blanckenburg, K. Kossert, and D. Jakob, *Nucl. Instrum. Methods Phys. Res., Sect. B* **268**, 192 (2010).
- [26] G. Korschinek, A. Bergmaier, T. Faestermann, U. C. Gerstmann, K. Knie, G. Rugel, A. Wallner, I. Dillmann, G. Dollinger, Ch. Lierse von Gostomski, K. Kossert, M. Maiti, M. Poutivtsev, and A. Remmert, *Nucl. Instrum. Methods Phys. Res., Sect. B* **268**, 187 (2010).
- [27] R. Bibron, R. Chesselet, G. Crozaz, G. Leger, and E. Picciotto, *Earth Planet. Sci. Lett.* **21**, 109 (1974).
- [28] D. Rodrigues, A. E. Negri, Ch. Balpardo, A. Arazi, T. Faestermann, J. O. Fernandez Niello, L. Fimiani, J. M. Gómez Guzmán, K. Hain, G. Korschinek, P. Ludwig, and G. V. Marti, *Appl. Radiat. Isot.* **140**, 342 (2018).
- [29] M. Poutivtsev, I. Dillmann, T. Faestermann, K. Knie, G. Korschinek, J. Lachner, A. Meier, G. Rugel, and A. Wallner, *Nucl. Instrum. Methods Phys. Res., Sect. B* **268**, 756 (2010).
- [30] D. Ackermann *et al.*, Cruise Report SO 141 „Hula 1”. Institut für Geowissenschaften, Universität Kiel, Vol. 8, 1999.
- [31] E. R. Flueh *et al.*, Cruise report SO 142 „Hula”. Institut für Geowissenschaften, Universität Kiel, 1999.

- [32] G. Friedrich and A. Schmitz-Wiechowsky, *Mar. Geol.* **37**, 71 (1980).
- [33] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.125.031101> for Supernova-produced ^{53}Mn on Earth.
- [34] G. J. Feldman and R. D. Cousins, *Phys. Rev. D* **57**, 3873 (1998).
- [35] C. Fitoussi, G. M. Raisbeck, K. Knie, G. Korschinek, T. Faestermann, S. Goriely, D. Lunney, M. Poutivtsev, G. Rugel, C. Waelbroeck, and A. Wallner, *Phys. Rev. Lett.* **101**, 121101 (2008).
- [36] J. Feige, A. Wallner, R. Altmeyer, L. K. Fifield, R. Golser, S. Merchel, G. Rugel, P. Steier, S. G. Tims, and S. R. Winkler, *Phys. Rev. Lett.* **121**, 221103 (2018).