Mechanics of Metric Frustration in Contorted Filament Bundles: From Local Symmetry to Columnar Elasticity

Daria W. Atkinson^{®*}

Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA

Christian D. Santangelo

Department of Physics, Syracuse University, Syracuse, New York 13210, USA

Gregory M. Grason

Department of Polymer Science and Engineering, University of Massachusetts Amherst, Amherst, Massachusetts 01003, USA

(Received 7 May 2021; revised 17 August 2021; accepted 22 October 2021; published 18 November 2021)

Bundles of filaments are subject to geometric frustration: certain deformations (e.g., bending while twisted) require longitudinal variations in spacing between filaments. While bundles are common—from protein fibers to yarns—the mechanical consequences of longitudinal frustration are unknown. We derive a geometrically nonlinear formalism for bundle mechanics, using a gaugelike symmetry under *reptations* along filament backbones. We relate force balance to orientational geometry and assess the elastic cost of frustration in twisted-toroidal bundles.

DOI: 10.1103/PhysRevLett.127.218002

Elastic zero modes are a ubiquitous feature of soft materials, from mechanical metamaterials [1,2] to liquid crystal elastomers [3]. Such systems can undergo large deformations with minimal strain, as geometrically coupled rotations and translations preserve local spacing between microscopic constituents. The smectic and columnar liquid crystalline phases provide paradigmatic examples of zero modes in soft elastic systems, permitting relative "sliding" of 2D layers and 1D columns, respectively. Smectic and columnar phases with zero-cost sliding displacements bear a striking resemblance to a much broader class of laminated and filamentous structures with largely inelastic, dissipative forces tangent to their layers or columns and without longitudinal correlations between material points, ranging from multilayer graphene materials [4] and stacked paper [5] to biopolymer bundles [6,7], nanotube yarns [8], and wire ropes [9].

While there are well-established frameworks which capture the geometric nonlinearities of smectic liquid crystals (i.e., the strain tensor accurately describes arbitrarily large and complex deformations) [10], no such framework exists for columnar and filamentous materials. The orientations of column backbones impose constraints on interfilament spacing, generating rich modes of geometric frustration without counterpart in smectic liquid crystals. In the simplest nontrivial case of helical bundles, predictions from a minimally nonlinear approximation of columnar elasticity [11] and tomographic analysis of elastic filament bundles [12] show that twist in straight bundles gives rise to nonuniform interfilament stress and spacing in transverse sections [see Fig. 1(a)]. Except for the restrictive

classes of straight, twisted bundles [13] and twist-free developable domains [14,15], bundle textures also generate *longitudinal* frustration, requiring local spacings to vary *along* a bundle [16]. Although deformations that introduce longitudinal frustration are the rule rather than the exception—for example, wire ropes or toroidal biopolymer condensates are both *twisted* and *bent* (e.g., Fig. 1)—existing frameworks of columnar elasticity fail to capture this effect.

In this Letter, we develop a fully geometrically nonlinear Lagrangian elasticity theory of columnar materials, which



FIG. 1. In (a), an equilibrium twisted bundle with straight, untwisted reference configuration, $\Omega R = 1$, and 2D Poisson ratio $\nu = 0.8$, colored by the local pressure $P = -S_{\alpha}^{\alpha}/2$. *Reptation* of the gold filament by σ along its contour, draws the filament (in the direction indicated by the gold arrow) above the planar cross-section of the bundle, but leaves the local separation $d\Delta$ unchanged. In (b), a twisted-toroidal bundle found by bending the same bundle to $\kappa_0 = 0.2/R$ and optimizing interfilament elastic cost.

completely captures the interplay between orientation and both lateral and longitudinal frustration of interfilament spacing. We construct this theory by imposing a gaugelike local symmetry under *reptations*, deformations that slide filaments along their contours without changing the interfilament spacing. The resultant equilibrium equations point to the role geometrical measures of nonequidistance play in bundles' mechanics. Within this framework, we compute the energetic costs of longitudinal frustration in twisted, toroidal bundles, and give evidence that (1) optimal configurations generically incorporate splay and (2) the bending cost that derives from nonuniform compression depends nonmonotonically on pretwist.

To construct the elastic theory, we divide space into points on filament backbones, labeled by two coordinates: **v**, a 2D label of filaments; and *s*, a length coordinate along filaments. Hence, the location of each point in the bundle is described by a function $\mathbf{r}(s, \mathbf{v})$, with $\partial_s \mathbf{r}(s, \mathbf{v})$ parallel to the tangent vector $\mathbf{t} = \partial_s \mathbf{r}/|\partial_s \mathbf{r}|$. Because they lack positional order along their backbone curves, filament bundles and columnar liquid crystals have a family of continuous zero modes, corresponding to *reptations* (Fig. 1),

$$\mathbf{r}'(s,\mathbf{v}) = \mathbf{r}(s+\sigma(s,\mathbf{v}),\mathbf{v}). \tag{1}$$

We assume that changes in local spacing can be described by an energy density function W, that depends only on the deformation gradient [17].

To account for reptation symmetry, we demand that \mathcal{W} depend on a modified deformation gradient, which transforms as a scalar under $\sigma(s, \mathbf{v})$, depends solely on the deformation \mathbf{r} of the material itself, and recovers the wellestablished 2D elasticity of developable bundles (i.e., parallel arrays) [14,15]. Specifically, we construct a covariant derivative $D_I \mathbf{r} = \nabla_I \mathbf{r} - \mathbf{A}_I$, where ∇_I is the usual covariant derivative on tensors in the material space and **r** determines A_I , such that if two configurations, **r** and **r**', are related by Eq. (1), then $D_I \mathbf{r} = D'_I \mathbf{r}'$. In order for $D_I \mathbf{r}$ to be reptation invariant, we must have that $-\mathbf{A}_{I}' + \mathbf{A}_{I} = \nabla_{I} \sigma \partial_{s} \mathbf{r}$. Therefore, $-\mathbf{A}'_{I} + \mathbf{A}_{I} = -(\mathbf{t} \cdot \nabla_{I} \mathbf{r}')\mathbf{t} + (\mathbf{t} \cdot \nabla_{I} \mathbf{r})\mathbf{t}$. To construct an elastic theory of columnar materials, we set $\mathbf{A}_{I} = (\mathbf{t} \cdot \nabla_{I} \mathbf{r}) \mathbf{t}$, which is manifestly reptation invariant, and leads to a deformation gradient that only measures deformations *transverse* to the local backbones,

$$D_I \mathbf{r} \equiv \nabla_I \mathbf{r} - (\mathbf{t} \cdot \nabla_I \mathbf{r}) \mathbf{t}.$$
 (2)

Notably, for two nearby filaments at $\mathbf{r}(\mathbf{v})$ and $\mathbf{r}(\mathbf{v} + d\mathbf{v})$ in material coordinates, the covariant derivative gives the local distance of closest approach $d\mathbf{\Delta} = dv^I D_I \mathbf{r}$, for which $d\mathbf{\Delta} \cdot \mathbf{t} = 0$ [see inset of Fig. 1(a)]. As shown explicitly in the Supplemental Material [18], this covariant deformation gradient captures the standard 2D deformation gradients of developable domains.

From this deformation gradient, we construct an effective metric $g_{IJ}^{\text{eff}} = D_I \mathbf{r} \cdot D_J \mathbf{r}$, which is naturally invariant under rotations of \mathbf{r} , and which encodes the metric inherited by the local 2D section transverse to the filaments in the bundle [21]. Because $D_s \mathbf{r} = 0$, the effective metric only has components for 2×2 block $I, J \neq s$, which we denote using index notation $\alpha, \beta \in \{1, 2\}$. We can now define the Green–Saint Venant strain tensor [17,25,26] as

$$\epsilon_{\alpha\beta} = \frac{1}{2} [D_{\alpha} \mathbf{r} \cdot D_{\beta} \mathbf{r} - g_{\alpha\beta}^{\text{tar}}], \qquad (3)$$

where $g_{\alpha\beta}^{\text{tar}}$ is the target metric corresponding to strain-free state, which for this Letter, we take to be developable with uniform spacing, so $g_{\alpha\beta}^{\text{tar}} = \delta_{\alpha\beta}$. For weak deflections from the uniform parallel state, Eq. (3) reduces to the small-tilt approximation to the nonlinear columnar strain [11,27,28], which captures the lowest-order dependence of spacing on orientation [18].

Assuming that strains are small though deformations may be large, we can treat the elastic energy as quadratic in the strain, so that

$$\mathcal{E}_s = \int dV \mathcal{W}(\epsilon) = \frac{1}{2} \int dV S^{\alpha\beta} \epsilon_{\alpha\beta}, \qquad (4)$$

where $S^{\alpha\beta} = (\partial \mathcal{W} / \partial \epsilon_{\alpha\beta}) = C^{\alpha\beta\gamma\delta} \epsilon_{\gamma\delta}$ is the nominal stress tensor, and $C^{\alpha\beta\gamma\delta}$ is a tensor of elastic constants which depends on both the crystalline symmetries of the underlying columnar order and the target metric, $g_{\alpha\beta}^{\text{tar}}$ [29]. Energetics of columnar materials also include other gauge-invariant costs, including the Frank-Oseen orientational free energy (particularly filament bending) and the cost of local density changes along columns [30]. Here, for clarity, we focus only on the energetics of columnar strain and detail the combined effects of other contributions elsewhere [36]. Given this gauge-invariant formulation of the columnar strain energy, we first illustrate the mechanical effects of orientational geometry on local forces. This follows from the bulk Euler-Lagrange equations of Eq. (4) (see Supplemental Material [18] for a complete derivation),

$$\frac{\delta \mathcal{E}_s}{\delta \mathbf{r}} = -\nabla_\alpha (S^{\alpha\beta} D_\beta \mathbf{r}) + \nabla_s \left(S^{\alpha\beta} D_\beta \mathbf{r} \frac{\mathbf{t} \cdot \nabla_\alpha \mathbf{r}}{|\nabla_s \mathbf{r}|} \right).$$
(5)

The bulk terms represent body forces generated by the columnar strains, which must be balanced by other internal stresses or external forces. To cast them in a more geometrical light, we consider separately the components tangential and perpendicular to \mathbf{t} , F_{\parallel} , and \mathbf{F}_{\perp} , respectively.

Making use of the identity, $\mathbf{t} \cdot \nabla_{\alpha} D_{\beta} \mathbf{r} = -\partial_{\alpha} \mathbf{t} \cdot \partial_{\beta} \mathbf{r}$, the tangential forces can be recast simply as

$$F_{\parallel} = S^{\alpha\beta} h_{\alpha\beta}, \tag{6}$$

where

$$h_{\alpha\beta} = \frac{1}{2} \left[\partial_{\alpha} \mathbf{t} \cdot \partial_{\beta} \mathbf{r} + \partial_{\beta} \mathbf{t} \cdot \partial_{\alpha} \mathbf{r} - \frac{\mathbf{t} \cdot \partial_{\alpha} \mathbf{r}}{|\partial_{s} \mathbf{r}|} \partial_{s} \mathbf{t} \cdot \partial_{\beta} \mathbf{r} - \frac{\mathbf{t} \cdot \partial_{\beta} \mathbf{r}}{|\partial_{s} \mathbf{r}|} \partial_{s} \mathbf{t} \cdot \partial_{\alpha} \mathbf{r} \right], \quad (7)$$

is the convective flow tensor that measures longitudinal variations in interfilament spacing [16]. Just as the second fundamental form measures gradients of a surface's normal vector [37], $h_{\alpha\beta}$ measure the symmetric gradients of **t** in its normal plane (i.e., its trace is the splay of filament tangents).

Here, we see that tangent forces couple nonequidistance to the stress tensor much like the Young-Laplace law couples in-plane stresses to normal forces in curved membranes [25]. This analogy becomes exact for zero twist textures: when $\mathbf{t} \cdot (\nabla \times \mathbf{t}) = 0$, filaments can be described by a set of surfaces normal to \mathbf{t} . In this case, it is possible to choose coordinates so that $\partial_{\alpha} \mathbf{r} \cdot \mathbf{t} = 0$, and tangential forces give the Young-Laplace force normal to each surface, with $h_{\alpha\beta}$ reducing to their second fundamental form. This illustrates the intuitive notion, shown schematically in Fig. 2, that columnar strain generates tangential body forces that push material points *toward* lower-stress locations in the array.

The bulk components of Eq. (5) perpendicular to \mathbf{t} give the transverse force

$$\mathbf{F}_{\perp} = -D_{\alpha}[S^{\alpha\beta}D_{\beta}\mathbf{r}] + D_{s}\left[S^{\alpha\beta}\frac{\mathbf{t}\cdot\partial_{\alpha}\mathbf{r}}{|\partial_{s}\mathbf{r}|}D_{\beta}\mathbf{r}\right].$$
 (8)

This form captures the divergence of stress in the planes perpendicular to backbones. The second term accounts for the corrections arising from material derivatives that lie along the backbone, such as twisted textures, when $\partial_{\alpha} \mathbf{r} \cdot \mathbf{t} \neq 0$. Thus, the longitudinal derivatives in \mathbf{F}_{\perp} are needed to capture transverse mechanics of even equidistant twisted bundles beyond the lowest-order geometric nonlinearity [11].

We now illustrate the energetics of longitudinal frustration by considering a prototypical nonequidistant geometry: twisted-toroidal bundles. Motivated in large part by the morphologies of condensed biopolymers [6,7],



FIG. 2. A 2D schematic of local forces in a filament bundle (with a uniform, parallel reference state), illustrating transverse stress S_{\perp}^{\perp} with regions under compression in blue and those under extension in red. The forces parallel and tangent to **t** are shown at two points, and push material points to regions of vanishing stress.

theoretical models of twisted toroids have focused on their orientational elasticity costs [38–40], ignoring the unavoidable frustration of interfilament spacing in this geometry. While satisfying force balance in nonequidistant bundles requires physical ingredients beyond the columnar strain energy, which we consider elsewhere [36], for the purposes of this Letter we take advantage of the full geometric nonlinearity of Eq. (5) to explore the specific costs of longitudinal gradients in spacing required by simultaneous twist and bend.

We construct twisted toroids from equilibrium twisted helical bundles of radius *R* and constant pitch, $2\pi/\Omega$ by bending them such that their central curve \mathbf{r}_0 is deformed from a straight line into a circle of radius κ_0^{-1} [see Fig. 1(b)]. We then define perturbative displacements $\mathbf{r}(s + \delta s, \rho + \delta \rho)$ $\phi + \delta \phi) \simeq \mathbf{r}_0 + \rho \hat{\rho} + \partial_s \mathbf{r}^{(0)} \delta s + \delta \rho \hat{\rho} + \delta \phi \hat{\phi}$ relative to the bent, pretwisted bundles, where ρ and ϕ describe the (Eulerian) distance from the central curve and the angular position relative to its normal in the plane perpendicular to its tangent \mathbf{t}_0 , and where $\partial_s \mathbf{r}^{(0)} = \mathbf{t}_0 + \Omega \rho \hat{\phi}$. The small- ρ limit of the force balance equations for the strain energy motivates the following displacements

$$\begin{pmatrix} \delta s \\ \delta \rho \\ \delta \phi \end{pmatrix} = \begin{pmatrix} a_s \Omega \kappa_0 \rho^3 \sin \phi \\ a_\rho \Omega^2 \kappa_0 \rho^4 \cos \phi \\ a_\phi \Omega^2 \kappa_0 \rho^3 \sin \phi \end{pmatrix}, \tag{9}$$

where a_s , a_ρ , and a_ϕ are variational parameters. Notably, to linear order in curvature, these parametrize the "almost equidistant" *Ansätze* considered previously, including splay-free [tr(h) = 0] [38] configurations and det(h) = 0 [16] *Ansätze*.

We expand the energy to quadratic order in κ_0 , holding the center of area at \mathbf{r}_0 , which constrains $a_s = a_\rho - a_\phi$, then minimize Eq. (4) with respect to the displacements for a given ΩR . Examples of the distribution of pressure $P = -S_{\alpha}^{\alpha}/2$ are shown in Fig. 1(b). Relative to the axisymmetric pressure induced by helical twist in the straight bundle, bending into a twisted toroid requires bunching (spreading) of the filaments at the inner (outer) positions in the toroid, leading to a polarization of the pressure toward the normal.

Because bending and twisting of bundles introduces longitudinal strain variation, bending the pretwisted bundle introduces additional stresses whose elastic cost can be characterized by an effective bending stiffness *B*, defined by $E_{\text{bend}} = (B/2) \int ds\kappa_0^2$, which derives purely from columnar strain, rather than intrafilament deformations. Figure 3 shows that longitudinal frustration leads to a bending cost that increases with small twist as $B \sim (\Omega R)^4$, but eventually gives way to remarkable nonmonotonic behavior at large pretwist. We note further that the bending cost grows with the 2D Poisson ratio ν of the columnar array, highlighting the importance of local compressional deformations in optimal twisted toroids.



FIG. 3. The effective bending modulus *B* which results from the frustration of constant spacing in twisted-toroidal bundles. The inset shows dimensionless measures of mean splay (Ψ) and biaxial splay (Γ) defined in the text, for fixed twist $\Omega R = 0.8$ for a range of 2D Poisson ratios, from $\nu = 0.2$ to 1.0. Approaching incompressibility ($\nu \rightarrow 1$), Ψ goes to zero, maintaining uniform areaper-filament at the expense of markedly increased strain energy.

We analyze the optimal modes of deformation via the convective flow tensor, in particular, the trace tr(h) (splay) and deviatoric components $h^{\text{dev}} = h_{\alpha\beta} - \text{tr}(h)\delta_{\alpha\beta}/2$ (biaxial splay) [41], which characterize longitudinal gradients of dilatory and shear stress in the columnar array. In contrast to a heuristic view that optimal packings should favor the uniform area per filament of splay-free textures, the inset of Fig. 3 instead shows that optimal toroids incorporate a mixture of both splay and biaxial splay where we define the respective measures of average splay and biaxial splay, $\Psi \equiv (1/\kappa_0^2 V) \int dV \operatorname{tr}(h)^2$ and $\Gamma \equiv (1/\kappa_0^2 V) \int dV \operatorname{tr}(h_{\text{dev}}^2)$. Only in the incompressible limit, as $\nu \rightarrow 1$, does the splay vanish, and only at the expense of additional biaxial splay and energetic cost, implying counterintuitively that splayed textures are in fact energetically favorable in longitudinally frustrated twisted toroids. Indeed, the energetic preference for splay in nonequidistant bundles can be traced to force balance conditions in this geometry [36].

As an elastic theory, our treatment of reptation invariant energetics neglects the nonconservative forces of tangential friction in 2D ordered bundles. Such a description will apply, by definition, to equilibria of any system with columnar order (i.e., without long-range interaxial correlations between filaments), conditions widely encountered in a range of molecular bundle-forming systems, from supermolecular fibers [42,43] to nanotube ropes [44–46]. For example, simulations of nanotube ropes [44,45], confirm that interfilament cohesion is sufficient to maintain dense, 2D order while columns reptate under highly contorted, twisted, and bent deformations. Condensed states of DNA and other polynucleotides, for example, exhibit columnar states [47], and notably exhibit twisted, contorted, and 2D ordered packings within viral capsids [7,48], although the existence of interhelical correlations in such structure and their effects on observed packing remains a matter of open investigation [49]. The question of when the reptation symmetric elastic description captures geometrically nonlinear deformations of macroscopic bundles, in which static friction between filaments may transmit axial shears, is more complex. While some recent experiments [12,50] with macroscopic bundles of densely packed, 2D ordered, and twisted bundles demonstrate that large degrees of distortion are possible without noticeable effects of sliding friction, prevailing theoretical models, backed up by finite-element simulations, of twisted cables, indicate that the effects of static friction lead to mechanical transitions between shear coupled and uncoupled regimes [51,52]. Notably, models of bending in pretwisted cables show that above a threshold bending (which increases with static friction and pretension), deformation transitions to a slipping regime [53,54]. In this context, we might speculate that deformations in the postslipping regimes of these models tends toward those captured by the minimal reptation-invariant elasticity. It may indeed be possible to explore this conjecture via a homogenization approach to a suitable constitutive model of fibered elastic composites [55-58], in which the interfiber matrix transmits strictly no elastic shears, but instead only an elastoplastic slipping of fibers determined by local transverse pressure (i.e., Coulomb friction).

In summary, we have shown that gauge-theoretic principles underlie the geometrically nonlinear theory of columnar elasticity, providing a means to quantify the cost of longitudinal frustration in the mechanics of bundles. Unlike phase field models of nonlinear elasticity (such as [59,60]), this description neither depends on the presence of a planar reference state, nor presupposes uniform crystalline order, allowing us to both accommodate the effective curvature of bundles of constant pitch helices [61], and provide a natural generalization to arbitrary target metrics [25]. Because this approach to elasticity relies only on the existence of local, continuous zero modes, we note that it can be generalized to other liquid crystals, like smectics, and anticipate that it may have applications beyond liquid crystals, including mechanical metamaterials.

The authors gratefully acknowledge useful discussions with R. Kusner, S. Zhou, and B. Davidovitch. M. Dimitriyev, D. Hall, and R. Kamien provided feedback on an early draft of this manuscript. This work was supported by the NSF under Grants No. DMR-1608862 and No. NSF DMR-1822638; D. A. received additional funding from the Penn Provost's postdoctoral fellowship and NSF MRSEC DMR-1720530.

atkinsod@sas.upenn.edu she/her or they/them

^[1] X. Mao, N. Xu, and T. C. Lubensky, Phys. Rev. Lett. 104, 085504 (2010).

^[2] K. Sun, A. Souslov, X. Mao, and T. C. Lubensky, Proc. Natl. Acad. Sci. U.S.A. 109, 12369 (2012).

- [3] M. Warner, P. Bladon, and E. M. Terentjev, J. Phys. II 4, 93 (1994).
- [4] Z. Xu, H. Sun, X. Zhao, and C. Gao, Adv. Mater. 25, 188 (2013).
- [5] S. Poincloux, T. Chen, B. Audoly, and P. Reis, Phys. Rev. Lett. 126, 218004 (2021).
- [6] N. V. Hud and K. H. Downing, Proc. Natl. Acad. Sci. U.S.A. 98, 14925 (2001).
- [7] A. Leforestier and F. Livolant, Proc. Natl. Acad. Sci. U.S.A. 106, 9157 (2009).
- [8] M. Zhang, K. R. Atkinson, and R. H. Baughman, Science 306, 1358 (2004).
- [9] G. A. Costello, *Theory of Wire Rope*, Mechanical Engineering Series (Springer-Verlag, New York, 1990).
- [10] G. Grinstein and R. A. Pelcovits, Phys. Rev. A 26, 915 (1982).
- [11] G. M. Grason, Phys. Rev. E 85, 031603 (2012).
- [12] A. Panaitescu, G. M. Grason, and A. Kudrolli, Phys. Rev. E 95, 052503 (2017).
- [13] G. M. Grason, Rev. Mod. Phys. 87, 401 (2015).
- [14] Y. Bouligand, J. Phys. 41, 1297 (1980).
- [15] M. Kléman, J. Phys. 41, 737 (1980).
- [16] D. W. Atkinson, C. D. Santangelo, and G. M. Grason, New J. Phys. 21, 062001 (2019).
- [17] P.G. Ciarlet, J. Elast. 78–79, 1 (2005).
- [18] See Supplemental Material at http://link.aps.org/supplemental/ 10.1103/PhysRevLett.127.218002 for details on the deformation gradient, force balence and energetic calculations for bending of twisted bundles, which includes [19,20].
- [19] M. Kléman and P. Oswald, J. Phys. 43, 655 (1982).
- [20] G. M. Grason, Soft Matter 9, 6761 (2013).
- [21] Equidistant bundles—where components of g_{IJ}^{eff} are uniform along *s*—are known as Riemannian fibrations, see, e.g., [22,23], and the transverse structure can be mapped on to single 2D base space. Nonequidistant bundles correspond to sub-Riemannian generalizations [24] where the inherited metric measures distances in the two planes perpendicular to **t** at each point.
- [22] B. O'Neill, Mich. Math. J. 13, 459 (1966).
- [23] D. Gromoll and G. Walschap, Submersions, foliations, and metrics, in *Metric Foliations and Curvature* (Birkhäuser Basel, Basel, 2009), pp. 1–44.
- [24] A. Agrachev, D. Barilari, and U. Boscain, A Comprehensive Introduction to Sub-Riemannian Geometry, Cambridge Studies in Advanced Mathematics (Cambridge University Press, Cambridge, England, 2019).
- [25] E. Efrati, E. Sharon, and R. Kupferman, J. Mech. Phys. Solids 57, 762 (2009).
- [26] M. A. Dias, J. A. Hanna, and C. D. Santangelo, Phys. Rev. E 84, 036603 (2011).
- [27] J. V. Selinger and R. F. Bruinsma, Phys. Rev. A 43, 2910 (1991).
- [28] G. M. Grason and R. F. Bruinsma, Phys. Rev. Lett. 99, 098101 (2007).
- [29] We raise and lower indices with $g^{\alpha\beta tar}$, the inverse of $g^{tar}_{\alpha\beta}$, rather than $g^{\alpha\beta eff}$, as a matter of convenience, and note that the difference in the elastic energy is higher order in the strain tensor [26].
- [30] The symmetry under local reptations in this theory is unlike a related treatment of both the nematic to smectic-*A* transition

[31,32] and its columnar analog [33–35] in which the gauge symmetry of a density wave is explicitly broken by, e.g., the splay elasticity of the underlying nematic order.

- [31] P.G. de Gennes, Solid State Commun. 10, 753 (1972).
- [32] T. C. Lubensky, G. Grinstein, and R. A. Pelcovits, Phys. Rev. B 25, 6022 (1982).
- [33] C. Giannessi, Phys. Rev. A 34, 705 (1986).
- [34] R. D. Kamien and D. R. Nelson, Phys. Rev. Lett. 74, 2499 (1995).
- [35] R. D. Kamien, J. Phys. II 6, 461 (1996).
- [36] D. W. Atkinson, C. D. Santangelo, and G. M. Grason (to be published).
- [37] R. S. Millman and G. D. Parker, *Elements of Differential Geometry* (Prentice-Hall, Englewood Cliffs, NJ, 1977).
- [38] I. M. Kulić, D. Andrienko, and M. Deserno, Europhys. Lett. 67, 418 (2004).
- [39] J. Charvolin and J. F. Sadoc, Eur. Phys. J. E 25, 335 (2008).
- [40] V. Koning, B. C. v. Zuiden, R. D. Kamien, and V. Vitelli, Soft Matter 10, 4192 (2014).
- [41] J. V. Selinger, Liq. Cryst. Rev. 6, 129 (2018).
- [42] T. Kato, T. Yasuda, Y. Kamikawa, and M. Yoshio, Chem. Commun. 729 (2009).
- [43] K. Yano, Y. Itoh, F. Araoka, G. Watanabe, T. Hikima, and T. Aida, Science 363, 161 (2019).
- [44] I. Ostanin, R. Ballarini, D. Potyondy, and T. Dumitric, J. Mech. Phys. Solids 61, 762 (2013).
- [45] I. Ostanin, R. Ballarini, and T. Dumitrică, J. Appl. Mech. 81, 061004 (2014).
- [46] Y. Wang, I. Ostanin, C. Gaidu, and T. Dumitric, Langmuir 31, 12323 (2015).
- [47] F. Livolant and A. Leforestier, Prog. Polym. Sci. 21, 1115 (1996), liquid Crystalline Polymers-Part 2.
- [48] S. L. Ilca, X. Sun, K. E. Omari, A. Kotecha, F. d. Haas, F. DiMaio, J. M. Grimes, D. I. Stuart, M. M. Poranen, and J. T. Huiskonen, Nature (London) 570, 252 (2019).
- [49] L. Barberi, F. Livolant, A. Leforestier, and M. Lenz, Nucleic Acids Res. 49, 3709 (2021).
- [50] A. Panaitescu, G. M. Grason, and A. Kudrolli, Phys. Rev. Lett. **120**, 248002 (2018).
- [51] D. Liu, S. Zheng, and Y. He, Math. Mech. Solids 24, 2154 (2019).
- [52] E. A. W. de Menezes and R. J. Marczak, Engineering structures 241, 112487 (2021).
- [53] K.-J. Hong, A. Der Kiureghian, and J. L. Sackman, J. Eng. Mech. 131, 500 (2005).
- [54] D. Zhang and M. Ostoja-Starzewski, J. Appl. Mech. 83, 031003 (2016).
- [55] K. D. Cherednichenko, V. P. Smyshlyaev, and V. V. Zhikov, Proc. R. Soc. Edinburgh, Sect. A 136, 87 (2006).
- [56] M. Bellieud, SIAM J. Math. Anal. 49, 2615 (2017).
- [57] D. Engl, C. Kreisbeck, and A. Ritorto, arXiv:2105.03971.
- [58] E. Davoli and C. Kreisbeck, arXiv:2108.00038.
- [59] P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, Cambridge, England, 1995).
- [60] O. Stenull and T. C. Lubensky (private communication).
- [61] I. R. Bruss and G. M. Grason, Proc. Natl. Acad. Sci. U.S.A. 109, 10781 (2012).