

Bose-Einstein Condensation of Efimovian Triples in the Unitary Bose GasS. Musolino ^{1,*}, H. Kurkjian ², M. Van Regemortel ³, M. Wouters ⁴, S. J. J. M. F. Kokkelmans ¹ and V. E. Colussi ^{5,†}¹*Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, Netherlands*²*Laboratoire de Physique Théorique, Université de Toulouse, CNRS, UPS, 31400 Toulouse, France*³*Joint Quantum Institute and The Institute for Research in Electronics and Applied Physics, University of Maryland, College Park, Maryland 20742, USA*⁴*TQC, Universiteit Antwerpen, Universiteitsplein 1, B-2610 Antwerp, Belgium*⁵*INO-CNR BEC Center and Dipartimento di Fisica, Università di Trento, 38123 Povo, Italy*

(Received 18 June 2021; revised 4 October 2021; accepted 1 December 2021; published 10 January 2022)

In an atomic Bose-Einstein condensate quenched to the unitary regime, we predict the sequential formation of a significant fraction of condensed pairs and triples. At short distances, we demonstrate the two-body and Efimovian character of the condensed pairs and triples, respectively. As the system evolves, their size becomes comparable to the interparticle distance, such that many-body effects become significant. The structure of the condensed triples depends on the size of Efimov states compared with density scales. Unexpectedly, we find universal condensed triples in the limit where these scales are well separated. Our findings provide a new framework for understanding dynamics in the unitary regime as the Bose-Einstein condensation of few-body composites.

DOI: [10.1103/PhysRevLett.128.020401](https://doi.org/10.1103/PhysRevLett.128.020401)

Introduction.—Bose-Einstein condensation (BEC) gives rise to such spectacular manifestations of quantum statistics as superfluidity, superconductivity, and supersolidity [1–4]. The paradigmatic theories of Bogoliubov and Bardeen-Cooper-Schrieffer (BCS) describe BECs of weakly coupled bosons and fermionic pairs, respectively, and have been applied in many fields of physics [4–10]. Here, the quantum statistics of the medium alters one-body dynamics, producing quasiparticles, and two-body dynamics, producing Cooper pairs, with the latter persisting even in the absence of a two-body bound state. These guiding concepts however must be reconsidered when describing strongly interacting systems such as liquid helium [11], ultracold gases [12], nuclear matter [13–16], and strongly coupled polarons [17–22]. The occurrence of a richer few-body physics including three-body bound Efimov states amongst these strongly interacting bosons and multicomponent fermions [23,24] necessitates the key shifting of many-body paradigms from two- to three-body correlations. Here, the fundamental question reemerges of whether bound states in vacuum (polymers: dimers, trimers, etc.) can be bound by the medium and converted into condensed few-body composites (n -tuples: pairs, triples, etc.) possessing long-range order.

Recently, the versatility of ultracold atomic platforms was utilized to shed new light on these open problems. Despite strong three-body losses, quasiequilibrated states were achieved in single-component Bose gases quenched to the unitary regime $n|a|^3 \gg 1$, with n the atomic density and a the s -wave scattering length [25–28]. Specifically, a macroscopic population of Efimov trimers was reported in

Ref. [26], following a second sweep of interactions to weak interactions. Historically, this technique was used to measure the condensation of Cooper pairs in the BCS-BEC crossover via their conversion into weakly bound dimers [29,30]. It is thus natural to ask whether the molecules measured in Ref. [26] reveal the existence of few-body condensates of pairs and triples in the unitary Bose gas. It is unknown whether the hypothesized universality of the medium [31], parametrized by the density (Fermi) scales $k_n = (6\pi^2 n)^{1/3}$, $E_n = \hbar^2 k_n^2 / 2m$, and $t_n = \hbar / E_n$, produces universal pairs and triples, or conversely whether a (non-universal) sensitivity to the Efimov effect and finite-range physics is preserved at the many-body level. Answering this question in such a nonequilibrium and strongly interacting quenched system requires a model both ergodic [32] and nonperturbative [33–37], which recovers the vacuum three-body spectrum [38–41]. Although widely used in statistical physics [42–44], the cumulant model was recently adapted to quantum gases and found to fulfill these requirements [45,46].

In this Letter, we study a uniform BEC quenched to the unitary regime and develop a general theory of simultaneous atomic, pair, and triple condensation in strongly interacting systems possessing the Efimov effect. Within the cumulant model, we construct generalized condensate wave functions and predict significant pair and triple condensation and associated off-diagonal long-range ordering (ODLRO) occurring between depleted atoms. We show that the Efimovian character of the triples is guaranteed at short distances; however at later times triples have a size comparable to the interparticle spacing. Remarkably even

when Fermi and Efimovian scales are well separated, medium effects lead to the persistent production of triple and pair BECs with universal populations and internal structures.

Model.—We model the system of N spinless bosons in a cubic volume V using a single-channel Hamiltonian with pairwise s -wave interactions

$$\hat{H} = \int d^3r \hat{\psi}^\dagger(\mathbf{r}) \left(-\frac{\hbar^2}{2m} \Delta_{\mathbf{r}} \right) \hat{\psi}(\mathbf{r}) + \frac{1}{2} \int d^3r d^3r' \hat{\psi}^\dagger(\mathbf{r}) \hat{\psi}^\dagger(\mathbf{r}') V(|\mathbf{r} - \mathbf{r}'|) \hat{\psi}(\mathbf{r}') \hat{\psi}(\mathbf{r}), \quad (1)$$

where $\hat{\psi}(\mathbf{r}) = (1/\sqrt{V}) \sum_{\mathbf{k}} \hat{a}_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}}$ are field operators and $\hat{a}_{\mathbf{k}}$ annihilates a boson of momentum $\hbar\mathbf{k}$. At unitarity ($|a| \rightarrow \infty$), the actual potential can be replaced by a simpler nonlocal separable potential $\hat{V} = g|\zeta\rangle\langle\zeta|$ with s -wave form factors $\langle\mathbf{k}|\zeta\rangle = \theta(\Lambda - |\mathbf{k}|)$, interaction strength $g = -\pi^3 \hbar^2 \bar{a}/m$, and cutoff $\Lambda = 2/\pi\bar{a}$, where $\bar{a} = 0.955 r_{\text{vdW}}$ is the mean scattering length and r_{vdW} is the van der Waals length for a particular atomic species [38,47–49]. This sets the three-body parameter $\kappa_* r_{\text{vdW}} \approx 0.211$, which is the wave number of the ground-state Efimov trimer at unitarity (see [37,38,45,50]).

We model the postquench many-body dynamics of an initially pure, noninteracting atomic condensate using the method of cumulants whose hierarchical structure reflects the sequential growth of intrinsically higher-order correlations [43,45,46,67–70]. Within the U(1) symmetry-breaking picture, we study the dynamics of the singlet $\langle\hat{a}_{\mathbf{k}}\rangle = \delta_{\mathbf{k}0} \sqrt{V} \psi_0$, which describes the atomic BEC in the $k = 0$ mode. In the frame rotating with the condensate phase θ_0 , we study also the doublets

$$n_{\mathbf{k}} = \langle\hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}}\rangle, \quad c_{\mathbf{k}} = e^{-2i\theta_0} \langle\hat{a}_{-\mathbf{k}} \hat{a}_{\mathbf{k}}\rangle, \quad (2)$$

describing the single-particle momentum distribution and pairing, respectively, and the triplets

$$M_{\mathbf{k},\mathbf{q}} = e^{i\theta_0} \langle\hat{a}_{\mathbf{k}-\mathbf{q}}^\dagger \hat{a}_{\mathbf{q}}^\dagger \hat{a}_{\mathbf{k}}\rangle, \quad R_{\mathbf{k},\mathbf{q}} = e^{-3i\theta_0} \langle\hat{a}_{\mathbf{q}-\mathbf{k}} \hat{a}_{\mathbf{k}} \hat{a}_{-\mathbf{q}}\rangle, \quad (3)$$

introducing ergodic processes and the Efimov effect [32,38,45,71]. Truncating the cumulant hierarchy can be justified at early times due to the sequential nature of correlation buildup [45,50]. However the increasing importance of quadruplets, in particular for energy conservation, limits our study to times $t \lesssim t_n$.

Off-diagonal long-range ordering.—The triplet model contains anomalous averages at the one-body level (ψ_0) in the atomic condensate and at the two- ($c_{\mathbf{k}}$) and three-body ($R_{\mathbf{k},\mathbf{q}}$) levels *within* the quantum depletion. These cumulants are intimately connected to the eigenfunctions of the reduced density matrices, signaling ODLRO and

condensation [72–74]. We begin from the spectral decomposition of the one-body density matrix

$$\rho^{(1)}(\mathbf{r}_1, \mathbf{r}_2; t) = \langle\hat{\psi}^\dagger(\mathbf{r}_2) \hat{\psi}(\mathbf{r}_1)\rangle = \sum_{\nu} N_{\nu}(t) \varphi_{\nu}^*(\mathbf{r}_2, t) \varphi_{\nu}(\mathbf{r}_1, t), \quad (4)$$

where φ_{ν} are orthogonal one-body eigenstates. Only one eigenvalue $N_{\nu=0}$ is assumed to be macroscopic such that φ_0 is responsible for ODLRO at the one-body level:

$$\lim_{|\mathbf{r}_1 - \mathbf{r}_2| \rightarrow \infty} \rho^{(1)}(\mathbf{r}_1, \mathbf{r}_2; t) = N_0(t) \varphi_0^*(\mathbf{r}_2, t) \varphi_0(\mathbf{r}_1, t). \quad (5)$$

Within the cumulant approach, the long-range part of $\rho^{(1)}$ is simply $|\psi_0|^2$, such that $N_0 = V|\psi_0|^2$ coincides with the condensate population $\langle\hat{a}_0^\dagger \hat{a}_0\rangle$ and fraction $n_0 = N_0/V$.

In the presence of one-body condensation, ODLRO occurs trivially at all higher orders [69,74]. We isolate therefore the atomic condensate from the fluctuations $\hat{\psi}(\mathbf{r}) = \psi_0 + \delta\hat{\psi}(\mathbf{r})$, satisfying $\langle\delta\hat{\psi}(\mathbf{r})\rangle = 0$. To study intrinsically few-body ODLRO amongst fluctuations, we adapt the treatment of Yang [74] and spectrally decompose the corresponding p -body density matrices

$$\langle\delta\hat{\psi}^\dagger(\mathbf{r}'_1) \dots \delta\hat{\psi}^\dagger(\mathbf{r}'_p) \delta\hat{\psi}(\mathbf{r}_p) \dots \delta\hat{\psi}(\mathbf{r}_1)\rangle = \sum_{\nu} N_{\nu}^{(p)}(t) \varphi_{\nu}^{(p)*}(\mathbf{r}'_1, \dots, \mathbf{r}'_p, t) \varphi_{\nu}^{(p)}(\mathbf{r}_1, \dots, \mathbf{r}_p, t), \quad (6)$$

where $\varphi_{\nu}^{(p)}$ and $N_{\nu}^{(p)}$ are the orthogonal p -body eigenstates and eigenvalues, respectively. Analogous to Eq. (5), when the p -body density matrix is nonzero in the long-range limit $|\sum_{i=1}^p \mathbf{r}_i - \mathbf{r}'_i|/p \rightarrow \infty$, there exists intrinsic p -body ODLRO. In the triplet model, this limit is dominated by the anomalous contraction $\langle\delta\hat{\psi}^\dagger \dots \delta\hat{\psi}^\dagger\rangle \langle\delta\hat{\psi} \dots \delta\hat{\psi}\rangle$, such that nonzero c or R cumulants generate ODLRO. The associated normalized pair and triple wave functions are

$$\varphi_0^{(2)}(\mathbf{r}, t) = \frac{c(\mathbf{r}, t)}{\sqrt{N_0^{(2)}(t)}}, \quad \varphi_0^{(3)}(\mathbf{r}, \boldsymbol{\rho}, t) = \frac{R(\mathbf{r}, \boldsymbol{\rho}, t)}{\sqrt{N_0^{(3)}(t)}}, \quad (7)$$

with $N_0^{(2)} = \sum_{\mathbf{k}} |c_{\mathbf{k}}|^2$, $N_0^{(3)} = \sum_{\mathbf{k}, \mathbf{q}} |R_{\mathbf{k}, \mathbf{q}}|^2$, and Jacobi vectors $\mathbf{r} \equiv \mathbf{r}_1 - \mathbf{r}_2$, $\boldsymbol{\rho} = \mathbf{r}_3 - (\mathbf{r}_1 + \mathbf{r}_2)/2$. We note that pair or triple condensation may generate trivial ODLRO in density matrices with $p \geq 4$. If one were to study, e.g., quadruple condensation, such contributions should be removed.

Condensate fractions.—Unlike one-body condensation, the macroscopic eigenvalues $N_0^{(p)}$ cannot be directly related to condensed fractions. To understand this, we construct composite operators annihilating condensed pairs and triples

$$\hat{b}_0^{(p)} = \frac{1}{\sqrt{p!}} \left[\prod_{i=1}^p \int d^3 r_i \delta\hat{\psi}(\mathbf{r}_i) \right] \varphi_0^{(p)*}(\mathbf{r}_1, \dots, \mathbf{r}_p). \quad (8)$$

Evaluating the quantum average of the commutators in the triplet model gives

$$\langle [\hat{b}_0^{(2)}, \hat{b}_0^{(2)\dagger}] \rangle = 1 + \frac{2}{N_0^{(2)}} \sum_{\mathbf{k}} |c_{\mathbf{k}}|^2 n_{\mathbf{k}}, \quad (9)$$

$$\langle [\hat{b}_0^{(3)}, \hat{b}_0^{(3)\dagger}] \rangle = 1 + \frac{3}{N_0^{(3)}} \sum_{\mathbf{k}, \mathbf{q}} |R_{\mathbf{k}, \mathbf{q}}|^2 n_{\mathbf{k}} (1 + n_{\mathbf{q}}), \quad (10)$$

which approximate the canonical relations only for weak excitations ($n_{\mathbf{k}} \ll 1$) or localized pairs and triples relative to the medium. In the opposite limit, composite bosons are created on top of densely populated Fourier modes, leading to Bose enhancement of atoms within the created composite and an overestimation of the condensed fraction [75]. Consequently, the rapid quantum depletion of the atomic condensate in the unitary regime yields $\hat{b}_0^{(p)}$ that are *approximately* bosonic only at $t \lesssim t_n$ (see the Supplemental Material [50]). The renormalization procedure,

$$\hat{B}_0^{(p)} = \frac{\hat{b}_0^{(p)}}{\sqrt{\langle [\hat{b}_0^{(p)}, \hat{b}_0^{(p)\dagger}] \rangle}}, \quad (11)$$

effectively ensures that the bosonic canonical relations are preserved *on average* at all times, which we use to compute the pair and triple condensate fractions as $n_0^{(p)}/n = \langle \hat{B}_0^{(p)\dagger} \hat{B}_0^{(p)} \rangle / (N/p)$.

The postquench dynamics of the condensate fractions are shown in Fig. 1. Compared with the doublet model, three-body processes in the triplet model lead to an accelerated depletion of the atomic condensate, reaching $n_0/n \approx 0.4$ by $t = t_n$. At early times, the formation of condensed triples follows the universal pair condensate growth sequentially, reflecting the hierarchical structure of the cumulant equations of motion (see Refs. [38,50]). At later times the dynamics depend strongly on the density regime, repeating log periodically with the density typical of the Efimov effect [39–41]. In the trimer regime [Fig. 1(a)], the ground-state Efimov trimer resonantly overlaps with the scale set by the density ($k_n/\kappa_* \sim 1$), and triple condensation dominates clearly at later times, becoming comparable to the atomic condensate fraction. Condensed atoms can be converted to pairs and triples via low energy two- and three-body scattering, respectively [71,76,77]. As this overlap becomes less resonant ($k_n/\kappa_* \lesssim 1$), the system enters the crossover regime where particle-number oscillations between pair and triple BECs visible in Fig. 1(b) are analogous to the atom-dimer coherences of Ref. [78]. In the universal regime [Fig. 1(c)], Efimovian and Fermi scales are well separated ($k_n/\kappa_* \ll 1$), and the oscillation




FIG. 1. Atomic (blue), pair (orange), and triple (green) condensate fraction dynamics in the (a) trimer, (b) crossover, and (c) universal density regimes. The gray dot-dashed lines indicate universal results in the doublet (Hartree-Fock Bogoliubov [80]) model. The comparison of Efimovian (κ_*) and Fermi (k_n) scales strongly determines the relative populations of the condensed few-body composites.

becomes increasingly faster relative to t_n . This is the characteristic dynamical signature of the Efimov effect [38,40,41,79]. At later times, pair condensation remains dominant while the nonuniversal oscillations fade and the condensate fractions converge universally, approaching $n_0^{(2)}/n \approx 0.2$ and $n_0^{(3)}/n \approx 0.1$ by $t = t_n$.

Short-range expansions.—We study now how the short-range behavior of the condensate wave functions $c(\mathbf{r}, t)$ and $R(\mathbf{r}, \boldsymbol{\rho}, t)$ are dictated by few-body physics. This can be understood from the corresponding cumulant equations of motion which are identical to few-body Schrödinger equations at large momenta compared with the many-body scales [1,38,45,50]. For distances larger than the short range of the potential ($r_{\text{vdW}} < r < k_n^{-1}, a$), the pair and triple condensate wave functions can be expanded in terms of the zero-energy few-body scattering wave functions

$$c(\mathbf{r}, t) \underset{r \rightarrow 0}{\sim} \frac{1}{4\pi} \Psi_0^{(2)}(t) \phi(\mathbf{r}), \quad (12)$$

$$R(\mathbf{r}, \boldsymbol{\rho}, t) \underset{R \rightarrow 0}{\sim} \frac{2^{3/2}}{3^{1/4} s_0} \Psi_0^{(3)}(t) \Phi(R, \boldsymbol{\Omega}), \quad (13)$$

which define the macroscopic order parameters $\Psi_0^{(p)}$ (see the Supplemental Material [50]). Here $\phi(\mathbf{r}) = 1/r - 1/a$ is the zero-energy two-body scattering state, and

$$\Phi(R, \boldsymbol{\Omega}) = \frac{1}{R^2} \sin \left[s_0 \log \frac{R}{R_t} \right] \frac{\phi_{is_0}(\boldsymbol{\Omega})}{\sqrt{\langle \phi_{is_0} | \phi_{is_0} \rangle}} \quad (14)$$

is the zero-energy three-body scattering state for hyper-radius $R = \sqrt{r^2/2 + 2\rho^2/3}$ and hyperangles $\mathbf{\Omega} = \{\hat{\rho}, \hat{\mathbf{r}}, \alpha = \arctan(r/\rho)\}$ [81]. Here, $s_0 \approx 1.00624$, $R_t = \sqrt{2} \exp\{\text{Im} \ln[\Gamma(1 + is_0)]/s_0\}/\kappa_*$, and Γ is the gamma function. The hyperangular function describing s -wave pairwise scatterings is $\phi_s(\mathbf{\Omega}) = (1 + \hat{P}_{13} + \hat{P}_{23}) \sin[s(\pi/2 - \alpha)]/\sin(2\alpha)\sqrt{4\pi}$ where \hat{P}_{ij} swaps particles i and j [82]. From Eqs. (12) and (13) we see then at unitarity that the pairs have a universal behavior $\sim 1/r$ at short distances, whereas the triples have an Efimovian character, diverging as $1/R^2$ and oscillating log periodically with the three-body parameter.

At short distances, the total probability to measure clustered pairs and triples is encoded in the two- and three-body contact densities \mathcal{C}_2 and \mathcal{C}_3 , respectively, central to a set of universal relations between system properties [81,83]. In the presence of pair and triple condensation, these clusters can be divided into contributions from the order parameters and higher-order cumulants

$$\mathcal{C}_2 = \frac{m^2 g^2}{\hbar^4} \langle (\hat{\psi}^\dagger)^2 \hat{\psi}^2 \rangle = |\Psi_0^{(2)}|^2 + \delta\mathcal{C}_2, \quad (15)$$

$$\mathcal{C}_3 = -\frac{m^2 g^2}{2\hbar^4 \Lambda^2} \left(H' + \frac{J'}{a\Lambda} \right) \langle (\hat{\psi}^\dagger)^3 \hat{\psi}^3 \rangle = |\Psi_0^{(3)}|^2 + \delta\mathcal{C}_3, \quad (16)$$

where H' and J' are log-periodic functions of Λ , $\hat{\psi} = \hat{\psi}(\mathbf{0})$ are local field operators, and $\delta\mathcal{C}_p$'s are contributions absent in the triplet model (see the Supplemental Material [50]). This establishes the square modulus of $\Psi_0^{(p)}$ as a probability density, analogous to ψ_0 at the one-body level. The dynamics shown in Fig. 2 can be understood from Refs. [33,40,41,45], namely, early-time growths $|\Psi_0^{(p)}(t)| \propto t^{(p-1)/2}$ with primary ($p=3$) and secondary ($p=2$) visibility of nonuniversal trimer oscillations in the crossover regime ($k_n/\kappa_* = 0.82$ and 0.61).

Internal structure.—We study now the longer-range internal structure of the pair and triple condensate wave functions, focusing on the interplay between Efimovian and Fermi scales. Figure 3 shows the triplet model results for the normalized pair and triple condensate wave functions at $t/t_n = 0.15, 0.5, 1$. To visualize the triple condensate wave function, we average $R_{\mathbf{k}, \mathbf{q}}$ over internal configurations at a fixed hypermomentum $K^2 = k^2 + q^2 + kq \cos \theta$ where $\cos \theta = \hat{\mathbf{k}} \cdot \hat{\mathbf{q}}$. The corresponding $\varphi_0^{(3)}(K)$ captures variations of the coherent tripling physics with changes in the overall three-body momentum scale [50].

At early stages of evolution ($t = 0.15t_n$), the pair and triple wave functions are relatively constant over the range of momenta considered consistent with the buildup of local correlations between nearby particles [33]. Accordingly, the small amount of clustered pairs and triples are dominated by few-body physics. This explains why their




FIG. 2. Macroscopic (a) pair and (b) triple order parameter dynamics over a range of densities within the triplet model. Black solid line: universal results within the doublet model. Inset: residual exponents $\Psi_0^{(p)} \propto \Lambda^{\gamma_p}$ evaluated at fixed $t = 0.05t_n$ converge as expected to 0 as the system becomes increasingly dilute with respect to the range of the interaction ($\Lambda/k_n \rightarrow \infty$).




FIG. 3. Internal structures of the normalized (a) pair and (b) triple condensate wave functions as the system evolves for densities within the crossover (orange) and universal (light blue) regimes. The filled circles in (a) indicate the three-body parameter. Insets: normalized pair and triple condensate wave functions in the universal regime at $t = t_n$ compared with the vacuum ground-state Efimov trimer $|\Psi_{3b}^{(0)}(K)|$ (black-dotted) for density $k_n/\kappa_* = 0.35$.

condensation dynamics shown in Fig. 1 follow the corresponding contact growth laws. We note that the local, structural origin of these laws was not recognized in Ref. [39].

At later times, both pair and triple wave functions become increasingly nonlocal. From the insets of Fig. 3 we see that as the system approaches the universal regime more deeply, both condensate wave functions acquire a universal form by $t = t_n$. Together with the universal behavior of the triple condensed fractions in Fig. 1(c), this remarkable finding suggests the existence of a condensate composed of *universal* Efimovian triples at later times despite strongly nonuniversal short-distance behavior [see Fig. 2(b)]. In the crossover regime results in Fig. 3(a), we find the development of a peak at momenta $k_n \sim \kappa_*$ reminiscent of the Cooper pair in the BEC-BCS crossover [84]. The absence of this peak in simulations of the doublet model and universal regime of the triplet model [Fig. 3(a)] ties it to the Efimov effect.

To study the presence and role of the ground-state Efimov trimer $|\Psi_{3b}^{(0)}\rangle$ in the triple condensate $|\varphi_0^{(3)}\rangle$, we evaluate the overlap

$$P_{3b}^{(0)}(t) = |\langle \varphi_0^{(3)} | \Psi_{3b}^{(0)} \rangle|^2, \quad (17)$$

as shown in Fig. 4 (see the Supplemental Material [50]). In the universal regime, the ground-state Efimov trimer is localized relative to the Fermi scales. At all times in this regime, $P_{3b}^{(0)}$ reflects therefore the short-range behavior of the triple condensate wave function encapsulated by $\Psi_0^{(3)}$, contributing the characteristic trimer oscillations visible in Fig. 4. After a small initial increase, the rapid decrease of $P_{3b}^{(0)}$ in this regime reflects the local to nonlocal transition of




FIG. 4. Evolution of the overlap [Eq. (17)] between ground-state Efimov trimer and triple condensate wave function for varying densities, with $P_{3b}^{(0)}(t=0)$ undefined. The numerical upper limit is comparable to the total probability ($\sim 82\%$) projected onto the Efimov channel in the quenched three-body problem with characteristic time scale $t_{3b}^{(0)} = m/\hbar\kappa_*^2$ (filled circles) [85]. The illustrations indicate the presence or absence of a trimer character in the triple condensate.

the triple condensate wave function, which bears little resemblance to the trimer as shown in the inset of Fig. 3(b). This is responsible also for the decreased visibility of the trimer oscillations in Fig. 1(c). Consequently, this local to nonlocal structural transition, generated by the medium effect of the strong quantum depletion, is the underlying mechanism by which macroscopic observables, such as the condensate fractions in Fig. 1(c), display a universal scale invariance. This occurs despite continued local sensitivity to the three-body parameter [Eq. (13)] which becomes less relevant as the coherent physics begins to occur predominantly on the Fermi scale. In the trimer regime, we find no decrease of $P_{3b}^{(0)}$ for $t \lesssim t_n$, and the early-time increase is more gradual. From Fig. 3(b) it is clear that although the coherent physics occurs predominantly on the Fermi scale at later times as before, when one has $k_n \sim \kappa_*$ the condensed triples are increasingly dominated by the ground-state Efimov trimer. In short, the nonuniversal, trimer character of the triple condensate increases at later times in the trimer regime, whereas it decreases in the universal regime where one finds triples *without* a vacuum equivalent.

Conclusion.—Using the cumulant model that includes three-body correlations, we have shown that novel types of few-body condensates are generated within the quantum depletion of a quenched unitary Bose gas. Crucially, the regime of universal pair and triple condensation demonstrates a strongly interacting many-body system behaving universally even in the presence of nonuniversal few-body physics such as the Efimov effect. We expect the molecular fractions produced following an interaction sweep back to weak interactions [26,29] to reflect the few-body composites present in the unitary regime. However the highlighted difficulties of counting composite bosons extended in the medium requires a precise modeling of the projection and remains the subject of future work [37,86]. Additionally, the tripling fluctuations discussed in this Letter raise interesting prospects for measuring non-Gaussian many-body states [87,88].

We thank Jinglun Li, Denise Ahmed-Braun, Paul Mestrom, Thomas Secker, and Gijs Groeneveld for fruitful discussions. M. V. R. acknowledges support by MURI-AFOSR FA9550-19-1-0399. M. W. acknowledges financial support from the FWO-Vlaanderen under the Project No. G016219N. V. E. C. acknowledges financial support from Provincia Autonoma di Trento and the Italian MIUR under the PRIN2017 project CEnTraL. S. M. and S. J. J. M. F. K. acknowledge financial support by the Netherlands Organisation for Scientific Research (NWO) under Grant No. 680-47-623.

*sil.musolino@gmail.com

†colussiv@gmail.com

[1] A. Leggett, *Quantum Liquids: Bose Condensation and Cooper Pairing in Condensed-Matter Systems*, Oxford Graduate Texts (Oxford University Press, Oxford, 2006).

- [2] L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation and Superfluidity*, International Series of Monographs on Physics (Oxford University Press, Oxford, 2016).
- [3] M. Tinkham, *Introduction to Superconductivity* (McGraw-Hill, New York, 1996).
- [4] S. A. Moskalenko and D. W. Snoke, *Bose-Einstein Condensation of Excitons and Biexcitons: And Coherent Nonlinear Optics with Excitons* (Cambridge University Press, Cambridge, England, 2000).
- [5] G. C. Strinati, P. Pieri, G. Röpke, P. Schuck, and M. Urban, The BCS–BEC crossover: From ultra-cold Fermi gases to nuclear systems, *Phys. Rep.* **738**, 1 (2018).
- [6] G. Röpke, A. Schnell, P. Schuck, and P. Nozières, Four-Particle Condensate in Strongly Coupled Fermion Systems, *Phys. Rev. Lett.* **80**, 3177 (1998).
- [7] H. Tajima, S. Tsutsui, T. M. Doi, and K. Iida, Cooper triples in attractive three-component fermions: Implication for Hadron-Quark crossover, [arXiv:2012.03627](https://arxiv.org/abs/2012.03627).
- [8] J. Makowiak and P. Tarasewicz, An extension of the Bardeen–Cooper–Schrieffer model of superconductivity, *Physica (Amsterdam)* **331C**, 25 (2000).
- [9] R. Sen'kov and V. Zelevinsky, Unified BCS-like model of pairing and alpha-correlations, *Phys. At. Nucl.* **74**, 1267 (2011).
- [10] P. Nozières and D. Saint James, Particle vs. pair condensation in attractive Bose liquids, *J. Phys.* **43**, 1133 (1982).
- [11] M. Kunitski, S. Zeller, J. Voigtsberger, A. Kalinin, L. P. H. Schmidt, M. Schöffler, A. Czasch, W. Schöllkopf, R. E. Grisenti, T. Jahnke, D. Blume, and R. Dörner, Observation of the Efimov state of the helium trimer, *Science* **348**, 551 (2015).
- [12] F. Chevy and C. Salomon, Strongly correlated Bose gases, *J. Phys. B* **49**, 192001 (2016).
- [13] V. Efimov, Weakly-bound states of 3 resonantly-interacting particles, *Sov. J. Nucl. Phys.* **12**, 589 (1971).
- [14] V. Efimov, Low-energy properties of three resonantly interacting particles, *Sov. J. Nucl. Phys.* **29**, 546 (1979).
- [15] Y. Nishida, New Type of Crossover Physics in Three-Component Fermi Gases, *Phys. Rev. Lett.* **109**, 240401 (2012).
- [16] H. Tajima and P. Naidon, Quantum chromodynamics (QCD)-like phase diagram with Efimov trimers and Cooper pairs in resonantly interacting SU(3) Fermi gases, *New J. Phys.* **21**, 073051 (2019).
- [17] Y. Nishida, Polaronic Atom-Trimer Continuity in Three-Component Fermi Gases, *Phys. Rev. Lett.* **114**, 115302 (2015).
- [18] L. A. Peña Ardila and S. Giorgini, Impurity in a Bose-Einstein condensate: Study of the attractive and repulsive branch using quantum Monte Carlo methods, *Phys. Rev. A* **92**, 033612 (2015).
- [19] M. Sun, H. Zhai, and X. Cui, Visualizing the Efimov Correlation in Bose Polarons, *Phys. Rev. Lett.* **119**, 013401 (2017).
- [20] N. G. Nygaard and N. T. Zinner, Efimov three-body states on top of a Fermi sea, *New J. Phys.* **16**, 023026 (2014).
- [21] J. Levinsen, M. M. Parish, and G. M. Bruun, Impurity in a Bose-Einstein Condensate and the Efimov Effect, *Phys. Rev. Lett.* **115**, 125302 (2015).
- [22] A. Christianen, J. I. Cirac, and R. Schmidt, From Efimov physics to the Bose polaron using gaussian states, [arXiv:2108.03175](https://arxiv.org/abs/2108.03175).
- [23] P. Naidon and S. Endo, Efimov physics: A review, *Rep. Prog. Phys.* **80**, 056001 (2017).
- [24] S. Taie, R. Yamazaki, S. Sugawa, and Y. Takahashi, An SU(6) Mott insulator of an atomic Fermi gas realized by large-spin Pomeranchuk cooling, *Nat. Phys.* **8**, 825 (2012).
- [25] P. Makotyn, C. E. Klauss, D. L. Goldberger, E. Cornell, and D. S. Jin, Universal dynamics of a degenerate unitary Bose gas, *Nat. Phys.* **10**, 116 (2014).
- [26] C. E. Klauss, X. Xie, C. Lopez-Abadia, J. P. D’Incao, Z. Hadzibabic, D. S. Jin, and E. A. Cornell, Observation of Efimov Molecules Created from a Resonantly Interacting Bose Gas, *Phys. Rev. Lett.* **119**, 143401 (2017).
- [27] C. Eigen, J. A. P. Glidden, R. Lopes, N. Navon, Z. Hadzibabic, and R. P. Smith, Universal Scaling Laws in the Dynamics of a Homogeneous Unitary Bose Gas, *Phys. Rev. Lett.* **119**, 250404 (2017).
- [28] C. Eigen, J. A. P. Glidden, R. Lopes, E. A. Cornell, R. P. Smith, and Z. Hadzibabic, Universal prethermal dynamics of Bose gases quenched to unitarity, *Nature (London)* **563**, 221 (2018).
- [29] C. A. Regal, M. Greiner, and D. S. Jin, Observation of Resonance Condensation of Fermionic Atom Pairs, *Phys. Rev. Lett.* **92**, 040403 (2004).
- [30] W. E. Zwirger, *The BCS-BEC Crossover and the Unitary Fermi Gas (Vol. 863 of Lecture Notes in Physics)*, edited by W. Zwirger (Springer-Verlag, Berlin, 2012).
- [31] T.-L. Ho, Universal Thermodynamics of Degenerate Quantum Gases in the Unitarity Limit, *Phys. Rev. Lett.* **92**, 090402 (2004).
- [32] M. Van Regemortel, H. Kurkjian, M. Wouters, and I. Carusotto, Prethermalization to thermalization crossover in a dilute Bose gas following an interaction ramp, *Phys. Rev. A* **98**, 053612 (2018).
- [33] A. G. Sykes, J. P. Corson, J. P. D’Incao, A. P. Koller, C. H. Greene, A. M. Rey, K. R. A. Hazzard, and J. L. Bohn, Quenching to unitarity: Quantum dynamics in a three-dimensional Bose gas, *Phys. Rev. A* **89**, 021601(R) (2014).
- [34] J. P. Corson and J. L. Bohn, Bound-state signatures in quenched Bose-Einstein condensates, *Phys. Rev. A* **91**, 013616 (2015).
- [35] C. Gao, M. Sun, P. Zhang, and H. Zhai, Universal Dynamics of a Degenerate Bose Gas Quenched to Unitarity, *Phys. Rev. Lett.* **124**, 040403 (2020).
- [36] A. Muñoz de las Heras, M. M. Parish, and F. M. Marchetti, Early-time dynamics of Bose gases quenched into the strongly interacting regime, *Phys. Rev. A* **99**, 023623 (2019).
- [37] S. Musolino, V. E. Colussi, and S. J. J. M. F. Kokkelmans, Pair formation in quenched unitary Bose gases, *Phys. Rev. A* **100**, 013612 (2019).
- [38] V. E. Colussi, S. Musolino, and S. J. J. M. F. Kokkelmans, Dynamical formation of the unitary Bose gas, *Phys. Rev. A* **98**, 051601(R) (2018).
- [39] J. P. D’Incao, J. Wang, and V. E. Colussi, Efimov Physics in Quenched Unitary Bose Gases, *Phys. Rev. Lett.* **121**, 023401 (2018).

- [40] V. E. Colussi, J. P. Corson, and J. P. D’Incao, Dynamics of Three-Body Correlations in Quenched Unitary Bose Gases, *Phys. Rev. Lett.* **120**, 100401 (2018).
- [41] V. E. Colussi, B. E. van Zwol, J. P. D’Incao, and S. J. J. M. F. Kokkelmans, Bunching, clustering, and the buildup of few-body correlations in a quenched unitary Bose gas, *Phys. Rev. A* **99**, 043604 (2019).
- [42] T. Bodineau, B. Derrida, V. Lecomte, and F. van Wijland, Long range correlations and phase transitions in non-equilibrium diffusive systems, *J. Stat. Phys.* **133**, 1013 (2008).
- [43] M. Kira and S. W. Koch, *Semiconductor Quantum Optics* (Cambridge University Press, Cambridge, England, 2011).
- [44] M. Tohyama and P. Schuck, Extension of time-dependent Hartree-Fock-Bogoliubov equations, *Phys. Rev. C* **91**, 034316 (2015).
- [45] V. E. Colussi, H. Kurkjian, M. Van Regemortel, S. Musolino, J. van de Kraats, M. Wouters, and S. J. J. M. F. Kokkelmans, Cumulant Theory of the Unitary Bose Gas: Prethermal and Efimovian Dynamics, *Phys. Rev. A* **102**, 063314 (2020).
- [46] M. Kira, Hyperbolic Bloch equations: Atom-cluster kinetics of an interacting Bose gas, *Ann. Phys. (Amsterdam)* **356**, 185 (2015).
- [47] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Feshbach resonances in ultracold gases, *Rev. Mod. Phys.* **82**, 1225 (2010).
- [48] V. V. Flambaum, G. F. Gribakin, and C. Harabati, Analytical calculation of cold-atom scattering, *Phys. Rev. A* **59**, 1998 (1999).
- [49] T. Köhler, T. Gasenzer, and K. Burnett, Microscopic theory of atom-molecule oscillations in a Bose-Einstein condensate, *Phys. Rev. A* **67**, 013601 (2003).
- [50] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.128.020401>, which includes Refs. [51–66], for additional details of our calculations.
- [51] L. D. Faddeev and S. P. Merkuriev, *Quantum Scattering Theory for Several Particle Systems*, Vol. 11 (Springer Science & Business Media, Netherlands, 2013).
- [52] W. Glöckle, *The Quantum Mechanical Few-Body Problem*, edited by W. Beiglböck (Springer Berlin Heidelberg, Berlin, 1983).
- [53] J. R. Taylor, *Scattering Theory: The Quantum Theory of Nonrelativistic Collisions* (Dover Publications, New York, 2006).
- [54] L. H. Thomas, The interaction between a neutron and a proton and the structure of H^3 , *Phys. Rev.* **47**, 903 (1935).
- [55] G. Skorniakov and K. Ter-Martirosian, Three body problem for short range forces. I. Scattering of low energy neutrons by deuterons, *Sov. Phys. JETP* **4** (1957).
- [56] C. H. Greene, P. Giannakeas, and J. Pérez-Ríos, Universal few-body physics and cluster formation, *Rev. Mod. Phys.* **89**, 035006 (2017).
- [57] J. Wang, J. P. D’Incao, B. D. Esry, and C. H. Greene, Origin of the Three-Body Parameter Universality in Efimov Physics, *Phys. Rev. Lett.* **108**, 263001 (2012).
- [58] P. Naidon, S. Endo, and M. Ueda, Physical origin of the universal three-body parameter in atomic Efimov physics, *Phys. Rev. A* **90**, 022106 (2014).
- [59] L. Salasnich, N. Manini, and A. Parola, Condensate fraction of a Fermi gas in the BCS-BEC crossover, *Phys. Rev. A* **72**, 023621 (2005).
- [60] H. Kurkjian, Y. Castin, and A. Sinatra, Phase operators and blurring time of a pair-condensed Fermi gas, *Phys. Rev. A* **88**, 063623 (2013).
- [61] F. Werner and Y. Castin, General relations for quantum gases in two and three dimensions: Two-component fermions, *Phys. Rev. A* **86**, 013626 (2012).
- [62] A. Leggett, Cooper pairing in spin-polarized fermi systems, *J. Phys. (Paris), Colloq.* **41**, C7 (1980).
- [63] M. Combescot, X. Leyronas, and C. Tanguy, On the N-exciton normalization factor, *Eur. Phys. J. B* **31**, 17 (2003).
- [64] M. Combescot and C. Tanguy, New criteria for bosonic behavior of excitons, *Europhys. Lett.* **55**, 390 (2001).
- [65] R. J. Fletcher, R. Lopes, J. Man, N. Navon, R. P. Smith, M. W. Zwierlein, and Z. Hadzibabic, Two- and three-body contacts in the unitary Bose gas, *Science* **355**, 377 (2017).
- [66] M. Sun, P. Zhang, and H. Zhai, High temperature virial expansion to universal quench dynamics, *Phys. Rev. Lett.* **125**, 110404 (2020).
- [67] J. Fricke, Transport equations including many-particle correlations for an arbitrary quantum system: A general formalism, *Ann. Phys. (N.Y.)* **252**, 479 (1996).
- [68] T. Köhler and K. Burnett, Microscopic quantum dynamics approach to the dilute condensed Bose gas, *Phys. Rev. A* **65**, 033601 (2002).
- [69] M. Kira, Excitation picture of an interacting Bose gas, *Ann. Phys. (Amsterdam)* **351**, 200 (2014).
- [70] M. Kira, Coherent quantum depletion of an interacting atom condensate, *Nat. Commun.* **6**, 6624 (2015).
- [71] T. Köhler, Three-body problem in a dilute Bose-Einstein condensate, *Phys. Rev. Lett.* **89**, 210404 (2002).
- [72] O. Penrose, CXXXVI. On the quantum mechanics of helium II, *London, Edinburgh Dublin Philos. Mag. J. Sci.* **42**, 1373 (1951).
- [73] O. Penrose and L. Onsager, Bose-Einstein condensation and liquid helium, *Phys. Rev.* **104**, 576 (1956).
- [74] C. N. Yang, Concept of off-diagonal long-range order and the quantum phases of liquid He and of superconductors, *Rev. Mod. Phys.* **34**, 694 (1962).
- [75] Y. H. Pong and C. K. Law, Bosonic characters of atomic Cooper pairs across resonance, *Phys. Rev. A* **75**, 043613 (2007).
- [76] P. F. Bedaque, H.-W. Hammer, and U. van Kolck, Renormalization of the Three-Body System with Short-Range Interactions, *Phys. Rev. Lett.* **82**, 463 (1999).
- [77] E. Braaten, H.-W. Hammer, and T. Mehen, Dilute Bose-Einstein Condensate with Large Scattering Length, *Phys. Rev. Lett.* **88**, 040401 (2002).
- [78] E. A. Donley, N. R. Claussen, S. T. Thompson, and C. E. Wieman, Atom-Molecule coherence in a Bose-Einstein condensate, *Nature (London)* **417**, 529 (2002).
- [79] E. Braaten, H.-W. Hammer, and M. Kusunoki, Efimov States in a Bose-Einstein Condensate near a Feshbach Resonance, *Phys. Rev. Lett.* **90**, 170402 (2003).
- [80] J. P. Blaizot and G. Ripka, *Quantum Theory of Finite Systems* (The MIT Press, Cambridge, Massachusetts, 1986).

- [81] F. Werner and Y. Castin, General relations for quantum gases in two and three dimensions. II. Bosons and mixtures, *Phys. Rev. A* **86**, 053633 (2012).
- [82] The normalization constant is $\langle \phi_s | \phi_s \rangle \equiv \int d\Omega |\phi_s(\Omega)|^2 = -(12\pi/s) \sin(s^*\pi/2) [\cos(s\pi/2) - (s\pi/2) \sin(s\pi/2) - (4\pi/3\sqrt{3}) \cos(s\pi/6)]$.
- [83] E. Braaten, D. Kang, and L. Platter, Universal Relations for Identical Bosons from Three-Body Physics, *Phys. Rev. Lett.* **106**, 153005 (2011).
- [84] M. M. Parish, B. Mihaila, E. M. Timmermans, K. B. Blagoev, and P. B. Littlewood, BCS-BEC crossover with a finite-range interaction, *Phys. Rev. B* **71**, 064513 (2005).
- [85] F. Werner, Atomes froids piégés en interaction résonnante: Gaz unitaire et problème à trois corps, Ph.D. thesis, Université Pierre et Marie Curie, 2008.
- [86] E. Altman and A. Vishwanath, Dynamic Projection on Feshbach Molecules: A Probe of Pairing and Phase Fluctuations, *Phys. Rev. Lett.* **95**, 110404 (2005).
- [87] A. Tenart, G. Herc, J.-P. Bureik, A. Dureau, and D. Clément, Observation of pairs of atoms at opposite momenta in an equilibrium interacting Bose gas, [arXiv:2105.05664](https://arxiv.org/abs/2105.05664).
- [88] T. Schweigler, V. Kasper, S. Erne, I. Mazets, B. Rauer, F. Cataldini, T. Langen, T. Gasenzer, J. Berges, and J. Schmiedmayer, Experimental characterization of a quantum many-body system via higher-order correlations, *Nature (London)* **545**, 323 (2017).