

Temperature-Dependent Periodicity of the Persistent Current in Strongly Interacting Systems

Ovidiu I. Păţu¹ and Dmitri V. Averin²

¹*Institute for Space Sciences, Bucharest-Măgurele R 077125, Romania*

²*Department of Physics and Astronomy, Stony Brook University, Stony Brook, New York 11794, USA*

 (Received 22 October 2021; accepted 14 February 2022; published 4 March 2022)

The persistent current in small isolated rings enclosing magnetic flux is the current circulating in equilibrium in the absence of an external excitation. While initially studied in superconducting and normal metals, recently, atomic persistent currents have been generated in ultracold gases spurring a new wave of theoretical investigations. Nevertheless, our understanding of the persistent currents in interacting systems is far from complete, especially at finite temperatures. Here we consider the fermionic one-dimensional Hubbard model and show that in the strong-interacting limit, the current can change its flux period and sign (diamagnetic or paramagnetic) as a function of temperature, features that cannot be explained within the single-particle or Luttinger liquid techniques. Also, the magnitude of the current can counterintuitively increase with temperature, in addition to presenting different rates of decay depending on the polarization of the system. Our work highlights the properties of the strongly interacting multicomponent systems that are missed by conventional approximation techniques, but can be important for the interpretation of experiments on persistent currents in ultracold gases.

DOI: [10.1103/PhysRevLett.128.096801](https://doi.org/10.1103/PhysRevLett.128.096801)

Introduction.—The existence of a persistent current (PC) in small metallic rings threaded by magnetic flux ϕ at low temperatures has been theoretically predicted since the early days of quantum mechanics [1] and superconductivity [2–5]. Following the publication of Ref. [6], PCs were the focus of intense theoretical investigations, and were experimentally confirmed in both individual and ensembles of metal rings [7–12]. Resurgent interest in the field is due to the generation of atomic PCs in ultracold gases of single-component bosons [13,14], spinor bosons [15], and very recently of spinful fermions [16].

The PC is a paradigmatic example of quantum coherence in mesoscopic systems and its magnitude is given by $I(\phi) = -\partial F(\phi)/\partial\phi$ with $F(\phi) = -k_B T \ln \mathcal{Z}(\phi)$ the free energy and $\mathcal{Z}(\phi)$ the canonical partition function [17,18]. Gauge invariance implies periodicity, $I(\phi + \phi_0) = I(\phi)$, with ϕ_0 the flux quantum $\phi_0 = h/e$ (h is Planck's constant and e the charge of the electron) and from time invariance we have $I(-\phi) = -I(\phi)$. In addition to the amplitude of the current, defined by $I_{\max} = \max_{\phi \in (0, \phi_0/2)} |I(\phi)|$, and periodicity, we are interested in the sign of the magnetic response: diamagnetic or paramagnetic. A system is diamagnetic (paramagnetic) if $F(\phi)$ has a local minimum (maximum) at $\phi = 0$. While there is a large body of work, mainly focused on free electrons with disorder, our understanding of PCs in interacting systems, especially its temperature dependence, is far from complete. Using a variational approach, Leggett conjectured [19] that the ground-state energy of N polarized interacting fermions is diamagnetic for odd N and paramagnetic for even N .

This conjecture was proved and extended to the case of small temperatures using Luttinger liquid (LL) methods [20]. At small temperatures and within the LL regime, the amplitude of the current decreases exponentially with temperature, but the periodicity and sign of the current remain unchanged. For interacting fermions with spin [N_\downarrow (N_\uparrow) electrons have spin-down (-up)] a general result valid at arbitrary temperature [21] is that $F(0) \leq F(\phi/\phi_0)$ for $N_{\downarrow,\uparrow}$ both odd, and $F(1/2) \leq F(\phi/\phi_0)$ for $N_{\downarrow,\uparrow}$ both even. This result does not preclude periodicities smaller than ϕ_0 or changes in the sign of the current with temperature, as we will show below.

The Hubbard model in a magnetic field.—We consider a system of N electrons of which N_\downarrow have spin-down on a ring lattice with L sites and repulsive interactions. The ring is threaded by an Aharonov-Bohm flux ϕ . The system is described by the Hubbard Hamiltonian [22–25]

$$\mathcal{H} = -t \sum_{j=1}^L \left[\sum_{\sigma} (e^{-ieA} c_{j+1,\sigma}^\dagger c_{j,\sigma} + \text{H.c.}) - \frac{U}{t} n_{j,\uparrow} n_{j,\downarrow} \right], \quad (1)$$

with $n_{j,\sigma} = c_{j,\sigma}^\dagger c_{j,\sigma}$ the number of electrons of spin $\sigma = \{\downarrow, \uparrow\}$ at site j . In Eq. (1), $A = 2\pi\phi/(L\phi_0)$ is the vector potential of the magnetic flux, t is the electron hopping integral, and $U > 0$ quantifies the strength of the repulsive interaction. In the following, we will measure the energies in units of t and set $\hbar = 1$ and $e = 1$. The Hamiltonian (1) is exactly solvable with the Bethe ansatz equations (BAEs)

[24]: $k_j L = 2\pi I_j + 2\pi\phi/\phi_0 - \sum_{\alpha=1}^{N_\downarrow} \theta[(\sin k_j - \lambda_\alpha)/u]$, $\sum_{j=1}^N \theta[(\lambda_\alpha - \sin k_j)/u] = 2\pi J_\alpha + \sum_{\beta=1}^{N_\downarrow} \theta[(\lambda_\alpha - \lambda_\beta)/(2u)]$ where $u \equiv U/4t$, $j = 1, \dots, N$, $\alpha = 1, \dots, N_\downarrow$, $\theta(x) = 2 \arctan(x)$, and $I_j = N_\downarrow/2 \pmod{1}$, i.e., I_j is integer or half-integer depending on whether N_\downarrow is even or odd, and similarly, $J_\alpha = (N - N_\downarrow + 1)/2 \pmod{1}$. The energy and momentum of a state are $E = -2 \sum_{j=1}^N \cos k_j$ and $P = [\sum_{j=1}^N k_j] \pmod{2\pi}$. To find the PC we need to solve the BAEs for the ground state (at $T = 0$) or for all the relevant excited states (at $T > 0$). We look first at some limiting cases.

The $U = 0$ case.—Because the particles are noninteracting, it is sufficient to consider spinless fermions. For fermions with spin, the PC is then given simply by the sum of the contributions from the two spin directions [26]. The momenta of M spinless fermions in the ring are $k_j = 2\pi(j + \phi/\phi_0)/L$, with j integer. This implies that at $\phi = 0$, the ground-state is degenerate for even M , but not for odd M , i.e., the PC depends on the parity of the number of particles. In terms of the ‘‘Fermi vector’’ $k_F(M) \equiv \pi M/L$ and the Fermi velocity $v_F(M) = 2 \sin k_F(M)$, the PC of spinless fermions is [27]: $I_{FF}(M, \phi) = -\{[I_0(M)]/(\sin \pi/L)\} \sin\{[(2\pi/L)(\phi/\phi_0)]\}$ for M odd, and $I_{FF}(M, \phi) = \{[I_0(M)]/(\sin \pi/L)\} \sin\{(\pi/L)[1 - (2|\phi|)/\phi_0]\} \text{sgn}\phi$ for M even with $I_0(M) = ev_F(M)/L$. Therefore, the PC for electrons with spin is $I(\phi) = I_{FF}(N_\downarrow, \phi) + I_{FF}(N_\uparrow, \phi)$. When $N_{\downarrow, \uparrow}$ are both odd, $I(\phi)$ is diamagnetic; when $N_{\downarrow, \uparrow}$ are both even, it is paramagnetic.

The $U = \infty$ case.—In this ‘‘impenetrable’’ limit [28–31] the BAEs for the k_j ’s become $k_j^\infty = 2\pi(I_j + \phi/\phi_0 + \sum_{\alpha=1}^{N_\downarrow} J_\alpha/N)/L$, which are equivalent to the result for

spinless fermions in a ring threaded by a magnetic flux $\phi/\phi_0 + \sum_{\alpha=1}^{N_\downarrow} J_\alpha/N$. In the ground state, I_j ’s fill an interval between I_{\min} and I_{\max} , resulting in the energy $E = -2\{[\sin(\pi N/L)]/[\sin(\pi/L)]\} \cos\{[(2\pi)/L][(\phi/\phi_0) + (1/N)\sum_{\alpha=1}^{N_\downarrow} J_\alpha + D]\}$ where $D \equiv (I_{\min} + I_{\max})/2$. This formula is valid for densities $n = N/L < 1$ (at half-filling, $n = 1$, the sine factor in E gives 0). The energy is minimized by choosing the set $\{J_\alpha\}_{\alpha=1}^{N_\downarrow}$ such that $\sum_{\alpha=1}^{N_\downarrow} J_\alpha = -p$ for $(p-1/2)/N < \phi/\phi_0 + D < (p+1/2)/N$. This implies that the PC for the not fully polarized impenetrable system at zero temperature is (i) periodic with a period of $1/N$ of the flux quantum [this remains valid in the case of $SU(\kappa)$ fermions with $\kappa > 2$ as shown in a recent study [31]], (ii) diamagnetic, and (iii) does not present parity effects [29,30]—see also Fig. 1. Note that for fully spin-polarized electrons, $N_\downarrow = 0$, the system is effectively noninteracting even for large U , and PC is described by the same expressions as for $U = 0$, e.g., has the period of one flux quantum. This abrupt change of the flux period (from 1 to $1/N$) between the polarized electrons and a system with even one flipped spin can be understood in terms of the change of the rotation period of the electron system in real space [32]. For polarized electrons, the rotation period is $1/N$ of the full rotation, while a spin flip changes this period to a full rotation.

First correction.—For u large but finite, the charge momenta with accuracy $1/u$ are [25,30,33] $k_j = k_j^\infty + \Delta k_j/u$ with $\Delta k_j = E_s \sum_{n=1}^{N_\downarrow} [\sin k_j^\infty - \sin k_n^\infty]/L$, where $E_s = -2 \sum_{\alpha=1}^{N_\downarrow} 1/[N(\Lambda_\alpha^2 + 1)]$ is the energy per lattice site of an antiferromagnetic XXX Heisenberg spin chain with spin rapidities $\{\Lambda_\alpha\}_{\alpha=1}^{N_\downarrow}$ satisfying the BAEs:




FIG. 1. Dependence of the ground-state energy (left columns; normalized by t) and current [right columns; in units of $I_0(N_\downarrow) + I_0(N_\uparrow)$] on the magnetic flux and strength of the interaction for $N = 8, N_\downarrow = 4$ and $N = 10, N_\downarrow = 5$. From top row to bottom the coupling strength is $U = 5000, 50, 5.5, 1, 0$ and the density is the same in both cases $n = 0.1$.

$N\theta(\Lambda_\alpha) = 2\pi J_\alpha + \sum_{\beta=1}^{N_\downarrow} \theta[(\Lambda_\alpha - \Lambda_\beta)/2]$. The energy with the same $1/u$ accuracy is $E = E^\infty + [(2E_s)/(Lu)][N \sum_{j=1}^N \sin^2 k_j^\infty - (\sum_{j=1}^N \sin k_j^\infty)^2]$ with $E^\infty = -2 \sum_{j=1}^N \cos k_j^\infty$. Note that the energy depends on ϕ via k_j^∞ . Thus, in the strong-coupling limit, the spin degrees of freedom are described by an antiferromagnetic Heisenberg spin chain with the coupling constant $\sim N_\downarrow/(Lu)$, while the charge degrees of freedom are similar to free fermions. This regime, in which the energies of the charge and spin sectors satisfy $E_{\text{charge}} \gg E_{\text{spin}}$, is called the spin-incoherent regime [34–40]. It is particular to multicomponent systems, and presents universal properties which are different from the LL case. For the Hubbard model, the lowest energy states in this regime are obtained [30] by considering the same distribution of I_j 's as in the infinite repulsion case and finding the states of the Heisenberg chain with the lowest energy for a given momentum $q = 2\pi \sum_{\alpha=1}^M J_\alpha/N$, which are the des-Cloizeaux-Pearson excitations [41].

The dependence of the ground-state energy and current on the strength of interaction at zero temperature is shown in Fig. 1 for “balanced” systems with $N = 8$ and $N = 10$. At very strong coupling ($U = 5000$) the current is diamagnetic and has other characteristic features of the impenetrable case: $1/N$ periodicity and no parity effects. For weaker interaction, the contribution of the spin sector becomes more pronounced, as manifested by the raising of the N parabolas. We can see that for $U = 50$ and $U = 5.5$, while the PC is still diamagnetic, the periodicity changes to $1/2$. At very weak interaction, the periodicity becomes 1, and for $U = 0$, one obtains a paramagnetic (diamagnetic) current for $N_{\downarrow,\uparrow}$ even (odd). Therefore, the strength of the interaction has a strong influence on the zero-temperature PC, changing its periodicity, amplitude, and even the sign (paramagnetic or diamagnetic). Also, the $1/2$ periodicity seen in Fig. 1 at strong and intermediary coupling ($U = 50$ and $U = 5.5$) is a particular case of the additional N_\downarrow/N periodicity characterizing the PC of the strongly interacting Hubbard model first discovered in Ref. [42] for $N_\downarrow \ll N$. Figure 1 provides a proof of this additional N_\downarrow/N periodicity in the microscopic regime of small N . From the point of view of the real-space rotations, this periodicity can be viewed as a manifestation of the antiferromagnetic order, which for the balanced electron system makes the rotation period $1/N_\downarrow$ of the full rotation, i.e., two times the period $1/N$ for spinless electrons.

In the mesoscopic regime ($N, L \gg 1$), and at zero temperature, the calculation of the persistent current is equivalent to the calculation of the finite size corrections to the energy due to the change of the boundary conditions from periodic to twisted [30]. For any value of U and $n < 1$, and assuming that the I_j 's and J_α 's are consecutive numbers, the corrections to the ground-state energy due to the magnetic flux are [30,43,44]

$$\Delta E(\phi) = \frac{2\pi v_c}{L} \left[Z_{cc} \left(D_c + \frac{\phi}{\phi_0} \right) + Z_{sc} D_s \right]^2 + \frac{2\pi v_s}{L} \left[Z_{cs} \left(D_c + \frac{\phi}{\phi_0} \right) + Z_{ss} D_s \right]^2, \quad (2)$$

where $Z_{cc}, Z_{cs}, Z_{sc}, Z_{ss}$ are the matrix elements of the dressed charge matrix (see Supplemental Material [45]), $v_{c,s}$ are the velocities of the charge and spin excitations, and D_c and D_s satisfy the constraints $I_{\max} - I_{\min} + 1 = N$, $I_{\max} + I_{\min} = 2D_c$, $J_{\max} - J_{\min} + 1 = N_\downarrow$, and $J_{\max} + J_{\min} = 2D_s$.

When $U \gg 1$ we have $Z_{cc} \sim 1$, $Z_{cs} \sim 0$, $Z_{sc} \sim N_\downarrow/N$, and Z_{ss} is given by a simple integral equation [47,48]. In the same limit the charge and spin velocities behave like $v_c \sim 2 \sin \pi n$, $v_s \sim 1/U$, which means that in the first approximation we can neglect the second term of Eq. (2)

$$\Delta E(\phi) = \frac{2\pi v_c}{L} \left[\left(D_c + \frac{\phi}{\phi_0} \right) + \frac{N_\downarrow}{N} D_s \right]^2. \quad (3)$$

When $N_\downarrow/N = 1/m$ with $m = 2, 3, \dots$, this expression shows that the current has a N_\downarrow/N periodicity. When N_\downarrow/N is not very close to $1/m$, the situation is more complicated (see Supplemental Material [45]).

Persistent current at finite temperature.—Computing thermodynamics of the Hubbard model, even in the thermodynamic limit, is a very difficult task, and it is sensible to assume that computing the $1/L$ corrections is outside the reach of analytical methods. In the strong coupling limit, however, one can take advantage of the fact that the energy of the spin sector is much smaller than the energy of the charge sector, which allows for direct computation of the canonical partition function at low temperatures by summing over all the spin eigenstates and only some of the charge excitations. For a dilute system ($n < 0.1$) the relevant temperature scales are $T_F = \pi^2 n^2$ for the charge degrees of freedom and $T_s = \pi^2 n^3/U$ for the spin degrees of freedom. For temperatures $T \ll T_F$ the partition function can be computed as $\mathcal{Z}(\phi) = \sum_{\text{relevant sets } I} \sum_{\text{all sets } J} \exp\{-E(\{k_j\}, \phi)/T\}$ and gives the PC. This approach requires the knowledge of the all $C_{N_\downarrow}^N$ states of the Heisenberg spin chain with N sites and N_\downarrow spins down, which can be found in Refs. [49–51]. Using this method we were able to investigate the PC for all systems with $N \leq 10$, $N_\downarrow \leq N/2$, and $T < 0.06T_F$. While below we focus on dilute systems, we note that our results remain valid for all densities $0 < n < 1$, if $U/n \gg 1$ (see Supplemental Material [45]).

The dependence of the PC on temperature in the strongly interacting Hubbard model is very complex with the polarization of the system playing an important role. For a system with $N = 8$ and $N_\downarrow = 2$, Fig. 2 shows that while at very low temperatures the current is diamagnetic with period $1/8$, at higher temperatures the periodicity changes




FIG. 2. Dependence of the free energy (left columns in units of τ) and current [right columns normalized by $I_0(N_\downarrow) + I_0(N_\uparrow)$] on the magnetic flux and temperature for $N = 8, N_\downarrow = 2$ and $N = 10, N_\downarrow = 2$. From top row to bottom the temperatures are $T/T_s = 1.01; 48.1; 582$ for $N = 8$ and $T/T_s = 1.01; 48.1; 455$ for $N = 10$. The density and interaction strength are $n = 0.01$ and $U = 100$.

to $1/4$. At even higher temperatures the current becomes paramagnetic with period 1. The evolution of the current with increasing temperature for a system with $N = 10$ and $N_\downarrow = 2$ is similar (but note the paramagnetic current at intermediate temperatures): diamagnetic with period $1/10$, paramagnetic with period $1/5$, and paramagnetic with period 1. Therefore, we see the following pattern: in the ground state, the current is diamagnetic with periodicity $1/N$, and is followed at higher temperatures by the current with N_\downarrow/N periodicity whose sign is the same as the one at zero temperature, which can be derived from Eq. (2). At very high temperatures, the current should have the same characteristics as for free fermions with spin, i.e., for both $N_{\downarrow,\uparrow}$ even, paramagnetic with period 1. This general pattern can be understood by noting that an increase in temperature is qualitatively similar to the decrease in U , and therefore the evolution of the current with T mimics the evolution of the current at $T = 0$, when interaction decreases. For instance, the doubling of the current period from $1/N$ to N_\downarrow/N in Fig. 2 can again be related to the change of the rotation symmetry of the electron system in real space, from full rotation at low temperatures, to half of the rotation at the intermediate temperatures, when the two spin-down electrons become located symmetrically in the system.

Qualitatively, the amplitude of the PC is reduced exponentially with increasing temperature. The quantitative temperature dependence of the amplitude is plotted in Fig. 3 for systems with $N = 6$ and $N = 10$, and shows that there are different rates of decay associated with the different ranges of the system parameters: Luttinger liquid, spin incoherent, and almost free. The fastest rate of decay is in the LL regime; in the spin-incoherent regime, the rate of decay depends strongly on the polarization of the system. A very interesting feature which can be seen in Fig. 3 is the presence of an interval of temperature in which the amplitude is *increasing* with T for $N_{\downarrow,\uparrow} = 3$ and $N_{\downarrow,\uparrow} = 5$. This counterintuitive feature is present at the

transition between the LL and the spin-incoherent regime and is due to the fact that the tail of the momentum distribution gets strongly suppressed as the temperature increases (for a similar phenomenon in spinless fermionic systems with nontrivial geometry or dissipation see Refs. [52,53]). This momentum reconstruction was first noticed in the case of the Gaudin-Yang model (obtained in the dilute limit of the Hubbard model) in Ref. [54]. In details, the PC is on the order of that produced by an electron at the Fermi level, $I = ev_F/L$. At finite temperature, the occupation probabilities of levels close in energy to the Fermi level, and producing positive and negative contributions to the current, are not very different, leading to suppression of the current associated with the broadening of the momentum distribution. At the LL-spin-incoherent transition, however, the fraction of particles with higher momenta decreases, resulting in a softer rate of decay for some polarizations, or even increase in the magnitude of the PC.




FIG. 3. Temperature dependence of the amplitudes [normalized by $I_0(N_\downarrow) + I_0(N_\uparrow)$] for $N = 6$ and $N = 10$. Note that for $N = 6, N_\downarrow = 3$ and $N = 10, N_\downarrow = 5$ there is an interval in which the amplitude is increasing with temperature. For all cases $n = 0.01$ and $U = 100$.

We expect that the temperature dependent periodicity and the different rates of decay of the PC to be general features of strongly interacting fermionic systems with spin-independent interactions. The reason is that such systems (integrable and nonintegrable) present both the LL and spin-incoherent regimes, as it can be seen more explicitly in the “Wigner-molecule” regime of the charged fermions.

Conclusions.—In summary, we calculated the persistent current in the strongly repulsive Hubbard model at finite temperatures from the Bethe ansatz equations. The current shows several notable characteristics, including the temperature-dependent period and the counterintuitive temperature dependence of the amplitude. To the best of our knowledge, this is the first example of a temperature-dependent period of the persistent current, despite a large number of previous studies of temperature-dependent PCs in many different models. It is quite unexpected, since the period is a fundamental quantum property of the system which should be statistics independent. An interesting future development would be an extension of our results to any value of temperature by the alternative (but considerably more computationally expensive) method of exact diagonalization and finding similar transport regimes in other, possibly nonintegrable, models of strongly interacting particles. We believe also that our findings will have considerable implications for the interpretation of experiments on persistent currents of multicomponent systems that can be generated and investigated in the present-day atomtronics circuits [55–62].

The work of O.I.P. has been supported from the LAPLAS 6 program of the Romanian National Authority for Scientific Research (CNCS-UEFISCDI). D.V.A. was supported by the U.S. NSF Grants No. EAGER 1836707 and No. 2104781.

[1] F. Hund, *Ann. Phys. (N.Y.)* **32**, 102 (1938).
 [2] F. Bloch, *Phys. Rev.* **137**, A787 (1965).
 [3] M. Schick, *Phys. Rev.* **166**, 404 (1968).
 [4] L. Gunther and Y. Imry, *Solid State Commun.* **7**, 1391 (1969).
 [5] I. O. Kulik, *Pis'ma Zh. Eksp. Teor. Fiz.* **11**, 407 (1970) [*JETP Lett.* **11**, 275 (1970)].
 [6] M. Buttiker, Y. Imry, and R. Landauer, *Phys. Lett.* **96A**, 365 (1983).
 [7] L. P. Lévy, G. Dolan, J. Dunsmuir, and H. Bouchiat, *Phys. Rev. Lett.* **64**, 2074 (1990).
 [8] V. Chandrasekhar, R. A. Webb, M. J. Brady, M. B. Ketchen, W. J. Gallagher, and A. Kleinsasser, *Phys. Rev. Lett.* **67**, 3578 (1991).
 [9] E. M. Q. Jariwala, P. Mohanty, M. B. Ketchen, and R. A. Webb, *Phys. Rev. Lett.* **86**, 1594 (2001).
 [10] R. Deblock, R. Bel, B. Reulet, H. Bouchiat, and D. Mailly, *Phys. Rev. Lett.* **89**, 206803 (2002).
 [11] H. Bluhm, N. C. Koshnick, J. A. Bert, M. E. Huber, and K. A. Moler, *Phys. Rev. Lett.* **102**, 136802 (2009).

[12] A. C. Bleszynski-Jayich, W. E. Shanks, B. Peaudecerf, E. Ginossar, F. von Oppen, L. Glazman, and J. G. E. Harris, *Science* **326**, 272 (2009).
 [13] C. Ryu, M. F. Andersen, P. Cladé, V. Natarajan, K. Helmerson, and W. D. Phillips, *Phys. Rev. Lett.* **99**, 260401 (2007).
 [14] K. C. Wright, R. B. Blakestad, C. J. Lobb, W. D. Phillips, and G. K. Campbell, *Phys. Rev. Lett.* **110**, 025302 (2013).
 [15] S. Beattie, S. Moulder, R. J. Fletcher, and Z. Hadzibabic, *Phys. Rev. Lett.* **110**, 025301 (2013).
 [16] Y. Cai, D. G. Allman, P. Sabharwal, and K. C. Wright, *arXiv:2104.02218*.
 [17] N. Byers and C. N. Yang, *Phys. Rev. Lett.* **7**, 46 (1961).
 [18] F. Bloch, *Phys. Rev. B* **2**, 109 (1970).
 [19] A. J. Leggett, in *Granular Nanoelectronics*, edited by D. K. Ferry, R. Barker, and C. Jacoboni, NATO ASI Ser. B Vol. 251 (Plenum, New York, 1991), p. 297.
 [20] D. Loss, *Phys. Rev. Lett.* **69**, 343 (1992).
 [21] X. Waintal, G. Fleury, K. Kazymyrenko, M. Houzet, P. Schmitteckert, and D. Weinmann, *Phys. Rev. Lett.* **101**, 106804 (2008).
 [22] E. H. Lieb and F. Y. Wu, *Phys. Rev. Lett.* **20**, 1445 (1968); **21**, 192(E) (1968).
 [23] A. A. Zvyagin, *Fiz. Tverd. Tela (Leningrad)* **32**, 1546 (1990) [*Sov. Phys. Solid State* **32**, 905 (1990)].
 [24] B. S. Shastry and B. Sutherland, *Phys. Rev. Lett.* **65**, 243 (1990).
 [25] F. H. L. Essler, H. Frahm, F. Göhmann, A. Klümper, and V. E. Korepin, *The One-Dimensional Hubbard Model* (Cambridge University Press, Cambridge, England, 2005).
 [26] D. Loss and P. Goldbart, *Phys. Rev. B* **43**, 13762(R) (1991).
 [27] H.-F. Cheung, Y. Gefen, E. K. Riedel, and W.-H. Shih, *Phys. Rev. B* **37**, 6050 (1988).
 [28] M. Ogata and H. Shiba, *Phys. Rev. B* **41**, 2326 (1990).
 [29] F. V. Kusmartsev, *J. Phys. Condens. Matter* **3**, 3199 (1991).
 [30] N. Yu and M. Fowler, *Phys. Rev. B* **45**, 11795 (1992).
 [31] W. J. Chetcuti, T. Haug, L.-C. Kwek, and L. Amico, *SciPost Phys.* **12**, 033 (2022).
 [32] D. V. Averin and C. Bruder, *Phys. Rev. B* **98**, 085304 (2018).
 [33] F. V. Kusmartsev, J. F. Weisz, R. Kishore, and M. Takahashi, *Phys. Rev. B* **49**, 16234 (1994).
 [34] A. Berkovich and J. H. Lowenstein, *Nucl. Phys.* **B285**, 70 (1987).
 [35] A. Berkovich, *J. Phys. A* **24**, 1543 (1991).
 [36] V. V. Cheianov and M. B. Zvonarev, *Phys. Rev. Lett.* **92**, 176401 (2004).
 [37] V. V. Cheianov and M. B. Zvonarev, *J. Phys. A* **37**, 2261 (2004).
 [38] K. A. Matveev, *Phys. Rev. Lett.* **92**, 106801 (2004).
 [39] G. A. Fiete and L. Balents, *Phys. Rev. Lett.* **93**, 226401 (2004).
 [40] G. A. Fiete, *Rev. Mod. Phys.* **79**, 801 (2007).
 [41] J. des Cloizeaux and J. J. Pearson, *Phys. Rev.* **128**, 2131 (1962).
 [42] F. V. Kusmartsev, *Phys. Rev. B* **52**, 14445 (1995).
 [43] F. Woyarovich, *J. Phys. A* **22**, 4243 (1989).
 [44] A. A. Zvyagin, *Finite Size Effects in Correlated Electron Models: Exact Results* (Imperial College Press, London, England, 2005).

- [45] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.128.096801> for the strong coupling limit of the BAEs, the definitions of the charge matrix and velocities, and a discussion on the N_{\downarrow}/N periodicity, which includes Ref. [46].
- [46] A. G. Izergin, V. E. Korepin, and N. Yu. Reshetikhin, *J. Phys. A* **22**, 2615 (1989).
- [47] H. Frahm and V. E. Korepin, *Phys. Rev. B* **42**, 10553 (1990).
- [48] H. Frahm and V. E. Korepin, *Phys. Rev. B* **43**, 5653 (1991).
- [49] M. Karbach and G. Muller, *Comput. Phys.* **11**, 36 (1997).
- [50] R. Hagemans and J.-S. Caux, *J. Phys. A* **40**, 14605 (2007).
- [51] W. Hao, R. I. Nepomechie, and A. J. Sommes, *Phys. Rev. E* **88**, 052113 (2013).
- [52] M. V. Moskalets and P. S. Deo, *Phys. Rev. B* **62**, 6920 (2000).
- [53] M. V. Moskalets, *Physica (Amsterdam)* **291B**, 75 (2000).
- [54] V. V. Cheianov, H. Smith, and M. B. Zvonarev, *Phys. Rev. A* **71**, 033610 (2005).
- [55] A. Ramanathan, K. C. Wright, S. R. Muniz, M. Zelan, W. T. Hill III, C. J. Lobb, K. Helmerson, W. D. Phillips, and G. K. Campbell, *Phys. Rev. Lett.* **106**, 130401 (2011).
- [56] L. Amico, A. Osterloh, and F. Cataliotti, *Phys. Rev. Lett.* **95**, 063201 (2005).
- [57] C.-C. Chien, S. Peotta, and M. Di Ventra, *Nat. Phys.* **11**, 998 (2015).
- [58] L. Amico *et al.*, *AVS Quantum Sci.* **3**, 039201 (2021).
- [59] L. Amico, D. Anderson, M. Boshier, J.-P. Brantut, L.-C. Kwek, A. Minguzzi, and W. von Klitzing, *arXiv*: 2107.08561.
- [60] G. Pecci, P. Naldesi, L. Amico, and A. Minguzzi, *Phys. Rev. Research* **3**, L032064 (2021).
- [61] G. Pecci, P. Naldesi, A. Minguzzi, and L. Amico, *arXiv*: 2105.10408.
- [62] W. J. Chetcuti, J. Polo, A. Osterloh, P. Castorina, and L. Amico, *arXiv*:2112.06950.