

Symmetry of Cosmological Observables, a Mirror World Dark Sector, and the Hubble Constant

Francis-Yan Cyr-Racine¹, Fei Ge², and Lloyd Knox²

¹*Department of Physics and Astronomy, University of New Mexico, Albuquerque, New Mexico 87106, USA*

²*Department of Physics and Astronomy, University of California, Davis, California 95616, USA*

 (Received 30 July 2021; revised 21 December 2021; accepted 6 April 2022; published 18 May 2022)

We find that a uniform scaling of the gravitational free-fall rates and photon-electron scattering rate leaves most dimensionless cosmological observables nearly invariant. This result opens up a new approach to reconciling cosmic microwave background and large-scale structure observations with high values of the Hubble constant H_0 : Find a cosmological model in which the scaling transformation can be realized without violating any measurements of quantities not protected by the symmetry. A “mirror world” dark sector allows for effective scaling of the gravitational free-fall rates while respecting the measured mean photon density today. Further model building might bring consistency with the two constraints not yet satisfied: the inferred primordial abundances of deuterium and helium.

DOI: [10.1103/PhysRevLett.128.201301](https://doi.org/10.1103/PhysRevLett.128.201301)

Introduction.—Different methodologies for determining the current rate of expansion, the Hubble constant H_0 , are leading to discrepant results. The most precise of the cosmological-model-dependent methods uses the *Planck* measurements of the cosmic microwave background (CMB). Assuming Lambda cold dark matter (Λ CDM), the result is $H_0 = (67.49 \pm 0.53)$ km/sec/Mpc [1–3]. The most precise of the more direct methods that are relatively independent of cosmological model assumptions comes from the SH₀ES team [4–7]. Using Cepheid-calibrated supernovae, they find $H_0 = (73.04 \pm 1.04)$ km/sec/Mpc ([8] hereafter R22), a 5σ difference from the above result.

The preference of Λ CDM-dependent methods for a low H_0 is remarkably robust to choice of datasets and details of applications; it remains for both the “inverse distance-ladder” methods [9–15] that make minimal use of CMB data, and methods that rely on large-scale structure observations instead of CMB data [16–21].

Likewise, various averages of the more direct methodologies [22–34], including those that exclude Cepheid-calibrated supernovae, also lead to $> 4\sigma$ discrepancies with the CMB- and Λ CDM-inferred value of H_0 [35]. On the other hand, there is no convergence of opinion yet among those using supernovae to measure H_0 . Most notably, the Carnegie-Chicago Hubble Program [36–45] finds some inconsistencies in SH₀ES estimates of distances to nearby supernovae. For a review of the observational situation, see Refs. [30,46].

A lot of recent theoretical work is inspired by the possibility that the tension arises from a failure of Λ CDM. For a wide-ranging discussion of possible avenues for a solution, see Ref. [47], for a summary of many of the proposed models, see Ref. [48], and for a ranking of their performance with respect to a common dataset,

see Ref. [49]. The search is difficult due to the variety of cosmological measurements, their sensitivity to the details of the cosmological models, their high precision, and their high degree of consistency with Λ CDM.

In this Letter, we provide new insight into the structure of cosmological models to help theorists navigate this challenging environment, as they search for solutions to the Hubble tension. In particular, we present a transformation that leaves all distance ratios and the statistical properties of fractional maps of CMB temperature anisotropy (i.e., $\Delta T/T$), CMB polarization, and galaxy number overdensity invariant. This symmetry transformation, under which all relevant length and timescales in the problem are rescaled by a constant scaling factor λ at all redshifts z [50], has its roots in the scale-free nature of primordial fluctuations.

A restricted version of this transformation was introduced in Ref. [51], in which only the gravitational timescales $1/\sqrt{G\rho_i}$, where ρ_i is the mean density of the i th component, were scaled. This leads to an approximate symmetry that is severely broken at small angular scales and in polarization on all scales. By extending their transformation to include a scaling of the photon scattering rate, we have found a symmetry of the above-mentioned observables that is exact in the limit of equilibrium recombination and zero neutrino mass. Crucially, we find that symmetry-breaking effects of real-world departures from these limits are mild.

We also report here on partial progress toward the use of this symmetry to solve the H_0 tension. Significant observational constraints prevent us from a straightforward scaling of gravitational rates by the scaling of mean densities. To evade these constraints and still reap the benefits of the scaling transformation, we are led quite naturally to the addition of a mirror world dark sector.

Dark sector models that contain a mirror (or “twin”) sector [52] that has the same particle content and gauge interactions as the standard model (SM), have been extensively studied in the literature (see, e.g., Refs. [56–86]) in the context of the little hierarchy problem. To scale the photon scattering rate, one can alter the primordial fraction of baryonic mass in helium Y_p , which is the method we use here. We show below that such models can, in principle, exploit the scaling symmetry to accommodate a higher value of H_0 with CMB observations.

A strict implementation of the scaling transformation, however, leads to a conflict between the measurements of Y_p and the primordial deuterium abundance with expectations from big bang nucleosynthesis (BBN) as well as the Y_p necessary for scaling up the scattering rate. Thus, we have essentially remapped the problem that has proven to be quite difficult of reconciling a high H_0 with phenomenologically complicated CMB data, to a problem of reconciling a high H_0 with the observationally inferred primordial abundances of helium and deuterium. Our work motivates the search for models that can solve this reframed problem [87]. Finally, although we focus here on inferences from CMB observations, *all* of the cosmological-model-dependent inferences of H_0 we referred to above, including those in Refs. [9–21], would be impacted similarly; i.e., replacing Λ CDM with a model that solves our reframed problem would also reconcile these inferences with a high H_0 due to the nature of the symmetry.

The scaling transformation.—Let us assume, for now, that recombination happens in equilibrium and neutrinos are massless. Then the only length scales in the linear perturbation evolution equations in the Λ CDM model written with the scale factor $a = 1/(1+z)$ as the independent timelike variable are the gravitational timescales of each of the $i = 1$ to N components $1/\sqrt{G\rho_i(a)}$ and the photon mean free path between electron scatters $1/[\sigma_T n_e(a)]$. As a result, if we consider the linear evolution of a single Fourier mode with wave number k , any fractional perturbation, such as $\delta\rho_m(\mathbf{k}, a)/\rho_m(a)$ satisfying the evolution equations will also satisfy them when transformed by a uniform scaling of all relevant (inverse) length scales (including k) by some factor λ . Since the initial conditions in Λ CDM do not introduce a length scale (the spectrum of initial perturbations is a power law), the statistical properties of fractional perturbations are independent of λ except for an overall amplitude. Dependence on the amplitude can be removed [51] by extending the scaling transformation to include $A_s \rightarrow A_s/\lambda^{(n_s-1)}$ where A_s is the amplitude of the primordial power spectrum at some fiducial value of k , and n_s is the spectral index of the power law power spectrum of initial density perturbations.

In Ref. [51], this transformation was introduced but without the photon scattering-rate scaling, which is critically important for our purposes. Including it, the transformation leads to an exact symmetry in the limit of

equilibrium recombination [88], massless neutrinos [89], and linearized equations [90] of distance ratios and the statistical properties of maps made in projection on the sky of quantities such as $\delta\rho/\rho$, and fractional CMB temperature and polarization anisotropies. These include galaxy clustering power spectra, shear power spectra, galaxy-shear cross-correlations, fractional CMB temperature and polarization power spectra, the temperature-polarization cross spectrum, and the CMB lensing spectrum. The invariance exists for the Λ CDM model, and any other model as long as additional length scales (if any, such as those related to mean curvature or neutrino mass) are properly scaled as well. Absent the introduction of new length scales, the full transformation can be written as

$$\begin{aligned} \sqrt{G\rho_i(a)} &\rightarrow \lambda\sqrt{G\rho_i(a)}, & \sigma_T n_e(a) &\rightarrow \lambda\sigma_T n_e(a), \\ \text{and } A_s &\rightarrow A_s/\lambda^{(n_s-1)}. \end{aligned} \quad (1)$$

Symmetry breaking.—The transformation given in Eq. (1) is severely constrained by observations that are sensitive to absolute densities of cosmological components. Most importantly, we know very precisely the mean energy density of the CMB today from measurements by FIRAS of its flux density across a broad range of wavelengths [93,94]. By anchoring ρ_γ , this measurement severely limits our ability to exploit the scaling transformation to raise H_0 [95].

Other important effects that break the above symmetry arise from departures from thermodynamic equilibrium, as emphasized in Ref. [51]. Unlike periods of equilibrium, during which we have no sensitivity to the rates of the reactions that are maintaining equilibrium, periods in which equilibrium is lost provide us with valuable sensitivity to the relevant reaction rates. If we then assume that such microphysical rates are known, we can gain sensitivity to the expansion rate. A prime example is BBN, where sensitivity of the yield of helium and deuterium to nuclear reaction rates allows one to infer, from measurements of helium and/or deuterium, the expansion rate during BBN, and thus, through the Friedmann equation, the mass or energy density of the Universe at that time. Similarly, hydrogen recombination is an out-of-equilibrium process which is sensitive to atomic reaction rates, and thus breaks the symmetry of the Eq. (1) transformation. We will see that the impact of this latter symmetry breaking on our parameter constraints is mild.

A mirror world dark sector and free Y_p .—Zahn and Zaldarriaga [51] did not provide a physical mechanism by which one could realize their transformation, other than by varying G , as their goal was only analytic understanding. We now introduce a physical mechanism that, while not allowing for the transformation as strictly written, permits a complete mimicry of its effects, so that the same invariance is achieved.

By extending the Λ CDM model to include a dark copy of the photons, baryons, and neutrinos (see, e.g., Refs. [97–117]), all with the same mean density ratios as in the visible sector, we can effectively mimic the $\sqrt{G\rho_i}$ part of the scaling transformation while evading the constraint from FIRAS. The dark photons (which have temperature T_D) are a replacement for the additional visible photons that would violate the FIRAS constraint. The dark baryons [implemented as “atomic dark matter” (ADM) [118–141]] allow us to scale up the total baryonlike density without changing the well-constrained (visible sector) baryon-to-photon ratio. The dark neutrinos allow us to scale up the effective number of free-streaming neutrino species from its Λ CDM value of $N_{\text{eff}}^{\text{FS}} = 3.046$ [142], preserving the well-constrained ratio of free-streaming to tightly coupled relativistic particle densities [144–146].

In our implementation, the new mirror dark sector interacts purely gravitationally with the visible sector and the CDM. Therefore, we can mimic a Λ CDM model with scaled-up densities if the perturbations in the mirror world evolve in the same way as what they replaced in the visible sector and thus provide the same source to the metric perturbations. For this to be the case, the dark photons must transition from tightly coupled to freely streaming when the visible photons do. We thus ensure that the ADM recombines at approximately the same time as regular hydrogen by keeping the ratio B_D/T_D fixed,

where B_D is the binding energy of the ADM, and by setting the dark fine structure constant and dark proton mass equal to those in the light sector. For simplicity, we assume there are no dark versions of helium or heavier nuclei. We can keep the Thomson scattering rate on the scaling trajectory by adjusting Y_p . At fixed baryon density, we have $n_e(z) \propto x_e(z)(1 - Y_p)$ where $x_e(z)$ is the fraction of free electrons. So to scale the scattering rate appropriately, approximating $x_e(z)$ as fixed, we send $(1 - Y_p) \rightarrow \lambda(1 - Y_p)$.

In Fig. 1, we show the temperature (TT) and polarization (EE) power spectra for the best-fit Λ CDM model given dataset 1 (defined below) as well as spectra for the models scaled from that one with $\lambda = 1.1$, with and without the photon-scattering-rate scaling included. The scaled spectra with no photon scattering scaling differ significantly from the Λ CDM spectra. The small differences between the fully scaled spectra and the Λ CDM spectra are primarily due to the symmetry-breaking effects of the atomic reaction rates affecting hydrogen recombination.

Results.—We now explore the parameter constraints in the framework just described. We define datasets 1 and 2 and model spaces *A*, *B*, and *C*, in Tables I and II. The results are shown in Fig. 2. In the upper left panel, we see the expected results from *A1*: The posterior is almost flat throughout the prior region [156]. This is the numerical manifestation of the exact symmetry we have presented.

For *B1*, the H_0 posterior remains quite broad. The soft symmetry-breaking effects of nonequilibrium recombination, evident in Fig. 1, are sufficiently degenerate with variation of other Λ CDM parameters to avoid significant constraint on H_0 . The tension with R21 has been completely eliminated. The very slight preference for $H_0 = 73.2$ km/s/Mpc in *B1* over the Λ CDM value is a parameter-volume effect; the best-fit Λ CDM model has a nearly identical χ^2 value as the best-fit *B* model constrained to $\lambda = 1.08$, lower by $\Delta\chi^2 = 0.2$.

In the other three panels of Fig. 2, we show how well departures away from scaling are constrained by the data if they are not prevented by fiat. The *B1* contours lie over the constraints given by model *C*; as expected, the scaling direction is preferred by the data. We also see that the region of high posterior probability density extends to parameter values far from the scaling solution, an indication




FIG. 1. CMB temperature (TT) and polarization (EE) power spectra with (red and green) and without (blue) scaling by $\lambda = 1.1$ from the best-fit Λ CDM model for dataset 1. Both $\lambda = 1.1$ cases are scaled by use of the mirror world dark sector, with (red) and without (green) scaling of the photon scattering rate. Data points are from *Planck* 2018 [147]. Fractional differences with the best-fit Λ CDM model are shown in the second and bottom panels.

TABLE I. Definition of datasets 1 and 2. We use the Plik TT + TE + EE, $\text{Low}l_T$, $\text{Low}l_E$, and lensing *Planck* likelihoods described in Ref. [147]. Baryon acoustic oscillations (BAO) datasets are 6dFGS [148], SDSS MGS [149], and BOSS DR12 [150]. R21 is the SH_0ES measurement [7], of which R22 is a recent update.

Label	Dataset
1	<i>Planck</i> TT, TE, EE, lensing + BAO
2	Dataset 1 plus R21 ($H_0 = 73.2 \pm 1.3$ km/s/Mpc)

TABLE II. Definition of model spaces A , B , and C . For model space A , the one non- Λ CDM parameter is λ . We restrict the additional components including Y_p to the scaling solution and (artificially) hold $x_e(z)$ fixed to its Λ CDM best-fit value for both the dark and light sectors. We scale Y_p from its BBN-consistent Λ CDM dataset 1 best-fit value of 0.2454. Model space B differs only in that we calculate the visible sector $x_e(z)$ using the atomic reaction rates (and the code RECFAST [151,152]) and the dark sector ionization evolution as in Ref. [128]. Model space C only differs from B in that we allow Y_p , the effective number of free-streaming neutrinos $N_{\text{eff}}^{\text{FS}}$, and the fraction of mirror world (or “atomic”) dark matter f_{ADM} to depart from their scaling values. For all model spaces, we adopt the uniform prior $1.00001 < \lambda < 1.3$, set the ratio of dark to light photon temperatures to $T_D/T_\gamma = (\lambda^2 - 1)^{1/4}$, and, although it introduces a new length scale, we take one of the neutrino species to have a mass of 0.06 eV [153]. We modified CAMB [154] to solve the relevant Einstein-Boltzmann equations and used COSMOMC [155] to calculate parameter posterior densities.

Label	Model space
A	Λ CDM + λ (scaling enforced), $x_e(z)$ fixed
B	Λ CDM + λ (scaling enforced), $x_e(z)$ calculated
C	Λ CDM + T_D + $N_{\text{eff}}^{\text{FS}}$ + f_{ADM} + Y_p

of some freedom that more detailed model building could exploit. For $C2$, we find that $f_{\text{ADM}} = 0.027 \pm 0.011$ and $T_D = (0.68 \pm 0.06)T_\gamma$. Meanwhile, the value of $\sigma_8 = 0.808 \pm 0.011$ is nearly unchanged (if not slightly lower) from its Λ CDM value, as expected.

In Fig. 2 we also see a problem: The Y_p values consistent with R21 are inconsistent with inferences from spectral observations of hot “metal-poor” gas such as the A21 finding of $Y_p = 0.2453 \pm 0.0034$. From $C2$ we have $Y_p = 0.170 \pm 0.025$, a 3.0σ difference. Compounding this trouble, the additional light relics, if we do not otherwise alter the standard thermal history, would increase the BBN-expected Y_p [158] in $C2$ to 0.2614 ± 0.0038 .

Discussion.—We have found important constraints on the project of turning the scaling transformation into a solution to the H_0 tension. The first of these, from FIRAS, can be accommodated with a mirror world dark sector. Constraints from light element abundances lead us to the articulation of two additional targets for model building: (i) a new mechanism for increasing the photon scattering rate and (ii) additional model features that would bring BBN predictions for helium and deuterium abundances in line with observations.

Item (i) follows from the fact that the Y_p required for consistency with R21 in models B and C is 3σ too low compared to the inference from observations in A21, hence, strongly suggesting that modifying Y_p is not a promising way to increase the photon scattering rate. Item (ii) follows since the extra relativistic species in the mirror sector required by R21 alter predictions of Y_p and deuterium.




FIG. 2. Constraints on parameters from datasets 1 and 2 given models A , B , and C (see Tables I and II). Top left panel: the (unnormalized) posterior probability density of H_0 . Other panels: the 68% and 95% contours of equal probability density in the $H_0 - Y_p$, $H_0 - f_{\text{ADM}}$, and $H_0 - T_D/T_\gamma$ planes. The gray band “R21” shows the 1σ and 2σ constraint on H_0 from R21. The purple band “A21” shows the same for Y_p from ([157] hereafter A21).

We have for $C2$ $\Delta N_{\text{eff}} = 1.3 \pm 0.34$. On the other hand, BBN consistency with Y_p and deuterium measurements leads to $\Delta N_{\text{eff}} = -0.17 \pm 0.28$ [87].

One idea worth exploring, to boost the photon scattering rate, is a heating of the baryons by a spectral distortion in the Wien tail at frequencies beyond FIRAS’s reach [159]. A time-varying electron mass [160,161] is another possible solution that has had some phenomenological success [160], and could even be well motivated by a supersymmetric gravity sector [162,163]. Successfully fitting the light element abundances could be achieved, for instance, by reheating the dark sector in the post-BBN era (see, e.g., Ref. [164]), or by possibly introducing interactions between the light and dark sectors that could affect the predicted yields; our inferred T_D/T_γ even puts decoupling of the dark and light sectors in the right ballpark for this to occur.

Our scenario requires a mirror sector that is phenomenologically close to the SM, albeit with a lower temperature. In its simplest realization, such a sector appears tightly constrained by particle collider data (see, e.g., Ref. [165]). Encouragingly however, preliminary work [166] indicates that such a mirror sector could be successfully built. More generally, our work opens the possibility of relaxing cosmological constraints (see, e.g., Refs. [68,72,78,167]) on such mirror scenarios, given mechanisms to adjust the photon scattering rate and light-element abundances.

We note that having $\sim 3\%$ of dark matter in ADM can lead to interesting astrophysical phenomenology such as exotic compact objects [168,169], dark stars [170,171], exotic gamma-bright point sources [172], and dark disks within galaxies [83,131–134,173–176].

Conclusions.—We have generalized a scaling transformation so that it now allows for large changes in the cosmological model while preserving the precisely measured and feature-rich CMB temperature and polarization spectra, as well as many other cosmological observables. Implementing this transformation, while evading constraints from FIRAS, leads us directly to a mirror world dark sector, a type of dark sector that has been proposed before for independent reasons.

While the scaling symmetry does not, by itself, provide an end-to-end solution to the H_0 tension, we have used it to single out the photon scattering rate and light-element abundances as players with important roles in the discrepancy. We have thus provided clear model-building targets for the community to explore.

F. G. and L. K. were partially supported by the U.S. Department of Energy Office of Science. F.-Y. C.-R is supported by the National Science Foundation (NSF) under Grant No. AST-2008696. We thank B. Fields, D. Green, D. Liu, and M. Luty for useful conversations and the UNM Center for Advanced Research Computing, supported in part by the NSF, for providing some of the computing resources used in this work. Part of this work was performed at the Aspen Center for Physics, which is supported by NSF Grant No. PHY-1607611.

-
- [1] L. Balkenhol *et al.* (SPT Collaboration), *Phys. Rev. D* **104**, 083509 (2021).
- [2] N. Aghanim *et al.* (Planck Collaboration), *Astron. Astrophys.* **641**, A6 (2020).
- [3] S. Aiola, E. Calabrese, L. Maurin, S. Naess, B. L. Schmitt, M. H. Abitbol, G. E. Addison, P. A. Ade, D. Alonso, M. Amiri *et al.*, *J. Cosmol. Astropart. Phys.* **12** (2020) 047.
- [4] A. G. Riess, L. Macri, S. Casertano, H. Lampeitl, H. C. Ferguson, A. V. Filippenko, S. W. Jha, W. Li, and R. Chornock, *Astrophys. J.* **730**, 119 (2011).
- [5] A. G. Riess *et al.*, *Astrophys. J.* **826**, 56 (2016).
- [6] A. G. Riess, S. Casertano, W. Yuan, L. M. Macri, and D. Scolnic, *Astrophys. J.* **876**, 85 (2019).
- [7] A. G. Riess, S. Casertano, W. Yuan, J. B. Bowers, L. Macri, J. C. Zinn, and D. Scolnic, *Astrophys. J. Lett.* **908**, L6 (2021).
- [8] A. G. Riess *et al.*, arXiv:2112.04510.
- [9] W. J. Percival, B. A. Reid, D. J. Eisenstein, N. A. Bahcall, T. Budavari, J. A. Frieman, M. Fukugita, J. E. Gunn, Ž. Ivezić, G. R. Knapp *et al.*, *Mon. Not. R. Astron. Soc.* **401**, 2148 (2010).
- [10] A. Heavens, R. Jimenez, and L. Verde, *Phys. Rev. Lett.* **113**, 241302 (2014).
- [11] É. Aubourg, S. Bailey, J. E. Bautista, F. Beutler, V. Bhardwaj, D. Bizyaev, M. Blanton, M. Blomqvist, A. S. Bolton, J. Bovy *et al.*, *Phys. Rev. D* **92**, 123516 (2015).
- [12] A. J. Cuesta, L. Verde, A. Riess, and R. Jimenez, *Mon. Not. R. Astron. Soc.* **448**, 3463 (2015).
- [13] J. L. Bernal, L. Verde, and A. G. Riess, *J. Cosmol. Astropart. Phys.* **10** (2016) 019.
- [14] L. Verde, J. L. Bernal, A. F. Heavens, and R. Jimenez, *Mon. Not. R. Astron. Soc.* **467**, 731 (2017).
- [15] P. Lemos, E. Lee, G. Efstathiou, and S. Gratton, *Mon. Not. R. Astron. Soc.* **483**, 4803 (2019).
- [16] T. M. C. Abbott *et al.* (DES Collaboration), *Mon. Not. R. Astron. Soc.* **480**, 3879 (2018).
- [17] G. D’Amico, J. Gleyzes, N. Kokron, K. Markovic, L. Senatore, P. Zhang, F. Beutler, and H. Gil-Marín, *J. Cosmol. Astropart. Phys.* **05** (2020) 005.
- [18] M. M. Ivanov, M. Simonović, and M. Zaldarriaga, *J. Cosmol. Astropart. Phys.* **05** (2020) 042.
- [19] T. Colas, G. D’Amico, L. Senatore, P. Zhang, and F. Beutler, *J. Cosmol. Astropart. Phys.* **06** (2020) 001.
- [20] O. H. E. Philcox, M. M. Ivanov, M. Simonović, and M. Zaldarriaga, *J. Cosmol. Astropart. Phys.* **05** (2020) 032.
- [21] P. Zhang, G. D’Amico, L. Senatore, C. Zhao, and Y. Cai, *J. Cosmol. Astropart. Phys.* **02** (2022) 036.
- [22] W. L. Freedman, B. F. Madore, V. Scowroft, C. Burns, A. Monson, S. E. Persson, M. Seibert, and J. Rigby, *Astrophys. J.* **758**, 24 (2012).
- [23] S. H. Suyu *et al.*, *Mon. Not. R. Astron. Soc.* **468**, 2590 (2017).
- [24] S. Birrer, T. Treu, C. E. Rusu, V. Bonvin, C. D. Fassnacht, J. H. H. Chan, A. Agnello, A. J. Shajib, G. C. F. Chen, M. Auger *et al.*, *Mon. Not. R. Astron. Soc.* **484**, 4726 (2019).
- [25] K. C. Wong *et al.*, *Mon. Not. R. Astron. Soc.* **498**, 1420 (2020).
- [26] C. D. Huang, A. G. Riess, W. Yuan, L. M. Macri, N. L. Zakamska, S. Casertano, P. A. Whitelock, S. L. Hoffmann, A. V. Filippenko, and D. Scolnic, *Astrophys. J.* **889**, 5 (2020).
- [27] E. Kourkchi, R. B. Tully, G. S. Anand, H. M. Courtois, A. Dupuy, J. D. Neill, L. Rizzi, and M. Seibert, *Astrophys. J.* **896**, 3 (2020).
- [28] M. J. Reid, D. W. Pesce, and A. G. Riess, *Astrophys. J. Lett.* **886**, L27 (2019).
- [29] W. L. Freedman, B. F. Madore, T. Hoyt, I. S. Jang, R. Beaton, M. G. Lee, A. Monson, J. Neeley, and J. Rich, *Astrophys. J.* **891**, 57 (2020).
- [30] W. L. Freedman, *Astrophys. J.* **919**, 16 (2021).
- [31] D. W. Pesce, J. A. Braatz, M. J. Reid, A. G. Riess, D. Scolnic, J. J. Condon, F. Gao, C. Henkel, C. M. V. Impellizzeri, C. Y. Kuo, and K. Y. Lo, *Astrophys. J. Lett.* **891**, L1 (2020).
- [32] N. Khetan *et al.*, *Astron. Astrophys.* **647**, A72 (2021).
- [33] J. P. Blakeslee, J. B. Jensen, C.-P. Ma, P. A. Milne, and J. E. Greene, *Astrophys. J.* **911**, 65 (2021).
- [34] S. Birrer *et al.*, *Astron. Astrophys.* **643**, A165 (2020).
- [35] E. Di Valentino, *Mon. Not. R. Astron. Soc.* **502**, 2065 (2021).
- [36] R. L. Beaton *et al.*, *Astrophys. J.* **832**, 210 (2016).

- [37] D. Hatt, R. L. Beaton, W. L. Freedman, B. F. Madore, I.-S. Jang, T. J. Hoyt, M. G. Lee, A. J. Monson, J. A. Rich, V. Scowcroft, and M. Seibert, *Astrophys. J.* **845**, 146 (2017).
- [38] D. Hatt, W. L. Freedman, B. F. Madore, R. L. Beaton, T. J. Hoyt, I. Sung Jang, M. Gyoon Lee, A. J. Monson, J. A. Rich, V. Scowcroft, and M. Seibert, *Astrophys. J.* **861**, 104 (2018).
- [39] D. Hatt, W. L. Freedman, B. F. Madore, I. Sung Jang, R. L. Beaton, T. J. Hoyt, M. Gyoon Lee, A. J. Monson, J. A. Rich, V. Scowcroft, and M. Seibert, *Astrophys. J.* **866**, 145 (2018).
- [40] C. R. Burns *et al.* (CSP Collaboration), *Astrophys. J.* **869**, 56 (2018).
- [41] T. J. Hoyt, W. L. Freedman, B. F. Madore, D. Hatt, R. L. Beaton, I. S. Jang, M. G. Lee, A. J. Monson, J. R. Neeley, J. A. Rich, and V. A. Mager, *Astrophys. J.* **882**, 150 (2019).
- [42] R. L. Beaton, M. Seibert, D. Hatt, W. L. Freedman, T. J. Hoyt, I. S. Jang, M. G. Lee, B. F. Madore, A. J. Monson, J. R. Neeley, J. A. Rich, and V. Scowcroft, *Astrophys. J.* **885**, 141 (2019).
- [43] W. L. Freedman, B. F. Madore, D. Hatt, T. J. Hoyt, I. S. Jang, R. L. Beaton, C. R. Burns, M. G. Lee, A. J. Monson, J. R. Neeley *et al.*, *Astrophys. J.* **882**, 34 (2019).
- [44] I. S. Jang, T. J. Hoyt, R. L. Beaton, W. L. Freedman, B. F. Madore, M. G. Lee, J. R. Neeley, A. J. Monson, J. A. Rich, and M. Seibert, *Astrophys. J.* **906**, 125 (2021).
- [45] T. J. Hoyt, R. L. Beaton, W. L. Freedman, I. S. Jang, M. G. Lee, B. F. Madore, A. J. Monson, J. R. Neeley, J. A. Rich, and M. Seibert, *Astrophys. J.* **915**, 34 (2021).
- [46] P. Shah, P. Lemos, and O. Lahav, *Astron. Astrophys. Rev.* **29**, 9 (2021).
- [47] L. Knox and M. Millea, *Phys. Rev. D* **101**, 043533 (2020).
- [48] E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D. F. Mota, A. G. Riess, and J. Silk, *Classical Quantum Gravity* **38**, 153001 (2021).
- [49] N. Schöneberg, G. F. Abellán, A. P. Sánchez, S. J. Witte, c. V. Poulin, and J. Lesgourgues, [arXiv:2107.10291](https://arxiv.org/abs/2107.10291).
- [50] Redshift is often used as a timelike variable in cosmology since the light that arrives here from a given event on our past light cone is stretched by the expansion by a factor of $1+z$, providing a monotonic relationship between time of emission and redshift z in a continuously expanding universe.
- [51] O. Zahn and M. Zaldarriaga, *Phys. Rev. D* **67**, 063002 (2003).
- [52] Models have also been proposed that contain multiple copies of the SM; see, e.g., Refs. [53–55].
- [53] G. Dvali, *Fortschr. Phys.* **58**, 528 (2010).
- [54] G. Dvali and M. Redi, *Phys. Rev. D* **80**, 055001 (2009).
- [55] N. Arkani-Hamed, T. Cohen, R. T. D’Agnolo, A. Hook, H. D. Kim, and D. Pinner, *Phys. Rev. Lett.* **117**, 251801 (2016).
- [56] Z. Chacko, H.-S. Goh, and R. Harnik, *Phys. Rev. Lett.* **96**, 231802 (2006).
- [57] Z. Chacko, Y. Nomura, M. Papucci, and G. Perez, *J. High Energy Phys.* **01** (2006) 126.
- [58] Z. Chacko, H.-S. Goh, and R. Harnik, *J. High Energy Phys.* **01** (2006) 108.
- [59] R. Barbieri, T. Gregoire, and L. J. Hall, [arXiv:hep-ph/0509242](https://arxiv.org/abs/hep-ph/0509242).
- [60] N. Craig and K. Howe, *J. High Energy Phys.* **03** (2014) 140.
- [61] N. Craig, A. Katz, M. Strassler, and R. Sundrum, *J. High Energy Phys.* **07** (2015) 105.
- [62] I. Garcia Garcia, R. Lasenby, and J. March-Russell, *Phys. Rev. D* **92**, 055034 (2015).
- [63] N. Craig and A. Katz, *J. Cosmol. Astropart. Phys.* **10** (2015) 054.
- [64] M. Farina, *J. Cosmol. Astropart. Phys.* **11** (2015) 017.
- [65] M. Farina, A. Monteux, and C. S. Shin, *Phys. Rev. D* **94**, 035017 (2016).
- [66] V. Prilepina and Y. Tsai, *J. High Energy Phys.* **09** (2017) 033.
- [67] R. Barbieri, L. J. Hall, and K. Harigaya, *J. High Energy Phys.* **11** (2016) 172.
- [68] N. Craig, S. Koren, and T. Trott, *J. High Energy Phys.* **05** (2017) 038.
- [69] J. Berger, K. Jedamzik, and D. G. E. Walker, *J. Cosmol. Astropart. Phys.* **11** (2016) 032.
- [70] Z. Chacko, N. Craig, P. J. Fox, and R. Harnik, *J. High Energy Phys.* **07** (2017) 023.
- [71] C. Csaki, E. Kuflik, and S. Lombardo, *Phys. Rev. D* **96**, 055013 (2017).
- [72] Z. Chacko, D. Curtin, M. Geller, and Y. Tsai, *J. High Energy Phys.* **09** (2018) 163.
- [73] G. Elor, H. Liu, T. R. Slatyer, and Y. Soreq, *Phys. Rev. D* **98**, 036015 (2018).
- [74] Y. Hochberg, E. Kuflik, and H. Murayama, *Phys. Rev. D* **99**, 015005 (2019).
- [75] A. Francis, R. J. Hudspith, R. Lewis, and S. Tulin, *J. High Energy Phys.* **12** (2018) 118.
- [76] K. Harigaya, R. McGehee, H. Murayama, and K. Schutz, *J. High Energy Phys.* **05** (2020) 155.
- [77] M. Ibe, A. Kamada, S. Kobayashi, T. Kuwahara, and W. Nakano, *Phys. Rev. D* **100**, 075022 (2019).
- [78] D. Dunsky, L. J. Hall, and K. Harigaya, *J. High Energy Phys.* **02** (2020) 078.
- [79] C. Csáki, C.-S. Guan, T. Ma, and J. Shu, *J. High Energy Phys.* **12** (2020) 005.
- [80] S. Koren and R. McGehee, *Phys. Rev. D* **101**, 055024 (2020).
- [81] J. Terning, C. B. Verhaaren, and K. Zora, *Phys. Rev. D* **99**, 095020 (2019).
- [82] L. Johns and S. Koren, [arXiv:2012.06591](https://arxiv.org/abs/2012.06591).
- [83] J.-S. Roux and J. M. Cline, *Phys. Rev. D* **102**, 063518 (2020).
- [84] A. C. Ritter and R. R. Volkas, *Phys. Rev. D* **104**, 035032 (2021).
- [85] D. Curtin and S. Gryba, *J. High Energy Phys.* **08** (2021) 009.
- [86] D. Curtin, S. Gryba, J. Setford, D. Hooper, and J. Scholtz, *Phys. Rev. D* **105**, 035033 (2022).
- [87] B. D. Fields, K. A. Olive, T.-H. Yeh, and C. Young, *J. Cosmol. Astropart. Phys.* **03** (2020) 010; **11** (2020) E02.
- [88] We could drop this caveat by extending the transformation further so that the atomic reaction rates scale appropriately as well. Doing so would also extend the challenge (significantly) of finding a model for which a direction in the parameter space corresponds to the scaling transformation.

- [89] The very mild symmetry breaking from nonzero neutrino masses could be removed by also scaling them by a factor of λ .
- [90] We include this caveat because we do not have proof that the symmetry holds in the full nonlinear theory. It might. We do know it holds to second order in perturbations; see Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.128.201301> for a discussion of the symmetry at first and second order in perturbation theory, which includes Refs. [51,91,92].
- [91] C.-P. Ma and E. Bertschinger, *Astrophys. J.* **455**, 7 (1995).
- [92] N. Bartolo, S. Matarrese, and A. Riotto, [arXiv:astro-ph/0703496](https://arxiv.org/abs/astro-ph/0703496).
- [93] D. J. Fixsen, E. S. Cheng, J. M. Gales, J. C. Mather, R. A. Shafer, and E. L. Wright, *Astrophys. J.* **473**, 576 (1996).
- [94] D. Fixsen, *Astrophys. J.* **707**, 916 (2009).
- [95] A similar point was made recently in Ref. [96].
- [96] M. M. Ivanov, Y. Ali-Haïmoud, and J. Lesgourgues, *Phys. Rev. D* **102**, 063515 (2020).
- [97] S. I. Blinnikov and M. Khlopov, *Sov. Astron.* **27**, 371 (1983).
- [98] L. Ackerman, M. R. Buckley, S. M. Carroll, and M. Kamionkowski, *Phys. Rev. D* **79**, 023519 (2009).
- [99] J. L. Feng, M. Kaplinghat, H. Tu, and H.-B. Yu, *J. Cosmol. Astropart. Phys.* **07** (2009) 004.
- [100] P. Agrawal, F.-Y. Cyr-Racine, L. Randall, and J. Scholtz, *J. Cosmol. Astropart. Phys.* **05** (2017) 022.
- [101] R. Foot and S. Mitra, *Phys. Rev. D* **66**, 061301(R) (2002).
- [102] R. Foot and R. R. Volkas, *Phys. Rev. D* **68**, 021304(R) (2003).
- [103] R. Foot, *Int. J. Mod. Phys. D* **13**, 2161 (2004).
- [104] R. Foot and R. R. Volkas, *Phys. Rev. D* **70**, 123508 (2004).
- [105] R. Foot, A. Kobakhidze, K. L. McDonald, and R. R. Volkas, *Phys. Rev. D* **77**, 035006 (2008).
- [106] R. Foot, *Phys. Lett. B* **711**, 238 (2012).
- [107] R. Foot, *Phys. Rev. D* **88**, 023520 (2013).
- [108] R. Foot, *Int. J. Mod. Phys. A* **29**, 1430013 (2014).
- [109] R. Foot and S. Vagnozzi, *Phys. Rev. D* **91**, 023512 (2015).
- [110] R. Foot and S. Vagnozzi, *J. Cosmol. Astropart. Phys.* **07** (2016) 013.
- [111] P. Ciarcelluti, *Int. J. Mod. Phys. D* **14**, 187 (2005).
- [112] P. Ciarcelluti, *Int. J. Mod. Phys. D* **14**, 223 (2005).
- [113] P. Ciarcelluti and A. Lepidi, *Phys. Rev. D* **78**, 123003 (2008).
- [114] P. Ciarcelluti, *Int. J. Mod. Phys. D* **19**, 2151 (2010).
- [115] P. Ciarcelluti and Q. Wallemacq, *Phys. Lett. B* **729**, 62 (2014).
- [116] P. Ciarcelluti and Q. Wallemacq, *Adv. High Energy Phys.* **2014**, 148319 (2014).
- [117] J.-R. Cudell, M. Y. Khlopov, and Q. Wallemacq, *Mod. Phys. Lett. A* **29**, 1440006 (2014).
- [118] H. Goldberg and L. J. Hall, *Phys. Lett. B* **174**, 151 (1986).
- [119] D. Fargion, M. Khlopov, and C. A. Stephan, *Classical Quantum Gravity* **23**, 7305 (2006).
- [120] M. Y. Khlopov, *Pis'ma Zh. Eksp. Teor. Fiz.* **83**, 3 (2006).
- [121] M. Y. Khlopov and C. Kouvaris, *Phys. Rev. D* **78**, 065040 (2008).
- [122] D. E. Kaplan, G. Z. Krnjaic, K. R. Rehermann, and C. M. Wells, *J. Cosmol. Astropart. Phys.* **05** (2010) 021.
- [123] M. Y. Khlopov, A. G. Mayorov, and E. Y. Soldatov, *Int. J. Mod. Phys. D* **19**, 1385 (2010).
- [124] D. E. Kaplan, G. Z. Krnjaic, K. R. Rehermann, and C. M. Wells, *J. Cosmol. Astropart. Phys.* **10** (2011) 011.
- [125] M. Y. Khlopov, *Mod. Phys. Lett. A* **26**, 2823 (2011).
- [126] S. R. Behbahani, M. Jankowiak, T. Rube, and J. G. Wacker, *Adv. High Energy Phys.* **2011**, 709492 (2011).
- [127] J. M. Cline, Z. Liu, and W. Xue, *Phys. Rev. D* **85**, 101302 (R) (2012).
- [128] F.-Y. Cyr-Racine and K. Sigurdson, *Phys. Rev. D* **87**, 103515 (2013).
- [129] J. M. Cline, Z. Liu, G. D. Moore, and W. Xue, *Phys. Rev. D* **89**, 043514 (2014).
- [130] F.-Y. Cyr-Racine, R. de Putter, A. Raccanelli, and K. Sigurdson, *Phys. Rev. D* **89**, 063517 (2014).
- [131] J. J. Fan, A. Katz, L. Randall, and M. Reece, *Phys. Rev. Lett.* **110**, 211302 (2013).
- [132] J. Fan, A. Katz, L. Randall, and M. Reece, *Phys. Dark Universe* **2**, 139 (2013).
- [133] M. McCullough and L. Randall, *J. Cosmol. Astropart. Phys.* **10** (2013) 058.
- [134] L. Randall and J. Scholtz, *J. Cosmol. Astropart. Phys.* **09** (2015) 057.
- [135] M. Y. Khlopov, *Int. J. Mod. Phys. A* **29**, 1443002 (2014).
- [136] L. Pearce, K. Petraki, and A. Kusenko, *Phys. Rev. D* **91**, 083532 (2015).
- [137] J. Choquette and J. M. Cline, *Phys. Rev. D* **92**, 115011 (2015).
- [138] K. Petraki, L. Pearce, and A. Kusenko, *J. Cosmol. Astropart. Phys.* **07** (2014) 039.
- [139] M. Cirelli, P. Panci, K. Petraki, F. Sala, and M. Taoso, *J. Cosmol. Astropart. Phys.* **05** (2017) 036.
- [140] K. Petraki, M. Postma, and J. de Vries, *J. High Energy Phys.* **04** (2017) 077.
- [141] D. Curtin and J. Setford, *J. High Energy Phys.* **03** (2021) 166.
- [142] We note that using the more recent estimate of $N_{\text{eff}}^{\text{FS}} = 3.044$ [143] would have minimal impact on our results.
- [143] J. Froustey, C. Pitrou, and M. C. Volpe, *J. Cosmol. Astropart. Phys.* **12** (2020) 015.
- [144] Z. Hou, R. Keisler, L. Knox, M. Millea, and C. Reichardt, *Phys. Rev. D* **87**, 083008 (2013).
- [145] B. Follin, L. Knox, M. Millea, and Z. Pan, *Phys. Rev. Lett.* **115**, 091301 (2015).
- [146] D. Baumann, D. Green, J. Meyers, and B. Wallisch, *J. Cosmol. Astropart. Phys.* **01** (2016) 007.
- [147] N. Aghanim *et al.* (Planck Collaboration), *Astron. Astrophys.* **641**, A5 (2020).
- [148] F. Beutler, C. Blake, M. Colless, D. H. Jones, L. Staveley-Smith, L. Campbell, Q. Parker, W. Saunders, and F. Watson, *Mon. Not. R. Astron. Soc.* **416**, 3017 (2011).
- [149] A. J. Ross, L. Samushia, C. Howlett, W. J. Percival, A. Burden, and M. Manera, *Mon. Not. R. Astron. Soc.* **449**, 835 (2015).
- [150] S. Alam *et al.* (BOSS Collaboration), *Mon. Not. R. Astron. Soc.* **470**, 2617 (2017).
- [151] S. Seager, D. D. Sasselov, and D. Scott, *Astrophys. J. Lett.* **523**, L1 (1999).
- [152] W. Y. Wong, A. Moss, and D. Scott, *Mon. Not. R. Astron. Soc.* **386**, 1023 (2008).

- [153] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz, and A. Zhou, *J. High Energy Phys.* **09** (2020) 178.
- [154] A. Lewis, A. Challinor, and A. Lasenby, *Astrophys. J.* **538**, 473 (2000).
- [155] A. Lewis and S. Bridle, *Phys. Rev. D* **66**, 103511 (2002).
- [156] This prior (see Table II caption) is made necessary by our use of a mirror dark sector to mimic a Λ CDM model with scaled-up densities. It ensures that the energy densities of dark photons and dark atoms are always positive.
- [157] E. Aver, D. A. Berg, K. A. Olive, R. W. Pogge, J. J. Salzer, and E. D. Skillman, *J. Cosmol. Astropart. Phys.* **03** (2021) 027.
- [158] R. Consiglio, P. F. de Salas, G. Mangano, G. Miele, S. Pastor, and O. Pisanti, *Comput. Phys. Commun.* **233**, 237 (2018).
- [159] J. Chluba (private communication).
- [160] T. Sekiguchi and T. Takahashi, *Phys. Rev. D* **103**, 083507 (2021).
- [161] L. Hart and J. Chluba, *Mon. Not. R. Astron. Soc.* **493**, 3255 (2020).
- [162] C. P. Burgess and F. Quevedo, *J. Cosmol. Astropart. Phys.* **04** (2022) 007.
- [163] C. P. Burgess, D. Dineen, and F. Quevedo, *J. Cosmol. Astropart. Phys.* **03** (2022) 064.
- [164] D. Aloni, A. Berlin, M. Joseph, M. Schmaltz, and N. Weiner, [arXiv:2111.00014](https://arxiv.org/abs/2111.00014).
- [165] G. Burdman, Z. Chacko, R. Harnik, L. de Lima, and C. B. Verhaaren, *Phys. Rev. D* **91**, 055007 (2015).
- [166] N. Blinov, G. Krnjaic, and S. W. Li, [arXiv:2108.11386](https://arxiv.org/abs/2108.11386).
- [167] S. Bansal, J. H. Kim, C. Kolda, M. Low, and Y. Tsai, [arXiv:2110.04317](https://arxiv.org/abs/2110.04317).
- [168] S. Shandera, D. Jeong, and Henry S. Grasshorn Gebhardt, *Phys. Rev. Lett.* **120**, 241102 (2018).
- [169] D. Singh, M. Ryan, R. Magee, T. Akhter, S. Shandera, D. Jeong, and C. Hanna, *Phys. Rev. D* **104**, 044015 (2021).
- [170] D. Curtin and J. Setford, *Phys. Lett. B* **804**, 135391 (2020).
- [171] D. Curtin and J. Setford, *J. High Energy Phys.* **03** (2020) 041.
- [172] P. Agrawal and L. Randall, *J. Cosmol. Astropart. Phys.* **12** (2017) 019.
- [173] E. D. Kramer and L. Randall, *Astrophys. J.* **829**, 126 (2016).
- [174] E. D. Kramer and L. Randall, *Astrophys. J.* **824**, 116 (2016).
- [175] K. Schutz, T. Lin, B. R. Safdi, and C.-L. Wu, *Phys. Rev. Lett.* **121**, 081101 (2018).
- [176] J. Buch, S. C. J. Leung, and J. Fan, *J. Cosmol. Astropart. Phys.* **04** (2019) 026.