

Anomalous Transport of Tracers in Active Baths

Omer Granek¹, Yariv Kafri¹, and Julien Tailleur²

¹*Department of Physics, Technion-Israel Institute of Technology, Haifa 3200003, Israel*

²*Université de Paris, Laboratoire Matière et Systèmes Complexes (MSC), UMR 7057 CNRS, F-75205 Paris, France*

 (Received 18 August 2021; revised 23 December 2021; accepted 21 June 2022; published 14 July 2022)

We derive the long-time dynamics of a tracer immersed in a one-dimensional active bath. In contrast to previous studies, we find that the damping and noise correlations possess long-time tails with exponents that depend on the tracer symmetry. For generic tracers, shape asymmetry induces ratchet effects that alter fluctuations and lead to superdiffusion and friction that grows with time when the tracer is dragged at a constant speed. In the singular limit of a completely symmetric tracer, we recover normal diffusion and finite friction. Furthermore, for small symmetric tracers, the active contribution to the friction becomes negative: active particles enhance motion rather than oppose it. These results show that, in low-dimensional systems, the motion of a passive tracer in an active bath cannot be modeled as a persistent random walker with a finite correlation time.

DOI: 10.1103/PhysRevLett.129.038001

Since Einstein and Smoluchowski, the motion of a tracer particle in a bath has been a topic of much interest [1]. The simplest textbook framework models the motion of the particle as a memoryless Brownian motion using an underdamped Langevin equation [2–4]. The momentum autocorrelation function then decays exponentially with a single timescale, signaling a transition between inertial and viscous regimes. This was, however, found to be oversimplistic: the conservation of momentum in the solvent instead leads to a power-law decay [5–7] and a host of interesting phenomena—especially in low dimensions—such as the breakdown of the Fourier law [8–10].

When compared with the equilibrium case, active fluids reveal a much richer physics, from the ratchet effects induced by asymmetric gears [11–14] and rectifiers [15–20] to the long-ranged forces and currents generated by asymmetric obstacles [20–24]. Over the past two decades, much activity has been devoted to studying passive tracers in active baths [25–63]. In the adiabatic limit in which the bath’s relaxation is much faster than the tracer’s response [64–71], the tracer’s dynamics is described by a generalized Langevin equation. In 1D, it reads as

$$\gamma_0 \dot{X}(t) + \int_0^t dt' \gamma(t-t') \dot{X}(t') = \mathcal{F}(t) + \eta(t), \quad (1)$$

where the interactions with the active particles lead to a stochastic force $\mathcal{F}(t)$ and a retarded friction $\int_0^t dt' \gamma(t-t') \dot{X}(t')$. Equation (1) also includes a memoryless viscous medium at temperature T that leads to the friction coefficient γ_0 and a Gaussian white noise $\eta(t)$ satisfying $\langle \eta(t)\eta(t') \rangle = 2\gamma_0 T \delta(t-t')$. Despite many efforts, a single unifying picture for the friction $\gamma(t)$ and

the force-force correlation functions $C_{\mathcal{F}} \equiv \langle \mathcal{F}(t)\mathcal{F}(0) \rangle_c$ does not emerge from the existing results.

First, a large class of experimental and numerical studies has suggested that the random, finite-duration encounters between the bath particles and the tracer lead to an exponential decay of $\gamma(t)$ over a short timescale [25–35]. Equation (1) then reduces to $(\gamma_0 + \gamma_T)\dot{X}(t) = \mathcal{F}(t)$, where $\gamma_T \equiv \int_0^\infty dt \gamma(t)$. In this case, similarly to an underdamped Brownian particle, the large-scale motion of the tracer is diffusive. This has been justified analytically in the simple case of a tracer connected by linear springs to a bath of active Ornstein-Uhlenbeck particles [72]—an active counterpart to the celebrated work of Vernon and Feynman [73–75].

In contrast, a second class of experiments and models on so-called wet-active matter suggests a more complex physics [37,41,43,45,50,55,59]. The long-ranged decay of hydrodynamic interactions can indeed turn $\gamma(t)$ and $C_{\mathcal{F}}(t)$ into power laws [37,45,59]. These may lead to anomalous diffusion on intermediate timescales but, ultimately, lead to long-time diffusion.

We note, however, that long-time tails are generic, even in the absence of hydrodynamic interactions. Indeed, the fluctuating density of active particles is a conserved quantity—and hence a slow field—so that the bath cannot have a single characteristic relaxation time. This leads to power-law memory and correlations, as already noted for equilibrium [6,7,76,77] and nonequilibrium [78,79] systems, including phoretic colloids [80] and driven tracers [81,82]. In low-dimensional systems, these tails may result in anomalous transport over long timescales [80,81]. Although thoroughly studied in other contexts, these effects were so far overlooked for tracers in dry active baths.




FIG. 1. (a) A large tracer and a bath of small active particles are immersed in a viscous medium inside a long narrow channel. (b) The short transverse dimension allows one to model the channel as a one-dimensional system where particles can bypass each other and the tracer, even though the transverse and orientational fluctuations of the tracer are lost in this one-dimensional description. Top: asymmetric tracer. Bottom: symmetric tracer.

In this Letter, to resolve this issue, we consider the simplest nontrivial system in which Eq. (1) can be systematically derived: a single tracer immersed in a dry one-dimensional active bath of run-and-tumble particles. To remain as close as possible to the phenomenology of an active bath in $d > 1$ dimensions, we allow particles to overtake each other and the tracer, hence modeling the latter by a soft repulsive potential $V(x)$; see Fig. 1. Starting from the coupled dynamics of the bath particles and tracer positions, $\{x_i(t), X(t)\}$, we determine explicitly the long-time behaviors of $\gamma(t)$ and $C_{\mathcal{F}}(t)$ as functions of the tracer shape and of the microscopic parameters of our model. To do so, we employ a *controlled* adiabatic expansion [83,84] valid in the large γ_0 limit in which the tracer dynamics can be described by Eq. (1). Our results show the emergence of long-time tails that lead to interesting and qualitatively different behaviors for symmetric and asymmetric tracers. For generic, *asymmetric* tracers, ratchet effects make $\gamma(t)$ and $C_{\mathcal{F}}(t)$ scale as $\sim t^{-1/2}$ in the long-time limit, leading to *superdiffusive* behavior around their mean displacements:

$$\langle X^2(t) \rangle_c \equiv \langle X^2(t) \rangle - \langle X(t) \rangle^2 \sim Kt^{3/2}. \quad (2)$$

When the tracer is towed at a constant velocity U , it experiences a friction force from the active particles that grows as

$$\frac{f_{\text{fric}}(t)}{U} \sim -\Gamma_T t^{1/2}. \quad (3)$$

We provide below explicit expressions for K and Γ_T in the presence of a soft asymmetric potential in a dilute active bath. In the singular limit of a *symmetric* tracer, $C_{\mathcal{F}}(t)$ and $\gamma(t)$ scale as $\sim t^{-3/2}$, similar to a tracer in a bath of equilibrium Brownian particles [76,85], which yields a *diffusive* behavior:

$$\langle X^2(t) \rangle_c \sim 2Dt. \quad (4)$$




FIG. 2. Simulation results (symbols) compared with our theoretical predictions for the long-time limit, without any fitting parameters, (dashed black lines): (a) mean squared displacement for symmetric and asymmetric tracers; (b) friction force exerted on an asymmetric tracer; (c) symmetric-tracer friction coefficient vs tracer size L_T . Simulation details and results for soft repulsive interactions are given in Sec. I of the Supplemental Material [87].

Towing the tracer at constant velocity U , the active particles exert a *finite* friction force:

$$\frac{f_{\text{fric}}(t)}{U} = -\gamma_T - \gamma_1 t^{-1/2} + \mathcal{O}(t^{-3/2}), \quad (5)$$

where $\gamma_T \equiv \int_0^\infty dt \gamma(t)$. Interestingly, for small tracer sizes, γ_T and γ_1 are negative: the active bath pushes the tracer in the towing direction. We provide perturbative expressions for D and γ_T and defer their systematic derivations for later work [86]. All our results are confirmed by microscopic simulations shown in Fig. 2. The derivation presented below suggests that the exponents are *universal* to any bath with long-time diffusive statistics. We confirm that they hold in the presence of soft repulsive interparticle forces in Sec. I of the Supplemental Material [87].

Model.—We consider bath particles moving with speed v and randomly switching their orientations with rate $\alpha/2$, leading to a persistence length $\ell_p = v/\alpha$. The tracer interacts with the active bath via a short-range potential V which vanishes outside $[0, L_T]$, such that the force on bath particle i is $f(x_i - X) = -\partial_{x_i} V(x_i - X)$ and the tracer size is L_T . We take $|\mu f(x)| < v$ so that particles are able to cross the tracer, which emulates the channel in Fig. 1(a). The tracer and bath-particle dynamics thus read as

$$\gamma_0 \dot{X}(t) = F_{\text{tot}}(t) \equiv -\sum_i f[x_i(t) - X(t)], \quad (6)$$

$$\dot{x}_i(t) = v\sigma_i(t) + \mu f[x_i(t) - X(t)], \quad (7)$$

where the $\sigma_i(t) \in \{\pm 1\}$ flip independently with rate $\alpha/2$ and μ is the bath-particle mobility. In Eqs. (6) and (7) we neglected the thermal noises acting on the tracer and bath particles, which are typically much weaker than the active and viscous forces [25,26,34,38,89]. (See Sec. V of the Supplemental Material [87] for a discussion of the $T \neq 0$ case.) In the analytical derivations below we consider a dilute bath of active particles, without interparticle forces, in either infinite systems or periodic ones of size $L \gg L_T, \ell_p$.

Theory.—The fluctuating force $F_{\text{tot}}(t)$ differs from the average force F exerted on a tracer held *fixed*. This is due to both the tracer's motion and the stochasticity of the active bath. The average correction due to the tracer motion is characterized by $\gamma(t)$ in Eq. (1). Within an adiabatic perturbation theory $\gamma(t)$ is defined as

$$\langle F_{\text{tot}}(t) \rangle - F \equiv -\int_0^t dt' \gamma(t-t') \dot{X}(t'), \quad (8)$$

where the average is conditioned on a given realization of $\dot{X}(t)$. The fluctuations of F_{tot} are then characterized through

$$\mathcal{F}(t) \equiv F_{\text{tot}}(t) + \int_0^t dt' \gamma(t-t') \dot{X}(t'). \quad (9)$$

Adiabatic perturbation theory tells us that, when γ_0 is large, the statistics of $\mathcal{F}(t)$ are identical to those of the force exerted on a tracer held fixed [84]. Furthermore, it relates $\gamma(t)$ and $\mathcal{F}(t)$ through an Agarwal-Kubo-type formula [83]:

$$\gamma(t-t') = \langle \mathcal{F}(t) \partial_{X_0} \ln \rho_s[x(t') - X_0, \sigma(t')] \rangle^s. \quad (10)$$

Here, $\rho_s(x - X_0, \sigma)$ is the steady-state density of bath particles with orientation σ and displacement $x - X_0$ from a tracer held fixed at X_0 . The brackets $\langle \cdot \rangle^s$ represent an average with respect to ρ_s . In the following, we set $X_0 = 0$ without loss of generality. For an equilibrium bath at temperature T , $\langle \mathcal{F}(t) \rangle^s = 0$ and Eq. (10) reduces to the fluctuation-dissipation theorem (FDT) $\gamma(t) = C_{\mathcal{F}}(t)/T$ where $C_{\mathcal{F}}(t) = \langle \mathcal{F}(t) \mathcal{F}(0) \rangle^s$. Outside equilibrium, these constraints need not hold.

To characterize the tracer dynamics, we compute independently $F = \langle \mathcal{F}(t) \rangle^s$, $C_{\mathcal{F}}(t)$, and $\gamma(t-t')$. To do so, we start from the expression for the steady state of non-interacting run-and-tumble particles in the presence of an external force $f(x)$ [90,91]:

$$\rho_s(x, \sigma) = \frac{\frac{1}{2}\rho_L}{1 + \sigma \frac{\mu}{v} f(x)} \exp \left\{ \beta_{\text{eff}} \int_0^x dy \frac{f(y)}{1 - [\frac{\mu}{v} f(y)]^2} \right\}, \quad (11)$$

where ρ_L is the particle density at $x = 0^-$, $T_{\text{eff}} = v^2/\mu\alpha$ is the effective temperature, and $\beta_{\text{eff}} = 1/T_{\text{eff}}$. The steady-state density is $\rho_s(x) = \sum_{\sigma} \rho_s(x, \sigma)$.

Asymmetric tracer.—For an asymmetric tracer, the densities of active particles ρ_R and ρ_L at the right and left ends of the tracer differ and are given by $\rho_R = 2\rho_0/[1 + \exp(\beta_{\text{eff}}\varepsilon)]$ and $\rho_L = 2\rho_0/[1 + \exp(-\beta_{\text{eff}}\varepsilon)]$, where $\varepsilon \equiv -\int dx f(x)/\{1 - [\mu f(x)/v]^2\}$. The density difference then leads to a nonvanishing average force $F = -\int dx f(x) \rho_s(x)$ exerted on the tracer [21,92,93], which is given by

$$F = -T_{\text{eff}}(\rho_R - \rho_L) = 2T_{\text{eff}}\rho_0 \tanh\left(\frac{\varepsilon}{2T_{\text{eff}}}\right), \quad (12)$$

where we have introduced the average background density $\rho_0 = (\rho_R + \rho_L)/2$. Note that Eq. (12) is consistent with the ideal gas law applied to the left and right sides of the tracer.

The long-time behavior of $C_{\mathcal{F}}(t)$ and $\gamma(t)$ can be derived from the knowledge of the propagator $p(x, \sigma, t|x', \sigma', 0)$. In the long-time limit, the dynamics of the active particles are diffusive so that the support of $p(x, \sigma, t|x', \sigma', 0)$ spreads over a region of length $2b(t)$ around x' , where $b(t) \sim (\pi D_{\text{eff}} t)^{1/2}$ is a diffusive propagating front. For any $x - x' \ll b(t)$, and to leading order in $b(t)$, $p(x, \sigma, t|x', \sigma', 0)$ has relaxed to the normalized steady-state distribution $\rho_s(x, \sigma)/\sum_{\sigma} \int_{-b(t)}^{b(t)} dx \rho_s(x, \sigma)$. For $L_T \ll 2b(t)$, one can neglect the region inside the tracer in the integral so that $\sum_{\sigma} \int_{-b(t)}^{b(t)} dx \rho_s(x, \sigma) \sim (\rho_R + \rho_L)b(t)$, up to corrections of order $\mathcal{O}(L^{-1})$. Since $b(t) \sim (\pi D_{\text{eff}} t)^{1/2}$ we get

$$p(x, \sigma, t|x', \sigma', 0) \sim \frac{\rho_s(x, \sigma)}{\rho_R + \rho_L} (\pi D_{\text{eff}} t)^{-1/2}. \quad (13)$$

This heuristic result can be derived exactly, within the adiabatic limit, and its subleading correction can be shown to scale as $\mathcal{O}(t^{-3/2})$ (See Sec. II of the Supplemental Material [87]).

On long times, $p(x, \sigma, t|x', \sigma', 0)$ is independent of the initial coordinate (x', σ') . Therefore, two-point correlations are factorized in this limit. Furthermore, for N noninteracting particles, the forces exerted by different particles on the tracer are uncorrelated so that $C_{\mathcal{F}}(t) = N\{\langle f(t)f(0) \rangle^s - [\langle f(t) \rangle^s]^2\}$, where $f(t)$ is the force due to a single bath particle. Since $\langle f(t) \rangle^s = F/N$, $N[\langle f(t) \rangle^s]^2$ only contributes a correction of order $\mathcal{O}(L^{-1})$ to $C_{\mathcal{F}}(t)$. Using Eq. (13), $C_{\mathcal{F}}(t)$ can then be evaluated as

$$C_{\mathcal{F}}(t) = \sum_{\sigma\sigma'} \int dx dx' f(x) p(x, \sigma, t | x', \sigma', 0) f(x') \rho_s(x', \sigma') \quad (14)$$

$$= \frac{F^2}{\rho_R + \rho_L} (\pi D_{\text{eff}} t)^{-1/2} + \mathcal{O}(t^{-3/2}). \quad (15)$$

Similarly, we obtain from Eqs. (10) and (12)

$$\gamma(t) = \sum_{\sigma\sigma'} \int dx dx' f(x) p(x, \sigma, t | x', \sigma', 0) \partial_{x'} \rho_s(x', \sigma') \quad (16)$$

$$= \beta_{\text{eff}} \frac{F^2}{\rho_R + \rho_L} (\pi D_{\text{eff}} t)^{-1/2} + \mathcal{O}(t^{-3/2}). \quad (17)$$

Remarkably, the long-time regime satisfies an effective FDT $\gamma(t) = \beta_{\text{eff}} C_{\mathcal{F}}(t) + \mathcal{O}(t^{-3/2})$. We also note that Eqs. (11)–(17) hold in the infinite-system-size limit. For large-but-finite systems, they are complemented by $\mathcal{O}(L^{-1})$ corrections, as discussed in Sec. III of the Supplemental Material [87].

Equations (15) and (17) immediately show that the asymmetric tracer undergoes anomalous dynamics on long times. Indeed, the noise and friction intensities, defined as $I = \int_0^\infty dt C_{\mathcal{F}}(t)$ and $\gamma_T = \int_0^\infty dt \gamma(t)$ are infinite, hence leading to an ill-defined diffusivity $D \equiv I/(\gamma_0 + \gamma_T)^2$. To characterize the anomalous dynamics of the tracer we first consider its free motion. We define the tracer's mobility $B(t)$ through $X(t) = \int_0^t dt' B(t-t') \mathcal{F}(t')$, which leads to

$$\langle X(t)^2 \rangle_c = 2 \int_0^t dt_1 \int_0^{t_1} dt_2 B(t_1) B(t_2) C_{\mathcal{F}}(t_1 - t_2). \quad (18)$$

Since we are working in the large γ_0 limit, $B(t) \sim 1/\gamma_0$ [94]. Using Eq. (15) for $C_{\mathcal{F}}(t)$ then gives Eq. (2), hence implying *superdiffusion*, with

$$K = \frac{4F^2}{3\rho_0\gamma_0^2\sqrt{\pi D_{\text{eff}}}}. \quad (19)$$

In addition to anomalous diffusion, the asymmetric tracer experiences friction that grows with time, as shown by the following towing experiment. Setting a constant velocity $\dot{X} = U$ in Eq. (1), the friction exerted by the active particles on the tracer can be measured as $f_{\text{fric}}(t) \equiv \langle F_{\text{tot}} \rangle - F$. From Eqs. (8) and (17), we get

$$f_{\text{fric}}(t) = -U \int_0^t dt' \gamma(t') \sim -U \frac{F^2}{T_{\text{eff}}\rho_0} \left(\frac{t}{\pi D_{\text{eff}}} \right)^{1/2}, \quad (20)$$

which yields Eq. (5) with $\Gamma_T = F^2(\pi D_{\text{eff}})^{-1/2}/T_{\text{eff}}\rho_0$.

Symmetric tracer.—For a symmetric tracer, $F = 0$. Equations (15) and (17) then imply that $\gamma(t)$, $C_{\mathcal{F}}(t) = \mathcal{O}(t^{-3/2})$. In this case, I and γ_T remain finite so




FIG. 3. Consider a symmetric tracer (blue potential) and an active particle located at its left end at position \hat{x} at $t = 0$. The particle is shown in orange and magenta for $\hat{\sigma} = \pm 1$ respectively. At late times, the particle position is distributed as a Gaussian centered around $x_c = \hat{x} + \hat{\sigma} \ell_p$. For $\hat{\sigma} = 1$, when $\ell_p \gg L_T$, the anticorrelation between $f(x')$ and $f(x)$ leads to a negative contribution to $C_{\mathcal{F}}(t)$. Conversely, a $\hat{\sigma} = -1$ particle leads to a positive contribution to $C_{\mathcal{F}}$. Due to the polarization against the potential, $\hat{\sigma} = \pm 1$ occur with different probabilities. This leads to an overall negative $C_{\mathcal{F}}(t)$ for small L_T and a positive one for large sizes.

that $D = I/(\gamma_0 + \gamma_T)^2$ is well defined and Eq. (4) holds. We now present heuristic discussions of $C_{\mathcal{F}}(t)$ and $\gamma(t)$ that account for two important features: their scaling as $t^{-3/2}$ and their sign changes for small tracers. These results can be derived exactly, within the adiabatic limit, for piecewise linear potentials [86].

Consider a symmetric tracer of length L_T whose potential is depicted in Fig. 3. While our results can be derived exactly [86], we present here a simple argument which holds in the limit in which the edges of the tracer have a small width d and small slopes $\pm f_0$. Consider first a single particle located at the left end of the tracer, at $\hat{x} \simeq 0$, moving in the direction $\hat{\sigma}$. At long times, the probability distribution of its position x is a Gaussian centered around $\hat{\sigma} \ell_p$, of variance $2D_{\text{eff}}t$ (see Fig. 3). The force-force correlation of this particle can be computed as

$$c(\hat{\sigma}, t) = \frac{f_0^2 d}{\sqrt{4\pi D_{\text{eff}} t}} \left[e^{-\frac{\ell_p^2}{4D_{\text{eff}} t}} - e^{-\frac{(L_T - \hat{\sigma} \ell_p)^2}{4D_{\text{eff}} t}} \right], \quad (21)$$

as can be inferred from Eq. (14) using $\rho_s(x', \sigma') = \delta(x') \delta_{\sigma', \hat{\sigma}}$. Note that the factor d comes from the integration over x in Eq. (14), which also leads to the two exponentials corresponding to $x \simeq 0$ and $x \simeq L_T$, respectively. This amounts to summing the contribution due to particles returning to the left end, such that $f(x)f(x') = f_0^2$, and that of particles crossing the tracer, such that $f(x)f(x') = -f_0^2$.

Let us return to the case of an active bath of density ρ_0 . We denote by m the polarization of particles around $x' = 0$ so that the local density of particles with orientation σ is $\rho_0[(1 + \sigma m)/2]$. The force-force correlation is then obtained from the single-particle result through $C_{\mathcal{F}}(t) = 2\rho_0\{[(1 + m)/2]c(1, t) + [(1 - m)/2]c(-1, t)\}$, where the factor 2 stems from the contributions of particles starting at $x' \simeq L_T$. Expanding the exponentials in Eq. (21) in the long-time limit, one finds the leading orders cancel,

yielding the $t^{-3/2}$ scaling of $C_{\mathcal{F}}(t)$. Using Eq. (11) leads to $m = \mu f_0/v$, which is consistent with the fact that active particles polarize against external potentials [95]. Straightforward algebra then gives

$$C_{\mathcal{F}}(t) \sim \frac{\rho_0(f_0 d L_T)^2}{4\pi^{1/2}(D_{\text{eff}} t)^{3/2}} G(\ell_p/L_T) \quad (22)$$

where $G(y) = 1 - (2\mu f_0/v)y$. Importantly, $C_{\mathcal{F}}(t)$ becomes negative when the size of the tracer is small, $L_T \leq 2\mu\ell_p f_0/v$. In the discussion above, we neglected $\mathcal{O}(f_0)$ corrections to the propagator and to the steady-state density due to the edges of the tracer. Including the f_0 corrections to all orders confirms the scaling [Eq. (22)], to order $\mathcal{O}(d^2)$, albeit with $G(y) = [1 - (2\mu f_0 y/v)^2]/[1 - (\mu f_0/v)^2]^2$ (see Sec. IV of the Supplemental Material [87]). This does not change the leading order estimate for the crossover length $\sim 2\mu f_0 \ell_p/v$. Negative autocorrelations have been reported in other contexts, in [7] and out [80] of equilibrium. Here, it is a direct consequence of the polarization against the potential. Setting $m = 0$ in the computation above always leads to $C_{\mathcal{F}}(t) > 0$.

We now turn to the long-time behavior of $\gamma(t)$. Inserting Eq. (11) in Eq. (10) leads to $\gamma = \gamma_p - \gamma_a$, with

$$\gamma_p(t-t') \equiv \beta_{\text{eff}} \left\langle \mathcal{F}(t) \frac{\mathcal{F}(t')}{1 - \frac{\mu}{v} \mathcal{F}(t')} \right\rangle_c, \quad (23)$$

$$\gamma_a(t-t') \equiv \beta_{\text{eff}} \left\langle \mathcal{F}(t) \frac{\sigma(t') \ell_p \partial_{x(t')} \mathcal{F}(t')}{1 - \sigma(t') \frac{\mu}{v} \mathcal{F}(t')} \right\rangle_c. \quad (24)$$

The heuristic argument developed above for $C_{\mathcal{F}}(t)$ directly extends to the correlators (23) and (24), showing that γ_a and γ_p both inherit the $t^{-3/2}$ scaling of $C_{\mathcal{F}}(t)$ at long times. Inspecting Eq. (23) shows that, to leading order in f_0 ,

$$\gamma_p(t) \sim \beta_{\text{eff}} C_{\mathcal{F}}(t) = \frac{\beta_{\text{eff}} \rho_0 (f_0 d L_T)^2}{4\pi^{1/2} (D_{\text{eff}} t)^{3/2}} G(\ell_p/L_T). \quad (25)$$

Equation (25) is nothing but an effective FDT for the passive tracer. Our results show that the FDT is only expected to hold for small f_0 and should be generically violated when γ_a is not negligible compared with γ_p .

The presence of $\sigma(t')$ in Eq. (24) makes the contributions of $\sigma' = \pm 1$ particle add up, instead of canceling, leading to $\gamma_a(t) > 0$ for all L_T and a long-time scaling $\gamma_a \sim \mathcal{O}(f_0^3) t^{-3/2}$. Therefore, to leading order in f_0 , $\gamma \sim \beta_{\text{eff}} C_{\mathcal{F}}(t)$. This suggests that $\gamma_T = \int_0^\infty dt \gamma(t)$ can also change sign and become *negative* for small tracers. Indeed, a perturbative calculation finds that

$$\gamma_T \sim \beta_{\text{eff}} v^{-1} \rho_0 (f_0 d)^2 \frac{L_T}{\ell_p} \left(1 - \frac{d^2 + 6\ell_p^2}{3dL_T} \right). \quad (26)$$

The derivations of this result and of the asymptotics of γ_a are not particularly illuminating; they are deferred to Sec. IV of the Supplemental Material [87]. Importantly, Eq. (26) implies that when a small symmetric tracer is dragged at velocity U , the active bath *enhances* its motion rather than resisting it.

Adiabatic limit.—Although Eq. (1) is a common framework to describe a tracer's dynamics, it relies on the assumption that the motion is slow. An important—but rarely debated—question is thus its range of validity. Here, this is set by the requirement that the tracer's response is much slower than the diffusive relaxation of the bath, i.e., $\langle X(t) \rangle, \langle X^2(t) \rangle_c^{1/2} \ll (D_{\text{eff}} t)^{1/2}$. For an asymmetric tracer, using $\langle X(t) \rangle \sim Ft/\gamma_0$ and Eq. (2), we find $t \ll \tau_1 \equiv D_{\text{eff}}(\gamma_0/F)^2$ and $t \ll \tau_2 \equiv (D_{\text{eff}}/K)^2$. Equation (19) implies $\tau_1 \ll \tau_2$ so that the adiabatic limit holds up to $t \ll \tau_1$. Beyond this timescale, which can be arbitrarily large, an asymmetric tracer in an active bath cannot be described by Eq. (1). Considering a finite system of size L , the diffusive relaxation time is $t = \tau_{\text{rel}} \sim L^2/D_{\text{eff}}$. Thus, the adiabatic limit for an asymmetric tracer in a finite system is valid for $FL \ll D_{\text{eff}}\gamma_0$, which can be achieved by designing the tracer shape to bound F or by using a small enough system. For a symmetric tracer, there is no temporal restriction, and the only requirement is $D \ll D_{\text{eff}}$, which can be fulfilled by setting $\gamma_0 \gg (I/D_{\text{eff}})^{1/2}$. For towing both asymmetric tracers and symmetric tracers at constant velocity U , the only requirement is $U \ll D_{\text{eff}}/L$.

Conclusion.—In this Letter, we have derived the long-time dynamics of a passive tracer in a dilute active bath under the sole assumption of an adiabatic evolution. We have revealed new regimes for both asymmetric and symmetric tracers. First, ratchet effects generically lead to the superdiffusion of asymmetric tracers, which also experience friction that grows with time when they are dragged at constant velocity U . For symmetric tracers, the long-time tail preserves the diffusive behavior, but negative active friction is observed for small tracers. The latter solely follows from the persistent motion of active particles and their polarization by external potentials, a mechanism that differs from previously studied cases with negative mobility [72,96,97]. We expect the tails for asymmetric and symmetric tracers to become $t^{-d/2}$ and $t^{-(d/2+1)}$ in d dimensions, respectively. This suggests, in two dimensions, that $\langle X(t)^2 \rangle_c \sim t \ln t$ for an asymmetric tracer, which remains to be verified. Our results stem from generic features of dry active particles and should thus hold generically. The exponents are expected to be universal, but the transport coefficients can be dressed, for instance, by interactions. Moreover, the mechanisms should lead to even richer behaviors for active suspensions in momentum-conserving fluids [37,45,50,59], or in the presence of phoresis [80].

We thank Yongjoo Baek, Bernard Derrida, and Xinpeng Xu for many useful discussions. O. G. and Y. K. are supported by Israel Science Foundation Grant No. 1331/17 and NSF-BSF Grant No. 2016624. J. T. is supported by the ANR grant THEMA. O. G. also acknowledges support from the Adams Fellowship Program of the Israeli Academy of Sciences and Humanities.

-
- [1] P. Hänggi and F. Marchesoni, *Chaos* **15**, 026101 (2005).
- [2] C. Gardiner, *Stochastic Methods: A Handbook for the Natural and Social Sciences*, Springer Series in Synergetics (Springer, Berlin Heidelberg, 2010).
- [3] P. Chaikin and T. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, Cambridge, England, 2000).
- [4] M. Kardar, *Statistical Physics of Fields* (Cambridge University Press, Cambridge, England, 2007).
- [5] B. J. Alder and T. E. Wainwright, *Phys. Rev. Lett.* **18**, 988 (1967).
- [6] Y. Pomeau and P. Résibois, *Phys. Rep.* **19**, 63 (1975).
- [7] H. van Beijeren, *Rev. Mod. Phys.* **54**, 195 (1982).
- [8] T. R. Kirkpatrick, D. Belitz, and J. V. Sengers, *J. Stat. Phys.* **109**, 373 (2002).
- [9] A. Dhar, *Adv. Phys.* **57**, 457 (2008).
- [10] H. Spohn, in *Lect. Notes Phys.* (Springer, Cham, 2016), Vol. 921, pp. 107–158.
- [11] R. Di Leonardo, L. Angelani, D. Dell’Arciprete, G. Ruocco, V. Iebba, S. Schippa, M. P. Conte, F. Mecarini, F. De Angelis, and E. Di Fabrizio, *Proc. Natl. Acad. Sci. U.S.A.* **107**, 9541 (2010).
- [12] A. Sokolov, M. M. Apodaca, B. A. Grzybowski, and I. S. Aranson, *Proc. Natl. Acad. Sci. U.S.A.* **107**, 969 (2010).
- [13] A. Kaiser, A. Peshkov, A. Sokolov, B. ten Hagen, H. Löwen, and I. S. Aranson, *Phys. Rev. Lett.* **112**, 158101 (2014).
- [14] C. Maggi, F. Saglimbeni, M. Dipalo, F. De Angelis, and R. Di Leonardo, *Nat. Commun.* **6**, 7855 (2015).
- [15] P. Galajda, J. Keymer, P. Chaikin, and R. Austin, *J. Bacteriol.* **189**, 8704 (2007).
- [16] M. B. Wan, C. J. Olson Reichhardt, Z. Nussinov, and C. Reichhardt, *Phys. Rev. Lett.* **101**, 018102 (2008).
- [17] J. Tailleur and M. E. Cates, *Europhys. Lett.* **86**, 60002 (2009).
- [18] J. Stenhammar, R. Wittkowski, D. Marenduzzo, and M. E. Cates, *Sci. Adv.* **2**, e1501850 (2016).
- [19] C. O. Reichhardt and C. Reichhardt, *Annu. Rev. Condens. Matter Phys.* **8**, 51 (2017).
- [20] T. Speck, *Soft Matter* **16**, 2652 (2020).
- [21] N. Nikola, A. P. Solon, Y. Kafri, M. Kardar, J. Tailleur, and R. Voituriez, *Phys. Rev. Lett.* **117**, 098001 (2016).
- [22] Y. Baek, A. P. Solon, X. Xu, N. Nikola, and Y. Kafri, *Phys. Rev. Lett.* **120**, 058002 (2018).
- [23] O. Granek, Y. Baek, Y. Kafri, and A. P. Solon, *J. Stat. Mech.* (2020) 063211.
- [24] T. Speck and A. Jayaram, *Phys. Rev. Lett.* **126**, 138002 (2021).
- [25] X.-L. Wu and A. Libchaber, *Phys. Rev. Lett.* **84**, 3017 (2000).
- [26] C. Maggi, M. Paoluzzi, N. Pellicciotta, A. Lepore, L. Angelani, and R. Di Leonardo, *Phys. Rev. Lett.* **113**, 238303 (2014).
- [27] A. Argun, A.-R. Moradi, E. Pinçe, G. B. Bağcı, A. Imparato, and G. Volpe, *Phys. Rev. E* **94**, 062150 (2016).
- [28] C. Maggi, M. Paoluzzi, L. Angelani, and R. Di Leonardo, *Sci. Rep.* **7**, 17588 (2017).
- [29] S. Chaki and R. Chakrabarti, *Physica (Amsterdam)* **511A**, 302 (2018).
- [30] S. Chaki and R. Chakrabarti, *Physica (Amsterdam)* **530A**, 121574 (2019).
- [31] L. Dabelow, S. Bo, and R. Eichhorn, *Phys. Rev. X* **9**, 021009 (2019).
- [32] K. Goswami, *Phys. Rev. E* **99**, 012112 (2019).
- [33] M. Knežević and H. Stark, *New J. Phys.* **22**, 113025 (2020).
- [34] S. Ye, P. Liu, F. Ye, K. Chen, and M. Yang, *Soft Matter* **16**, 4655 (2020).
- [35] S. Belan and M. Kardar, *J. Chem. Phys.* **154**, 024109 (2021).
- [36] G. Soni, B. Jaffar Ali, Y. Hatwalne, and G. Shivashankar, *Biophys. J.* **84**, 2634 (2003).
- [37] D. T. N. Chen, A. W. C. Lau, L. A. Hough, M. F. Islam, M. Goulian, T. C. Lubensky, and A. G. Yodh, *Phys. Rev. Lett.* **99**, 148302 (2007).
- [38] D. Loi, S. Mossa, and L. F. Cugliandolo, *Phys. Rev. E* **77**, 051111 (2008).
- [39] P. T. Underhill, J. P. Hernandez-Ortiz, and M. D. Graham, *Phys. Rev. Lett.* **100**, 248101 (2008).
- [40] A. W. C. Lau and T. C. Lubensky, *Phys. Rev. E* **80**, 011917 (2009).
- [41] K. C. Leptos, J. S. Guasto, J. P. Gollub, A. I. Pesci, and R. E. Goldstein, *Phys. Rev. Lett.* **103**, 198103 (2009).
- [42] J. Dunkel, V. B. Putz, I. M. Zaid, and J. M. Yeomans, *Soft Matter* **6**, 4268 (2010).
- [43] H. Kurtuldu, J. S. Guasto, K. A. Johnson, and J. P. Gollub, *Proc. Natl. Acad. Sci. U.S.A.* **108**, 10391 (2011).
- [44] G. Miño, T. E. Mallouk, T. Darnige, M. Hoyos, J. Dauchet, J. Dunstan, R. Soto, Y. Wang, A. Rousselet, and E. Clement, *Phys. Rev. Lett.* **106**, 048102 (2011).
- [45] I. M. Zaid, J. Dunkel, and J. M. Yeomans, *J. R. Soc. Interface* **8**, 1314 (2011).
- [46] G. Foffano, J. S. Lintuvuori, K. Stratford, M. E. Cates, and D. Marenduzzo, *Phys. Rev. Lett.* **109**, 028103 (2012).
- [47] G. L. Miño, J. Dunstan, A. Rousselet, E. Clément, and R. Soto, *J. Fluid Mech.* **729**, 423 (2013).
- [48] T. V. Kasyap, D. L. Koch, and M. Wu, *Phys. Fluids* **26**, 081901 (2014).
- [49] A. Morozov and D. Marenduzzo, *Soft Matter* **10**, 2748 (2014).
- [50] J.-L. Thiffeault, *Phys. Rev. E* **92**, 023023 (2015).
- [51] A. E. Pateson, A. Gopinath, P. K. Purohit, and P. E. Arratia, *Soft Matter* **12**, 2365 (2016).
- [52] A. Suma, L. F. Cugliandolo, and G. Gonnella, *J. Stat. Mech.* (2016) 054029.
- [53] E. W. Burkholder and J. F. Brady, *Phys. Rev. E* **95**, 052605 (2017).
- [54] M. J. Y. Jerez, M. N. P. Confesor, M. V. Carpio Bernido, and C. C. Bernido, in *AIP Conference Proceedings* (American Institute of Physics, College Park, 2017), Vol. 1871, p. 050004 10.1063/1.4996528.

- [55] T. Kurihara, M. Aridome, H. Ayade, I. Zaid, and D. Mizuno, *Phys. Rev. E* **95**, 030601(R) (2017).
- [56] P. Pietzonka and U. Seifert, *J. Phys. A* **51**, 01LT01 (2018).
- [57] E. W. Burkholder and J. F. Brady, *J. Chem. Phys.* **150**, 184901 (2019).
- [58] R. Chatterjee, N. Segall, C. Merrigan, K. Ramola, B. Chakraborty, and Y. Shokef, *J. Chem. Phys.* **150**, 144508 (2019).
- [59] K. Kanazawa, T. G. Sano, A. Cairoli, and A. Baule, *Nature (London)* **579**, 364 (2020).
- [60] J. Reichert and T. Voigtmann, arXiv:2010.13769.
- [61] L. Abbaspour and S. Klumpp, *Phys. Rev. E* **103**, 052601 (2021).
- [62] J. Katuri, W. E. Usual, M. N. Popescu, and S. Sánchez, *Sci. Adv.* **7**, eabd0719 (2021).
- [63] J. Reichert, L. F. Granz, and T. Voigtmann, *Eur. Phys. J. E* **44**, 27 (2021).
- [64] M. S. Green, *J. Chem. Phys.* **20**, 1281 (1952).
- [65] H. Mori, *Prog. Theor. Phys.* **33**, 423 (1965).
- [66] R. Kubo, *Rep. Prog. Phys.* **29**, 255 (1966).
- [67] N. Van Kampen, *Phys. Rep.* **124**, 69 (1985).
- [68] N. Van Kampen and I. Oppenheim, *Physica (Amsterdam)* **138A**, 231 (1986).
- [69] U. Seifert, *Rep. Prog. Phys.* **75**, 126001 (2012).
- [70] C. Maes and S. Steffenoni, *Phys. Rev. E* **91**, 022128 (2015).
- [71] M. Krüger and C. Maes, *J. Phys. Condens. Matter* **29**, 064004 (2017).
- [72] C. Maes, *Phys. Rev. Lett.* **125**, 208001 (2020).
- [73] R. Feynman and F. Vernon, *Ann. Phys. (N.Y.)* **24**, 118 (1963).
- [74] G. W. Ford, M. Kac, and P. Mazur, *J. Math. Phys. (N.Y.)* **6**, 504 (1965).
- [75] A. Caldeira and A. Leggett, *Physica (Amsterdam)* **121A**, 587 (1983).
- [76] S. Hanna, W. Hess, and R. Klein, *J. Phys. A* **14**, L493 (1981).
- [77] J. Boon and S. Yip, *Molecular Hydrodynamics* (Dover Publications, New York, 1991).
- [78] I. Goychuk and P. Hänggi, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 3552 (2002).
- [79] A. W. C. Lau, B. D. Hoffmann, A. Davies, J. C. Crocker, and T. C. Lubensky, *Phys. Rev. Lett.* **91**, 198101 (2003).
- [80] R. Golestanian, *Phys. Rev. Lett.* **102**, 188305 (2009).
- [81] O. Bénichou, A. Bodrova, D. Chakraborty, P. Illien, A. Law, C. Mejía-Monasterio, G. Oshanin, and R. Voituriez, *Phys. Rev. Lett.* **111**, 260601 (2013).
- [82] P. Illien, O. Bénichou, C. Mejía-Monasterio, G. Oshanin, and R. Voituriez, *Phys. Rev. Lett.* **111**, 038102 (2013).
- [83] L. D'Alessio, Y. Kafri, and A. Polkovnikov, *J. Stat. Mech.* (2016) 023105.
- [84] P. Weinberg, M. Bukov, L. D'Alessio, A. Polkovnikov, S. Vajna, and M. Kolodrubetz, *Phys. Rep.* **688**, 1 (2017).
- [85] S. Hanna, W. Hess, and R. Klein, *Physica (Amsterdam)* **111A**, 181 (1982).
- [86] O. Granek, Y. Kafri, and J. Tailleur (to be published).
- [87] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.129.038001>, which cites Ref. [88], for simulation and computation details.
- [88] L. Angelani, A. Costanzo, and R. Di Leonardo, *Europhys. Lett.* **96**, 68002 (2011).
- [89] M. E. Cates, *Rep. Prog. Phys.* **75**, 042601 (2012).
- [90] K. Kitahara, W. Horsthemke, and R. Lefever, *Phys. Lett.* **70A**, 377 (1979).
- [91] A. P. Solon, Y. Fily, A. Baskaran, M. E. Cates, Y. Kafri, M. Kardar, and J. Tailleur, *Nat. Phys.* **11**, 673 (2015).
- [92] L. Angelani and R. D. Leonardo, *New J. Phys.* **12**, 113017 (2010).
- [93] S. A. Mallory, C. Valeriani, and A. Cacciuto, *Phys. Rev. E* **90**, 032309 (2014).
- [94] The diverging behavior of γ_T sets an upper bound on the time-scale for which this approximation holds, as discussed at the end of the Letter.
- [95] M. Enculescu and H. Stark, *Phys. Rev. Lett.* **107**, 058301 (2011).
- [96] See, e.g., J. Cividini, D. Mukamel, and H. A. Posch, *J. Phys. A* **51**, 085001 (2018) and references therein.
- [97] P. K. Ghosh, P. Hänggi, F. Marchesoni, and F. Nori, *Phys. Rev. E* **89**, 062115 (2014).