

Minimal Massive Supergravity

Nihat Sadik Deger^{1,2,3,*}, Marc Geiller^{4,†}, Jan Rosseel^{5,6,3,‡} and Henning Samtleben^{4,7,§}

¹*Department of Mathematics, Bogazici University, Bebek, 34342 Istanbul, Turkey*

²*Feza Gursey Center for Physics and Mathematics, Bogazici University, Kandilli, 34684 Istanbul, Turkey*

³*Erwin Schrödinger International Institute for Mathematics and Physics, University of Vienna, Boltzmannngasse 9, 1090, Vienna, Austria*

⁴*ENSL, CNRS, Laboratoire de physique, F-69342 Lyon, France*

⁵*Faculty of Physics, University of Vienna, Boltzmannngasse 5, 1090 Vienna, Austria*

⁶*Division of Theoretical Physics, Rudjer Bošković Institute, Bijenička 54, 10000 Zagreb, Croatia*

⁷*Institut Universitaire de France (IUF), 75005 Paris, France*



(Received 7 June 2022; revised 2 August 2022; accepted 22 September 2022; published 18 October 2022)

Minimal massive gravity in three dimensions propagates a single massive spin-2 mode around an anti-de Sitter vacuum. It is distinguished by allowing for vacua with positive central charges of the asymptotic conformal algebra and a bulk graviton of positive energy. We present a new action for the model (and its higher order extensions) in terms of a dreibein and an independent spin connection. From this, we construct its supersymmetric extension. Surprisingly, all vacua complying with bulk and boundary unitarity appear to break supersymmetry spontaneously. In contrast, all supersymmetric vacua have a negative central charge whenever the bulk graviton has positive energy.

DOI: [10.1103/PhysRevLett.129.171601](https://doi.org/10.1103/PhysRevLett.129.171601)

Three-dimensional (3D) gravity has long been established as a distinguished testing ground in order to develop, examine, and challenge fundamental issues of quantum gravity, black hole physics, and holography, building on the seminal works [1–5]. In what is now understood as a precursor of the anti-de Sitter/conformal field theory (AdS/CFT) correspondence, Brown and Henneaux discovered that Einstein gravity on a 3D AdS space has an infinite-dimensional asymptotic symmetry algebra forming two copies of the Virasoro algebra with nonvanishing central charges [3]. This observation has been at the origin of fundamental precision computations [6,7] of the entropy of the Bañados-Teitelboim-Zanelli black hole [8], obtained as a discrete quotient of 3D AdS.

Over the years, various 3D gravitational models have been constructed featuring massive spin-2 excitations around Minkowski and AdS spaces, including in particular topologically massive gravity (TMG) [9], new massive gravity [10], minimal massive gravity (MMG) [11], and higher order extensions thereof, such as the exotic massive gravity [12].

In this Letter, we will focus on the MMG model (and its higher order extensions). This model propagates a single massive spin-2 mode around an AdS background, and is

distinguished by the fact that its parameter space allows for a region in which the massive spin-2 mode is neither ghost nor tachyonic, while maintaining both Brown-Henneaux central charges positive. In this sense, MMG evades the bulk-boundary unitarity clash from which most other 3D massive gravity models suffer, elevating the model to a viable holographic dual of a putative unitary 2D CFT.

To date, none of the 3D bosonic models complying with bulk and boundary unitarity [11,13] have been embedded into a supersymmetric theory, despite the fact that supersymmetry typically provides an invaluable set of tools to establish and corroborate consistent holographic scenarios. The construction of a supersymmetric extension of MMG is one of the main results of this Letter and gives rise to some surprising observations regarding its vacuum structure, notably an apparent clash between unbroken supersymmetry and unitarity for its AdS vacua.

MMG is defined as a deformation of TMG by terms quadratic in the Riemann tensor according to

$$\frac{1}{\mu} C_{\mu\nu} + \bar{\sigma} G_{\mu\nu} + \bar{\Lambda}_0 g_{\mu\nu} = \frac{\gamma}{2\mu^2} \epsilon_{\mu\kappa\lambda} \epsilon_{\nu\sigma\tau} S^{\kappa\sigma} S^{\lambda\tau}. \quad (1)$$

Here, $G_{\mu\nu}$ is the Einstein tensor associated with the metric $g_{\mu\nu}$, $S_{\mu\nu}$ is the associated Schouten tensor, and $C_{\mu\nu}$ is its Cotton tensor

$$\begin{aligned} G_{\mu\nu} &= R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R, & S_{\mu\nu} &= R_{\mu\nu} - \frac{1}{4} g_{\mu\nu} R, \\ C_{\mu\nu} &= \epsilon_{\mu\rho\sigma} \nabla^\rho S^\sigma{}_\nu. \end{aligned} \quad (2)$$

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

In the limit $\gamma \rightarrow 0$, Eqs. (1) reduce to those of TMG. The MMG equations (1) cannot be derived from a standard action principle of the metric alone. In particular, the fact that the right-hand side has zero divergence (as required for consistency) is not automatic, but follows on-shell from iterating Eq. (1) itself—a mechanism dubbed “third way consistency” in [11]. On the other hand, in [11] Eqs. (1) could be derived by variation of a first order Lagrangian with auxiliary fields, in the region of parameter space where [14]

$$\mu^2(1 + \gamma\bar{\sigma})^2 > \gamma^3\bar{\Lambda}_0. \quad (3)$$

The absence of a standard action functional has among other things hampered the construction of the supersymmetric extension of (1). As a main result of this Letter, we will employ a new action for MMG (and higher order extensions thereof), in order to construct the supersymmetric extension of (1). As it turns out, the extension is not unique, and for a given set of coupling constants $\{\gamma, \mu, \bar{\sigma}, \bar{\Lambda}_0\}$ we find (up to four) different supersymmetric extensions of the model. The underlying supersymmetric structures provide additional tools for the vacuum analysis of the model.

As a starting point, we consider the following class of actions, depending on a dreibein e_μ^a and an independent (torsionful) spin connection ϖ_μ^a as

$$\begin{aligned} \mathcal{L}[e, \varpi] = & \mathcal{L}_0[e] + \tau \epsilon^{\mu\nu\rho} e_\mu^a D[\varpi]_\nu e_{\rho a} \\ & + \kappa \epsilon^{\mu\nu\rho} \left(\varpi_\mu^a \partial_\nu \varpi_{\rho a} + \frac{1}{3} \epsilon_{abc} \varpi_\mu^a \varpi_\nu^b \varpi_\rho^c \right), \end{aligned} \quad (4)$$

with local Lorentz indices a, b, \dots (for our conventions see Ref. [15]). To begin with, $\mathcal{L}_0[e]$ is an arbitrary gravitational Lagrangian, depending only on the dreibein e_μ^a . The constants τ and κ denote the coupling constants of the torsion and the Chern-Simons term for ϖ , respectively. Variation of the Lagrangian (4) with respect to the connection ϖ yields an equation for its curvature

$$R[\varpi]_{\mu\nu, a} = -\frac{\tau}{\kappa} \epsilon_{abc} e_\mu^b e_\nu^c. \quad (5)$$

Variation of (4) with respect to the dreibein e_μ^a on the other hand determines the torsion $T[\varpi]_{\mu\nu, a}$ of the connection ϖ by

$$0 = 2\mathbb{G}_a^\mu + \tau \epsilon^{\mu\nu\rho} T[\varpi]_{\nu\rho, a}, \quad (6)$$

where \mathbb{G}_a^μ is defined as

$$\delta\mathcal{L}_0 = \sqrt{-g} \mathbb{G}_a^\mu \delta e_\mu^a. \quad (7)$$

Diffeomorphism and Lorentz symmetry imply that the tensor $\mathbb{G}_{\mu\nu} \equiv \mathbb{G}_\mu^a e_{a\nu}$ is symmetric and divergence free

$$\mathbb{G}_{\mu\nu} = \mathbb{G}_{\nu\mu}, \quad \nabla^\mu \mathbb{G}_{\mu\nu} = 0. \quad (8)$$

The contorsion of the connection ϖ is defined as

$$\begin{aligned} K[\varpi]_\mu^a &= \varpi_\mu^a - \omega_\mu^a \\ &= \frac{1}{2} \epsilon^{\rho\sigma\tau} \left(e_\mu^b e_\rho^a - \frac{1}{2} e_\mu^a e_\rho^b \right) T[\varpi]_{\sigma\tau, b}, \end{aligned} \quad (9)$$

in terms of the torsionless Levi-Civita connection ω_μ^a . The field equations (6) can then be rewritten as

$$K[\varpi]_\mu^a = -\frac{1}{\tau} \left(\mathbb{G}_\mu^a - \frac{1}{2} e_\mu^a \mathbb{G} \right) \equiv -\frac{1}{\tau} \mathbb{S}_\mu^a, \quad (10)$$

where $\mathbb{G} \equiv \mathbb{G}_\mu^a e_\mu^a$. Finally, using the general relation between curvature and contorsion

$$\begin{aligned} R[\varpi]_{\mu\nu}^a &= R[\omega]_{\mu\nu}^a + 2D[\omega]_{[\mu} K[\varpi]_{\nu]}^a \\ &+ \epsilon^{abc} K[\varpi]_{\mu b} K[\varpi]_{\nu c}, \end{aligned} \quad (11)$$

and combining this with (5) and (10), yields the equation

$$\epsilon_{\mu\sigma\tau} \nabla^\sigma \mathbb{S}^\tau{}_\nu - \tau \mathbb{G}_{\mu\nu} + \frac{\tau^2}{\kappa} g_{\mu\nu} = \frac{1}{2\tau} \epsilon_{\mu\sigma\tau} \epsilon_{\nu\kappa\lambda} \mathbb{S}^{\sigma\kappa} \mathbb{S}^{\tau\lambda}, \quad (12)$$

exclusively formulated in terms of the dreibein, with $\mathbb{S}_{\mu\nu}$ defined via (10), (7). Just as in (1), these equations are “third-way consistent” in that the vanishing of the divergence of the right-hand side follows from iterating Eq. (12) itself [together with the relations (8)]. A key feature of this construction is that the final equations (12) do not arise directly among the Euler-Lagrange equations, but only after combination with the integrability conditions (11) [16].

Let us also point out that consistent matter couplings to (12) are straightforwardly implemented in the Lagrangian (4) by

$$\begin{aligned} \mathcal{L}_0[e] &\rightarrow \mathcal{L}_0[e] + \mathcal{L}_{\text{matter}}[e, \dots], \\ &\Rightarrow \mathbb{G}_{\mu\nu} \rightarrow \mathbb{G}_{\mu\nu} + T_{\mu\nu}, \end{aligned} \quad (13)$$

with the standard (covariantly conserved) energy-momentum tensor $T_{\mu\nu}$, reproducing the (on-shell) results of [12,17].

In the following, we specialize to MMG, by setting

$$\mathcal{L}_0[e] = \frac{1}{G_3} \epsilon^{\mu\nu\rho} e_\mu^a R_{\nu\rho, a} + \lambda \epsilon^{\mu\nu\rho} \epsilon_{abc} e_\mu^a e_\nu^b e_\rho^c, \quad (14)$$

where for simplicity we will set the gravitational constant $G_3 = 1$. The resulting Eqs. (12) then reproduce (1), with the (bijective up to rescaling) translation of parameters according to

$$\frac{\mu}{\gamma} = \tau, \quad \mu\bar{\Lambda}_0 = \frac{9\lambda^2}{4\tau} + \frac{\tau^2}{\kappa}, \quad \mu\bar{\sigma} = -\tau - \frac{3\lambda}{2\tau}. \quad (15)$$

Equivalently, (14) can be replaced by the first order Palatini Lagrangian in terms of an independent connection ω ,

$$\mathcal{L}_0[e, \omega] = \varepsilon^{\mu\nu\rho} (e_\mu^a R[\omega]_{\nu\rho,a} + \lambda \varepsilon_{abc} e_\mu^a e_\nu^b e_\rho^c), \quad (16)$$

such that the final Lagrangian (4) is given by the sum of the so-called ‘‘standard’’ and the ‘‘exotic’’ action of 2 + 1 gravity [5], however with both actions carrying different spin connections ω , and ϖ , respectively [18]. In this first order formulation, and after redefinition

$$\omega_\mu^a = \Omega_\mu^a + \alpha h_\mu^a, \quad \varpi_\mu^a = \Omega_\mu^a + \frac{\sqrt{-\kappa\tau}}{\kappa} e_\mu^a, \quad (17)$$

the Lagrangian (4), written as $\mathcal{L}[e, \Omega, h]$, reproduces the first-order Lagrangian of [11]. The condition $\kappa\tau < 0$ precisely defines the region in parameter space (3) in which the Lagrangian of [11] exists.

We will now use the Lagrangian (4), (14) in its second order form as a starting point for the construction of supersymmetric extensions of the model. Separate supersymmetrization of the two parts of (4) is known in terms of super-Chern-Simons theories, with in particular (14) admitting a general $\mathcal{N} = (p, q)$ supersymmetric extension [4]. Again, the nontrivial structure here arises since both parts of (4) share the same dreibein e_μ^a while carrying independent spin connections ω , and ϖ . For simplicity, we will only attempt to impose minimal [$\mathcal{N} = (1, 0)$] supersymmetry.

Our ansatz for the fermionic sector of the model carries two gravitino fields, ψ_μ and χ_μ , sharing only one local supersymmetry, with spinor parameter ε . Reminiscent of the first order formulation of TMG [19,20], this allows the first order fermionic Lagrangian [expected for the supersymmetrization of (4), (14)] to consistently accommodate a massive spin-3/2 mode as a superpartner to the massive spin-2 mode. Indeed, this turns out to be the correct structure. Leaving the technical details for [21], let us only spell out the result. To quadratic order in the fermions, their Lagrangian is given by

$$\begin{aligned} \mathcal{L}_{\text{ferm}} = & \frac{1}{\zeta^2} \varepsilon^{\mu\nu\rho} [(1 + \zeta)\bar{\psi}_\mu + \bar{\chi}_\mu] D[\omega]_\nu [(1 - \zeta)\psi_\rho + \chi_\rho] \\ & + \frac{1}{2\zeta^2} \varepsilon^{\mu\nu\rho} K[\varpi]_\nu^a (\bar{\psi}_\mu + \bar{\chi}_\mu) \gamma_a (\psi_\rho + \chi_\rho) \\ & - \tau \varepsilon^{\mu\nu\rho} \bar{\psi}_\mu \gamma_\nu \chi_\rho + \frac{1}{2} \tau \varepsilon^{\mu\nu\rho} \bar{\chi}_\mu \gamma_\nu \chi_\rho \\ & + \frac{1}{4} \left((\zeta^2 - 2)\tau + \frac{1}{\zeta^2 \kappa} \right) \varepsilon^{\mu\nu\rho} \bar{\psi}_\mu \gamma_\nu \psi_\rho, \end{aligned} \quad (18)$$

with the couplings determined as functions of a new parameter ζ . The full Lagrangian can be shown to be

invariant under the supersymmetry transformations (also given to lowest order in the fermions)

$$\begin{aligned} \delta e_\mu^a &= \frac{1}{2} \bar{\psi}_\mu \gamma^a \varepsilon, \\ \delta \varpi_\mu^a &= -\frac{1}{2\zeta^2 \kappa} (\bar{\psi}_\mu + \bar{\chi}_\mu) \gamma^a \varepsilon - \frac{1}{2} D[\varpi]_\mu (\bar{\chi}_\nu \varepsilon e^{\nu a}), \\ \delta \psi_\mu &= D[\omega]_\mu \varepsilon - \frac{1}{4} \left(\zeta^2 \tau + \frac{1}{\zeta^2 \kappa} \right) \gamma_\mu \varepsilon, \\ \delta \chi_\mu &= \frac{1}{2} K[\varpi]_\mu^a \gamma_a \varepsilon - \frac{1}{4} \left(\zeta^2 \tau - \frac{1}{\zeta^2 \kappa} \right) \gamma_\mu \varepsilon, \end{aligned} \quad (19)$$

up to quartic terms in the fermions, which can be removed by higher order fermion contributions to the Lagrangian and transformation rules [22]. Details will appear elsewhere [21].

The final result is thus given by the sum of the bosonic Lagrangian (4), (14) and the fermionic Lagrangian (18). Supersymmetry requires the following relation

$$\lambda = \frac{1 + \zeta^2 \kappa \tau [2\zeta^2 + (\zeta^2 - 4)\zeta^4 \kappa \tau + 4]}{12\zeta^4 \kappa^2}, \quad (20)$$

between the parameter ζ parametrizing the fermionic couplings, and the coupling constants $\{\lambda, \kappa, \tau\}$ of the bosonic model (4), (14). Supersymmetrizability of the MMG model thus translates into the existence of real roots (for ζ) of (20). A necessary and sufficient set of conditions for the existence of such real roots is

$$\begin{aligned} \text{(i)} \quad & \frac{3\lambda}{\tau^2} - \frac{1}{\kappa\tau} + 1 \geq 0, \\ \text{(ii)} \quad & \text{either } \kappa\tau \geq -1 \quad \text{or} \quad \frac{3\lambda}{\tau^2} \geq -\frac{2}{\sqrt{-\kappa\tau}}. \end{aligned} \quad (21)$$

There are in general up to eight real roots (pairwise related by the flip $\zeta \rightarrow -\zeta$). While this may appear to place strong constraints on the model, remarkably our analysis below reveals that every MMG model (1) admitting an AdS vacuum also admits a supersymmetric extension.

The bosonic MMG equations (1) are obtained from the second order Lagrangian (4), (14), after elimination of the connection ϖ by its field equations. We can now carry out the analog construction in the fermionic sector. The fermionic field equations, obtained from variation of (18)

$$D[\omega]_{[\mu} \psi_{\nu]} = \frac{1}{4} \left(\zeta^2 \tau + \frac{1}{\zeta^2 \kappa} \right) \gamma_{[\mu} \psi_{\nu]} - \tau \gamma_{[\mu} \chi_{\nu]}, \quad (22)$$

can be solved algebraically for χ_μ . Plugging this solution back into the remaining fermionic field equations, we find after some computation (still to lowest order in fermions)

$$\begin{aligned} \tau C^\rho &= \frac{1}{8} \nu(2) \nu(-2) R^\rho - \frac{1}{4} \nu(0) \epsilon^{\mu\nu\rho} \gamma_\sigma \psi_\nu S_\mu^\sigma \\ &\quad - \frac{1}{32} \nu(0) \nu(2) \nu(-2) \epsilon^{\mu\nu\rho} \gamma_\mu \psi_\nu - \frac{1}{2} R^\mu G_\mu^\rho \\ &\quad - \frac{1}{2} \epsilon^{\rho\mu\nu} \gamma^\sigma R_\mu G_{\nu\sigma} - \frac{1}{2} \epsilon^{\rho\mu\nu} \gamma_\mu R^\sigma G_{\nu\sigma}, \end{aligned} \quad (23)$$

where we have introduced the gravitino curvature

$$R^\mu = \epsilon^{\mu\nu\rho} D[\omega]_\nu \psi_\rho, \quad (24)$$

and the ‘‘Cottino vector spinor’’ [23]

$$C^\mu = \gamma^\rho \gamma^{\mu\nu} D[\omega]_\nu R_\rho - \epsilon^{\mu\nu\rho} S_{\rho\sigma} \gamma^\sigma \psi_\nu, \quad (25)$$

and moreover defined the functions

$$\nu(n) \equiv (\zeta + n)^2 \tau + \frac{1}{\zeta^2 \kappa}. \quad (26)$$

We thus obtain the second-order fermionic field equation, entirely expressed in terms of the dreibein e_μ^a and the gravitino ψ_μ . Equation (23) constitutes the ‘‘superpartner’’ to the bosonic MMG equations (1). In the process, the latter receive additional source terms bilinear in the fermions. In order not to spoil consistency of the field equations (1), these source terms have to satisfy certain consistency conditions which in turn are implied by (23). Details will appear in [21].

Let us also note that in the TMG limit, where the right-hand side of (1) vanishes, Eq. (23) reduces to

$$C^\rho = -2\mu\bar{\sigma}R^\rho - \frac{\mu\bar{\sigma}}{\sqrt{-\bar{\Lambda}_0}} \gamma^{\rho\nu} \psi_\nu. \quad (27)$$

This is precisely the super-TMG equation from [23] (where $\bar{\sigma} = 1$ was assumed).

Let us now explore the landscape of (A)dS vacua of the super-MMG model and in particular localize the bulk-boundary unitary AdS vacua discovered in [11]. The bosonic MMG equations (1) admit maximally symmetric vacua $G_{\mu\nu} = -\Lambda g_{\mu\nu}$, provided that

$$\mu^2 \bar{\sigma}^2 \geq \gamma \bar{\Lambda}_0. \quad (28)$$

With the translation of parameters (15), this turns out to precisely coincide with the first condition in (21). The cosmological constants are given by

$$\Lambda_\pm = -\tau^2 \left((1 \pm \Gamma)^2 + \frac{1}{\kappa\tau} \right), \quad \Gamma \equiv \sqrt{\frac{3\lambda}{\tau^2} - \frac{1}{\kappa\tau}} + 1. \quad (29)$$

As a consequence, every supersymmetrizable MMG model admits maximally symmetric vacua. Evaluating the

super-MMG field equations (1) with (15) and (20), we obtain the values of their cosmological constants as

$$\begin{aligned} \Lambda_{\text{susy}} &= -\frac{(1 + \zeta^4 \kappa\tau)^2}{4\zeta^4 \kappa^2} \equiv -\frac{1}{\ell^2}, \\ \Lambda_{\text{ns}} &= -\frac{1 + \zeta^2 \kappa\tau [8 + 2\zeta^2 + \zeta^2 (\zeta^2 - 4)^2 \kappa\tau]}{4\zeta^4 \kappa^2}. \end{aligned} \quad (30)$$

The first vacuum in (30) is AdS (or Minkowski) and preserves part of the supersymmetry with the Killing spinor defined by

$$D[\omega]_\mu \epsilon - \frac{1}{2\ell} \gamma_\mu \epsilon = 0. \quad (31)$$

From (19), it then follows that $\delta\psi_\mu = 0$ is satisfied as usual for AdS, whereas $\delta\chi_\mu = 0$ holds identically, as a consequence of (10). On the other hand, for the nonsupersymmetric vacuum Λ_{ns} in (30), the Killing spinor equations for ψ_μ and χ_μ (19) cannot be solved simultaneously.

Linearizing and diagonalizing the bosonic Lagrangian around the supersymmetric AdS vacuum Λ_{susy} (with \tilde{e}_μ^a as background dreibein) yields

$$\begin{aligned} \mathcal{L}_{\text{lin}} &= \epsilon^{\mu\nu\rho} \alpha_+ f_\mu^{(+a)} (\partial_\nu f_{\rho,a}^{(+)} - \ell^{-1} \epsilon_{abc} f_\nu^{(+b)} \tilde{e}_\rho^c) \\ &\quad + \epsilon^{\mu\nu\rho} \alpha_- f_\mu^{(-a)} (\partial_\nu f_{\rho,a}^{(-)} + \ell^{-1} \epsilon_{abc} f_\nu^{(-b)} \tilde{e}_\rho^c) \\ &\quad - \epsilon^{\mu\nu\rho} \alpha_0 p_\mu^a (\partial_\nu p_{\rho,a} + M \epsilon_{abc} p_\nu^b \tilde{e}_\rho^c), \end{aligned} \quad (32)$$

exhibiting two massless and one massive spin-2 mode, $f^{(\pm)}$ and p , respectively, with coefficients factorizing in terms of ζ in an intriguing pattern as

$$\begin{aligned} \alpha_+ &= \kappa(1 - \zeta^2)(1 + \zeta^4 \kappa\tau), \\ \alpha_- &= \kappa(1 + \zeta^2 \kappa\tau)(1 + \zeta^4 \kappa\tau), \\ \alpha_0 &= \kappa(1 - \zeta^2)(1 + \zeta^2 \kappa\tau), \end{aligned} \quad (33)$$

and with the mass given by

$$M\ell = -\frac{(1 + \zeta^2 \kappa\tau) + \zeta^2 (1 - \zeta^2) \kappa\tau}{1 + \zeta^4 \kappa\tau}. \quad (34)$$

The no-tachyon condition $M^2 \ell^2 > 1$ for the massive spin-2 mode translates into

$$\kappa\tau(1 - \zeta^2)(1 + \zeta^2 \kappa\tau) > 0. \quad (35)$$

An analysis following [11,24] quickly shows that imposing no-ghost and no-tachyon unitarity conditions on the massive spin-2 mode necessarily implies negative central charges. Remarkably, all supersymmetric AdS vacua thus exhibit the clash between bulk and boundary unitarity.




FIG. 1. Different regions in parameter space. As long as the MMG model admits an AdS vacuum, i.e., outside of the gray and yellow areas, it admits at least one supersymmetric extension. The AdS vacua evading the bulk-boundary unitarity clash [11] are the nonsupersymmetric AdS vacua in region V.

We can finally map out the full landscape of (A)dS vacua in order to reconcile these results with the earlier findings of [11]. To this end, we combine the conditions on supersymmetrizability (21) with the values of the cosmological constants (29), (30) in order to identify the various regions in parameter space as depicted in Fig. 1.

Region 0.—In this region

$$\frac{1}{\kappa\tau} > 1 + \frac{3\lambda}{\tau^2}, \quad (36)$$

the first condition in (21) is violated. According to the above discussion, no (A)dS vacua exist in this region and the model is not supersymmetrizable.

Region I.—This region is bounded by region 0 and the parabola

$$\frac{1}{\kappa\tau} = -\frac{9}{4} \left(\frac{\lambda}{\tau^2} \right)^2, \quad (37)$$

with $(\lambda/\tau^2) < -\frac{2}{3}$. The second condition in (21) is violated, thus the model is not supersymmetrizable. With (29), it follows that both vacua Λ_{\pm} are of dS type.

Region II.—This region is bounded from above by the parabola (37). Λ_+ is an AdS vacuum, Λ_- is dS. There are two different solutions to (20), i.e., two supersymmetric extensions of the bosonic model, in both of which Λ_+ is

supersymmetric, but exhibits the bulk-boundary unitarity clash.

Region III.—This region is not covered by the Lagrangian of [11], since $\kappa\tau > 0$. Both vacua Λ_{\pm} are of AdS type. Equation (20) admits two different solutions

$$\zeta_{\pm}^2 = (1 \pm \Gamma) + \sqrt{(1 \pm \Gamma)^2 + \frac{1}{\kappa\tau}}. \quad (38)$$

There are thus two supersymmetric extensions of the bosonic model, satisfying

$$\Lambda_{\text{susy}}(\zeta_{\pm}) = \Lambda_{\pm} = \Lambda_{\text{ns}}(\zeta_{\mp}). \quad (39)$$

That is, each of the vacua (29) of the bosonic model is supersymmetric in one of the supersymmetric extensions and nonsupersymmetric in the other. Both vacua exhibit the bulk-boundary unitarity clash.

Region IV.—Both vacua Λ_{\pm} are AdS. There are four supersymmetric extensions of the model with a structure similar to (39), i.e., each of the AdS vacua of the bosonic model is supersymmetric in some supersymmetric extension(s) of the model. Again, both vacua exhibit the bulk-boundary unitarity clash.

Region V.—Both vacua Λ_{\pm} are AdS. There are two solutions of Eqs. (20)

$$\zeta_{\pm}^2 = (1 + \Gamma) \pm \sqrt{(1 + \Gamma)^2 + \frac{1}{\kappa\tau}}, \quad (40)$$

i.e., two supersymmetric extensions of the model. In both of these, the vacuum Λ_+ is supersymmetric, whereas Λ_- is not. A careful translation of parameters shows that this region is hosting all the vacua identified in [11] as evading the bulk-boundary clash. Specifically, these are the non-supersymmetric Λ_- vacua, as is also consistent with the above analysis of supersymmetric vacua.

Along the border lines separating the different regions in Fig. 1, there is always one Minkowski vacuum together with an (A)dS vacuum. The red line is the so-called merger line [17] in which both (A)dS vacua of the model coincide. Note that the two regions $\kappa\tau < 0$ and $\kappa\tau > 0$ of the parameter space are not connected, as the model (4) degenerates for $\kappa\tau = 0$ (or $(1/\kappa\tau) = 0$).

As anticipated above, the analysis shows that MMG admits supersymmetric extensions in all regions (II–V) that admit AdS vacua. In particular, the bulk-boundary unitary AdS vacua discovered in [11] are all situated in region V and can be embedded into supersymmetric models. Moreover, the analysis shows that in all these vacua supersymmetry is spontaneously broken.

The supersymmetric extensions of the MMG model thus offer new perspectives on the AdS vacuum analysis of the bosonic model and it will be most interesting to explore its repercussions in the context of other solutions of the model,

such as [17]. The product pattern of the coefficients (33) in terms of the supersymmetry parameter ζ indicates the location of the chiral points at which one of the central charges vanishes. The underlying supersymmetric structure will also be of great value for the unitarity analysis at these special points. Further interesting research directions include the construction of possible supersymmetric matter couplings to MMG, as well as the supersymmetrization of higher order extensions of the model [12,25]—given that our action (4) naturally accommodates all such generalizations. A superspace formulation of our construction would be highly desirable to address these issues.

Interestingly, we have identified different minimal supersymmetric extensions of the same bosonic MMG in which different vacua of the bosonic model appear supersymmetric. This may be read as a hint of an underlying structure of extended supersymmetry into which these models could be embedded, as is typical for such twin supergravities [26]. As a technical challenge this would require embedding the single massive bosonic degree of freedom into some extended multiplet structure.

Perhaps the most surprising result of our analysis is the observed clash between supersymmetry and bulk-boundary unitarity. It is precisely the AdS vacua in which supersymmetry is spontaneously broken which reconcile positive central charges with a positive energy bulk graviton. It would be very interesting to understand if this observed clash of unbroken supersymmetry and unitarity goes back to some more fundamental principle and has deeper implications for holography. The minimal massive supergravity constructed in this Letter provides a natural starting point for studying aspects of holography around non-supersymmetric vacua. The simultaneous (and unavoidable) presence of a supersymmetric AdS vacuum with bulk-boundary unitarity clash and a second AdS vacuum avoiding the clash but breaking supersymmetry allows us to probe such issues in a single model.

A more detailed version of the presented results will appear elsewhere [21].

We would like to thank Olaf Hohm and Mehmet Ozkan for useful discussions and comments. J. R. and N. S. D. are grateful to the Erwin Schrödinger Institute (ESI), Vienna where part of this work was done in the framework of the “Research in Teams” Programme. N. S. D. also wishes to thank ENS de Lyon for hospitality during the course of this work.

*sodik.deger@boun.edu.tr

†marc.geiller@ens-lyon.fr

‡jan.rosseel@univie.ac.at

§henning.samtleben@ens-lyon.fr

[1] S. Deser, R. Jackiw, and G. 't Hooft, Three-dimensional Einstein gravity: dynamics of flat space, *Ann. Phys. (N.Y.)* **152**, 220 (1984).

- [2] S. Deser and R. Jackiw, Three-dimensional cosmological gravity: dynamics of constant curvature, *Ann. Phys. (N.Y.)* **153**, 405 (1984).
- [3] J. D. Brown and M. Henneaux, Central charges in the canonical realization of asymptotic symmetries: An example from three-dimensional gravity, *Commun. Math. Phys.* **104**, 207 (1986).
- [4] A. Achucarro and P. K. Townsend, A Chern-Simons action for three-dimensional anti-de Sitter supergravity theories, *Phys. Lett. B* **180**, 89 (1986).
- [5] E. Witten, (2+1)-dimensional gravity as an exactly soluble system, *Nucl. Phys.* **B311**, 46 (1988).
- [6] A. Strominger, Black hole entropy from near horizon microstates, *J. High Energy Phys.* **02** (1998) 009.
- [7] D. Birmingham, I. Sachs, and S. Sen, Entropy of three-dimensional black holes in string theory, *Phys. Lett. B* **424**, 275 (1998).
- [8] M. Banados, C. Teitelboim, and J. Zanelli, The Black Hole in Three-Dimensional Space-Time, *Phys. Rev. Lett.* **69**, 1849 (1992).
- [9] S. Deser, R. Jackiw, and S. Templeton, Three-Dimensional Massive Gauge Theories, *Phys. Rev. Lett.* **48**, 975 (1982).
- [10] E. A. Bergshoeff, O. Hohm, and P. K. Townsend, Massive Gravity in Three Dimensions, *Phys. Rev. Lett.* **102**, 201301 (2009).
- [11] E. Bergshoeff, O. Hohm, W. Merbis, A. J. Routh, and P. K. Townsend, Minimal massive 3D gravity, *Classical Quantum Gravity* **31**, 145008 (2014).
- [12] M. Özkan, Y. Pang, and P. K. Townsend, Exotic massive 3D gravity, *J. High Energy Phys.* **08** (2018) 035.
- [13] E. A. Bergshoeff, S. de Haan, O. Hohm, W. Merbis, and P. K. Townsend, Zwei-Dreibein Gravity: A Two-Frame-Field Model of 3D Massive Gravity, *Phys. Rev. Lett.* **111**, 111102 (2013); **111**, 259902(E) (2013).
- [14] This inequality immediately follows from expressing the parameters of the MMG equations (1) in terms of the parameters $\{\sigma, \alpha, \mu\}$ of the first-order Lagrangian of [11]: $\mu^2(1 + \gamma\bar{\sigma})^2 - \gamma^3\Lambda_0 = \mu^2(1 + \sigma\alpha)^{-2} > 0$ (recall that $\sigma^2 = 1$ in [11]).
- [15] The conventions used in this Letter are as follows. We use the $(- + +)$ signature convention. The Levi-Civita tensor density and tensor are denoted by $\varepsilon^{\mu\nu\rho}$ and $\epsilon^{\mu\nu\rho}$, respectively. All spinors are Majorana. The covariant derivatives with respect to ω and ϖ are denoted by $D[\omega]_\mu$ and $D[\varpi]_\mu$. On a Lorentz vector X^a and spinor Ψ , one has $D[\omega]_\mu X^a = \partial_\mu X^a + \varepsilon^{abc}\omega_{\mu b}X_c$ and $D[\omega]_\mu \Psi = \partial_\mu \Psi + \frac{1}{2}\omega_\mu^a \gamma_a \Psi$ and analogously for $D[\varpi]_\mu X^a$ and $D[\varpi]_\mu \Psi$. The torsion and curvature for ω (and mutatis mutandis for ϖ) are defined as $T[\omega]_{\mu\nu}^a = 2D[\omega]_{[\mu}e_{\nu]}^a$ and $R[\omega]_{\mu\nu}^a = 2\partial_{[\mu}\omega_{\nu]}^a + \varepsilon^{abc}\omega_{\mu b}\omega_{\nu c}$. We have $\gamma_{\mu\nu} = \epsilon_{\mu\nu\rho}\gamma^\rho$ and spinor bilinears obey $\bar{\psi}\chi = \bar{\chi}\psi$ and $\bar{\psi}\gamma_\mu\chi = -\bar{\chi}\gamma_\mu\psi$.
- [16] A similar mechanism has been employed in N. S. Deger and H. Samtleben, A note on the third way consistent deformation of Yang-Mills theory, *Phys. Lett. B* **833**, 137275 (2022) for a description of the so-called third way consistent deformation of Yang-Mills theory in terms of a gauged scalar sigma model.
- [17] A. S. Arvanitakis, A. J. Routh, and P. K. Townsend, Matter coupling in 3D minimal massive gravity, *Classical Quantum Gravity* **31**, 235012 (2014).

- [18] The appearance of two independent spin connections is responsible for the presence of a massive degree of freedom in this formulation of MMG. For $\omega = \varpi$, the Lagrangian (4) would merely correspond to a reformulation of 3D (topological) gravity, such as studied in E. W. Mielke and P. Baekler, Topological gauge model of gravity with torsion, *Phys. Lett. A* **156**, 399 (1991); , Chern-Simons formulation of three-dimensional gravity with torsion and nonmetricity, *J. Geom. Phys.* **56**, 2523 (2006).
- [19] E. Sezgin and Y. Tani, Witten-Nester energy in topologically massive gravity, *Classical Quantum Gravity* **26**, 235005 (2009).
- [20] A. Routh, Hamiltonian form of topologically massive supergravity, *Phys. Rev. D* **88**, 024022 (2013).
- [21] N. S. Deger, M. Geiller, J. Rosseel, and H. Samtleben (to be published).
- [22] Thus pushing potential obstacles to sixth order in the fermions.
- [23] G. Gibbons, C. Pope, and E. Sezgin, The general supersymmetric solution of topologically massive supergravity, *Classical Quantum Gravity* **25**, 205005 (2008).
- [24] E. A. Bergshoeff, W. Merbis, and P. K. Townsend, On asymptotic charges in 3D gravity, *Classical Quantum Gravity* **37**, 035003 (2020).
- [25] H. R. Afshar and N. S. Deger, Exotic massive 3D gravities from truncation, *J. High Energy Phys.* **11** (2019) 145.
- [26] D. Roest and H. Samtleben, Twin supergravities, *Classical Quantum Gravity* **26**, 155001 (2009).