

Reconciling Conflicting Approaches for the Tunneling Time Delay in Strong Field Ionization

M. Klaiber¹, Q. Z. Lv¹, S. Sukiasyan¹, D. Bakucz Canário¹, K. Z. Hatsagortsyan^{1,*}, and C. H. Keitel¹
Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany



(Received 29 March 2020; revised 24 May 2021; accepted 12 October 2022; published 9 November 2022)

Several recent attoclock experiments have investigated the fundamental question of a quantum mechanically induced time delay in tunneling ionization via extremely precise photoelectron momentum spectroscopy. The interpretations of those attoclock experimental results were controversially discussed, because the entanglement of the laser and Coulomb field did not allow for theoretical treatments without undisputed approximations. The method of semiclassical propagation matched with the tunneled wave function, the quasistatic Wigner theory, the analytical R-matrix theory, the backpropagation method, and the under-the-barrier recollision theory are the leading conceptual approaches put forward to treat this problem, however, with seemingly conflicting conclusions on the existence of a tunneling time delay. To resolve the contradicting conclusions of the different approaches, we consider a very simple tunneling scenario which is not plagued with complications stemming from the Coulomb potential of the atomic core, avoids consequent controversial approximations and, therefore, allows us to unequivocally identify the origin of the tunneling time delay.

DOI: [10.1103/PhysRevLett.129.203201](https://doi.org/10.1103/PhysRevLett.129.203201)

Time delay in tunneling is a fascinating fundamental quantum problem, most recently measured in an experiment with cold atoms [1]. In particular, there has been intense and often controversial discussion about time delay in strong field tunneling ionization [2–39], confirming or disputing the interpretation of the experimental attoclock results [2–7]. The main difficulty stems from the fact that in such an experiment the photoelectron momentum distribution (PMD) is measured, rather than the tunneling time directly. This time is retrieved using time-to-angle mapping for photoelectrons tunnel-ionized in a laser field of elliptical polarization close to circular [2]. This mapping straightforwardly follows from the so-called simple man model [40]. According to this the photoelectron emission angle is determined by the direction of the laser vector potential at the moment of the electron appearing in the continuum. However, in a real physical situation the extraction of information on the tunneling time from PMD is not straightforward, because the Coulomb field of the atomic core induces a similar effect in PMD as the tunneling time delay (with respect to the peak of the laser field), and this effect is difficult to account for quantum mechanically with high accuracy. For this reason a

semiclassical method was proposed [2–4], where the tunneling was treated quantum mechanically, but the further electron motion in the continuum under the simultaneous action of the laser and Coulomb fields, classically. Moreover, the Coulomb field effect essentially depends on the tunnel exit coordinate, which in the quasistatic regime of ionization was calculated including tunnel ionization in parabolic coordinates with induced dipole and Stark shift also referred to as TIPIS model [4]. The semiclassical method was further improved, deriving the initial conditions of the classical propagation via the quantum mechanical Wigner trajectory emerging from the tunneling region [6,17]. However, non-adiabaticity of the tunneling ionization renders the quasistatic Wigner theory and related matched quantum-classical model inaccurate at large Keldysh parameters [41].

The numerical solutions of the time-dependent Schrödinger equation (TDSE) [7,13,42–46] for the attoclock reproduce the experimental results, yielding confidence that the attoclock PMD features have a single electron origin. However, the numerical results do not contribute much to our understanding of the tunneling time. Not long ago the backpropagation method was proposed to deduce the tunneling time delay from the numerical solution of TDSE [22–24]. In this method, the asymptotic numerical solution of TDSE is simulated by a classical ensemble, which is backpropagated classically up to the tunnel exit. With the backpropagation method a negligible time delay is obtained in weak fields, though nonnegligible negative time delays up to several atomic units (a.u.) at high fields. However, a conclusion was drawn that the tunneling

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Open access publication funded by the Max Planck Society.

time delay is negligible, assuming intuitively that the origin of the negative time cannot be tunneling. The TDSE numerical results have been also compared to the analytical R-matrix (ARM) theory [13], which is the state-of-the-art theory of the Coulomb-corrected strong field approximation (SFA) [47–51]. The comparison revealed that TDSE has a negative time delay with respect to ARM at large laser fields which, however, has been interpreted as a consequence of the bound state depletion and frustrated ionization [52]. The problem of the accurate description of subtle features of PMD has been addressed in Ref. [18] within SFA. A new type of quantum orbit was identified there, corresponding to under-the-barrier recollisions, where interference with the direct ionization path induces a gentle modification of the asymptotic PMD.

There are basically two opinions in interpreting the attoclock experiment, one claiming negligible (or zero) [7,13,23–26], and other nonnegligible asymptotic tunneling time delay (ATD) [4–6,17,18]. While ATD is read out from the asymptotic PMD, some also considered theoretically the near exit time delay (ETD) [15–17,27], observable in a Gedanken experiment with a so-called virtual detector near the tunnel exit [53,54]. These two faces of the concept of the tunneling time delay should be clearly distinguished. ATD is defined by the time delay of the classical trajectory, which is classically backpropagated from the peak of the photoelectron asymptotic wave function up to the exact point of the tunnel exit. The classical backpropagation is physically relevant, as the electron dynamics in the continuum is quasiclassical. However, the dynamical information can be extracted more accurately from the full quantum mechanical wave function. In particular, the so-called Wigner trajectory is deduced from the latter, which traces the time-dependent peak of the current density via the electron wave function during the tunneling ionization and defines ETD. The Wigner trajectory is in accordance with the classical backpropagation trajectory a few de-Broglie wavelengths away from the tunnel exit. However, near the tunnel exit the Wigner trajectory deviates strongly from the classical one and shows a positive ETD [17,38]. We note that ATD and ETD characterize the tunneling dynamics from different perspectives: While ATD is related to the attoclock protocol, ETD describes how the classical trajectory emerges from the quantum dynamics of the laser driven atomic electron.

This Letter is devoted to resolving conflicting conclusions of theories on the tunneling time delay and clarifying the difference between ATD and ETD. We judiciously consider a very basic tunneling scenario which is not plagued with complications stemming from Coulomb effects and the depletion of the bound state, is applicable in the adiabatic as well as in the nonadiabatic regimes, avoids consequent controversial approximations, and allows for analytical results. We consider ionization of a one-dimensional (1D) atom bound with a zero-range

potential driven by an half-cycle laser field. The calculation using SFA [41,55,56] is carried out fully analytically which facilitates a qualitative comparison of all differing models. Our results are confirmed by the numerical solutions of TDSE in 1D as well as in 3D with linearly and circularly polarized laser pulses. The reasons for the conclusions deviating from Refs. [7,22–26] are all analyzed.

We consider ionization of an electron bound in a 1D zero-range potential $V(x) = -\kappa\delta(x)$ [57], in a half-cycle laser pulse with electric field $E(t) = -E_0 \cos^2(\omega t)$, where $\omega = 0.05$ a.u., $\kappa = \sqrt{2I_p} = 1$ a.u., and I_p is the ionization potential. The Keldysh parameter is $\gamma = \tilde{\omega}\kappa/E_0$, with the effective frequency $\tilde{\omega} \equiv \sqrt{2}\omega$ related to the \cos^2 pulse. Atomic units are used throughout. We employ SFA, with incorporated low-frequency approximation (LFA) for a more accurate treatment (beyond the Born approximation) of the recollision [58–60]. The LFA validity is justified as the laser frequency $\omega \ll \varepsilon_r$ [58], with the recollision energy $\varepsilon_r \sim 1$ a.u. The asymptotic momentum distribution, $w(p) = |m(p)|^2 = |m_D(p) + m_R(p)|^2$ (see Fig. 1) is




FIG. 1. (a) Asymptotic PMD as function of $\delta p \equiv p + A(0)$ for $E_0 = 0.25$ a.u., where the grid line at $\delta p = 0$ shows the PMD peak with only the direct ionization amplitude m_D . (b) Tunneling time distribution using the backpropagation method of the asymptotic PMD. (c) The ratio of the rescattering amplitude to the direct ionization one $|m_R/m_D|$. (d) The shift of the PMD peak (δp) due to the tunneling time delay vs the laser field. For all panels (green-solid) is via the full SFA $m = m_D + m_R$, including sub-barrier direct and rescattered paths, (green-dashed) via the SFA direct amplitude, (black) via the numerical TDSE solution, (red) TDSE with the bound state depletion subtracted via the backpropagation method (see Sec. XIII.E.3 in the Supplemental Material [63]), (brown) via the static Wigner trajectory.

determined by interference of the direct ionization amplitude

$$m_D(p) = -i \int dt \langle \psi_p^V(t) | H_i(t) | \phi(t) \rangle, \quad (1)$$

and the ionization amplitude with rescattering, described by a second-order SFA [60,61]:

$$m_R(p) = - \int dt \int_t ds \int dq \langle \psi_p^V(s) | T[p + A(s)] | \psi_q^V(s) \rangle \times \langle \psi_q^V(t) | H_i(t) | \phi(t) \rangle. \quad (2)$$

Here $\phi(x, t) = \sqrt{\kappa} \exp(-\kappa|x| + ik^2/2t)$ is the bound state wave function, $\psi_p^V(x, t)$ the Volkov wave function [62], $H_i(t) = xE(t)$ the electron interaction Hamiltonian with the laser field, and $A(t) = - \int_{t_f}^t ds E(s)$. In the considered half-cycle laser field the rescattering takes place during the under-the-barrier dynamics, which is in LFA described with the exact laser-free scattering T matrix: $\langle p | T(p) | q \rangle = -(\kappa/2\pi)/(1 - ik/|p|)$ [60]. The time integration in m_R is carried out via 3D saddle-point approximation [63]. We have also calculated asymptotic PMD via numerical TDSE solution [63], which is in accordance with the analytical result and shows a momentum shift with respect to the zero-time delay case (PMD via the direct SFA amplitude m_D) corresponding to a negative ATD $\delta t = -\delta p/E_0$; see Fig. 1.

We retrieve the distribution of ATD from the asymptotic PMD [see Fig. 1(b)], using the backpropagating method [22]. The classical backpropagation up to the tunnel exit, where the longitudinal velocity is vanishing $p + A(t_e) = 0$, is carried out using the photoelectron asymptotic wave function $\psi(x, t) = \int m(p) \exp(ipx) dp$, with the total amplitude $m(p) = m_D + m_R$ [63]. The interference of the direct and the under-the-barrier rescattered trajectories, which is governed by the ratio of amplitudes m_R/m_D , induces a visible shift in the asymptotic PMD with respect to the case of the PMD based on the direct trajectory only, although $|m_R/m_D| \approx 0.13$. For instance, the momentum shift is $\delta p \sim 0.3$ a.u., which is equivalent to the negative ATD $t_e \sim -1$ a.u., at $E_0 = 0.25$ a.u.; see Figs. 1(b) and 1(c). The exponential suppression of ATD is governed by the parameter E_0/E_{th} [18], with the threshold field E_{th} of over-the-barrier ionization (OTBI) [63–65]. It is larger near the OTBI threshold [17,66] (the shorter the barrier length, the larger is the tunneling time [67]).

In Fig. 1(d) the dependence of the momentum shift δp due to the tunneling time delay on the laser field amplitude is shown. Thus, the calculations with our basic tunneling scenario show that the peak of asymptotic PMD can have a time delay up to the order of 1 a.u. [see Fig. 1(d)], due to interference of the direct and rescattered paths. The averaging over PMD decreases the time delay. We can give a simple estimation of the latter property. From Ref. [18], the negative

time delay is proportional to the Keldysh exponent $t_e(t) \approx t_0 \exp\{-2\kappa^3/[3|E(t)|]\}$, with the maximum of the time delay $t_{\max} = t_0 \exp\{-2\kappa^3/(3E_0)\}$, and $t_0 \sim 1/\kappa^2$, such that the averaged time delay can be estimated as $\langle t_e \rangle \sim \int_{-t_f}^{t_f} t_e(t) w(t) dt / \int_{-t_f}^{t_f} w(t) dt \sim 0.7 t_{\max}$, with the tunneling ionization probability $w(t) \sim \exp\{-2\kappa^3/[3|E(t)|]\}$.

The time delay calculated from the second-order SFA in the adiabatic regime $\gamma \ll 1$ is closely related to the static Wigner time delay. This is illustrated in Fig. 1(d) (see brown line), where the estimation of the static Wigner time delay is used; see Refs. [18,63]. In strong fields, the regime of ionization is adiabatic and the quasistatic Wigner theory is relevant [in Fig. 1(d) at $\gamma \lesssim 0.5$]. Note that in Ref. [6], the deviation of the experimental data from the quasistatic estimation takes place at $\gamma \gtrsim 0.6$.

In Ref. [23] the average tunneling time delay is calculated for helium ($\kappa = 1.345$). In this case the time delay is by a factor of $\kappa_{He}^2/\kappa^2 \approx 1.8$ smaller than in our $\kappa = 1$ a.u. case, since $t_e \sim 1/\kappa^2$ [68]. We may compare qualitatively our 1D case of Fig. 1(d) ($I_p = 0.5$) with the helium result of Ref. [23] at the same ratio of $E_0/E_{th}^{1D} = E_0/E_{th}^{He}$, using for helium $E_{th}^{He} = 0.24$ a.u. [63]. The scaled data of Ref. [23] provide the average negative tunneling time delay of the order up to 1 a.u. ($E_0 \lesssim 0.25$ a.u.), which qualitatively is in accordance with our model.

With the SFA time-dependent amplitude $m(p, t)$, we retrieve the time-dependent SFA wave function $\Psi(x, t)$ for the ionized electron:

$$\Psi(x, t) = \phi(x, t) + \int_{-\infty}^{\infty} dp \int_{-\infty}^t dt' \psi_p^V(x, t') m(p, t'); \quad (3)$$

and derive ETD from the latter as the peak of the full electron current density $j(x_e)$ at the tunnel exit x_e : see also Refs. [17,68,69]. The quantum mechanical description allows one to find the physical time delay at the exit which is read out as the peak of the time-dependent electron current density near the tunnel exit; see the current density distribution in Fig. 2, which shows that the most probable ETD is positive. This is also observed in the static tunneling case via the Wigner trajectory $t_e^W \sim 1/E_0^{2/3}$, yielding also a nonvanishing group velocity at the tunnel exit $v_W \sim E_0^{1/3}$ [17,68]. The SFA wave function in Fig. 2 is calculated via the direct ionization path. The difference between the TDSE and SFA results is due to the contribution of the recolliding trajectory. Thus, Fig. 2 shows that ETD is mostly determined by the direct ionization path. The inclusion of the recollision path disturbs ETD only slightly. Thus, the following picture of tunneling ionization emerges: the Wigner trajectory via the virtual detector (peak of the current density) shows a positive ETD. A few de-Broglie wavelengths away from the tunnel exit the most probable classical trajectory emerges from the Wigner one. The classical trajectory shows a negative ATD, which is




FIG. 2. The electron time-dependent current density $j(x_e)$ near the tunnel exit during tunneling ionization: (green-solid) via the full SFA, and (green-dashed) via the first-order SFA, (black) via the TDSE numerical solution, (red) via the TDSE numerical solution with depletion subtracted [63]. The ETD is positive as exhibited by the SFA curve, as well as by the TDSE with subtracted depletion.

vanishing in the deep tunneling regime $E_0 \ll E_{\text{th}}$; however, ETD is largest in this regime.

In Ref. [26] a “trajectory-free” method is proposed to address the tunneling time problem. It is based on the assumption that the (real part of the) saddle point of the time integrand of the ionization amplitude determines the ionization time (ATD). We question this assumption. Our line of reasoning is the following. From one side, with our simple SFA model we show that in the near OTBI regime the asymptotic momentum distribution is shifted with respect to the zero ATD case, i.e., demonstrating a nonvanishing ATD. From another side, we apply the “trajectory-free” method to the SFA wave function [63], i.e., represent the SFA wave function as a time integral, calculate the time saddle-point, and obtain that the saddle-point of the ionization wave function is zero in the same case, which shows a shift in the momentum distribution corresponding to the nonzero ATD; see Fig. 3. This demonstrates that the time saddle point and ATD are not equivalent. We interpret it as follows. The time saddle point indicates the complex time when the electron ionization path starts at the atomic core. Both amplitudes m_D and m_R start at $\text{Re}\{t_s\} = 0$ at the core, and each amplitude generates a momentum distribution with the peak corresponding to vanishing ATD. However, their interference causes a deviation of the momentum distribution peak from the no-time-delay model. In short, the time saddle point is the ionization starting point at the core, and it does not coincide with the time delay because the origin of the latter is the interference of two paths.

Finally, we have to comment on the numerically calculated vanishing ATD in Ref. [7] for an attoclock in the case of a short-range Yukawa potential $V(r) = -(Z/r)\exp(-r)$, with $Z = 1.908$. In Ref. [7] it is argued that the attoclock time delay in atoms is fully due to the Coulomb effect, and when the Coulomb field is removed in the case of a Yukawa potential, the ATD will be




FIG. 3. The time integrand of the ionization amplitude via SFA in the complex plane in the case of the most probable momentum $p = p_{\text{max}}$ for the total amplitude $m = m_D + m_R$. The saddle point of the total amplitude (see the cross) is vanishing: $\text{Re}\{t_s\} = 0$, while $\text{Re}\{t_s\} < 0$ for m_D and m_R , see Fig. 2 in the Supplemental Material [63]. The total amplitude has a shifted peak in momentum due to interference, however, the real part of the saddle point of the total amplitude for the peak momentum is still vanishing.

vanishing. However, the result of vanishing ATD in Ref. [7] is due to the fact that the range of the field strength of the calculation is not high enough for $E_0 \lesssim 0.075$. The threshold field of the applied Yukawa potential is $E_{\text{th}}^{\text{Yukawa}} \approx 0.1855$; it is smaller than that of the 1D short-range potential $E_{\text{th}}^{\text{Yukawa}}/E_{\text{th}}^{\text{1D}} \sim 0.7$. From the latter it is expectable to have a sizable ATD, for instance of the order of 1 a.u., near the threshold at $E_0 = 0.175$, while in the short-range potential it is observed at a larger value $E_0 = 0.23$; see Fig. 1(d). More detailed E_0/E_{th} scaling is different in 1D and 3D cases, which stems from the wave packet spreading factor in the transverse direction [18]. Furthermore, with the given high charge $Z \approx 2$, the momentum transfer at the tunnel exit due to the atomic potential is not negligible at high field strengths, when the tunnel exit is close to the core [63]. The attoclock angular offset angle due to the atomic potential corresponds to the positive time delay, which counteracts and reduces the observable negative ATD. We have carried out calculations of the ATD for the given Yukawa potential in a large range of field strengths using numerical 3D TDSE solutions for attoclock scenarios [43,44,63]. In Fig. 4 we show the total time delay calculated via the backpropagation from the 3D TDSE wave function, as well as the time delay after the subtraction of the depletion contribution (via BP method, see Sec. XII.E.3 of the Supplemental Material [63]). The attoclock offset angle is vanishingly small when the field is not large $E_0 \lesssim 0.075$ (up to the laser intensity 4×10^{14} W/cm²), in agreement with the calculation of




FIG. 4. Time delay calculated by the classical backpropagation (BP) of 3D TDSE results with a Yukawa potential $V(r) = -(Z/r)\exp(-r)$, with $Z = 1.908$ ($E_0 \leq 0.125$ a.u.) and $Z = 1.908183$ ($E_0 \geq 0.15$ a.u.) in a circularly polarized laser pulse: (black circles) total time shift, δt_{total} (BP), (red squares) final time shift δt_{ATD} (BP) with the depletion subtracted via the backpropagation method (see Sec. XIII.E.3 in the Supplemental Material [63]). The field strengths E_0 are shown up to $E_{\text{th}}^{\text{Yukawa}}$.

Ref. [7], but increases at high field strengths showing a negative time delay up to 1.7 a.u.

Concluding, we have analyzed the tunneling time delay in strong field ionization employing a simple tunneling scenario where leading theoretical approaches with seemingly conflicting conclusions could be compared without critical approximations. This way we have demonstrated that the peak of the tunneling wave packet emerging from the barrier around the tunnel exit significantly deviates from the most probable classical backpropagated trajectory, featuring a positive ETD. It however asymptotically merges with the backpropagated trajectory, which itself shows negative ATD, originating from the interference of the direct and recolliding sub-barrier paths. Finally, in explaining the absence of tunneling times from the remaining other methods, we have clarified that there are indeed no conflicts among the various approaches. In particular, we have shown that the attoclock offset angle cannot arise solely due to the Coulomb effect of the atomic core, because it exists also in the case of a Yukawa potential for strong fields. Furthermore, the starting point of ionization at the atom without time delay with respect to the peak of the laser field does not exclude nonvanishing ATD.

*k.hatsagortsyan@mpi-hd.mpg.de

[1] R. Ramos, D. Spierings, I. Racicot, and A. M. Steinberg, Measurement of the time spent by a tunnelling atom within the barrier region, *Nature (London)* **583**, 529 (2020).

- [2] P. Eckle, M. Smolarski, F. Schlup, J. Biegert, A. Staudte, M. Schöffler, H. G. Müller, R. Dörner, and U. Keller, Attosecond angular streaking, *Nat. Phys.* **4**, 565 (2008).
- [3] P. Eckle, A. N. Pfeiffer, C. Cirelli, A. Staudte, R. Dörner, H. G. Müller, M. Büttiker, and U. Keller, Attosecond ionization and tunneling delay time measurements in helium, *Science* **322**, 1525 (2008).
- [4] A. N. Pfeiffer, C. Cirelli, M. Smolarski, D. Dimitrovski, M. Abu-samha, L. B. Madsen, and U. Keller, Attoclock reveals natural coordinates of the laser-induced tunnelling current flow in atoms, *Nat. Phys.* **8**, 76 (2012).
- [5] A. S. Landsman, M. Weger, J. Maurer, R. Boge, A. Ludwig, S. Heuser, C. Cirelli, L. Gallmann, and U. Keller, Ultrafast resolution of tunneling delay time, *Optica* **1**, 343 (2014).
- [6] N. Camus, E. Yakaboylu, L. Fechner, M. Klaiber, M. Laux, Y. Mi, K. Z. Hatsagortsyan, T. Pfeifer, C. H. Keitel, and R. Moshhammer, Experimental Evidence for Wigner's Tunneling Time, *Phys. Rev. Lett.* **119**, 023201 (2017).
- [7] U. S. Sainadh, H. Xu, X. Wang, A. Atia-Tul-Noor, W. C. Wallace, N. Douguet, A. Bray, I. Ivanov, K. Bartschat, A. Kheifets, R. T. Sang, and I. V. Litvinyuk, Attosecond angular streaking and tunnelling time in atomic hydrogen, *Nature (London)* **568**, 75 (2019).
- [8] A. Czirják, R. Kopold, W. Becker, M. Kleber, and W. Schleich, The Wigner function for tunneling in a uniform static electric field, *Opt. Commun.* **179**, 29 (2000).
- [9] M. Lein, Streaking analysis of strong-field ionisation, *J. Mod. Opt.* **58**, 1188 (2011).
- [10] G. Orlando, C. R. McDonald, N. H. Protik, G. Vampa, and T. Brabec, Tunnelling time, what does it mean?, *J. Phys. B* **47**, 204002 (2014).
- [11] G. Orlando, C. R. McDonald, N. H. Protik, and T. Brabec, Identification of the Keldysh time as a lower limit for the tunneling time, *Phys. Rev. A* **89**, 014102 (2014).
- [12] A. S. Landsman and U. Keller, Tunnelling time in strong field ionisation, *J. Phys. B* **47**, 204024 (2014).
- [13] L. Torlina, F. Morales, J. Kaushal, I. Ivanov, A. Kheifets, A. Zielinski, A. Scrinzi, H. G. Müller, S. Sukiasyan, M. Ivanov, and O. Smirnova, Interpreting attoclock measurements of tunnelling times, *Nat. Phys.* **11**, 503 (2015).
- [14] A. S. Landsman and U. Keller, Attosecond science and the tunnelling time problem, *Phys. Rep.* **547**, 1 (2015).
- [15] N. Teeny, E. Yakaboylu, H. Bauke, and C. H. Keitel, Ionization Time and Exit Momentum in Strong-Field Tunnel Ionization, *Phys. Rev. Lett.* **116**, 063003 (2016).
- [16] N. Teeny, C. H. Keitel, and H. Bauke, Virtual-detector approach to tunnel ionization and tunneling times, *Phys. Rev. A* **94**, 022104 (2016).
- [17] E. Yakaboylu, M. Klaiber, H. Bauke, K. Z. Hatsagortsyan, and C. H. Keitel, Relativistic features and time delay of laser-induced tunnel ionization, *Phys. Rev. A* **88**, 063421 (2013).
- [18] M. Klaiber, K. Z. Hatsagortsyan, and C. H. Keitel, Under-the-Tunneling-Barrier Recollisions in Strong-Field Ionization, *Phys. Rev. Lett.* **120**, 013201 (2018).
- [19] T. Zimmermann, S. Mishra, B. R. Doran, D. F. Gordon, and A. S. Landsman, Tunneling Time and Weak Measurement in Strong Field Ionization, *Phys. Rev. Lett.* **116**, 233603 (2016).

- [20] J. Liu, Y. Fu, W. Chen, Z. Lü, J. Zhao, J. Yuan, and Z. Zhao, Offset angles of photocurrents generated in few-cycle circularly polarized laser fields, *J. Phys. B* **50**, 055602 (2017).
- [21] Y. Song, Y. Yang, F. Guo, and S. Li, Revisiting the time-dependent ionization process through the bohmian-mechanics method, *J. Phys. B* **50**, 095003 (2017).
- [22] H. Ni, U. Saalmann, and J.-M. Rost, Tunneling Ionization Time Resolved by Backpropagation, *Phys. Rev. Lett.* **117**, 023002 (2016).
- [23] H. Ni, U. Saalmann, and J.-M. Rost, Tunneling exit characteristics from classical backpropagation of an ionized electron wave packet, *Phys. Rev. A* **97**, 013426 (2018).
- [24] H. Ni, N. Eicke, C. Ruiz, J. Cai, F. Oppermann, N. I. Shvetsov-Shilovski, and L.-W. Pi, Tunneling criteria and a nonadiabatic term for strong-field ionization, *Phys. Rev. A* **98**, 013411 (2018).
- [25] A. S. Kheifets, The attoclock and the tunneling time debate, *J. Phys. B* **53**, 072001 (2020).
- [26] N. Eicke and M. Lein, Trajectory-free ionization times in strong-field ionization, *Phys. Rev. A* **97**, 031402(R) (2018).
- [27] N. Douguet and K. Bartschat, Dynamics of tunneling ionization using Bohmian mechanics, *Phys. Rev. A* **97**, 013402 (2018).
- [28] A. W. Bray, S. Eckart, and A. S. Kheifets, Keldysh-Rutherford Model for the Attoclock, *Phys. Rev. Lett.* **121**, 123201 (2018).
- [29] B. Baytaş, M. Bojowald, and S. Crowe, Canonical tunneling time in ionization experiments, *Phys. Rev. A* **98**, 063417 (2018).
- [30] X. Ren, Y. Wu, L. Wang, and Y. Zheng, Entangled trajectories during ionization of an h atom driven by n-cycle laser pulse, *Phys. Lett. A* **382**, 2662 (2018).
- [31] J. Tan, Y. Zhou, M. He, Y. Chen, Q. Ke, J. Liang, X. Zhu, M. Li, and P. Lu, Determination of the Ionization Time Using Attosecond Photoelectron Interferometry, *Phys. Rev. Lett.* **121**, 253203 (2018).
- [32] D. Sokolovski and E. Akhmatskaya, No time at the end of the tunnel, *Commun. Phys.* **1**, 47 (2018).
- [33] W. Quan, V. V. Serov, M. Z. Wei, M. Zhao, Y. Zhou, Y. L. Wang, X. Y. Lai, A. S. Kheifets, and X. J. Liu, Attosecond Molecular Angular Streaking with All-Ionic Fragments Detection, *Phys. Rev. Lett.* **123**, 223204 (2019).
- [34] N. Douguet and K. Bartschat, Attoclock setup with negative ions: A possibility for experimental validation, *Phys. Rev. A* **99**, 023417 (2019).
- [35] C. Hofmann, A. S. Landsman, and U. Keller, Attoclock revisited on electron tunnelling time, *J. Mod. Opt.* **66**, 1052 (2019).
- [36] V. V. Serov, A. W. Bray, and A. S. Kheifets, Numerical attoclock on atomic and molecular hydrogen, *Phys. Rev. A* **99**, 063428 (2019).
- [37] R. Wang, Q. Zhang, D. Li, S. Xu, P. Cao, Y. Zhou, W. Cao, and P. Lu, Identification of tunneling and multiphoton ionization in intermediate Keldysh parameter regime, *Opt. Express* **27**, 6471 (2019).
- [38] M. Han, P. Ge, Y. Fang, X. Yu, Z. Guo, X. Ma, Y. Deng, Q. Gong, and Y. Liu, Unifying Tunneling Pictures of Strong-Field Ionization with an Improved Attoclock, *Phys. Rev. Lett.* **123**, 073201 (2019).
- [39] M. Yuan, Direct probing of tunneling time in strong-field ionization processes by time-dependent wave packets, *Opt. Express* **27**, 6502 (2019).
- [40] P. B. Corkum, Plasma Perspective on Strong Field Multiphoton Ionization, *Phys. Rev. Lett.* **71**, 1994 (1993).
- [41] L. V. Keldysh, Ionization in the field of a strong electromagnetic wave, *Zh. Eksp. Teor. Fiz.* **47**, 1945 (1964) [*Sov. Phys. JETP* **20**, 1307 (1965)].
- [42] I. A. Ivanov and A. S. Kheifets, Strong-field ionization of he by elliptically polarized light in attoclock configuration, *Phys. Rev. A* **89**, 021402(R) (2014).
- [43] H. G. Muller, An efficient scheme for the time-dependent Schrödinger equation in the velocity gauge, *Laser Phys.* **9**, 138 (1999).
- [44] D. Bauer and P. Koval, Qprop: A Schrödinger-solver for intense laser-atom interaction, *Comput. Phys. Commun.* **174**, 396 (2006).
- [45] S. Patchkovskii and H. G. Muller, Simple, accurate, and efficient implementation of 1-electron atomic time-dependent schrödinger equation in spherical coordinates, *Comput. Phys. Commun.* **199**, 153 (2016).
- [46] V. P. Majety and A. Scrinzi, Absence of electron correlation effects in the helium attoclock setting, *J. Mod. Opt.* **64**, 1026 (2017).
- [47] S. V. Popruzhenko, G. G. Paulus, and D. Bauer, Coulomb-corrected quantum trajectories in strong-field ionization, *Phys. Rev. A* **77**, 053409 (2008).
- [48] S. V. Popruzhenko and D. Bauer, Strong field approximation for systems with Coulomb interaction, *J. Mod. Opt.* **55**, 2573 (2008).
- [49] L. Torlina and O. Smirnova, Time-dependent analytical *R*-matrix approach for strong-field dynamics. I. One-electron systems, *Phys. Rev. A* **86**, 043408 (2012).
- [50] L. Torlina, J. Kaushal, and O. Smirnova, Time-resolving electron-core dynamics during strong-field ionization in circularly polarized fields, *Phys. Rev. A* **88**, 053403 (2013).
- [51] J. Kaushal and O. Smirnova, Nonadiabatic coulomb effects in strong-field ionization in circularly polarized laser fields, *Phys. Rev. A* **88**, 013421 (2013).
- [52] T. Nubbemeyer, K. Gorling, A. Saenz, U. Eichmann, and W. Sandner, Strong-Field Tunneling without Ionization, *Phys. Rev. Lett.* **101**, 233001 (2008).
- [53] B. Feuerstein and U. Thumm, On the computation of momentum distributions within wavepacket propagation calculations, *J. Phys. B* **36**, 707 (2003).
- [54] X. Wang, J. Tian, and J. H. Eberly, Extended Virtual Detector Theory for Strong-Field Atomic Ionization, *Phys. Rev. Lett.* **110**, 243001 (2013).
- [55] F. H. M. Faisal, Multiple absorption of laser photons by atoms, *J. Phys. B* **6**, L89 (1973).
- [56] H. R. Reiss, Effect of an intense electromagnetic field on a weakly bound system, *Phys. Rev. A* **22**, 1786 (1980).
- [57] K. Krajewska, J. Kamiński, and K. Wódkiewicz, Zero-range interaction in arbitrary dimensions and in the presence of external forces, *Opt. Commun.* **283**, 843 (2010).
- [58] N. M. Kroll and K. M. Watson, Charged-particle scattering in the presence of a strong electromagnetic wave, *Phys. Rev. A* **8**, 804 (1973).

- [59] A. Čerkić, E. Hasović, D. B. Milošević, and W. Becker, High-order above-threshold ionization beyond the first-order born approximation, *Phys. Rev. A* **79**, 033413 (2009).
- [60] D. B. Milošević, Low-frequency approximation for above-threshold ionization by a laser pulse: Low-energy forward rescattering, *Phys. Rev. A* **90**, 063423 (2014).
- [61] W. Becker, F. Grasbon, R. Kopold, D. B. Milošević, G. G. Paulus, and H. Walther, Above-threshold ionization: From classical features to quantum effects, *Adv. At. Mol. Opt. Phys.* **48**, 35 (2002).
- [62] D. M. Wolkow, Über eine Klasse von Lösungen der Diracschen Gleichung, *Z. Phys.* **94**, 250 (1935).
- [63] See Supplemental Materials at <http://link.aps.org/supplemental/10.1103/PhysRevLett.129.203201> for the details.
- [64] S. Augst, D. Strickland, D. D. Meyerhofer, S. L. Chin, and J. H. Eberly, Tunneling Ionization of Noble Gases in a High-Intensity Laser Field, *Phys. Rev. Lett.* **63**, 2212 (1989).
- [65] N. B. Delone and V. P. Krainov, *Multiphoton Processes in Atoms*, Springer Series on Atoms+Plasmas (Springer, Berlin, 2000).
- [66] E. Yakaboylu, M. Klaiber, and K. Z. Hatsagortsyan, Wigner time delay for tunneling ionization via the electron propagator, *Phys. Rev. A* **90**, 012116 (2014).
- [67] A. Peres, Measurement of time by quantum clocks, *Am. J. Phys.* **48**, 552 (1980).
- [68] M. Klaiber, E. Yakaboylu, H. Bauke, K. Z. Hatsagortsyan, and C. H. Keitel, Under-the-Barrier Dynamics in Laser-Induced Relativistic Tunneling, *Phys. Rev. Lett.* **110**, 153004 (2013).
- [69] D. Bakucz Canário, M. Klaiber, K. Z. Hatsagortsyan, and C. H. Keitel, Role of reflections in the generation of a time delay in strong-field ionization, *Phys. Rev. A* **104**, 033103 (2021).