

New Direct Limit on Neutrinoless Double Beta Decay Half-Life of ^{128}Te with CUORE

D. Q. Adams,¹ C. Alduino,¹ K. Alfonso,² F. T. Avignone III,¹ O. Azzolini,³ G. Bari,⁴ F. Bellini,^{5,6} G. Benato,⁷ M. Beretta,⁸ M. Biassoni,⁹ A. Branca,^{10,9} C. Brofferio,^{10,9} C. Bucci,⁷ J. Camilleri,¹¹ A. Caminata,¹² A. Campani,^{13,12} L. Canonica,^{14,7} X. G. Cao,¹⁵ S. Capelli,^{10,9} C. Capelli,¹⁶ L. Cappelli,⁷ L. Cardani,⁶ P. Carniti,^{10,9} N. Casali,⁶ E. Celi,^{17,7} D. Chiesa,^{10,9} M. Clemenza,^{10,9} S. Copello,^{13,12} O. Cremonesi,⁹ R. J. Creswick,¹ A. D'Addabbo,⁷ I. Dafinei,⁶ F. Del Corso,⁴ S. Dell'Oro,^{10,9} S. Di Domizio,^{13,12} S. Di Lorenzo,⁷ V. Dompè,^{5,6} D. Q. Fang,¹⁵ G. Fantini,^{5,6} M. Faverzani,^{10,9} E. Ferri,⁹ F. Ferroni,^{17,6} E. Fiorini,^{9,10} M. A. Franceschi,¹⁸ S. J. Freedman,^{16,8*} S. H. Fu,¹⁵ B. K. Fujikawa,¹⁶ S. Ghislandi,^{17,7} A. Giachero,^{10,9} A. Gianvecchio,^{10,9} L. Gironi,^{10,9} A. Giuliani,¹⁹ P. Gorla,⁷ C. Gotti,⁹ T. D. Gutierrez,²⁰ K. Han,²¹ E. V. Hansen,⁸ K. M. Heeger,²² R. G. Huang,⁸ H. Z. Huang,² J. Johnston,¹⁴ G. Keppel,³ Yu. G. Kolomensky,^{8,16} R. Kowalski,²³ R. Liu,²² L. Ma,² Y. G. Ma,¹⁵ L. Marini,^{17,7} R. H. Maruyama,²² D. Mayer,¹⁴ Y. Mei,¹⁶ S. Morganti,⁶ T. Napolitano,¹⁸ M. Nastasi,^{10,9} J. Nikkel,²² C. Nones,²⁴ E. B. Norman,^{25,26} A. Nucciotti,^{10,9} I. Nutini,^{10,9} T. O'Donnell,¹¹ M. Olmi,⁷ J. L. Ouellet,¹⁴ S. Pagan,²² C. E. Pagliarone,^{7,27} L. Pagnanini,⁷ M. Pallavicini,^{13,12} L. Pattavina,⁷ M. Pavan,^{10,9} G. Pessina,⁹ V. Pettinacci,⁶ C. Pira,³ S. Pirro,⁷ S. Pozzi,^{10,9} E. Previtali,^{10,9} A. Puiu,^{17,7} S. Quitadamo,^{17,7} A. Ressa,^{5,6} C. Rosenfeld,¹ S. Sangiorgio,²⁵ B. Schmidt,¹⁶ N. D. Scielzo,²⁵ V. Sharma,¹¹ V. Singh,⁸ M. Sisti,⁹ D. Speller,²³ P. T. Surukuchi,²² L. Taffarello,²⁸ F. Terranova,^{10,9} C. Tomei,⁶ K. J. Vetter,^{8,16} M. Vignati,^{5,6} S. L. Wagaarachchi,^{8,16} B. S. Wang,^{25,26} B. Welliver,^{8,16} J. Wilson,¹ K. Wilson,¹ L. A. Winslow,¹⁴ S. Zimmermann,²⁹ and S. Zucchelli^{30,4}

(CUORE Collaboration)

¹*Department of Physics and Astronomy, University of South Carolina, Columbia, South Carolina 29208, USA*

²*Department of Physics and Astronomy, University of California, Los Angeles, California 90095, USA*

³*INFN—Laboratori Nazionali di Legnaro, Legnaro (Padova) I-35020, Italy*

⁴*INFN—Sezione di Bologna, Bologna I-40127, Italy*

⁵*Dipartimento di Fisica, Sapienza Università di Roma, Roma I-00185, Italy*

⁶*INFN—Sezione di Roma, Roma I-00185, Italy*

⁷*INFN—Laboratori Nazionali del Gran Sasso, Assergi (L'Aquila) I-67100, Italy*

⁸*Department of Physics, University of California, Berkeley, California 94720, USA*

⁹*INFN—Sezione di Milano Bicocca, Milano I-20126, Italy*

¹⁰*Dipartimento di Fisica, Università di Milano-Bicocca, Milano I-20126, Italy*

¹¹*Center for Neutrino Physics, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, USA*

¹²*INFN—Sezione di Genova, Genova I-16146, Italy*

¹³*Dipartimento di Fisica, Università di Genova, Genova I-16146, Italy*

¹⁴*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

¹⁵*Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), Institute of Modern Physics, Fudan University, Shanghai 200433, China*

¹⁶*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

¹⁷*Gran Sasso Science Institute, L'Aquila I-67100, Italy*

¹⁸*INFN—Laboratori Nazionali di Frascati, Frascati (Roma) I-00044, Italy*

¹⁹*Universit Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France*

²⁰*Physics Department, California Polytechnic State University, San Luis Obispo, California 93407, USA*

²¹*INPAC and School of Physics and Astronomy, Shanghai Jiao Tong University;*

Shanghai Laboratory for Particle Physics and Cosmology, Shanghai 200240, China

²²*Wright Laboratory, Department of Physics, Yale University, New Haven, Connecticut 06520, USA*

²³*Department of Physics and Astronomy, The Johns Hopkins University,*

3400 North Charles Street Baltimore, Maryland 21211, USA

²⁴*IRFU, CEA, Universit Paris-Saclay, F-91191 Gif-sur-Yvette, France*

²⁵*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*

²⁶*Department of Nuclear Engineering, University of California, Berkeley, California 94720, USA*

²⁷*Dipartimento di Ingegneria Civile e Meccanica, Università degli Studi di Cassino e del Lazio Meridionale, Cassino I-03043, Italy*

²⁸*INFN—Sezione di Padova, Padova I-35131, Italy*

²⁹*Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

³⁰*Dipartimento di Fisica e Astronomia, Alma Mater Studiorum—Università di Bologna, Bologna I-40127, Italy*



(Received 6 May 2022; revised 22 July 2022; accepted 3 October 2022; published 23 November 2022)

The Cryogenic Underground Observatory for Rare Events (CUORE) at Laboratori Nazionali del Gran Sasso of INFN in Italy is an experiment searching for neutrinoless double beta ($0\nu\beta\beta$) decay. Its main goal is to investigate this decay in ^{130}Te , but its ton-scale mass and low background make CUORE sensitive to other rare processes as well. In this Letter, we present our first results on the search for $0\nu\beta\beta$ decay of ^{128}Te , the Te isotope with the second highest natural isotopic abundance. We find no evidence for this decay, and using a Bayesian analysis we set a lower limit on the ^{128}Te $0\nu\beta\beta$ decay half-life of $T_{1/2} > 3.6 \times 10^{24}$ yr (90% CI). This represents the most stringent limit on the half-life of this isotope, improving by over a factor of 30 the previous direct search results, and exceeding those from geochemical experiments for the first time.

DOI: [10.1103/PhysRevLett.129.222501](https://doi.org/10.1103/PhysRevLett.129.222501)

Double beta ($\beta\beta$) decay is a rare second-order Fermi interaction in which a nucleus (A, Z) transforms into its isobar ($A, Z + 2$) by the simultaneous transmutation of two neutrons into two protons. This standard model process occurs with the emission of two electrons and two electron antineutrinos in the final state ($2\nu\beta\beta$ decay), such that the lepton number (L) conservation holds; this process has been measured for 11 nuclei [1], with half-lives in the range of 10^{18} – 10^{22} years. A second decay mode, neutrinoless double beta ($0\nu\beta\beta$) decay, has been hypothesized but never observed. This process would consist of a nucleus $\beta\beta$ decaying into its daughter with the emission of two electrons and no antineutrinos in the final state, thus violating L by two units. The experimental signature of this process is a peak in the two-electron total energy spectrum at the Q value ($Q_{\beta\beta}$) of the transition. The search for $0\nu\beta\beta$ decay addresses one of the most relevant open questions in neutrino physics: its observation would establish that L is not a symmetry of nature and neutrinos are Majorana fermions, providing a clear signature of physics beyond the standard model [2,3]. This would provide significant input for the explanation of the matter-antimatter asymmetry in the Universe via leptogenesis [4,5], as well as constraints on the absolute mass scale and ordering of neutrinos complementing other approaches [3,6].

CUORE is a ton-scale array of 988 TeO_2 crystals designed to search for $0\nu\beta\beta$ decay of ^{130}Te . Besides having the world leading sensitivity for this process [7,8] due to its very large mass—742 kg of TeO_2 —and low background, CUORE is also a powerful detector for other rare processes, in particular other Te decay channels [9–11]. In this Letter we report on a new direct search for ^{128}Te $0\nu\beta\beta$ decay. The CUORE array is grown from material with natural isotopic composition, which given the natural abundance of 31.75% [12] contains 188 kg of ^{128}Te . Despite this high abundance, the direct search is challenging due to the low $Q_{\beta\beta}$ value of (866.7 ± 0.7) keV [13] which lies in a region of the energy spectrum dominated by $2\nu\beta\beta$ decay of ^{130}Te and γ backgrounds from other natural radioactivity. The most recent ^{128}Te $0\nu\beta\beta$ decay half-life limit from a direct search experiment, $T_{1/2}^{0\nu} > 1.1 \times 10^{23}$ yr, was set by MiDBD in 2003 [14]. More stringent limits than this have

been set by indirect geochemical measurements (see Ref. [15] for a review), which evaluate the presence of the $\beta\beta$ decay products accumulated in geological mineral samples of known age via the assessment of the parent-daughter nuclei ratio.

The geochemical studies are not sensitive to the $\beta\beta$ decay mode but rather to the sum of all the possible decays ($2\nu\beta\beta$ or $0\nu\beta\beta$, to the ground or excited states), although the dominant contribution is expected to be the two-neutrino mode. The direct search result reported in this Letter improves by more than 30-fold the previous best direct search limit for this isotope and surpasses—for the first time—the indirect geochemical results.

Before reporting the details of our direct search for ^{128}Te $0\nu\beta\beta$ decay, we present an updated evaluation of the half-life value for ^{128}Te $\beta\beta$ decay based on the ratio $T_{1/2}(^{130}\text{Te})/T_{1/2}(^{128}\text{Te}) = (3.52 \pm 0.11) \times 10^{-4}$ [16] from ion-counting mass spectrometry of Xe in ancient Te samples. Using the most recent ^{130}Te $2\nu\beta\beta$ decay half-life measurement, $7.71_{-0.06}^{+0.08}(\text{stat})_{-0.15}^{+0.12}(\text{syst}) \times 10^{20}$ yr [9], we obtain $T_{1/2}^{2\nu}(^{128}\text{Te}) = (2.19 \pm 0.07) \times 10^{24}$ yr. This result replaces and is in agreement with the previously published value of $T_{1/2}^{2\nu}(^{128}\text{Te}) = (2.25 \pm 0.09) \times 10^{24}$ yr [1], which used the weighted average of the ^{130}Te $2\nu\beta\beta$ decay half-lives from CUORE-0 [17] and CUORE [18].

The CUORE detector comprises 19 towers of 52 crystals each. The basic unit is a $5 \times 5 \times 5$ cm³ TeO_2 crystal operated as an individual cryogenic calorimeter. Each crystal is equipped with a neutron transmutation doped (NTD) Ge thermistor [19], used as a temperature sensor, and a Si resistor to inject controlled heat pulses for thermal gain stabilization. The crystal is coupled through polytetrafluoroethylene and Cu supports to the coldest stage of a dilution refrigerator operating at a temperature of ~ 10 mK [20]. Any particle interaction in a TeO_2 absorber crystal produces an energy deposition that is converted into heat (phonons) and measured via the temperature sensor. A large and novel cryogenic infrastructure has been developed to provide the needed cooling power [8]. The CUORE cryostat is designed to meet the CUORE background specifications [21], and provides a low thermal noise

environment, minimizing vibration and thermal dissipation on the cryogenic calorimeters [20,22–24]. CUORE is the most advanced realization of the cryogenic calorimetric technology, developed over 30 years using TeO_2 -based detectors [25].

We acquire data in day-long periods called runs, which in turn are grouped into ~ 40 – 60 days collections called datasets. A typical dataset consists of 4–5 days of calibration runs, followed by 30–50 days of so-called physics runs, and finally another 4–5 days of final calibration to check the energy scale stability within a dataset. Calibration runs are performed using γ -ray sources of ^{232}Th and ^{60}Co to illuminate the detectors.

The procedure for the data acquisition and processing is described in [7]. We apply a digital optimum trigger algorithm [26,27] to the acquired continuous data stream and evaluate the amplitude of the triggered waveform by applying a frequency-based optimum filter that weights the Fourier components of the signal, exploiting the noise power spectrum to reduce the impact of noisy frequencies. We compensate for thermal gain variations in the crystals due to small fluctuations in their operating temperature with two independent methods. The first utilizes heat pulses of fixed amplitude injected regularly (every 570 s) via the Si heaters affixed to the crystals. For crystals with nonfunctional heaters we use the 2615 keV γ events from ^{208}Tl in calibration data as a reference. We use the data from calibration runs to convert the thermal amplitudes to units of energy. We exploit the granularity of the CUORE detector to perform a coincidence study and determine if signals in different crystals within a short time and spatial distance (typically of 10 ms and 150 mm) are attributed to the same physical interaction. We refer to these as coincident signals, to which we assign a multiplicity number \mathcal{M}_n , where n corresponds to the number of crystals simultaneously involved in the interaction (e.g., two events in different crystals due to Compton scattering of the 2615 keV ^{208}Tl line are labeled as \mathcal{M}_2), with single-crystal interactions labeled as \mathcal{M}_1 . We apply a pulse shape analysis (PSA) algorithm to identify and discriminate pulses due to particle energy depositions from nonphysical signals (e.g., noise spikes, abrupt baseline disturbances, pileup events).

The present analysis includes five datasets for a total TeO_2 exposure of 309.33 kg yr or 78.56 kg yr of ^{128}Te . These are the same data we used to measure the ^{130}Te $2\nu\beta\beta$ decay half-life [9]. However, the latter exposure is marginally lower (300.72 kg yr) due to stricter selection criteria on the energy scale calibration in both the β/γ (< 3 MeV) and α (> 3 MeV) regions for the $2\nu\beta\beta$ decay result. In contrast, this analysis requires only good performance in the β/γ region.

In the following, we provide a detailed description of the analysis technique used to search for ^{128}Te $0\nu\beta\beta$ decay, whose signature is a monoenergetic peak at

$Q_{\beta\beta} = (866.7 \pm 0.7)$ keV in the summed energy of the two emitted electrons. In the great majority of the cases the two electrons are absorbed by the same crystal: we therefore select \mathcal{M}_1 events only, within a region of interest (ROI) of (820–890) keV.

The signal efficiency is the product of the containment efficiency and the total analysis efficiency. We define the containment efficiency (ϵ_{MC}) as the fraction of ^{128}Te $0\nu\beta\beta$ decay events that release their full energy, i.e., $Q_{\beta\beta}$, in a single crystal [28]. We evaluate ϵ_{MC} by simulating 10^8 events in the CUORE crystals [21], obtaining $\epsilon_{\text{MC}} = 97.59 \pm 0.01\%$. The total analysis efficiency (exposure-weighted average $\epsilon_{\text{cut}} = 87.74 \pm 0.19\%$) is the product of the total reconstruction efficiency, the anticoincidence efficiency and the PSA efficiency. The first term is the probability that an event with a given energy is triggered, its energy is correctly reconstructed, and it is not rejected as a pileup event by the analysis cuts applied during the data processing; the anticoincidence efficiency is the probability that a single-hit event is not assigned the wrong multiplicity due to a random accidental coincidence with an unrelated event; the PSA efficiency is the probability that events passing the base pileup cuts also survive the PSA cut. We refer to [7] for a more detailed description of the computation methods of these efficiency terms.

To avoid introducing bias when choosing the fit model of the present analysis, we choose the ROI based on the CUORE background model simulations, particularly taking into account backgrounds close to $Q_{\beta\beta}$ for ^{128}Te (Fig. 1). Based on this, we choose an ROI of (820–890) keV.

Multiple peaks populate this energy window: the closest expected structure to $Q_{\beta\beta}$ is a γ line at 860.6 keV from ^{208}Tl , a ^{232}Th chain element. A prominent peak at 834.8 keV due




FIG. 1. \mathcal{M}_1 spectrum from the CUORE background model simulations in the proximity of the ^{128}Te $0\nu\beta\beta$ decay $Q_{\beta\beta}$. From left to right: ^{54}Mn γ (834.8 keV), ^{208}Tl γ (860.6 keV), and ^{228}Ac γ (911.2 keV). The ROI for this analysis is denoted by the dashed green box, and includes the ^{54}Mn and ^{208}Tl lines.

to a ^{54}Mn γ line is also identified: the presence of ^{54}Mn stems from the cosmogenic activation of copper [17,29]. The visible peak to the right of $Q_{\beta\beta}$ is the 911.2 keV γ line from ^{228}Ac , another element of the ^{232}Th chain. In addition, we observe a continuous background contribution mainly induced by the $2\nu\beta\beta$ decay of ^{130}Te and by multiple Compton scattering of the various γ rays from environmental radioactivity and cosmic radiation. The choice of the ROI is driven by the need for the energy window to fully contain the events of the posited $0\nu\beta\beta$ peak, while being large enough to include and constrain the background structures, allowing us to evaluate the signal rate correctly. The ROI contains the ^{54}Mn and ^{208}Tl peaks, while the ^{228}Ac line is excluded as it is 45 keV ($> 5\sigma$ with FWHM energy resolution of ~ 4.3 keV in the ROI) away from $Q_{\beta\beta}$.

We perform a simultaneous binned Bayesian fit on the five included datasets. The fit is performed with the Bayesian Analysis Toolkit (BAT) [30], that samples from the posterior probability density by performing a Markov Chain Monte Carlo (MCMC) using the Metropolis-Hastings algorithm. We fit the CUORE \mathcal{M}_1 spectrum over the chosen ROI; the lower limit on the $0\nu\beta\beta$ decay rate is taken as the rate corresponding to 90% of the marginalized posterior.

We fit the CUORE \mathcal{M}_1 spectrum over the chosen ROI with a likelihood that includes the posited signal peak plus the background structures present in the ROI, namely the ^{54}Mn peak, the ^{208}Tl peak, and a continuum distribution. We model the latter with a linear function, that describes the decreasing trend over the fit region. This simpler effective model is consistent with the full CUORE background model [9]. The binned likelihood for each dataset is the product of Poisson terms, and the total likelihood is

$$\mathcal{L} = \prod_{ds} \prod_i^{N_{\text{bins}}} \frac{\mu_i^{n_i} e^{-\mu_i}}{n_i!}, \quad (1)$$

where ds indexes the dataset, and the index i runs over the 140 bins (0.5 keV/bin). In the approximation of small bin width, the number of expected counts μ_i in the i th bin can be taken as the value of the model function at the center of the bin:

$$\mu_i = S f_S^{ds}(i) + C_{\text{Mn}} f_{\text{Mn}}^{ds}(i) + C_{\text{Tl}} f_{\text{Tl}}^{ds}(i) + f_{\text{linear}}^{ds}(i), \quad (2)$$

where S , C_{Mn} , and C_{Tl} are the number of counts at the signal, ^{54}Mn and ^{208}Tl peaks, while $f_S^{ds}(i)$, $f_{\text{Mn}}^{ds}(i)$, $f_{\text{Tl}}^{ds}(i)$, and $f_{\text{linear}}^{ds}(i)$ are the values at the i th bin of the probability density functions used to model the shape of each component.

We model the shape of each peak as the sum of three Gaussian distributions based on the CUORE detector response function, corrected for the energy dependence of the detector response (energy-resolution scaling and energy reconstruction bias) studied in Ref. [7]. The definition of

each component of Eq. (2) is detailed in the following. We implement all terms as parameters of the fit.

The $0\nu\beta\beta$ decay rate $\Gamma_{0\nu}$ is connected to the expected number of signal counts S for a given dataset through the formula

$$S = \Gamma_{0\nu} \frac{N_A}{A_{\text{TeO}_2}} \eta_{128} (M\Delta t)_{ds} \epsilon_{ds}^{\text{cut}} \epsilon_{\text{MC}}, \quad (3)$$

where N_A is Avogadro's constant, A_{TeO_2} is the TeO_2 molar mass, η_{128} is the ^{128}Te natural isotopic abundance, $(M\Delta t)_{ds}$ is the dataset exposure (in units of kg yr), $\epsilon_{ds}^{\text{cut}}$ is the dataset total analysis efficiency, and ϵ_{MC} is the containment efficiency. The decay rate $\Gamma_{0\nu}$ in the model is a global parameter common to all the datasets. We make a statistical inference on this parameter of interest.

The ^{54}Mn originates from cosmogenic activation of Cu, which occurred before the CUORE cryostat and detector structure components were moved underground at Laboratori Nazionali del Gran Sasso. This element has a half-life of 312.2 days; the analyzed data were taken over a period of ~ 2 years, thus we expect the number of events due to ^{54}Mn decay to decrease over time. To account for this reduction, we include a multiplicative factor in the definition of the number of expected ^{54}Mn events in each dataset, resulting from the integration of the exponential decay over the dataset duration:

$$C_{\text{Mn}} = \Gamma_{\text{Mn}} \tau \left(e^{-\frac{t_{ds}^{\text{in}}}{\tau}} - e^{-\frac{t_{ds}^{\text{fin}}}{\tau}} \right) \frac{(M\Delta t)_{ds}}{t_{ds}^{\text{fin}} - t_{ds}^{\text{in}}} \epsilon_{ds}^{\text{cut}}, \quad (4)$$

where t_{ds}^{in} and t_{ds}^{fin} , respectively, refer to the start time and the end time of the dataset with respect to the beginning of the data taking. The live time fraction $[\Delta t_{ds} / (t_{ds}^{\text{fin}} - t_{ds}^{\text{in}})]$ accounts for the dead times—few time intervals of data taking that are removed from the analysis, for example noisy periods due to short maintenance interruptions, activities in the local laboratory or earthquakes—over the integration time interval. The ^{54}Mn rate Γ_{Mn} [units of counts/(kg yr)] is a nuisance parameter of the fit common to all the datasets.

^{208}Tl belongs to the naturally occurring ^{232}Th chain. Given that the amplitude of the observed higher intensity ^{208}Tl γ peaks are constant in time across the datasets, we assume the 860.6 keV rate to be stable. We then define the expected number of events at this ^{208}Tl line in the ROI for a given dataset as

$$C_{\text{Tl}} = \Gamma_{\text{Tl}} (M\Delta t)_{ds} \epsilon_{ds}^{\text{cut}}, \quad (5)$$

where the ^{208}Tl decay rate Γ_{Tl} is expressed in units of counts/(kg yr). As with the ^{54}Mn rate, this represents a nuisance parameter of the fit.

We model the continuous background distribution as a linear function of energy according to the following expression:

$$f_{\text{linear}}^{ds}(i) = C_b^{ds} + m_{ds}(E_i - E_{1/2}), \quad (6)$$

where C_b^{ds} and m_{ds} are the expected number of background events and the background slope for a given dataset, E_i is the energy at the center of the i th bin, and $E_{1/2}$ is the energy corresponding to center of the ROI. We define the expected number of events C_b^{ds} in each dataset as

$$C_b^{ds} = \text{BI}_{ds}(M\Delta t)_{ds}w_i, \quad (7)$$

where BI_{ds} is the background index of dataset ds in units of counts/(keV kg yr) and w_i is the bin width, which is constant across the energy spectrum. The slope and the background index are also nuisance parameters in the fit and are dataset-dependent quantities.

We adopt a uniform prior for each parameter of the fit for several reasons. Because of the 100-fold increase in exposure, CUORE's sensitivity on $\Gamma_{0\nu}$ is expected to be factor of ~ 10 better with respect to the past direct limit. The absence of knowledge on $\Gamma_{0\nu}$ at the range that CUORE can probe justifies the choice of a uniform prior for $\Gamma_{0\nu} \geq 0$ according to the principle of indifference, which assigns equal probabilities to all the possible values up to a maximum that can be greater than the past limit. The CUORE background model can provide information on some nuisance parameters, however, it is constructed through a fit on the same data that are used for the present analysis, and including such information would bias the result. Thus, in the absence of independent measurements, we use a uniform prior for all the nuisance parameters. The signal, ^{54}Mn and ^{208}Tl rates and the BI are constrained to nonnegative physical values only, while for the background slope m both negative and positive values are allowed. We run the Bayesian fit on the data, and find no evidence for ^{128}Te $0\nu\beta\beta$ decay. From the marginalized posterior distribution of the signal rate, we extract a 90% CI limit of

$$\Gamma_{0\nu} < 1.9 \times 10^{-25} \text{ yr}^{-1}. \quad (8)$$

This lower limit corresponds to a 90% CI upper limit on the ^{128}Te $0\nu\beta\beta$ decay half-life of

$$T_{1/2}^{0\nu} > 3.6 \times 10^{24} \text{ yr}. \quad (9)$$

This result is the most stringent limit on the $0\nu\beta\beta$ decay of ^{128}Te to date, representing a more than 30-fold improvement over the previous limit [14] from direct searches, and exceeds for the first time the combined $0\nu\beta\beta$ and $2\nu\beta\beta$ decay half life obtained by geochemical measurements. The fit result and the total ROI spectrum are shown in Fig. 2.




FIG. 2. Top: data spectrum in the ROI, together with the best-fit curve (red solid) and the best-fit curve with the signal rate set at the 90% CI limit (blue dashed). Bottom: residual plot with fit, compatible with 0 (intercept at -1.1 ± 2.6 counts/keV, $\chi^2/\text{dof} = 82/69$).

We extract the median exclusion sensitivity to ^{128}Te $0\nu\beta\beta$ decay by repeating the statistical only Bayesian fit on 10^4 toy MC simulations of the experiment. We produce the toy MCs using the global mode values of the background parameters from a Bayesian fit without the signal component on the CUORE data. The median exclusion sensitivity is the median of the distribution of the 90% CI limits on $T_{1/2}^{0\nu}$, each resulting from a signal plus background fit to one of the 10^4 background-only toy MCs. This distribution is shown in Fig. 3, and its median is $\hat{T}_{1/2}^{0\nu} = 2.2 \times 10^{24}$ yr. The probability to obtain a more stringent limit than the one observed with the CUORE data is 8.8%. We also repeat the fit on the data, allowing the signal rate to assume non-physical negative values. In this case, the global mode of $\Gamma_{0\nu}$ is $(-2.4 \pm 1.8) \times 10^{-25} \text{ yr}^{-1}$, resulting in an under-fluctuation with a statistical significance of $\sim 1.4\sigma$, which is




FIG. 3. Distribution of the 90% CI limits on $T_{1/2}^{0\nu}$ extracted from repeating the analysis on the 10^4 background-only pseudoexperiments. The solid line corresponds to the median exclusion sensitivity, while the dashed one shows the 90% CI limit from the analysis of the CUORE data.

TABLE I. Systematic effects on the ^{128}Te $0\nu\beta\beta$ decay signal rate 90% limit. The first row refers to the BAT intrinsic uncertainty, due to the stochastic nature of the MCMC. The efficiencies and the ^{128}Te isotopic abundance were treated as nuisance parameters in the fit, while an alternative approach was adopted for the uncertainties on the ^{128}Te $Q_{\beta\beta}$ and the detector response function parameters. The dominant systematic effect is the value on $Q_{\beta\beta}$.

Systematic	Prior	Effect on $\Gamma_{0\nu}^{90\%}$
BAT stat only fit	...	0.3%
Bayesian approach		
Containment efficiency	Gaussian	0.4%
^{128}Te isotopic abundance	Gaussian	< 0.3% (0.05%)
Analysis cut efficiency	Gaussian	< 0.3% (0.1%)
Repeated fit approach		
^{128}Te $Q_{\beta\beta}$	Gaussian	7.0%
Energy reconstruction bias	Multivariate	< 0.3% (0.1%)
Energy resolution scaling	Multivariate	< 0.3% (0.1%)

compatible with the 8.8% underfluctuation obtained from the sensitivity study.

We summarize in Table I a series of systematic uncertainties affecting our limit. For this study, we run the fit without the constraint $\Gamma_{0\nu} \geq 0$, to access the full range $\Gamma_{0\nu}$ marginalized posterior. We adopt a fully Bayesian approach to evaluate the effect due to the uncertainties on the containment efficiency, the analysis cut efficiency, and the ^{128}Te natural isotopic abundance. We implement these as independent nuisance parameters in the likelihood with a Gaussian prior, whose mean and sigma are equal to the respective central value and associated error. We thus repeat the Bayesian fit activating one nuisance parameter at the time to allow its value to vary according to the corresponding prior.

We treat the systematics due to the uncertainty on the ^{128}Te $Q_{\beta\beta}$ and on the detector response function parameters (namely the energy reconstruction bias and resolution scaling) using an alternative approach, which we refer to as the repeated fit approach, because of the excessive computation time required to treat them as nuisance parameters in the fit. This method consists of repeating the fit for a series of discrete values of the systematic parameter under study, covering a $\pm 3\sigma$ region around its prior mean value. We then sum the $\Gamma_{0\nu}$ marginalized posteriors obtained from each fit weighting by the prior probability of the parameter considered as systematic, and take the signal rate corresponding to the 90% quantile of the obtained distribution. We take additional care when treating the detector response parameter systematics. It was previously observed in CUORE [7] that both the bias on the energy reconstruction and the resolution scaling exhibit an energy dependence which we model with two independent second order polynomial functions. As a consequence, a set of three correlated parameters describes the energy bias and another set of three

exists for the resolution scaling. The correlations among these parameters are taken into account using multidimensional priors. The dominant systematic is $Q_{\beta\beta}$, which has an effect of 7.0% on the limit. We expect this due to the relatively large error on its literature value (866.7 ± 0.7) keV [13]. All the other systematics affect the limit by less than 1%; the ^{128}Te isotopic abundance, the analysis cut efficiency, and the detector response function parameters result in values below the intrinsic BAT uncertainty due to the MCMC stochastic behavior (0.3%).

Several standard model extended theories include mechanisms that try to explain how the $0\nu\beta\beta$ decay takes place [31–33]. Among these models, the exchange of a light Majorana neutrino is the most favored [34]. However, a positive signal for the $0\nu\beta\beta$ decay of a single isotope would not determine the mechanism of this process [2]. Discriminating among the existing models [35] and possibly testing the calculations of nuclear matrix elements for double beta decay [36], would be possible via the comparison of results from different isotopes. It has been pointed out [35] that the study of the $0\nu\beta\beta$ decay of ^{128}Te can be particularly useful for such model discrimination. In this Letter, we present the first results on the ^{128}Te $0\nu\beta\beta$ decay search with the CUORE experiment. With a binned Bayesian fit of the CUORE data with a total exposure of 309.33 kg yr (78.6 kg yr of ^{128}Te), we find no evidence for ^{128}Te $0\nu\beta\beta$ decay, and we set a 90% CI limit on the half-life of this process at $T_{1/2}^{0\nu} > 3.6 \times 10^{24}$ yr. This represents the most stringent limit in literature, improving by over a factor 30 the previous limit from a direct search experiment, and exceeding those from indirect geochemical measurements for the first time. From the analyzed exposure, the CUORE median exclusion sensitivity to this decay is $\hat{T}_{1/2}^{0\nu} = 2.2 \times 10^{24}$ yr, giving an 8.8% probability to obtain a stronger limit. The dominant systematic, affecting the result at the level of 7.0%, is due to the uncertainty on $Q_{\beta\beta}$.

The analysis presented in this Letter has been carried out with about one-tenth of the final exposure scheduled for CUORE, corresponding to ~ 3.7 ton y. We plan to update these results with such an unprecedented amount of data collected with TeO_2 crystals.

The CUORE Collaboration thanks the directors and staff of the Laboratori Nazionali del Gran Sasso and the technical staff of our laboratories. This work was supported by the Istituto Nazionale di Fisica Nucleare (INFN); the National Science Foundation under Grants No. NSF-PHY-0605119, No. NSF-PHY-0500337, No. NSF-PHY-0855314, No. NSF-PHY-0902171, No. NSF-PHY-0969852, No. NSF-PHY-1307204, No. NSF-PHY-1314881, No. NSF-PHY-1401832, and No. NSF-PHY-1913374; and Yale University. This material is also based upon work supported by the U.S. Department of Energy (DOE) Office of Science under Contracts No. DE-AC02-05CH11231 and




FIG. 4. Linear fit on the mean reconstructed signal rate as a function of the injected one. The intercept and slope are compatible with 0 and 1, respectively.

No. DE-AC52-07NA27344; by the DOE Office of Science, Office of Nuclear Physics under Contracts No. DE-FG02-08ER41551, No. DE-FG03-00ER41138, No. DE-SC0012654, No. DE-SC0020423, No. DE-SC0019316; and by the EU Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreement No. 754496. This research used resources of the National Energy Research Scientific Computing Center (NERSC). This work makes use of both the DIANA data analysis and APOLLO data acquisition software packages, which were developed by the CUORICINO, CUORE, LUCIFER and CUPID-0 Collaborations.

Appendix.—We develop and optimize the fit strategy on toy MC spectra. We generate the toy MCs according to the signal-plus-background model, extracting the values of Γ_{Mn} , Γ_{Tl} , BI, and m from the CUORE background model. We refer to Ref. [37] for a more detailed discussion of the method. We take advantage of the toy MCs to verify that the fit correctly reconstructs the simulated background components and to inspect if a bias is introduced in the $0\nu\beta\beta$ decay rate reconstruction when a signal contribution is added in the toy MC. We generate 10^4 toy MCs with no signal and run the fit independently on each of them. We then construct the distributions of the best-fit values from all the toy MCs for each parameter, in order to compare the extracted and simulated values. As expected, these distributions are centered at the values used to produce the toy MC. Thanks to the large number of toy MCs, we are able to identify a small bias in the reconstruction of the BI and the slope corresponding to a $< 0.15\%$ underestimation and a $\leq 1.6\%$ overestimation, respectively. No correlations are seen between these two parameters. The reconstructed values of the ^{54}Mn and ^{208}Tl rates are compatible with the injected values. To test the signal rate

TABLE II. Parameter ranges for the fit parameters. All parameters are assigned a uniform prior; the signal, Mn and Tl rates and the BI are constrained to nonnegative physical values only, while both positive and negative values are allowed for the background slope.

Parameter	Prior range
$\Gamma_{0\nu}$	$[0, 1.74 \times 10^{-24}] \text{ yr}^{-1}$
BI_1	$[1.1634, 1.73] \text{ cts}/(\text{keV kg yr})$
BI_2	$[1.188, 1.6513] \text{ cts}/(\text{keV kg yr})$
BI_3	$[1.2453, 1.7374] \text{ cts}/(\text{keV kg yr})$
BI_4	$[1.2204, 1.7412] \text{ cts}/(\text{keV kg yr})$
BI_5	$[1.0221, 1.4536] \text{ cts}/(\text{keV kg yr})$
m_1	$[-1, 1] \text{ 1/keV}$
m_2	$[-1, 1] \text{ 1/keV}$
m_3	$[-1, 1] \text{ 1/keV}$
m_4	$[-1, 1] \text{ 1/keV}$
m_5	$[-1, 1] \text{ 1/keV}$
Γ_{Mn}	$[0, 44.58] \text{ cts}/(\text{kg yr})$
Γ_{Tl}	$[0, 6.16] \text{ cts}/(\text{kg yr})$

reconstruction, we repeat the fit on five sets of 2000 toy MCs, injecting a different signal amplitude in the range $(2-10) \times 10^{-25} \text{ yr}^{-1}$ in each set. This range includes the signal rate corresponding to the CUORE sensitivity of $3.2 \times 10^{-25} \text{ yr}^{-1}$ obtained from pure toy MC, i.e., without including real data.

Figure 4 shows the mean reconstructed signal rate as a function of the injected one. The relation between the two is well described by a linear function: the intercept is compatible with 0 at a $\sim 1.3\sigma$, and the slope is compatible with 1 within 1σ . These results allow us to conclude that no bias is introduced by the Bayesian fit in the signal rate reconstruction.

We also study the intrinsic stability of the BAT fit, by repeating it 2×10^3 times on the same toy MC populated only with the background components, obtaining a 0.3%

TABLE III. Best-fit values for all parameters of the fit on CUORE data. The signal rate is allowed to take nonnegative values only.

Parameter	Fit result	Units
$\Gamma_{0\nu}$	0	yr^{-1}
BI_1	1.48 ± 0.02	$\text{cts}/(\text{keV kg yr})$
BI_2	1.43 ± 0.02	$\text{cts}/(\text{keV kg yr})$
BI_3	1.49 ± 0.02	$\text{cts}/(\text{keV kg yr})$
BI_4	1.48 ± 0.02	$\text{cts}/(\text{keV kg yr})$
BI_5	1.26 ± 0.02	$\text{cts}/(\text{keV kg yr})$
m_1	-0.07 ± 0.03	keV^{-1}
m_2	-0.06 ± 0.03	keV^{-1}
m_3	-0.08 ± 0.03	keV^{-1}
m_4	-0.04 ± 0.03	keV^{-1}
m_5	-0.12 ± 0.03	keV^{-1}
Γ_{Mn}	15.9 ± 0.7	$\text{cts}/(\text{kg yr})$
Γ_{Tl}	0.5 ± 0.2	$\text{cts}/(\text{kg yr})$




FIG. 5. Top: marginalized posterior of the signal rate obtained from the official fit, which allows only physical values. Bottom: marginalized posterior of the signal rate obtained from the alternative fit, which allows also nonphysical values of $\Gamma_{0\nu}$. A $\sim 1.4\sigma$ significance underfluctuation, compatible with the results of the sensitivity studies, is observed. Both distributions are normalized to one.

root mean square on the distribution of the $\Gamma_{0\nu}$ limits at 90% credibility interval (CI).

Table II reports the ranges for all fit parameters. All parameters proportional to a number of counts, namely the signal, Mn, and Tl rates, are allowed to assume only nonnegative values. The BI of each dataset is further constrained according to a preliminary estimation of the number of background counts. The background slopes are allowed to assume also negative values.

Table III reports the value at the global mode for all parameters of the fit to the data.

Figure 5 shows the posterior distribution for $\Gamma_{0\nu}$ obtained from the reference fit, and from the alternative fit performed with the signal rate allowed to artificially assume non-physical negative values.

*Deceased.

- [1] A. S. Barabash, *Universe* **6**, 159 (2020).
 [2] S. M. Bilenky and C. Giunti, *Int. J. Mod. Phys. A* **30**, 1530001 (2015).

- [3] M. J. Dolinski, A. W. P. Poon, and W. Rodejohann, *Annu. Rev. Nucl. Part. Sci.* **69**, 219 (2019).
 [4] L. Canetti, M. Drewes, and M. Shaposhnikov, *New J. Phys.* **14**, 095012 (2012).
 [5] M. A. Luty, *Phys. Rev. D* **45**, 455 (1992).
 [6] S. Dell’Oro, S. Marocco, and F. Vissani, *Phys. Rev. D* **100**, 073003 (2019).
 [7] D. Q. Adams *et al.* (CUORE Collaboration), *Phys. Rev. Lett.* **124**, 122501 (2020).
 [8] D. Q. Adams *et al.* (CUORE), *Nature (London)* **604**, 53 (2022).
 [9] D. Q. Adams *et al.* (CUORE Collaboration), *Phys. Rev. Lett.* **126**, 171801 (2021).
 [10] D. Q. Adams *et al.* (CUORE Collaboration), *Eur. Phys. J. C* **81**, 567 (2021).
 [11] D. Q. Adams *et al.*, *Phys. Rev. C* **105**, 065504 (2022).
 [12] M. A. Fehr, M. Rehkamper, and A. N. Halliday, *Int. J. Mass Spectrom.* **232**, 83 (2004).
 [13] M. Wang, W. J. Huang, F. G. Kondev, G. Audi, and S. Naimi, *Chin. Phys. C* **45**, 030003 (2021).
 [14] C. Arnaboldi *et al.*, *Phys. Lett. B* **557**, 167 (2003).
 [15] A. Campani, V. Dompè, and G. Fantini, *Universe* **7**, 212 (2021).
 [16] T. Bernatowicz, J. Brannon, R. Brazzle, R. Cowsik, C. Hohenberg, and F. Podosek, *Phys. Rev. Lett.* **69**, 2341 (1992).
 [17] C. Alduino *et al.* (CUORE Collaboration), *Eur. Phys. J. C* **77**, 13 (2017).
 [18] I. Nutini *et al.*, *J. Low Temp. Phys.* **199**, 519 (2020).
 [19] E. E. Haller, N. P. Palaio, M. Rodder, W. L. Hansen, and E. Kreysa, *NTD germanium: A novel material for low temperature bolometers*, in *Neutron Transmutation Doping of Semiconductor Materials* (Springer, Boston, 1984), pp. 21–36.
 [20] C. Alduino *et al.*, *Cryogenics* **102**, 9 (2019).
 [21] C. Alduino *et al.* (CUORE Collaboration), *Eur. Phys. J. C* **77**, 543 (2017).
 [22] A. D’Addabbo, C. Bucci, L. Canonica, S. Di Domizio, P. Gorla, L. Marini, A. Nucciotti, I. Nutini, C. Rusconi, and B. Welliver, *Cryogenics* **93**, 56 (2018).
 [23] V. Dompè *et al.*, *J. Low Temp. Phys.* **200**, 286294 (2020).
 [24] D. Adams *et al.*, *Prog. Part. Nucl. Phys.* **122**, 103902 (2022).
 [25] C. Brofferio and S. Dell’Oro, *Rev. Sci. Instrum.* **89**, 121502 (2018).
 [26] S. Di Domizio, F. Orio, and M. Vignati, *J. Instrum.* **6**, P02007 (2011).
 [27] A. Campani *et al.*, *J. Low Temp. Phys.* **200**, 321 (2020).
 [28] C. Alduino *et al.* (CUORE Collaboration), *Phys. Rev. C* **93**, 045503 (2016).
 [29] M. Laubenstein and G. Heusser, *Appl. Radiat. Isot.* **67**, 750 (2009).
 [30] A. Caldwell, D. Kollar, and K. Kroninger, *Comput. Phys. Commun.* **180**, 2197 (2009).
 [31] B. Pontecorvo, *Phys. Lett.* **26B**, 630 (1968).
 [32] M. Mitra, G. Senjanovic, and F. Vissani, *Nucl. Phys.* **B856**, 26 (2012).

- [33] V. Tello, M. Nemevsek, F. Nesti, G. Senjanovic, and F. Vissani, *Phys. Rev. Lett.* **106**, 151801 (2011).
- [34] S. Dell’Oro, S. Marcocci, M. Viel, and F. Vissani, *Adv. High Energy Phys.* **2016**, 2162659 (2016).
- [35] F. Deppisch and H. Pas, *Phys. Rev. Lett.* **98**, 232501 (2007).
- [36] S. M. Bilenky and J. A. Grifols, *Phys. Lett. B* **550**, 154 (2002).
- [37] V. Dompè, Search for neutrinoless double beta decay of ^{128}Te with the CUORE experiment, Ph.D. thesis, Gran Sasso Science Institute, 2021.